

Astronomy Question Bank

These questions were carefully written or selected to prepare students for the astronomy team selection process in Brazil. Some questions are not original; those are marked with () before the statement. The question-bank template is the same one used by Professor Kevin Zhou. His work is invaluable, and many ideas in this set can be found in his handouts.

1 Optics and Telescopes

- [3] **Problem 1.** A Keplerian telescope has two converging lenses. The first has a radius of curvature of both faces equal to $R = 2$ m, and the second has both radii of curvature equal to $r = 0.5$ m. Considering that both lenses have negligible thickness and are made of a material with refractive index $n = 1.4$, calculate the diameter and the magnification of the telescope, knowing that this telescope is an $f/12$.
- [5] **Problem 2.** In this problem, we will become familiar with one of the most powerful tools in optics, *geometric optics*. The goal of this tool is to model lenses and mirrors in the form of matrices. This is very useful for solving problems involving combinations of several lenses, and it is a method for solving geometric-optics problems (practically) without using geometry. For that, pay attention to the following definitions in the figure below.




Figure 1: Diagram 1

All dashed lines are parallel to the optical axis, represented by the horizontal arrow. P_k denotes the point where the light passes from one medium to another, and y_k is the vertical coordinate of the point P_k . The subscripts i and t are used to refer to incident and transmitted, respectively. Thus, for example, $\alpha_{i,k}$ is the angle that the incident light ray makes with the horizontal at the point P_k , while the angle α_k is the angle between the point P_k and the center of lens k . For example, in the figure, the angle α_1 is represented by $V_1\hat{C}P_1$.

For this problem, we will assume that all angles of interest are small, so that $\tan \theta \approx \sin \theta \approx \theta$.

- a) Starting from Snell's law, find a relation between $n_{i,1}$, $n_{t,1}$, α_1 , $\alpha_{i,1}$, and $\alpha_{t,1}$.

- b) Find a relation for (1) $n_{t,1}\alpha_{t,1}$ and (2) $y_{t,1}$. Leave your answers in terms of $n_{i,1}$, $\alpha_{i,1}$, $y_{i,1}$, and $\mathcal{D}_1$, where

$$\mathcal{D}_1 = \frac{n_{t,1} - n_{i,1}}{R_1}$$

Now for one more definition. Let the vector $\mathbf{r}_{t,1} = (n_{t,1}\alpha_{t,1}, y_{t,1})$ and $\mathbf{r}_{i,1} = (n_{i,1}\alpha_{i,1}, y_{i,1})$.

- c) The two equations found in the previous item can be written in matrix form, as

$$\mathbf{r}_{t,1} = \mathcal{R}_1 \mathbf{r}_{i,1}$$

Find the 2×2 matrix equivalent to $\mathcal{R}_1$.

Our interest now is to find the relations between the points P_2 and P_1 .

- d) Find an equation for $n_{i,2}$ and $y_{i,2}$. Using the reasoning of the previous item, leave your answer in the form

$$\mathbf{r}_{i,2} = \Gamma_{2,1} \mathbf{r}_{t,1}$$

where $\Gamma_{2,1}$ is also a 2×2 matrix. Leave your answer in terms of the lens thickness, $d_{2,1}$.

- e) We define the lens matrix, $\mathcal{A}_{2,1}$, by the following equation:

$$\mathbf{r}_{t,2} = \mathcal{A}_{2,1} \mathbf{r}_{i,1}$$

so that

$$\mathcal{A}_{2,1} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

Find $\mathcal{A}_{2,1}$ explicitly. This is extremely important, because it describes the light ray that leaves the lens in terms of the ray that enters.

- f) Among all the properties of the lens matrix, the most curious is that the term a_{12} is proportional to $-1/f$, where f is the focal length of the lens. Prove this result. (Hint: you will be able to write $-a_{12}$ with a well-known equation from optics.)

Now let us get to work and make practical use of matrix optics.

- g) Consider the following diagram:




Figure 2: Diagram 2

Find y_i in terms of y_0 and the data in the figure.

- h) If the lens is between two different media, $n_{i,1} \neq n_{t,2}$, we have the following relation:

$$a_{12} = -\frac{n_{i,1}}{f_0} = -\frac{n_{t,2}}{f_I}$$

Using this, redo the previous item, considering that there is water with $n_w = 4/3$ between the two lenses.

- i) A very common lens arrangement is the Tessar objective, found in many cameras because of its efficiency in reducing aberration and astigmatism. The arrangement has the following form:




Figure 3: Tessar objective diagram

Find the equivalent lens matrix, $\mathcal{A}_{1,7}$, in terms of $\mathcal{R}_i$ and $\Gamma_{i,j}$.

- [2] **Problem 3.** The wave theory of light shows that a perfect focus is not possible, because of diffraction effects associated with the finite aperture of the lens. This lack of a perfect focus prevents objects that are very close together from being distinguished. This problem can be studied from two different points of view:

The wave theory of light predicts that a lens of diameter D cannot focus a parallel beam of light of wavelength λ into an angle smaller than the diffraction limit:

$$\theta_m \approx 1.22 \frac{\lambda}{D} \quad (1)$$

Now consider a quantum approach: the photons that are focused by the lens. These photons are known to have passed somewhere within a radius of the center of the lens. The uncertainty in the position x is associated with an uncertainty in the x component of the photon's momentum. Consequently, a photon that, in the absence of this uncertainty, would have been brought onto the optical axis of the focal plane may now be deflected by an angle $\theta \ll 1$.

Consider the de Broglie wavelength $\lambda = \frac{h}{p}$. Find a limit for θ .

Hint: Heisenberg's uncertainty principle relates the imprecision between measurements of the momentum and the position of a particle by

$$\Delta p_x \Delta x = \frac{\hbar}{2}$$

- [3] **Problem 4.** (Magna notes) Prove that the quantity $n_i R_i \sin \theta_i$, for spherical shells adjacent to one another, is an invariant, where n_i is the refractive index of the i -th spherical shell, θ_i is the angle that the light ray makes with the normal to the i -th spherical shell, and R_i is the radius of the i -th spherical shell.

- [4] **Problem 5.** The refractive index of a planet's atmosphere is given by

$$n(h) = \frac{n_0}{1 + \epsilon h}$$

where n_0 and ϵ are constants.

- a) A light ray reaches the atmosphere parallel to the surface, at a height $h' \ll R$. What is the trajectory?
- b) Given the trajectory of the light ray, calculate the radius of the planet.

Hint: You may find the following relation useful,

$$\int \frac{t}{\sqrt{a - bt^2}} dt = -\frac{1}{b} \sqrt{a - bt^2} + C$$