

# Assignment 7: GLMs (Linear Regressios, ANOVA, & t-tests)

Maia Griffith

Fall 2023

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on generalized linear models.

## Directions

1. Rename this file `<FirstLast>_A07_GLMs.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

## Set up your session

1. Set up your session. Check your working directory. Load the tidyverse, agricolae and other needed packages. Import the *raw* NTL-LTER raw data file for chemistry/physics (NTL-LTER\_Lake\_ChemistryPhysics\_Raw.csv). Set date columns to date objects.
2. Build a ggplot theme and set it as your default theme.

```
#1 Setting up my session.
```

```
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
```

```
## v dplyr      1.1.3      v readr      2.1.4
```

```
## v forcats    1.0.0      v stringr   1.5.0
```

```
## v ggplot2    3.4.3      v tibble    3.2.1
```

```
## v lubridate  1.9.3      v tidyr     1.3.0
```

```
## v purrr      1.0.2
```

```
## -- Conflicts ----- tidyverse_conflicts() --
```

```
## x dplyr::filter() masks stats::filter()
```

```
## x dplyr::lag()     masks stats::lag()
```

```
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors
```

```
library(agricolae)
```

```
library(corrplot)
```

```
## corrplot 0.92 loaded
```

```
library(dplyr)
library(here)

## here() starts at /home/guest/ENV872/maia-g/EDE_Fall2023

here()

## [1] "/home/guest/ENV872/maia-g/EDE_Fall2023"

#Load the correct dataset
lake.chem.phys <- read.csv(here("Data/Raw/NTL-LTER_Lake_ChemistryPhysics_Raw.csv"),
                           stringsAsFactors = TRUE)

#Change into date format
lake.chem.phys$sampdate <- as.Date(lake.chem.phys$sampdate , format = "%m/%d/%Y")

#2 Build and set new theme.
MGtheme <- theme_light(base_size = 11) +
  theme(axis.text = element_text(color = "darkgray"),
        legend.position = "right")
theme_set(MGtheme)
```

## Simple regression

Our first research question is: Does mean lake temperature recorded during July change with depth across all lakes?

3. State the null and alternative hypotheses for this question: > Answer: H0: July mean lake temperature across all lakes does not change with depth. Ha: July mean lake temperature across all lakes changes with depth.
4. Wrangle your NTL-LTER dataset with a pipe function so that the records meet the following criteria:
  - Only dates in July.
  - Only the columns: lakename, year4, daynum, depth, temperature\_C
  - Only complete cases (i.e., remove NAs)
5. Visualize the relationship among the two continuous variables with a scatter plot of temperature by depth. Add a smoothed line showing the linear model, and limit temperature values from 0 to 35 °C. Make this plot look pretty and easy to read.

```
#4 Pipe to wrangle the data
Jul.lake.chem.phys <-
  lake.chem.phys %>% # Calling the data frame
  mutate(Month = month(sampdate)) %>% # Creating a month col
  filter(Month == 7 ) %>% # Filtering by July (month number 7)
  select(lakename, year4, daynum, depth, temperature_C) %>% # Selecting cols
  na.omit() # Removing ALL NA values from ALL cols

dim(Jul.lake.chem.phys)
```

```
## [1] 9728    5
```


```
#5 Scatter plot of temp by depth (x = depth, y = temp) with line showing linear model, temp from 0 to 3
temp_depth_plot <-
  ggplot(Jul.lake.chem.phys, aes(x = depth, # Creating the plot
                                y = temperature_C)) +

  geom_point() + # Scatterplot
  geom_smooth(method = "lm") + # Adding line of best fit
  labs(y = "Temperature (C)",
       x = "Depth (meters)",
       title = "Lake Temperature by Depth") + # Editing axis titles
  ylim(0,35) # Limiting temp values

temp_depth_plot
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```

```
## Warning: Removed 24 rows containing missing values ('geom_smooth()').
```



6. Interpret the figure. What does it suggest with regards to the response of temperature to depth? Do the distribution of points suggest about anything about the linearity of this trend?

Answer: The plot suggests that there is a negative relationship between temperature and depth, meaning with increasing depth there is decreasing temperature. The distribution of points suggests that the trend may not be simply linear, but have a curve of some kind in which temperature decreases strongly within the first 10 meters, and then starts to level out.

7. Perform a linear regression to test the relationship and display the results

```
#7
tempdepth.regr <-
  lm(Jul.lake.chem.phys$temperature_C ~
      Jul.lake.chem.phys$depth)

summary(tempdepth.regr)

##
## Call:
## lm(formula = Jul.lake.chem.phys$temperature_C ~ Jul.lake.chem.phys$depth)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.5173 -3.0192  0.0633  2.9365 13.5834
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    21.95597     0.06792   323.3  <2e-16 ***
## Jul.lake.chem.phys$depth -1.94621     0.01174  -165.8  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.835 on 9726 degrees of freedom
## Multiple R-squared:  0.7387, Adjusted R-squared:  0.7387
## F-statistic: 2.75e+04 on 1 and 9726 DF, p-value: < 2.2e-16
```

8. Interpret your model results in words. Include how much of the variability in temperature is explained by changes in depth, the degrees of freedom on which this finding is based, and the statistical significance of the result. Also mention how much temperature is predicted to change for every 1m change in depth.

Answer: The Adjusted R-squared value is 0.7387, which means that about 73.8% of the variability in temperature is explained by changes in depth. There are 9726 degrees of freedom, with a high statistical significance with a p-value of less than 0.05 (p-value: < 2.2e-16). For every 1m change in depth, the temperature decreases by about 1.9 degrees C.

---

## Multiple regression

Let's tackle a similar question from a different approach. Here, we want to explore what might the best set of predictors for lake temperature in July across the monitoring period at the North Temperate Lakes LTER.

9. Run an AIC to determine what set of explanatory variables (year4, daynum, depth) is best suited to predict temperature.
10. Run a multiple regression on the recommended set of variables.

```

#9 AIC for year4, daynum, and depth for predicted temp
Jul.lakes.AIC <- lm(data = Jul.lake.chem.phys,
                    temperature_C ~ year4 + daynum + depth)

step(Jul.lakes.AIC)

## Start:  AIC=26065.53
## temperature_C ~ year4 + daynum + depth
##
##           Df Sum of Sq    RSS   AIC
## <none>                 141687 26066
## - year4    1         101 141788 26070
## - daynum   1         1237 142924 26148
## - depth    1       404475 546161 39189

##
## Call:
## lm(formula = temperature_C ~ year4 + daynum + depth, data = Jul.lake.chem.phys)
##
## Coefficients:
## (Intercept)      year4      daynum      depth
##   -8.57556      0.01134      0.03978     -1.94644

# Stepwise AIC shows the initial AIC is best, with each step making it worse.
# It also shows that removing depth has the largest impact.
# The variables year4 has very little impact, and daynum has a small impact.

#10 New regression using all 3 variables to explain temperature.
temp.depthdaynumyear.regr <- lm(data = Jul.lake.chem.phys,
                                temperature_C ~ year4 + depth + daynum)
summary(temp.depthdaynumyear.regr)

##
## Call:
## lm(formula = temperature_C ~ year4 + depth + daynum, data = Jul.lake.chem.phys)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6536 -3.0000  0.0902  2.9658 13.6123
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) -8.575564   8.630715  -0.994   0.32044
## year4        0.011345   0.004299   2.639   0.00833 **
## depth       -1.946437   0.011683 -166.611 < 2e-16 ***
## daynum       0.039780   0.004317   9.215 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.817 on 9724 degrees of freedom
## Multiple R-squared:  0.7412, Adjusted R-squared:  0.7411
## F-statistic: 9283 on 3 and 9724 DF,  p-value: < 2.2e-16

```

11. What is the final set of explanatory variables that the AIC method suggests we use to predict temperature in our multiple regression? How much of the observed variance does this model explain? Is this an improvement over the model using only depth as the explanatory variable?

Answer: The AIC method suggests we use year4 (low impact), depth (high impact) and daynum (some impact) in our multiple regression. This new model explains 74.1% of the variability in temperature, which is a small improvement over only using depth.

---

## Analysis of Variance

12. Now we want to see whether the different lakes have, on average, different temperatures in the month of July. Run an ANOVA test to complete this analysis. (No need to test assumptions of normality or similar variances.) Create two sets of models: one expressed as an ANOVA models and another expressed as a linear model (as done in our lessons).

```
#12
# Format ANOVA as aov
Jul.Temp.anova <- aov(data = Jul.lake.chem.phys, temperature_C ~ lakename)
summary(Jul.Temp.anova)

##              Df Sum Sq Mean Sq F value Pr(>F)
## lakename      8  21642   2705.2     50 <2e-16 ***
## Residuals    9719 525813     54.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

# Format ANOVA as linear model (lm)
Jul.Temp.anova.lm <- lm(data = Jul.lake.chem.phys, temperature_C ~ lakename)
summary(Jul.Temp.anova.lm)

##
## Call:
## lm(formula = temperature_C ~ lakename, data = Jul.lake.chem.phys)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -10.769  -6.614  -2.679   7.684  23.832
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    17.6664    0.6501  27.174 < 2e-16 ***
## lakenameCrampton Lake    -2.3145    0.7699  -3.006 0.002653 **
## lakenameEast Long Lake   -7.3987    0.6918 -10.695 < 2e-16 ***
## lakenameHummingbird Lake  -6.8931    0.9429  -7.311 2.87e-13 ***
## lakenamePaul Lake       -3.8522    0.6656  -5.788 7.36e-09 ***
## lakenamePeter Lake      -4.3501    0.6645  -6.547 6.17e-11 ***
## lakenameTuesday Lake    -6.5972    0.6769  -9.746 < 2e-16 ***
## lakenameWard Lake       -3.2078    0.9429  -3.402 0.000672 ***
## lakenameWest Long Lake  -6.0878    0.6895  -8.829 < 2e-16 ***
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7.355 on 9719 degrees of freedom
## Multiple R-squared:  0.03953,    Adjusted R-squared:  0.03874
## F-statistic:    50 on 8 and 9719 DF,  p-value: < 2.2e-16
```

13. Is there a significant difference in mean temperature among the lakes? Report your findings.

Answer: Yes, there is a significant difference in mean temperature among the lakes, because the p-value (in both models) is less than 0.05 (p-value: < 2.2e-16).

14. Create a graph that depicts temperature by depth, with a separate color for each lake. Add a `geom_smooth` (method = "lm", se = FALSE) for each lake. Make your points 50 % transparent. Adjust your y axis limits to go from 0 to 35 degrees. Clean up your graph to make it pretty.

```
#14.
temp_depth_ANOVA_plot <-
  ggplot(Jul.lake.chem.phys, aes(x = depth,
                                y = temperature_C,
                                color = lakename)) +


  geom_point(alpha = 0.5) +
  geom_smooth(method = "lm",
             se = FALSE) +
  labs(y = "Temperature (C)",
       x = "Depth (meters)",
       color = "Lake Name",
       title = "Comparison of Temperature by Depth in Each Lake") +
  ylim(0,35)

temp_depth_ANOVA_plot
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```

```
## Warning: Removed 73 rows containing missing values ('geom_smooth()').
```

## Comparison of Temperature by Depth in Each Lake



15. Use the Tukey's HSD test to determine which lakes have different means.

#15

```
TukeyHSD(Jul.Temp.anova)
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = temperature_C ~ lakename, data = Jul.lake.chem.phys)
##
## $lakename
##
```

	diff	lwr	upr	p adj
Crampton Lake-Central Long Lake	-2.3145195	-4.7031913	0.0741524	0.0661566
East Long Lake-Central Long Lake	-7.3987410	-9.5449411	-5.2525408	0.0000000
Hummingbird Lake-Central Long Lake	-6.8931304	-9.8184178	-3.9678430	0.0000000
Paul Lake-Central Long Lake	-3.8521506	-5.9170942	-1.7872070	0.0000003
Peter Lake-Central Long Lake	-4.3501458	-6.4115874	-2.2887042	0.0000000
Tuesday Lake-Central Long Lake	-6.5971805	-8.6971605	-4.4972005	0.0000000
Ward Lake-Central Long Lake	-3.2077856	-6.1330730	-0.2824982	0.0193405
West Long Lake-Central Long Lake	-6.0877513	-8.2268550	-3.9486475	0.0000000
East Long Lake-Crampton Lake	-5.0842215	-6.5591700	-3.6092730	0.0000000
Hummingbird Lake-Crampton Lake	-4.5786109	-7.0538088	-2.1034131	0.0000004
Paul Lake-Crampton Lake	-1.5376312	-2.8916215	-0.1836408	0.0127491
Peter Lake-Crampton Lake	-2.0356263	-3.3842699	-0.6869828	0.0000999
Tuesday Lake-Crampton Lake	-4.2826611	-5.6895065	-2.8758157	0.0000000



```
## Ward Lake-Crampton Lake      -0.8932661 -3.3684639  1.5819317 0.9714459
## West Long Lake-Crampton Lake -3.7732318 -5.2378351 -2.3086285 0.0000000
## Hummingbird Lake-East Long Lake  0.5056106 -1.7364925  2.7477137 0.9988050
## Paul Lake-East Long Lake      3.5465903  2.6900206  4.4031601 0.0000000
## Peter Lake-East Long Lake      3.0485952  2.2005025  3.8966879 0.0000000
## Tuesday Lake-East Long Lake    0.8015604 -0.1363286  1.7394495 0.1657485
## Ward Lake-East Long Lake      4.1909554  1.9488523  6.4330585 0.0000002
## West Long Lake-East Long Lake  1.3109897  0.2885003  2.3334791 0.0022805
## Paul Lake-Hummingbird Lake     3.0409798  0.8765299  5.2054296 0.0004495
## Peter Lake-Hummingbird Lake    2.5429846  0.3818755  4.7040937 0.0080666
## Tuesday Lake-Hummingbird Lake  0.2959499 -1.9019508  2.4938505 0.9999752
## Ward Lake-Hummingbird Lake     3.6853448  0.6889874  6.6817022 0.0043297
## West Long Lake-Hummingbird Lake 0.8053791 -1.4299320  3.0406903 0.9717297
## Peter Lake-Paul Lake           -0.4979952 -1.1120620  0.1160717 0.2241586
## Tuesday Lake-Paul Lake         -2.7450299 -3.4781416 -2.0119182 0.0000000
## Ward Lake-Paul Lake            0.6443651 -1.5200848  2.8088149 0.9916978
## West Long Lake-Paul Lake       -2.2356007 -3.0742314 -1.3969699 0.0000000
## Tuesday Lake-Peter Lake        -2.2470347 -2.9702236 -1.5238458 0.0000000
## Ward Lake-Peter Lake           1.1423602 -1.0187489  3.3034693 0.7827037
## West Long Lake-Peter Lake      -1.7376055 -2.5675759 -0.9076350 0.0000000
## Ward Lake-Tuesday Lake         3.3893950  1.1914943  5.5872956 0.0000609
## West Long Lake-Tuesday Lake    0.5094292 -0.4121051  1.4309636 0.7374387
## West Long Lake-Ward Lake       -2.8799657 -5.1152769 -0.6446546 0.0021080
```

*#Then find groupings of lakes*

```
Temp.groups <- HSD.test(Jul.Temp.anova, "lakename", group = TRUE)
Temp.groups
```

```
## $statistics
##      MSerror  Df      Mean      CV
##      54.1016 9719 12.72087 57.82135
##
## $parameters
##      test  name.t ntr StudentizedRange alpha
##      Tukey lakename  9          4.387504 0.05
##
## $means
##               temperature_C      std      r      se Min  Max   Q25   Q50
## Central Long Lake      17.66641 4.196292  128 0.6501298 8.9 26.8 14.400 18.40
## Crampton Lake          15.35189 7.244773  318 0.4124692 5.0 27.5  7.525 16.90
## East Long Lake         10.26767 6.766804  968 0.2364108 4.2 34.1  4.975  6.50
## Hummingbird Lake       10.77328 7.017845  116 0.6829298 4.0 31.5  5.200  7.00
## Paul Lake              13.81426 7.296928 2660 0.1426147 4.7 27.7  6.500 12.40
## Peter Lake             13.31626 7.669758 2872 0.1372501 4.0 27.0  5.600 11.40
## Tuesday Lake           11.06923 7.698687 1524 0.1884137 0.3 27.7  4.400  6.80
## Ward Lake              14.45862 7.409079  116 0.6829298 5.7 27.6  7.200 12.55
## West Long Lake         11.57865 6.980789 1026 0.2296314 4.0 25.7  5.400  8.00
##
##               Q75
## Central Long Lake 21.000
## Crampton Lake    22.300
## East Long Lake    15.925
## Hummingbird Lake 15.625
## Paul Lake         21.400
## Peter Lake        21.500
```

```
## Tuesday Lake      19.400
## Ward Lake         23.200
## West Long Lake    18.800
##
## $comparison
## NULL
##
## $groups
##           temperature_C groups
## Central Long Lake    17.66641      a
## Crampton Lake        15.35189     ab
## Ward Lake            14.45862     bc
## Paul Lake            13.81426      c
## Peter Lake           13.31626      c
## West Long Lake       11.57865      d
## Tuesday Lake         11.06923     de
## Hummingbird Lake     10.77328     de
## East Long Lake       10.26767      e
##
## attr(,"class")
## [1] "group"
```

16. From the findings above, which lakes have the same mean temperature, statistically speaking, as Peter Lake? Does any lake have a mean temperature that is statistically distinct from all the other lakes?

Answer: The HSD.test shows that Paul Lake and Ward Lake have a mean temperature that is statistically the same as Peter Lake. It does not look like any lake is statistically distinct from all other lakes.

17. If we were just looking at Peter Lake and Paul Lake. What's another test we might explore to see whether they have distinct mean temperatures?

Answer: A two-sample t-test could be run since we want to test the hypothesis that the means of the two lakes are equivalent. A low p-value of less than 0.05 would tell us that the means ARE statistically different from each other.

18. Wrangle the July data to include only records for Crampton Lake and Ward Lake. Run the two-sample T-test on these data to determine whether their July temperature are same or different. What does the test say? Are the mean temperatures for the lakes equal? Does that match your answer for part 16?

```
# Wrangle the data
Cram.Ward.combo <- Jul.lake.chem.phys %>%
  filter(lakename %in% c("Crampton Lake", "Ward Lake"))

# Used the following to make sure the filter worked properly
#summary(Cram.Ward.combo$lakename)
#summary(Jul.lake.chem.phys$lakename)

# Two-sample T-test (null: the lake means are equal; alt: means are not equal)
temp.twosample <-
  t.test(Cram.Ward.combo$temperature_C ~ Cram.Ward.combo$lakename)
temp.twosample
```

```
##
## Welch Two Sample t-test
##
## data: Cram.Ward.combo$temperature_C by Cram.Ward.combo$lakename
## t = 1.1181, df = 200.37, p-value = 0.2649
## alternative hypothesis: true difference in means between group Crampton Lake and group Ward Lake is not equal to 0
## 95 percent confidence interval:
## -0.6821129 2.4686451
## sample estimates:
## mean in group Crampton Lake      mean in group Ward Lake
##                15.35189                14.45862
```

Answer: The results give a p-value  $> 0.05$ , therefore we fail to reject the null hypothesis, and can conclude that there is not a significant difference in the mean temperatures between the two lakes. This does match my number 16, HSD.test resulted in a grouping of Crampton and Ward lakes, which also meant that there was not a significant difference between the temperatures of those lakes.