

On the Behavior of the Fundamental Solution of the Heat Equation with Variable Coefficients*

S. R. S. VARADHAN

1. Introduction

The solution $q(t, x, y)$ of the equation

$$(1.1) \quad \frac{\partial q}{\partial t} = \frac{1}{2} \sum_{i=1}^k \frac{\partial^2 q}{\partial x_i^2}$$

with the boundary condition $q(t, x, y) \rightarrow \delta_x(y)$ as $t \rightarrow 0$ can of course be written explicitly as

$$(1.2) \quad q(t, x, y) = (2\pi t)^{-k/2} \exp \left\{ -\frac{1}{2t} \|x - y\|^2 \right\},$$

where $\|x - y\|$ denotes the Euclidean distance. Looking at (1.2) one sees immediately that

$$(1.3) \quad \lim_{t \rightarrow 0} [-2t \log q(t, x, y)] = \|x - y\|^2.$$

We shall consider the analogue of (1.1). Let $p(t, x, y)$ be the solution of the equation

$$(1.4) \quad \frac{\partial p}{\partial t} = \frac{1}{2} \sum_{i,j=1}^k a_{ij}(x) \frac{\partial^2 p}{\partial x_i \partial x_j}$$

with the boundary condition $p(t, x, y) \rightarrow \delta_x(y)$ as $t \rightarrow 0$. We prove a formula similar to (1.3):

$$(1.5) \quad \lim_{t \rightarrow 0} [-2t \log p(t, x, y)] = d^2(x, y),$$

where $d(x, y)$ is the distance induced by a Riemannian metric derived from the coefficients $\{a_{ij}(x)\}$.

* This paper represents results obtained at the Courant Institute of Mathematical Sciences, New York University, under the sponsorship of the Office of Naval Research, Contract Nonr-285(46), and a Sloan Foundation Grant for Probability and Statistics. Reproduction in whole or in part is permitted for any purpose of the United States Government.

If G is a region with a boundary B , we consider the solution $\phi(x, \lambda)$ of the equation

$$\frac{1}{2} \sum a_{ij}(x) \frac{\partial^2 \phi}{\partial x_i \partial x_j} = \lambda \phi \quad \text{for } x \in G$$

with the boundary value $\phi = 1$ on B . We prove that

$$(1.6) \quad \lim_{\lambda \rightarrow \infty} [-(2\lambda)^{-1/2} \log \phi(x, \lambda)] = d(x, B),$$

where x is any point of G and $d(x, B)$ is the shortest distance to the boundary B from x .

The behavior (1.5) of $p(t, x, y)$ is derived from the behavior (1.6) of $\phi(x, \lambda)$ which is proved first. Although these results are closely related to certain properties of Markov processes, the probabilistic connections will be explored separately, [3].

Section 2 covers the preliminaries. The assumptions and the main theorems are stated there. The later sections cover the actual proof.

2. Preliminaries

L stands for the following differential operator acting on smooth functions on R_k :

$$(2.1) \quad Lf = \frac{1}{2} \sum a_{ij}(x) \frac{\partial^2 f}{\partial x_i \partial x_j}.$$

The matrix $\{a(x)\}$ of coefficients is of course symmetric and positive definite and the following assumptions are made concerning $a(x)$:

(i) Uniform Hölder condition, i.e., for all i and j we have

$$|a_{ij}(x) - a_{ij}(y)| \leq M \|x - y\|^\alpha \text{ for all } x, y, \quad \alpha > 0.$$

(2.2) (ii) Uniform ellipticity condition, i.e., there exist constants α and A such that for any vector ξ_1, \dots, ξ_k , and x

$$A \sum \xi_i^2 \geq \sum a_{ij}(x) \xi_i \xi_j \geq \alpha \sum \xi_i^2.$$

We then consider the equation

$$(2.3) \quad \frac{\partial p}{\partial t} = Lp$$

with the boundary condition $p(t, x, y) \rightarrow \delta_x(y)$ as $t \rightarrow 0$.

The following theorem concerning the existence of a solution $p(t, x, y)$ for (2.3) can be found in [1], [2].

THEOREM 2.1. Under the hypothesis (2.2) on the coefficients there exists a solution $p(t, x, y)$ for equation (2.3) and it has the following properties:

- (i) $p(t, x, y) \geq 0$,
- (ii) $\int p(t, x, y)p(s, y, z) dy = p(t + s, x, z)$,
- (iii) $p(t, x, y)$ is continuous in t, x , and y ,
- (iv) there are constants M and α such that

$$p(t, x, y) \leq Mt^{-k/2} \exp \left\{ -\frac{\alpha}{2t} \|x - y\|^2 \right\},$$

- (v) there are positive constants $\alpha_1, \alpha_2, M_1, M_2, \lambda$ such that

$$p(t, x, y) \geq M_1 t^{-k/2} \exp \left\{ -\frac{\alpha_1}{2t} \|x - y\|^2 \right\} \\ - M_2 t^{-k/2+\lambda} \exp \left\{ -\frac{\alpha_2}{2t} \|x - y\|^2 \right\}.$$

Further $p(t, x, y)$ is unique.

We now introduce the metric d . Let $p(\tau)$, $0 \leq \tau \leq 1$, be a smooth path in R_k . Then the length of such a path is defined as

$$l(p) = \int_0^1 [\dot{p}(\tau) a^{-1}(p(\tau)) \dot{p}(\tau)]^{1/2} d\tau,$$

where $\dot{p}(\tau)$ stands for $dp(\tau)/d\tau$, $a^{-1}(p(\tau))$ for the matrix inverse to $a(p(\tau))$ and $(\theta a \theta)$ for the quadratic form $\sum a_{ij} \theta_i \theta_j$; $l(p)$ is the natural length in a metric defined locally as

$$ds^2 = \sum a^{ij}(x) dx_i dx_j.$$

The operator L modified by a first order term is Laplacian in this metric. The global distance $d(x, y)$ induced by this metric is defined as

$$(2.4) \quad d(x, y) = \inf_{p: \substack{p(0)=x \\ p(1)=y}} l(p).$$

We can now state the first main theorem concerning the behavior of $p(t, x, y)$ introduced in Theorem 2.1 for small t .

THEOREM 2.2.

$$\lim_{t \rightarrow 0} [-2t \log p(t, x, y)] = d^2(x, y)$$

uniformly over x and y such that $d(x, y)$ is bounded.

We consider an open set G in R_k . Let B be the boundary of G . It is assumed that every point b on B is a limit point of the exterior of G .

Let $\phi(x, \lambda)$ be the solution of the equation

$$(2.5) \quad \begin{aligned} L\phi &= \lambda\phi & \text{for } x \in G, \\ \phi &= 1 & \text{for } b \in B. \end{aligned}$$

The solution exists for $\lambda \geq 0$, if the boundary value is properly interpreted. The next theorem is concerned with the behavior of the solution $\phi(x, \lambda)$ for large λ .

THEOREM 2.3. *With the assumptions stated above,*

$$\lim_{\lambda \rightarrow \infty} \left[-\frac{1}{\sqrt{2\lambda}} \log \phi(x, \lambda) \right] = d(x, B)$$

uniformly over compact subsets of $G \cup B$.

We state some notation and a few elementary consequences concerning the distance $d(x, y)$. The distance $[(x - y) \Lambda (x - y)]^{1/2}$ is denoted by $\|x - y\|_\Lambda$, where Λ is a positive definite matrix. If Λ is chosen as $a^{-1}(z)$, then $\|x - y\|$ is denoted by $\|x - y\|_z$; $\|x - y\|$ is of course the Euclidean distance. It is clear from the assumptions that $d, \| \cdot \|, \| \cdot \|_z$ are all equivalent in the sense that the ratio of any two is bounded by a universal constant. We shall denote by $\Delta(x, y)$ any one of these metrics. If a and b are positive definite matrices, we shall mean by $a \geq b$ that $a - b$ is positive semidefinite. The following is a consequence of the Hölder condition in (2.2).

LEMMA 2.4. *There exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ such that if $\Delta(x, y) \leq \varepsilon$, then*

$$\begin{aligned} [1 - \delta(\varepsilon)]a(x) &\leq a(y) \leq [1 + \delta(\varepsilon)]a(x), \\ [1 - \delta(\varepsilon)]a^{-1}(x) &\leq a^{-1}(y) \leq [1 + \delta(\varepsilon)]a^{-1}(x). \end{aligned}$$

From Lemma 2.4 and the definition of $d(x, y)$ we deduce

LEMMA 2.5. *There exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ such that if $\Delta(x, y) \leq \varepsilon$ and $\Delta(x, z) \leq \varepsilon$, then*

$$[1 - \delta(\varepsilon)] \|x - y\|_z \leq d(x, y) \leq [1 + \delta(\varepsilon)] \|x - y\|_z.$$

3. Elliptic Equation

In this section we prove Theorem 2.3. The method consists of comparing our solution with the solution of an equation having constant coefficients in a small domain and then piecing together the estimate with the help of the maximum principle.

Let y be an arbitrary point in R_k . Let the set S_ε and its boundary B_ε be defined as follows:

$$S_\varepsilon = [x: \|x - y\|_y < \varepsilon],$$

$$B_\varepsilon = [x: \|x - y\|_y = \varepsilon].$$

We denote by L_y the operator

$$L_y f = \frac{1}{2} \sum a_{ij}(y) \frac{\partial^2 f}{\partial x_i \partial x_j}.$$

It is an operator with constant coefficients. Let $\psi_\varepsilon(x, \lambda)$ be the solution to the equation

$$(3.1) \quad \begin{aligned} L_y \psi_\varepsilon &= \lambda \psi_\varepsilon & \text{for } x \in S_\varepsilon, \\ \psi_\varepsilon &= 1 & \text{for } x \in B_\varepsilon. \end{aligned}$$

THEOREM 3.1. For every $\lambda > 0$, $\psi(x, \lambda)$ is explicitly given by the formula below for $x \in S_\varepsilon$:

$$\begin{aligned} \psi_\varepsilon(x, \lambda) &= \frac{\cosh(\sqrt{2\lambda} r)}{\cosh(\sqrt{2\lambda} \varepsilon)} & \text{for } k = 1, \\ \psi_\varepsilon(x, \lambda) &= \frac{\int_0^\pi \cosh(\sqrt{2\lambda} r \cos \theta) \sin^{k-2} \theta d\theta}{\int_0^\pi \cosh(\sqrt{2\lambda} \varepsilon \cos \theta) \sin^{k-2} \theta d\theta} & \text{for } k \geq 2, \end{aligned}$$

where $r = \|x - y\|_y$. For all k , $\psi_\varepsilon(x, \lambda)$ is a convex function of x .

Proof: The proof is elementary and consists of direct verification.

THEOREM 3.2. For every $\rho > 0$ there exists a constant $M_\rho < \infty$ depending only on the dimension k such that, for all y, ε and λ ,

$$\psi_\varepsilon(y, \lambda) \leq M_\rho \exp \{-\varepsilon(1 - \rho)\sqrt{2\lambda}\}.$$

Proof: When $x = y$ or $r = 0$ the numerator in Theorem 3.1 reduces to a dimension constant. As for the denominator, it can be estimated from below by limiting the range of integration to the region $0 \leq \theta \leq \cos^{-1}(1 - \rho)$.

We now consider the sets

$$T_\varepsilon = [x: d(x, y) < \varepsilon],$$

$$D_\varepsilon = [x: d(x, y) = \varepsilon].$$

Then in view of Lemma 2.5, there exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ such that the set

$$S = [x: \|x - y\|_y < \varepsilon(1 - \delta(\varepsilon))]$$

is contained in T_ε . Let B be the boundary of S :

$$B = [x: \|x - y\|_y = \varepsilon(1 - \delta(\varepsilon))].$$

Obviously B is contained in $T_\varepsilon \cup D_\varepsilon$.

Let $f_\varepsilon(x, \lambda)$, $g_\varepsilon(x, \lambda)$, and $h_\varepsilon(x, \lambda)$ denote the solutions of the following equations:

$$(3.2) \quad \begin{aligned} Lf &= \lambda f & \text{for } x \in T_\varepsilon, \\ f &= 1 & \text{for } x \in D_\varepsilon, \end{aligned}$$

$$(3.3) \quad \begin{aligned} Lg &= \lambda g & \text{for } x \in S, \\ g &= 1 & \text{for } x \in B, \end{aligned}$$

$$(3.4) \quad \begin{aligned} L_y h &= \lambda h & \text{for } x \in S, \\ h &= 1 & \text{for } x \in B. \end{aligned}$$

LEMMA 3.3.

$$f(y, \lambda) \leq g(y, \lambda).$$

Proof: From equation (3.2) it is obvious that $f \leq 1$ in $T_\varepsilon \cup D_\varepsilon$. Since $B \subset T_\varepsilon \cup D_\varepsilon$, it follows that $f \leq 1$ on B . But f and g satisfy the same equation in S and on the boundary B of S , $g = 1$ and $f \leq 1$. Hence $f \leq g$ everywhere in S . Therefore in particular, $f(y, \lambda) \leq g(y, \lambda)$.

LEMMA 3.4. There exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ such that

$$g(y, \lambda) \leq h(y, \lambda(1 - \delta(\varepsilon))).$$

Proof: By Theorem 3.1, h is convex and by Lemma 2.4

$$a(x) \leq \frac{1}{1 - \delta(\varepsilon)} a(y).$$

Therefore,

$$\begin{aligned} \frac{1}{2} \sum a_{ij}(x) \frac{\partial^2 h}{\partial x_i \partial x_j} &\leq \frac{1}{2(1 - \delta(\varepsilon))} \sum a_{ij}(y) \frac{\partial^2 h}{\partial x_i \partial x_j} \\ &= \frac{1}{(1 - \delta(\varepsilon))} L_y h \\ &= \frac{\lambda}{(1 - \delta(\varepsilon))} h, \end{aligned}$$

or

$$Lh(x, \lambda(1 - \delta(\varepsilon))) \leq \lambda h \quad \text{in} \quad S.$$

But

$$Lg(x, \lambda) = \lambda g \quad \text{in} \quad S,$$

and h and g have the same boundary value on B . Hence the solution $g(x, \lambda)$ is smaller than $h(x, \lambda(1 - \delta(\varepsilon)))$ in S . In particular,

$$g(y, \lambda) \leq h(y, \lambda(1 - \delta(\varepsilon))).$$

THEOREM 3.5. *The solution $f_\varepsilon(x, \lambda)$ of equation (3.2) has the following property. There exist a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ and a constant $M_\rho < \infty$ for every $\rho > 0$ such that, for all ε, y and λ ,*

$$f_\varepsilon(y, \lambda) \leq M_\rho \exp \{-(1 - \delta(\varepsilon))(1 - \rho)\varepsilon\sqrt{2\lambda}\}.$$

The proof is immediate from Theorem 3.2 and Lemmas 3.3 and 3.4.

Let G be any arbitrary region with a boundary B . Let $\phi(x, \lambda)$ be the solution to the equation

$$\begin{aligned} L\phi &= \lambda\phi & \text{for} & \quad x \in G, \\ \phi &= 1 & \text{for} & \quad x \in B. \end{aligned} \tag{3.5}$$

If G is unbounded, we determine ϕ by the additional restriction that $0 \leq \phi \leq 1$. If G is bounded, it is automatically true that $0 \leq \phi \leq 1$. Let

$$d(x, B) = \inf_{b \in B} d(x, b). \tag{3.6}$$

We then have the following estimate concerning $\phi(x, \lambda)$.

THEOREM 3.6. *For every $\rho > 0$, there exists a function $C_\rho(x)$ such that*

$$\phi(x, \lambda) \leq C_\rho(x) \exp \{-(1 - \rho)d(x, B)\sqrt{2\lambda}\}.$$

$C_\rho(x)$ depends only on $d(x, B)$ and is uniformly bounded whenever x varies in such a way that $d(x, B)$ remains bounded.

Proof: Let $x \in G$ be arbitrary. Then the set

$$S_R = [y: d(x, y) < d(x, B)]$$

is obviously contained in G and its boundary is

$$B_R = [y: d(x, y) = d(x, B)].$$

Hence if $\psi(x, \lambda)$ is the solution of

$$(3.7) \quad \begin{aligned} L\psi &= \lambda\psi & \text{for } y \in S_R, \\ \psi &= 1 & \text{for } y \in B_R, \end{aligned}$$

then

$$\phi(x, \lambda) \leq \psi(x, \lambda).$$

To estimate $\psi(x, \lambda)$ we choose ε small so that $n\varepsilon = d(x, B)$ and define the sets S_1, S_2, \dots, S_n as follows:

$$S_j = [y: d(x, y) < j\varepsilon],$$

$$B_j = [y: d(x, y) = j\varepsilon].$$

Let ψ_j be the solution to the equation

$$(3.8) \quad \begin{aligned} L\psi_j &= \lambda\psi_j & \text{for } y \text{ in } S_j, \\ \psi_j &= 1 & \text{for } y \text{ on } B_j. \end{aligned}$$

We estimate ψ_{j+1} on B_j . It is obvious that if $b_j \in B_j$, the set $S = [y: d(b_j, y) < \varepsilon]$ is contained in S_{j+1} and hence

$$\psi_{j+1}(b_j, \lambda) \leq f_\varepsilon(b_j, \lambda),$$

where f_ε refers to the function defined in equation (3.2). Therefore,

$$(3.9) \quad \psi_{j+1}(b_j, \lambda) \leq M_\rho \exp \{ -[1 - \delta(\varepsilon)](1 - \rho)\sqrt{2\lambda} \varepsilon \}.$$

Now we can use the inequality

$$\phi(b_j, \lambda) \leq \psi_{j+1}(b_j, \lambda) \sup_{b_{j+1} \in B_{j+1}} \phi(b_{j+1}, \lambda)$$

which with (3.9) leads to

$$(3.10) \quad \phi(x, \lambda) \leq [M_\rho]^n \exp \{ -[1 - \delta(\varepsilon)](1 - \rho)n\varepsilon\sqrt{2\lambda} \}.$$

Since $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ and since $n\varepsilon = d(x, B)$, this leads immediately to the theorem.

For the rest of the section we shall be concerned with obtaining a lower bound.

Let y be an arbitrary point in R_k , let S be the set

$$S = [x: \|x - y\|_v < \varepsilon]$$

and B its boundary

$$B = [x: \|x - y\|_v = \varepsilon].$$

Γ is that part of the boundary B defined by

$$\Gamma = [x: \|x - y\|_y = \varepsilon, \|x - z\|_y < \varepsilon\sqrt{2\eta}] ,$$

where z is a point of B and η is a positive number. So Γ is a small portion of the boundary around z .

Let $\phi(x, \lambda)$ be the solution to the equation

$$\begin{aligned} (3.11) \quad L\phi &= \lambda\phi \quad \text{for } x \text{ in } S, \\ \phi &= 1 \quad \text{for } x \text{ on } \Gamma, \\ \phi &= 0 \quad \text{for } x \text{ on } B - \Gamma. \end{aligned}$$

We have the following theorem giving us a lower bound on $\phi(y, \lambda)$.

THEOREM 3.7. *There exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ and a constant M_ρ for every $\rho > 0$ such that for all y and λ*

$$\phi(y, \lambda) \geq \exp \{-\varepsilon(1 + \delta(\varepsilon))\sqrt{2\lambda}\} [1 - M_\rho \exp \{-\varepsilon(\eta - \rho - \delta(\varepsilon))\sqrt{2\lambda}\}] .$$

Proof: Let θ be a vector in R_k . For any z in R_k , θz is the scalar product $\sum \theta_i z_i$. For x in S we have

$$\begin{aligned} Le^{\theta(x-y)} &= \frac{1}{2} [\theta a(x) \theta] e^{\theta(x-y)} \\ &\geq \frac{1}{2} (1 - \delta(\varepsilon)) (\theta \Lambda \theta) e^{\theta(x-y)}, \end{aligned}$$

where $\Lambda = a(y)$. Putting $\lambda = \frac{1}{2}(1 - \delta(\varepsilon))[\theta \Lambda \theta]$, we have

$$Le^{\theta(x-y)} \geq \lambda e^{\theta(x-y)} \quad \text{for } x \text{ in } S.$$

Hence if $g(x, \lambda)$ is the solution to

$$\begin{aligned} (3.12) \quad Lg &= \lambda g \quad \text{in } S, \\ g &= e^{\theta(x-y)} \quad \text{for } x \text{ on } B, \end{aligned}$$

then

$$(3.13) \quad e^{\theta(x-y)} \leq g(x, \lambda) .$$

Let $\psi(x, \lambda)$ be the solution to

$$\begin{aligned} (3.14) \quad L\psi &= \lambda\psi \quad \text{for } x \text{ in } S, \\ \psi &= 1 \quad \text{for } x \text{ on } B. \end{aligned}$$

Then from Theorem 3.2, there is, for every $\rho > 0$, a constant M_ρ such that

$$(3.15) \quad \psi(y, \lambda) \leq M_\rho \exp \{-\varepsilon(1 - \rho)\sqrt{2\lambda}\} .$$

Let us now take the vector $\theta = k\Lambda^{-1}(z - y)$, where the constant k is to be chosen later, such that

$$(3.16) \quad \lambda = \frac{1}{2}(1 - \delta(\varepsilon))[\theta\Lambda\theta] .$$

The following estimates are easily obtained:

$$\begin{aligned} \sup_{x \in B} \theta(x - y) &= k\varepsilon^2, \\ \sup_{x \in B - \Gamma} \theta(x - y) &= k\varepsilon^2(1 - \eta) . \end{aligned}$$

Hence on B ,

$$(3.17) \quad e^{\theta(x-y)} \leq e^{k\varepsilon^2(1-\eta) + k\varepsilon^2} \chi_{\Gamma}(x) ,$$

where $\chi_{\Gamma}(x)$ is the indicator function of Γ . Let us recall that $\phi(x, \lambda)$ satisfies the equation

$$(3.18) \quad \begin{aligned} L\phi &= \lambda\phi & \text{for } x &\text{ in } S, \\ \phi &= \chi_{\Gamma}(x) & \text{for } x &\text{ on } B, \end{aligned}$$

and compare the equations (3.12), (3.14) and (3.18). The functions g , ψ , and ϕ satisfy the same equation in S but with different boundary values $e^{\theta(x-y)}$, 1, and $\chi_{\Gamma}(x)$, respectively, on B . Since (3.17) is valid on the boundary, the following inequality is valid throughout S :

$$(3.19) \quad g(x, \lambda) \leq e^{k\varepsilon^2(1-\eta)}\psi(x, \lambda) + e^{k\varepsilon^2}\phi(x, \lambda) .$$

Setting $x = y$ in (3.19) and using (3.13), we obtain

$$e^{k\varepsilon^2(1-\eta)}\psi(y, \lambda) + e^{k\varepsilon^2}\phi(y, \lambda) \geq 1 .$$

Using now the estimate (3.15) for $\psi(y, \lambda)$, we get

$$(3.20) \quad \phi(y, \lambda) \geq e^{-k\varepsilon^2} - e^{-k\varepsilon^2} \eta M_{\rho} e^{-\varepsilon(1-\rho)\sqrt{2\lambda}} .$$

Now (3.16) leads to the value $\varepsilon k = [1 + \delta(\varepsilon)]\sqrt{2\lambda}$. Substitution of this value of k in (3.20), a little simplification and replacement of $\delta(\varepsilon)$ by a larger function which again tends to zero as $\varepsilon \rightarrow 0$ yields

$$\phi(y, \lambda) \geq e^{-\varepsilon(1+\delta(\varepsilon))\sqrt{2\lambda}} (1 - M_{\rho} e^{-\varepsilon(\eta-\rho-\delta(\varepsilon))\sqrt{2\lambda}}) .$$

This completes the proof.

Let y and z be two arbitrary points in R_k with $d(x, y) = \varepsilon$. Let S be the set

$$S = [x: d(x, z) \leq 2\varepsilon\sqrt{\eta}] .$$

We consider the solution $\phi(x, \lambda)$ of the equation

$$(3.21) \quad \begin{aligned} L\phi &= \lambda\phi \quad \text{for } x \text{ in the exterior of } R_k - S, \\ \phi &= 1 \quad \text{for } x \text{ on the boundary } B \text{ of } S. \end{aligned}$$

We further assume that η is small enough so that y is in the exterior of S . Then we have

THEOREM 3.8. *There exists a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$, and a constant M_ρ for every $\rho > 0$ such that, for all y and z ,*

$$\phi(y, \lambda) \geq \exp \{-\varepsilon(1 + \delta(\varepsilon))\sqrt{2\lambda}\} [1 - M_\rho \exp \{-(\eta - \rho - \delta(\varepsilon))\sqrt{2\lambda}\}].$$

Proof: Let $\varepsilon' = \|y - z\|_y$. Choose η' such that the set

$$E = [x: \|x - z\|_y \leq \varepsilon'\sqrt{2\eta'}]$$

is contained in S . Let F denote the boundary

$$F = [x: \|x - z\|_y = \varepsilon'\sqrt{2\eta'}]$$

of S . Let $\psi(x, \lambda)$ be the solution to the equation

$$(3.22) \quad \begin{aligned} L\psi &= \lambda\psi \quad \text{for } x \text{ in the exterior of } E, \\ \psi &= 1 \quad \text{for } x \text{ on } F. \end{aligned}$$

Let S' be the set

$$S' = [x: \|x - y\|_y < \varepsilon']$$

with boundary

$$B' = [x: \|x - y\|_y = \varepsilon']$$

and Γ' the set $\Gamma' = B' \cap E$. Let $g(x, \lambda)$ be the solution of

$$(3.23) \quad \begin{aligned} Lg &= \lambda g \quad \text{for } x \text{ in } S', \\ g &= \chi_{\Gamma'}(x) \quad \text{for } x \text{ on } B'. \end{aligned}$$

Since $E \subset S$ and ψ is the solution to (3.22) in the exterior of E , ψ is defined on the boundary B of S and $0 \leq \psi \leq 1$ on B . But ϕ is the solution of (3.21) in the exterior of S and therefore for x in the exterior of S

$$(3.24) \quad \phi(x, \lambda) \geq \psi(x, \lambda).$$

Let A be the set $S' \cap (\text{exterior of } E)$. Its boundary is given by $[B' \cap (\text{exterior of } E)] \cup (F \cap S')$. Then g and ψ are both solutions of the same equation $Lf = \lambda f$ in A . As for their boundary values,

$$\begin{aligned} \text{on } B' \cap (\text{exterior of } E), \quad g &= 0, \quad \psi \geq 0, \\ \text{on } F \cap S', \quad g &\leq 1, \quad \psi = 1. \end{aligned}$$

Hence throughout A , $\psi \geq g$. Since y is in A , we have with (3.24)

$$\phi(y, \lambda) \geq g(y, \lambda).$$

Using Theorem 3.7 to estimate $g(y, \lambda)$, we have

$$g(y, \lambda) \geq \exp \{-\varepsilon'(1 + \delta(\varepsilon'))\sqrt{2\lambda}\} [1 - M_\rho \exp \{-(\eta' - \rho - \delta(\varepsilon'))\sqrt{2\lambda}\}].$$

It is easy to estimate using Lemma 2.5:

$$\varepsilon' \leq \varepsilon(1 + \delta(\varepsilon))$$

and

$$\eta' \geq 2\eta \left(\frac{1 - \delta(\varepsilon)}{1 + \delta(\varepsilon)} \right)^2.$$

Therefore for ε small enough, $\eta' \geq \eta$ and this proves the theorem.

Let y, z be two points with $d(y, z) = d$. Let n be an integer such that $n\varepsilon = d$. Let G be the sphere around z of radius $4\varepsilon\sqrt{\eta}$:

$$G = [x: d(x, z) \leq 4\varepsilon\sqrt{\eta}].$$

Let $\phi(x, \lambda)$ be the solution of the equation

$$\begin{aligned} L\phi &= \lambda\phi \quad \text{for } x \text{ in the exterior of } G, \\ \phi &= 1 \quad \text{on the boundary of } G. \end{aligned} \tag{3.25}$$

THEOREM 3.9. *For the solution ϕ of equation (3.25) there exist a function $\delta(\varepsilon) \downarrow 0$ as $\varepsilon \downarrow 0$ and a constant M_ρ for every $\rho > 0$ such that*

$$\begin{aligned} \phi(y, \lambda) &\geq \exp \{-d(1 + 4\sqrt{\eta})(1 + \delta(\varepsilon))\sqrt{2\lambda}\} \\ &\times [1 - M_\rho \exp \{-\varepsilon(1 - 4\sqrt{\eta})(\eta - \rho - \delta(\varepsilon))\sqrt{2\lambda}\}]^n. \end{aligned}$$

Proof: Let the points $y = y_0, y_1, y_2, \dots, y_n = z$ be chosen such that $d(y_i, y_{i+1}) = \varepsilon$; $n\varepsilon$ being equal to d . Let $\varepsilon > 0$. Construct the sets C_i successively as follows:

$$\begin{aligned} C_1 &= \overline{S(y_1, 2\varepsilon\sqrt{\eta})}, \\ C_2 &= \overline{\bigcup_{z_1 \in C_1} S(y_2, 2\sqrt{\eta} d(z_1, y_2))}, \\ C_{j+1} &= \overline{\bigcup_{z_j \in C_j} S(y_{j+1}, 2\sqrt{\eta} d(z_j, y_{j+1}))}, \end{aligned}$$

and let

$$\begin{aligned}\Delta_j &= \sup_{z_j \in C_j} d(z_j, y_j), \\ \rho_{j-1} &= \sup_{z_{j-1} \in C_{j-1}} d(z_{j-1}, y_j), \\ \mu_{j-1} &= \inf_{z_{j-1} \in C_{j-1}} d(z_{j-1}, y_j).\end{aligned}$$

Then

$$\begin{aligned}\rho_{j-1} &\leq \varepsilon + \Delta_{j-1}, \\ \mu_{j-1} &\geq \varepsilon - \Delta_{j-1}, \\ \Delta_j &\leq 2\sqrt{\eta} \rho_{j-1}, \\ \rho_0 &= \varepsilon, \quad \Delta_0 = 0.\end{aligned}$$

It follows from these equations that

$$\begin{aligned}\Delta_j &\leq \varepsilon[2\sqrt{\eta} + (2\sqrt{\eta})^2 + \cdots + (2\sqrt{\eta})^j] \\ &\leq \varepsilon 2\sqrt{\eta} \frac{1}{1 - 2\sqrt{\eta}} \\ &\leq 4\varepsilon\sqrt{\eta} \quad \text{for} \quad 2\sqrt{\eta} \leq \frac{1}{2}, \quad j = 1, 2, \cdots, n.\end{aligned}$$

If ε_j is a typical distance from C_j to y_{j+1} , then

$$\varepsilon - \Delta_j \leq \varepsilon_j \leq \varepsilon + \Delta_j,$$

so that

$$\varepsilon(1 - 4\sqrt{\eta}) \leq \varepsilon_j \leq \varepsilon(1 + 4\sqrt{\eta}).$$

Now we can apply Theorem 3.8 and obtain a uniform estimate for x in C_{n-1} . Repeated application of Theorem 3.8 leads to

$$\begin{aligned}\phi(y, \lambda) &\geq \exp\{-d(1 + 4\sqrt{\eta})(1 + \delta(\varepsilon))\sqrt{2\lambda}\} \\ &\quad \times [1 - M_\rho \exp\{-\varepsilon(1 - 4\sqrt{\eta})(\eta - \rho - \delta(\varepsilon))\sqrt{2\lambda}\}]^n.\end{aligned}$$

THEOREM 3.10. *Let z be a point in R_k , S the sphere $S = [x: d(x, z) < \delta]$. Let $\phi(x, \lambda)$ be the solution to the equation $L\phi = \lambda\phi$ in the exterior of S with $\phi = 1$ on the boundary of S . Then*

$$\liminf_{\lambda \rightarrow \infty} \frac{1}{\sqrt{2\lambda}} \log \phi(x, \lambda) \geq -d(x, z)$$

uniformly over all x, z and δ , provided δ remains bounded away from zero and $d(x, z)$ remains bounded.

The proof follows at once from Theorem 3.9. One chooses η first and then ρ and ε suitably; n remains bounded for any $\varepsilon > 0$ if d is bounded.

Let G be a region with exterior E . B is the boundary of G . It is assumed that B is also the boundary of E .

LEMMA 3.11. *Let $x \in G$ and let $d = d(x, B)$. Then for any $\varepsilon > 0$ there exists a $\delta > 0$ and a point $z \in E$ such that*

$$(i) \quad d(x, z) \leq d + \varepsilon,$$

$$(ii) \quad [y: d(y, z) < \delta] \subset E.$$

Moreover δ can be chosen uniformly when x varies over a bounded subset of G .

Proof: Let $x \in G \cup B$. Then there exists a y on B such that $d(x, B) = d(x, y)$. The sphere around y of radius $\frac{1}{2}\varepsilon$ has a non-empty intersection with E and there exists a sphere around some point z of some radius δ completely contained inside the intersection. Obviously $d(x, z) \leq d(x, y) + \frac{1}{2}\varepsilon$. To show uniformity we point out that if x' is in $G \cup B$ with $d(x, x') < \frac{1}{2}\varepsilon$, then

$$\begin{aligned} d' = d(x', B) &\geq d(x, B) - d(x, x') \\ &\geq d - \frac{1}{2}\varepsilon \\ &\geq d(x, z) - \frac{1}{2}\varepsilon - \frac{1}{2}\varepsilon \\ &= d(x, z) - \varepsilon, \end{aligned}$$

or

$$d(x, z) \leq d' + \varepsilon.$$

In other words, the sphere of radius δ around z serves not only for x but for a neighborhood around x as well. A routine compactness argument using finite subcovers proves the lemma for compact subsets of $G \cup B$ or for bounded subsets of G .

Now we can prove the main theorem:

THEOREM 3.12. *Let G be any region with boundary B . B is assumed to be the boundary of the exterior E of G as well. If $\phi(x, \lambda)$ is the solution of the equation*

$$\begin{aligned} L\phi &= \lambda\phi \quad \text{for } x \text{ in } G, \\ \phi &= 1 \quad \text{on } B, \end{aligned}$$

then

$$\lim_{\lambda \rightarrow \infty} \left[-\frac{1}{\sqrt{2\lambda}} \log \phi(x, \lambda) \right] = d(x, B)$$

uniformly for x varying over bounded subsets of G .

The proof follows from Theorems 3.6, 3.10, and Lemma 3.11.

4. The Fundamental Solution

Let $F(t)$ be a nondecreasing function of t for $t \geq 0$ with $0 \leq F(t) \leq 1$ and $F(0) = 0$. Let $\phi(\lambda)$ be the Laplace transform

$$(4.1) \quad \phi(\lambda) = \int_0^\infty e^{-\lambda\tau} dF(\tau).$$

We assume that the $\{\phi_\alpha(\lambda)\}$ are Laplace transforms of $\{F_\alpha(t)\}$ and that

$$(4.2) \quad \lim_{\lambda \rightarrow \infty} \left[-\frac{1}{\sqrt{2\lambda}} \log \phi_\alpha(\lambda) \right] = d_\alpha,$$

where the d_α are such that $0 \leq d_\alpha \leq D < \infty$. We want to conclude that if the limit in (4.2) is uniform with respect to α , then

$$(4.3) \quad \lim_{t \rightarrow 0} \left[-2t \log F_\alpha(t) \right] = d_\alpha^2$$

uniformly in α .

Initially we assume that (4.2) holds uniformly and that in addition $0 < d \leq d_\alpha \leq D < \infty$. The following lemmas are conclusions regarding $F_\alpha(t)$.

LEMMA 4.1. *For every $\varepsilon > 0$ there exists a constant $A_\varepsilon < \infty$ such that, for all $t \geq 0$,*

$$F_\alpha(t) \leq A_\varepsilon \exp \left\{ -\frac{(1-\varepsilon)}{2t} d_\alpha^2 \right\}.$$

Proof: Since $0 < d \leq d_\alpha \leq D < \infty$ we can conclude, from the fact that (4.2) holds uniformly, that for every $\varepsilon > 0$ there exists a constant A_ε such that for all $\lambda \geq 0$

$$(4.4) \quad \phi_\alpha(\lambda) \leq A_\varepsilon \exp \{ -\sqrt{1-\varepsilon} \sqrt{2\lambda} d_\alpha \}.$$

It follows from (4.1) that

$$\phi_\alpha(\lambda) \geq e^{-\lambda t} F_\alpha(t),$$

or

$$(4.5) \quad F_\alpha(t) \leq e^{\lambda t} \phi_\alpha(\lambda).$$

Choosing $\lambda = (1-\varepsilon)d_\alpha^2/2t$ and combining (4.5) and (4.4) we get

$$F_\alpha(t) \leq A_\varepsilon \exp \left\{ -\frac{(1-\varepsilon)}{2t} d_\alpha^2 \right\}.$$

COROLLARY 4.2.

$$\limsup_{t \rightarrow 0} 2t \log F_\alpha(t) \leq -d_\alpha^2$$

uniformly in α .LEMMA 4.3. For every $k > 1$,

$$\int_{kt}^{\infty} \exp \left\{ -\frac{d_\alpha^2 \tau}{2t^2} \right\} F_\alpha(\tau) d\tau \leq \frac{2A_\varepsilon t^2}{\varepsilon d^2} \exp \left\{ -\frac{(1-\varepsilon)d_\alpha^2}{2t} - \frac{\rho_k}{t} \right\},$$

where $\rho_k = \frac{1}{2}d^2(k + 1/k - 2)$.

Proof: Using Lemma 4.1, we have

$$\begin{aligned} \int_{kt}^{\infty} \exp \left\{ -\frac{d_\alpha^2 \tau}{2t} \right\} F_\alpha(\tau) d\tau &\leq A_\varepsilon \int_{kt}^{\infty} \exp \left\{ -\frac{d_\alpha^2 \tau}{2t^2} \right\} \exp \left\{ -\frac{(1-\varepsilon)d_\alpha^2}{2t} \right\} d\tau \\ &= A_\varepsilon t \int_k^\infty \exp \left\{ -\frac{1}{2t} d_\alpha^2 \left[u + \frac{(1-\varepsilon)}{u} \right] \right\} du \\ &= A_\varepsilon t \int_k^\infty \exp \left\{ -\frac{\varepsilon d_\alpha^2}{2t} \right\} \exp \left\{ -\frac{(1-\varepsilon)d_\alpha^2}{2t} \left[u + \frac{1}{u} \right] \right\} du \\ &\leq \frac{2A_\varepsilon t^2}{\varepsilon d^2} \exp \left\{ -\frac{(1-\varepsilon)d_\alpha^2}{t} - \frac{\rho_k}{t} \right\}. \end{aligned}$$

LEMMA 4.4. For every $k' < 1$,

$$\int_0^{k't} \exp \left\{ -\frac{d_\alpha^2 \tau}{2t^2} \right\} F_\alpha(\tau) d\tau \leq \frac{2A_\varepsilon t^2}{\varepsilon d^2} \exp \left\{ -\frac{(1-\varepsilon)d_\alpha^2}{t} - \frac{\rho_{k'}}{t} \right\},$$

where $\rho_{k'} = \frac{1}{2}d^2[k' + 1/k' - 2]$.

The proof is similar to that of Lemma 4.3.

LEMMA 4.5.

$$\lim_{t \rightarrow 0} \left[-2t \log F_\alpha(t) \right] = d_\alpha^2$$

uniformly in α .Proof: Because of (4.2), there exists, for every $\varepsilon > 0$, a constant B_ε such that for all $\lambda \geq 0$

$$(4.6) \quad \phi_\alpha(\lambda) \geq B_\varepsilon \exp \{ -(1+\varepsilon)\sqrt{2\lambda} d_\alpha \}.$$

However,

$$\begin{aligned}
 \phi_\alpha(\lambda) &= \lambda \int_0^\infty e^{-\lambda\tau} F_\alpha(\tau) d\tau \\
 (4.7) \quad &= \lambda \left[\int_0^{k't} + \int_{k't}^{kt} + \int_{kt}^\infty \right] e^{-\lambda\tau} F_\alpha(\tau) d\tau.
 \end{aligned}$$

Therefore, substituting $\lambda = d_\alpha^2/2t^2$ and combining (4.6) and (4.7) with Lemmas 4.3 and 4.4, we have

$$(4.8) \quad \int_{k't}^{kt} \exp \left\{ -\frac{d_\alpha^2\tau}{2t^2} \right\} F_\alpha(\tau) d\tau \geq C_\varepsilon t^2 \exp \left\{ \frac{(1+\varepsilon)d_\alpha^2}{t} \right\} \left(1 - k_\varepsilon \exp \left\{ \frac{2\varepsilon D^2 - \rho}{t} \right\} \right),$$

where $\rho = \min(\rho_k, \rho_{k'})$.

For a given k and k' with $k' < 1 < k$ one can find ε small enough so that $2\varepsilon D^2 - \rho < 0$. Therefore, we conclude from (4.8) that, for every k, k' with $k' < 1 < k$,

$$(4.9) \quad \liminf_{t \rightarrow 0} t \log \left[\int_{k't}^{kt} \exp \left\{ -\frac{d_\alpha^2}{2t^2} \right\} F_\alpha(\tau) d\tau \right] \geq -d_\alpha^2$$

uniformly in α .

On the other hand,

$$\int_{k't}^{kt} \exp \left\{ -\frac{d_\alpha^2\tau}{2t^2} \right\} F_\alpha(\tau) d\tau \leq F_\alpha(kt) \exp \left\{ -\frac{k'd_\alpha^2}{2t} \right\}$$

which with (4.9) yields

$$\liminf_{t \rightarrow 0} t \log F_\alpha(kt) \geq -d_\alpha^2 + \frac{1}{2}k'd_\alpha^2,$$

or

$$\liminf_{t \rightarrow 0} 2t \log F_\alpha(t) \geq -2kd_\alpha^2 + kk'd_\alpha^2$$

uniformly in α . Since k and k' can be chosen close to 1, we have

$$\liminf_{t \rightarrow 0} 2t \log F_\alpha(t) \geq -d_\alpha^2$$

uniformly in α . This is one part of the lemma and the other part is Corollary 4.2.

Now we drop the assumption that $d > 0$. We assume only that $0 \leq d_\alpha \leq D$.

LEMMA 4.6. *Lemma 4.5 still holds if we assume only that $0 \leq d_\alpha \leq D$.*

Proof: It is of course enough to show that if, for $\lambda \geq \lambda_0$,

$$-\varepsilon \leq -\frac{1}{\sqrt{2\lambda}} \log \phi_\alpha(\lambda) \leq \varepsilon,$$

then there exists a t_0 and a function $\eta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ such that

$$-\eta \leq -2t \log F_\alpha(t) \leq \eta.$$

Since $0 \leq \phi_\alpha(\lambda) \leq 1$ and $0 \leq F_\alpha(t) \leq 1$, one side is trivial. If

$$\phi_\alpha(\lambda) \geq e^{-\varepsilon\sqrt{2\lambda}} \quad \text{for} \quad \lambda \geq \lambda_0,$$

we shall show that for ε small enough there exist η and t_0 such that

$$F_\alpha(t) \geq e^{-\eta/t} \quad \text{for} \quad t \leq t_0.$$

To prove this we notice that

$$\phi_\alpha(\lambda) \leq F_\alpha(t) + e^{-\lambda t},$$

or

$$\begin{aligned} F_\alpha(t) &\geq \phi_\alpha(\lambda) - e^{-\lambda t} \\ &\geq e^{-\varepsilon\sqrt{2\lambda}} - e^{-\lambda t} \quad \text{for} \quad \lambda \geq \lambda_0. \end{aligned}$$

Choose $\lambda = 2/t^2$. If $t \leq t_0 = \sqrt{2/\lambda_0}$ and $\varepsilon < \frac{1}{2}$, then

$$f_\alpha(t) \geq e^{-2\varepsilon/t}(1 - e^{-1/t})$$

which concludes the proof.

We have, therefore, proved the following theorem:

THEOREM 4.7. Let $\phi_\alpha(\lambda) = \int_0^\infty e^{-\lambda t} dF_\alpha(t)$. If

$$\lim_{\lambda \rightarrow \infty} \left[-\frac{1}{\sqrt{2\lambda}} \log \phi_\alpha(\lambda) \right] = d_\alpha$$

uniformly in α and if $0 \leq d_\alpha \leq D < \infty$, then

$$\lim_{t \rightarrow 0} \left[-2t \log F_\alpha(t) \right] = d_\alpha^2$$

uniformly in α .

Let $G \subset R_k$ be a region with a boundary B . It is assumed that B is the boundary of the exterior of G as well. Let S be the cylinder in $R_k \times (0, \infty)$ with base G .

That is,

$$S = G \times (0, \infty).$$

The boundary of S is $\{B \times (0, \infty)\} \cup \{G \times \{0\}\}$. We consider the solution $u(x, t)$ to the equation

$$(4.10) \quad \begin{aligned} u_t - Lu &= 0 & \text{for } (x, t) \in G \times (0, \infty), \\ u(x, 0) &= 0 & \text{for } x \in G, \\ u(x, t) &= 1 & \text{for } x \in B, \quad t > 0. \end{aligned}$$

The solution to the equation exists and has the properties:

$u(x, t)$ is nondecreasing in t for $t \geq 0$,

$0 \leq u(x, t) \leq 1$.

THEOREM 4.8.

$$\limsup_{t \rightarrow 0} 2t \log u(x, t) \leq -d^2(x, B)$$

uniformly over all regions G and points x in G so long as $d(x, B)$ remains bounded.

Proof: If $\phi(x, \lambda) = \int_0^\infty e^{-\lambda\tau} u(x, d\tau)$, then $\phi(x, \lambda)$ satisfies the equation

$$(4.11) \quad \begin{aligned} L\phi &= \lambda\phi & \text{for } x \in G, \\ \phi &= 1 & \text{for } x \in B. \end{aligned}$$

From Theorem 3.6, there exists for every $\rho > 0$, a $C_\rho < \infty$ such that

$$\phi(x, \lambda) \leq C_\rho \exp \{-(1 - \rho)\sqrt{2\lambda} d(x, B)\}$$

so long as $d(x, B)$ remains bounded. From equation (4.5), we have

$$u(x, t) \leq e^{\lambda t} \phi(x, \lambda)$$

for every $\lambda \geq 0$. Choosing $\lambda = (1 - \rho)^2 d^2(x, B)/2t^2$,

$$u(x, t) \leq C_\rho \exp \left\{ \frac{(1 - \rho)^2 d^2(x, B)}{2t} \right\}$$

which proves the theorem.

THEOREM 4.9. The fundamental solution $p(t, x, y)$ of Theorem 2.1 has the property

$$\limsup_{t \rightarrow 0} 2t \log p(t, x, y) \leq -d^2(x, y)$$

uniformly over all x, y such that $d(x, y)$ is bounded.

Proof: Let $d(x, y) = d$. Define, for any $\varepsilon > 0$,

$$G = [z: d(x, z) < d - \varepsilon],$$

$$B = [z: d(x, z) = d - \varepsilon].$$

Then $p(t, x, y)$ satisfies the equation

$$\begin{aligned} p_t - Lp &= 0 & \text{for } (x, t) &\text{ in } G \times (0, \infty), \\ (4.12) \quad p(0, x, y) &= 0 & \text{for } x &\in G, \\ p(t, x, y) &= p(t, b, y) & \text{for } b &\in B. \end{aligned}$$

If $u(x, t)$ is the solution to

$$\begin{aligned} (4.13) \quad u_t - Lu &= 0 & \text{for } (x, t) &\text{ in } G \times (0, \infty), \\ u &= 0 & \text{for } x &\in G \text{ and } t = 0, \\ u &= 1 & \text{for } x &\in B \text{ and } t > 0, \end{aligned}$$

then clearly

$$(4.14) \quad p(t, x, y) \leq \left[\sup_{\substack{0 \leq \tau < \infty \\ b \in B}} p(\tau, b, y) \right] u(x, t).$$

Since $d(b, y) \geq \varepsilon$, we have from property (iv) of Theorem 2.1

$$(4.15) \quad \sup_{0 < \tau < \infty} \sup_{b \in B} p(\tau, b, y) \leq M_\varepsilon < \infty.$$

Hence, combining (4.14) and (4.15) with Theorem 4.8, we obtain

$$\lim_{t \rightarrow 0} 2t \log p(t, x, y) \leq -(d - \varepsilon)^2.$$

Since ε is arbitrary, this proves the theorem. However, apparently the uniformity holds only in regions of the form $0 < \varepsilon \leq d(x, y) \leq D < \infty$. But the direct estimate in Theorem 2.1,

$$p(t, x, y) \leq Ct^{-k/2} \exp \left\{ \frac{-\alpha \|x - y\|^2}{2t} \right\},$$

establishes the uniformity over all x, y such that $d(x, y)$ remains bounded.

We now proceed to obtain a lower bound on $p(t, x, y)$ for small t .

THEOREM 4.10. *There exists a function $\psi(\varepsilon)$ such that $\psi(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ and*

$$\liminf_{t \rightarrow 0} \inf_{x, y: d(x, y) \leq \varepsilon} 2t \log p(t, x, y) \geq -\psi(\varepsilon).$$

Proof: Let $s > 0$. Define G_s and its boundary B_s by

$$\begin{aligned} G_s &= [z: d(y, z) > s], \\ B_s &= [z: d(y, z) = s]. \end{aligned}$$

Let $u_s(x, t)$ be the solution to the equation

$$\begin{aligned} (4.16) \quad u_t - Lu &= 0 & \text{in } G_s \times (0, \infty), \\ u &= 1 & \text{on } B_s \times (0, \infty), \\ u &= 0 & \text{on } G_s \times \{0\}. \end{aligned}$$

Let $\phi_s(x, \lambda) = \int_0^\infty e^{\lambda \tau} u_s(x, d\tau)$. Then $\phi_s(x, \lambda)$ satisfies the equation

$$\begin{aligned} (4.17) \quad L\phi &= \lambda\phi & \text{in } G_s, \\ \phi &= 1 & \text{on } B_s. \end{aligned}$$

From Theorem 3.9,

$$\begin{aligned} (4.18) \quad \phi_s(x, \lambda) &\geq \exp \{-d(1 + 4\sqrt{\eta})(1 + \delta(\varepsilon))\sqrt{2\lambda}\} \\ &\quad \times [1 - M_\rho \exp \{-\varepsilon(1 - 4\sqrt{\eta})(\eta - \rho - \delta(\varepsilon))\sqrt{2\lambda}\}]^n, \end{aligned}$$

where $d = d(x, y)$, ε, η, ρ are arbitrary, $n\varepsilon = d$, with the restriction that $s \geq 4\varepsilon\sqrt{\eta}$. Since

$$u_s(x, t) \geq \phi_s(x, \lambda) - e^{-\lambda t}$$

for every λ , let us choose $\lambda = 8/t^2$, $\varepsilon = s$ and η small enough. We further put $s = \sqrt{t}$. Then using (4.18) we get

$$(4.19) \quad u_{\sqrt{t}}(x, t) \geq \exp \left\{ -\frac{4d}{t} (1 + \theta) \right\} (1 - M_\rho e^{-C/\sqrt{t}})^{d/\sqrt{t}} - e^{-8/t},$$

where θ is a small number which can be chosen arbitrarily small. This function is defined for x such that $d(x, y) \geq s = \sqrt{t}$. It is clear from (4.19) that if $\varepsilon \leq 1$

$$(4.20) \quad \liminf_{t \rightarrow 0} \inf_{\substack{x, y: d(x, y) \geq \sqrt{t} \\ d(x, y) \leq \varepsilon}} t \log u_{\sqrt{t}}(x, t) \geq -4\varepsilon.$$

We now consider the solution $u_s(x, t_0, t)$ to the equation

$$\begin{aligned} (4.21) \quad u_t - Lu &= 0 & \text{in } G_s \times (0, \infty), \\ u &= 0 & \text{on } B_s \times (0, t_0), \\ u &= 1 & \text{on } B_s \times (t_0, \infty), \\ u &= 0 & \text{on } G_s \times \{0\}. \end{aligned}$$

It is clear that

$$\begin{aligned} u_s(x, t_0, t) &= 0 & \text{for } t \leq t_0, \\ u_s(x, t_0, t) &= u_s(x, t - t_0) & \text{for } t > t_0. \end{aligned}$$

We are interested in $u_{\sqrt{t}}(x, \frac{1}{2}t, t) = u_{\sqrt{t}}(x, \frac{1}{2}t)$. From (4.20) we deduce

$$(4.22) \quad \liminf_{t \rightarrow 0} \inf_{\substack{(x, y): d(x, y) \geq \sqrt{t} \\ d(x, y) \leq \varepsilon}} t \log u_{\sqrt{t}}(x, \frac{1}{2}t, t) \geq -8\varepsilon.$$

Going back to $p(t, x, y)$ we see that it satisfies an equation similar to (4.21) in $G_s \times (0, \infty)$ with different boundary values. However, since the solution $u_s(x, t_0, t)$ depends only on the boundary values on $B_s \times (0, \infty)$ for $0 < \tau < t$ we conclude by direct comparison that

$$p(t, x, y) \geq u_s(x, t_0, t) \inf_{\substack{b \in B_s \\ t_0 \leq \tau \leq t}} p(\tau, b, y).$$

Since t_0 and t are arbitrary, we choose $t_0 = \frac{1}{2}t$ and $s = \sqrt{t}$. Therefore,

$$(4.23) \quad p(t, x, y) \geq u_{\sqrt{t}}(x, \frac{1}{2}t, t) \inf_{\substack{b \in B_{\sqrt{t}} \\ \frac{1}{2}t \leq \tau \leq t}} p(\tau, b, y).$$

For t sufficiently small we have from property (v) of Theorem 2.1

$$\inf_{\substack{b \in B_{\sqrt{t}} \\ \frac{1}{2}t \leq \tau \leq t}} p(\tau, b, y) \geq \gamma > 0.$$

This with (4.22) and (4.23) yields

$$\liminf_{t \rightarrow 0} \inf_{\substack{x, y: d(x, y) \leq \varepsilon \\ d(x, y) \geq \sqrt{t}}} 2t \log p(t, x, y) \geq -8\varepsilon.$$

On the other hand, if $d(x, y) \leq \sqrt{t}$, again from property (v) of Theorem 2.1,

$$\liminf_{t \rightarrow 0} \inf_{x, y: d(x, y) \leq \sqrt{t}} 2t \log p(t, x, y) \geq 0.$$

This completes the proof.

THEOREM 4.11. Let $\varepsilon > 0$ and let G_ε be the set $\{z: d(y, z) < \varepsilon\}$. If $k(x, t)$ is defined as

$$k(x, t, y) = \int_{G_\varepsilon} p(t, x, z) dz,$$

then

$$\liminf_{t \rightarrow 0} 2t \log k(x, t, y) \geq -d^2(x, y)$$

uniformly over all x, y such that $d(x, y)$ is bounded.

Proof: Let

$$H_y = [z: d(y, z) > \tfrac{1}{2}\varepsilon],$$

$$B_y = [z: d(y, z) = \tfrac{1}{2}\varepsilon].$$

From property (iv) of Theorem 2.1 it follows that

$$(4.24) \quad k(x, t, y) \rightarrow 1 \quad \text{as} \quad t \rightarrow 0$$

uniformly for x on B_y .

Assume that $u(x, t)$ solves the equation

$$(4.25) \quad \begin{aligned} u_t - Lu &= 0 & \text{in} & \quad H_y \times (0, \infty), \\ u &= 1 & \text{on} & \quad B_y \times (0, \infty), \\ u &= 0 & \text{on} & \quad H_y \times \{0\}. \end{aligned}$$

Then the Laplace transform $\phi(x, \lambda)$ solves the equation

$$\begin{aligned} L\phi &= \lambda\phi & \text{for } x & \text{ in } H_y, \\ \phi &= 1 & \text{on } B_y. \end{aligned}$$

From Theorem 3.12 it follows that

$$\lim_{t \rightarrow \infty} \left[-\frac{1}{\sqrt{2\lambda}} \log \phi(x, \lambda) \right] = d(x, B_y).$$

Moreover, from Theorems 3.6 and 3.10 it is clear that the above limit is uniform whenever $d(x, y)$ remains bounded. Of course $\phi(x, \lambda)$ is defined only when $d(x, y) \geq \tfrac{1}{2}\varepsilon$. Now from Theorem 4.7 we obtain

$$(4.26) \quad \lim_{t \rightarrow \infty} 2t \log u(x, t) \geq -d^2(x, B_y) \geq -d^2(x, y)$$

uniformly whenever $d(x, y)$ remains bounded. Standard comparison of $k(x, t, y)$ which satisfies an equation similar to (4.25) with $u(x, t)$ shows that

$$(4.27) \quad k(x, t, y) \geq u(x, t) \inf_{\substack{0 \leq \tau \leq t \\ b \in B_y}} k(b, \tau, y).$$

Inequalities (4.26) and (4.27) complete the proof insofar as the region $d(x, y) \geq \tfrac{1}{2}\varepsilon$ is concerned. However, property (iv) of Theorem 2.1 shows that

$$k(x, t, y) \rightarrow 1 \quad \text{as} \quad t \rightarrow 0$$

uniformly over the set $d(x, y) \leq \tfrac{1}{2}\varepsilon$.

THEOREM 4.12.

$$\liminf_{t \rightarrow 0} 2t \log p(t, x, y) \geq -d^2(x, y)$$

uniformly over all x, y such that $d(x, y)$ is bounded.

Proof: If G_y is the set $[z: d(y, z) \leq \varepsilon]$, then

$$\begin{aligned} p(t, x, y) &= \int p([1 - \delta]t, x, z) p(\delta t, z, y) dz \\ &\geq \int_{G_y} p([1 - \delta]t, x, z) p(\delta t, z, y) dz \\ &\geq k(x, [1 - \delta]t, y) \inf_{z \in G_y} p(\delta t, z, y). \end{aligned}$$

Therefore, for every $\varepsilon > 0$ and $\delta > 0$, applying Theorems 4.10 and 4.11 we get

$$\liminf_{t \rightarrow 0} 2t \log p(t, x, y) \geq - \left[\frac{d^2(x, y)}{1 - \delta} + \frac{\psi(\varepsilon)}{\delta} \right].$$

Since δ is arbitrary, let us choose

$$\delta = \frac{[\psi(\varepsilon)]^{1/2}}{d(x, y) + [\psi(\varepsilon)]^{1/2}}.$$

Then

$$\liminf_{t \rightarrow 0} 2t \log p(t, x, y) \geq -[d(x, y) + [\psi(\varepsilon)]^{1/2}].$$

Since ε is arbitrary and $\psi(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$,

$$\liminf_{t \rightarrow 0} 2t \log p(t, x, y) \geq -d^2(x, y).$$

Since $d(x, y)$ is bounded, the only possible trouble regarding uniformity can occur when $\delta \rightarrow 1$ or $d(x, y)$ is very small. But then Theorem 4.10 can be used directly to show that it is uniform even around $d(x, y) = 0$.

We have finally proved

THEOREM 4.13.

$$\lim_{t \rightarrow 0} \left[-2t \log p(t, x, y) \right] = d^2(x, y)$$

uniformly in x, y such that $d(x, y)$ is bounded.

Remarks. The assumption in Theorem 3.12 that B is the common boundary of G and of its exterior can be weakened further. It suffices to assume that, for a

dense set $B_0 \subset B$, every sufficiently small sphere S around $b_0 \in B_0$ is disconnected by B .

We can consider equation (2.3) in a domain $G \subset R_k$ with the extra condition that the solution is zero on the boundary. Then there is a formula similar to Theorem 2.2, where $d(x, y)$ is replaced by $d_G(x, y)$ which is the length of the shortest path from x to y not leaving G .

These results are best proved using probabilistic methods and are therefore left to [3].

Acknowledgement. The methods used by the author in an earlier version of this paper were heavily probabilistic. The author expresses his thanks to Prof. L. Nirenberg whose suggestions were useful in replacing the probabilistic methods by analytical ones.

Bibliography

- [1] Friedman, A., *Partial Differential Equations of Parabolic Type*, Prentice-Hall, Englewood Cliffs, N.J., 1964.
- [2] Miranda, C., *Equazioni alle Derivate Parziali di Tipo Ellittico*, Springer, Berlin, 1955.
- [3] Varadhan, S. R. S., *Diffusion processes in a small time interval*, to appear in Comm. Pure. Appl. Math., Vol. 20, 1967.

Received August, 1966.