

ENVY-FREE CHORE ALLOCATION WITH UNIT SUBSIDIES UNDER NEGATIVE DICHOTOMOUS VALUATIONS

Mainak Sarkar^{*1} and Soumyarup Sadhukhan^{†1}

¹Indian Institute of Technology Kanpur

Abstract

We study the allocation of indivisible chores with subsidies under negative dichotomous valuations, where the marginal disutility of every chore is either zero or one. This is the chore analogue of the dichotomous goods model of Barman et al. [BKNS22], for which a subsidy of at most one unit per agent is known to suffice for envy-freeness.

We show that the same optimal guarantee holds for chores, under no structural assumption beyond binary marginals. For every instance with negative dichotomous valuations, there exists a complete allocation $\mathcal{A}$ and a subsidy vector $p \in \{0, 1\}^n$ with $\sum_{i \in N} p_i \leq n - 1$ such that $(\mathcal{A}, p)$ is envy-free. Moreover, the allocation $\mathcal{A}$ is EF1 even before subsidies are paid. The bound is tight, and the allocation and subsidies can be computed in polynomial time in the value-oracle model. We further show that the unit-subsidy guarantee cannot, in general, be maintained if Pareto optimality is also required.

Our algorithm starts from an envy-free partial allocation produced by the algorithm of Tao et al. [TWYZ25], assigns the remaining chores to distinct agents in a tail strongly connected component selected at termination, and determines the subsidised agents through a backward closure in the associated equality graph.

1 Introduction

Discrete fair division studies the allocation of indivisible resources among agents with heterogeneous preferences, and lies at the interface of mathematical economics and computer science [BCE⁺16, End17]. A central fairness criterion is *envy-freeness* [Fol67, Var74]: no agent prefers another agent’s bundle to her own. For indivisible items, however, an envy-free allocation need not exist, as the standard one-item two-agent instance illustrates. This has motivated a large literature on relaxations of envy-freeness, including envy-freeness up to one good (EF1) [Bud11, LMMS04] and the stronger notion of envy-freeness up to any good (EFX) [CKM⁺19].

^{*}Contact: mainaks23@iitk.ac.in

[†]Contact: soumyarup.sadhukhan@gmail.com

A second and complementary line of work retains exact envy-freeness by allowing monetary subsidies. In addition to her allocated bundle A_i , each agent i receives a subsidy $p_i \geq 0$ from a third party, and the outcome is envy-free if

$$v_i(A_i) + p_i \geq v_i(A_j) + p_j \quad \text{for all } i, j.$$

Halpern and Shah [HS19] show that every instance admits an envy-freeable allocation and study the minimum subsidy required to achieve envy-freeness. Under the standard normalization that each item value is at most one,¹ they show that, for a given envy-freeable allocation, a total subsidy of at most $(n - 1)m$ may be required in the worst case, and conjecture that a total subsidy of $n - 1$ always suffices when the allocation can also be chosen. Brustle et al. [BDN⁺20] prove this conjecture for additive valuations, obtaining an allocation that is also EF1 and balanced, and show that for general monotone valuations a subsidy of at most $2(n - 1)$ per agent suffices.

Barman et al. [BKNS22] obtain a substantially stronger guarantee for dichotomous valuations. When every marginal $v_i(S \cup \{g\}) - v_i(S)$ belongs to $\{0, 1\}$, they show that there exists an allocation admitting subsidies in $\{0, 1\}^n$, with total subsidy at most $n - 1$, and that such an outcome can be computed in polynomial time in the value-oracle model. This bound is tight: for n agents and a single good valued at one by every agent, the $n - 1$ agents who do not receive the good each require one unit of subsidy.

In this paper, we consider the corresponding problem for indivisible chores, where receiving an additional item weakly decreases an agent’s utility. Chore-allocation problems arise naturally in applications such as shift assignment, on-call duties, teaching and refereeing loads, committee service, and maintenance tasks. Monetary compensation is also natural in such settings, where it may be interpreted as compensation for undertaking an undesirable task or additional workload.

We study *negative dichotomous* valuations: for every agent i , every $S \subseteq M$, and every $g \in M \setminus S$,

$$v_i(S \cup \{g\}) - v_i(S) \in \{0, -1\}.$$

Thus, every item is a chore for every agent, and adding a chore changes an agent’s utility by either zero or minus one. This domain is the sign analogue of the dichotomous goods model of [BKNS22].

Binary marginals capture settings in which an additional task either creates a new unit of burden or can be absorbed by an existing commitment. For example, an agent already covering an on-call window may incur no additional cost from another incident during that window, or a driver already routed through a district may absorb an additional stop at no extra cost. Since whether a chore is costly may depend on the rest of the assigned bundle, the model naturally allows non-additive valuations, including valuations that are not subadditive.

Although chore allocation is formally related to goods allocation by reversing the direction of preferences, this correspondence does not generally extend to algorithmic arguments. Several works have observed that algorithms and structural results for goods need not carry over directly to chores [ARSW17, SCD23, BSV20]. The same difficulty arises in our setting. The algorithm of Barman et al. [BKNS22] for dichotomous goods is

¹Such a normalization is necessary when stating absolute bounds on monetary subsidies.

incremental: it allocates the goods one at a time while maintaining an envy-freeable partial allocation whose required subsidies lie in $\{0, 1\}^n$. For chores, one might naturally try to preserve the same invariant by assigning each new chore to an agent who currently receives no subsidy. This approach, however, may not succeed. There are negative dichotomous instances in which a partial allocation admits an envy-free solution with subsidies in $\{0, 1\}^n$, but after introducing one additional chore, no assignment of that chore to an agent preserves the unit-subsidy guarantee; this remains true even if the previously allocated bundles are allowed to be permuted before assigning the new chore. We give such an example in Appendix 1. Thus, the one-item extension property underlying the goods-side construction fails for chores, and the argument of Barman et al. [BKNS22] does not directly carry over.

A general subsidy bound for our setting follows from the work of Kawase et al. [KMS⁺25a]. They show that, for doubly monotone valuations with marginal values bounded in absolute value by one, an EF1 allocation can be converted into an envy-free solution with a subsidy of at most $n - 1$ per agent and at most $n(n - 1)/2$ in total. Negative dichotomous valuations form a subclass of this domain. Thus, Kawase et al. [KMS⁺25a] already imply, for our setting, a subsidy bound that is independent of the number of chores: at most $n - 1$ per agent and $n(n - 1)/2$ in total. The question is whether this can be strengthened to the unit-subsidy guarantee known for dichotomous goods. Our main result answers this question affirmatively.

Theorem. *For every instance with negative dichotomous valuations, there exists a complete allocation $\mathcal{A}$ and a subsidy vector $p \in \{0, 1\}^n$ such that $(\mathcal{A}, p)$ is envy-free and*

$$\sum_{i \in N} p_i \leq n - 1.$$

Moreover, $\mathcal{A}$ is EF1 even before subsidies are paid, and such an allocation and subsidy vector can be computed in polynomial time in the value-oracle model.

The total subsidy bound is tight, even for identical binary additive costs. Consider n agents and $n - 1$ chores with $c_i(S) = |S|$ for every agent i . In every complete allocation, some agent receives the empty bundle. Comparing each agent with such an agent in the envy-freeness inequalities gives $p_i \geq |A_i|$, and therefore

$$\sum_{i \in N} p_i \geq \sum_{i \in N} |A_i| = n - 1.$$

Our proof follows a different route from the incremental construction of Barman et al. [BKNS22]. We start from the polynomial-time algorithm of Tao et al. [TWYZ25], which returns an envy-free partial allocation while leaving at most $n - 1$ chores unassigned. At termination, the remaining chores are assigned to distinct agents in a suitable tail strongly connected component of the associated equality graph. We then identify the subsidised agents through a backward-closure operation in this graph. The resulting complete allocation is EF1, and assigning one unit of subsidy to the agents in this closure makes it exactly envy-free.

The unit-subsidy guarantee cannot, in general, be strengthened to require Pareto optimality: even with two agents and identical binary submodular costs, every Pareto-optimal allocation may require a subsidy of 2 for some agent (see Proposition 1).

Section 2 introduces the model and the necessary preliminaries. Section 3 presents the algorithm and proves the main result.

1.1 Related work

The literature on envy-freeness with subsidies has been developed primarily for goods. Halpern and Shah [HS19] establish the basic characterisation of envy-freeable allocations and obtain the optimal total subsidy bound of $n - 1$ for binary additive valuations. Goko et al. [GIK⁺24] extend the unit-subsidy guarantee to binary submodular valuations, together with strategy-proofness, while Barman et al. [BKNS22] show that no structure beyond binary marginals is required. The result most directly relevant to our setting is due to Kawase et al. [KMS⁺25a], who consider doubly monotone valuations and show that any EF1 allocation can be converted into an envy-free solution using at most $n - 1$ subsidy per agent and $n(n - 1)/2$ in total. Since negative dichotomous valuations form a subclass of doubly monotone valuations, this provides the previously known m -independent benchmark for our problem.

Fair allocation of indivisible chores without subsidies has also received considerable attention. Bhaskar et al. [BSV21] give polynomial-time algorithms for EF1 allocations of chores and doubly monotone instances, and show that deciding the existence of an exactly envy-free chore allocation is NP-complete already for binary additive costs. Tao et al. [TWYZ25] study binary-marginal chores and prove the existence of an EFX and Pareto-optimal allocation for binary additive costs. The additive assumption cannot be dropped in general: Lin et al. [LLTZ26] construct binary XOS and binary supermodular instances for which no complete EFX allocation exists. Our result is different in that it guarantees exact envy-freeness after subsidies throughout the full binary-marginal chore domain, while the underlying allocation is always EF1.

Subsidies have also been studied directly for chore allocation. For proportionality under additive costs, tight bounds are known for equal entitlements [WXZ25], and weighted variants have been studied in [WZ24, GSW25]. More recently, Lu et al. [LMS26] show that for additive instances containing both goods and chores, one unit of subsidy per agent suffices and the total bound of $n - 1$ is tight. Their result, however, does not subsume the present setting: negative dichotomous valuations need not be additive. Conversely, additive valuations need not be dichotomous. Thus, prior to the present work, the unit-subsidy guarantee was known for additive chores [LMS26] and for dichotomous goods [BKNS22], but not for non-additive dichotomous chores.

To the best of our knowledge, this is the first unit-subsidy guarantee for envy-freeness under negative dichotomous valuations.

2 Notation and Preliminaries

We study the allocation of m indivisible items among n agents, with subsidies. Write $N = [n]$ for the set of agents and M for the set of items, $|M| = m$. Each agent $i \in N$ has a valuation $v_i : 2^M \rightarrow \mathbb{R}$ with $v_i(\emptyset) = 0$, and we represent an instance by the tuple $\langle N, M, \{v_i\}_{i \in N} \rangle$. Algorithms operate in the standard *value-oracle* model: for each v_i we assume only an oracle

that returns $v_i(S)$ for a queried set $S \subseteq M$, and running time is measured in n, m and the number of oracle calls.

An allocation $\mathcal{A} = (A_1, \dots, A_n)$ is an ordered tuple of pairwise disjoint *bundles*, with A_i assigned to agent i . The allocation is *complete* if $\bigcup_{i \in N} A_i = M$ and *partial* otherwise. The ordering matters throughout: much of the theory below concerns permuting a fixed family of bundles among the agents, and for a permutation $\sigma \in \mathbb{S}_n$ we write $\mathcal{A}_\sigma := (A_{\sigma(1)}, \dots, A_{\sigma(n)})$ for the allocation in which agent i receives $A_{\sigma(i)}$. The utilitarian welfare of $\mathcal{A}$ is $\text{SW}(\mathcal{A}) = \sum_{i \in N} v_i(A_i)$.

Definition 1 (Envy-Freeness). *An allocation $\mathcal{A} = (A_1, \dots, A_n)$ is envy-free (EF) iff $v_i(A_i) \geq v_i(A_j)$ for all agents $i, j \in N$.*

Envy-free allocations of indivisible items need not exist, so we allow a third party to supplement the allocation with money. Agents have quasi-linear utilities: agent i receiving bundle A_i and subsidy p_i enjoys utility $v_i(A_i) + p_i$, with money entering linearly and at the same exchange rate for every agent.

Definition 2 (Envy-Free Solution). *An allocation $\mathcal{A} = (A_1, \dots, A_n)$ and a subsidy vector $p = (p_1, \dots, p_n) \in \mathbb{R}_+^n$ constitute an envy-free solution $(\mathcal{A}, p)$ iff $v_i(A_i) + p_i \geq v_i(A_j) + p_j$ for all $i, j \in N$.*

An allocation $\mathcal{A}$ is *envy-freeable* if some $p \in \mathbb{R}_+^n$ makes $(\mathcal{A}, p)$ an envy-free solution; this is a property of the allocation alone. Following [BKNS22] we write $M(p) := \{i \in N \mid p_i \geq p_j \text{ for all } j \in N\}$ for the set of maximally subsidised agents.

We will also use the standard relaxation of envy-freeness up to one item (EF1). Since our framework allows chores, we use the two-sided form of EF1, in which one is allowed to remove at most one item from either of the two bundles being compared [KMS⁺25b]. Formally:

Definition 3 (EF1, two-sided form). *An allocation $\mathcal{A}$ is envy-free up to one item (EF1) iff for all $i, j \in N$ there exists $X \subseteq A_i \cup A_j$ with $|X| \leq 1$ such that $v_i(A_i \setminus X) \geq v_i(A_j \setminus X)$.*

For an all-goods instance, removing an item from the envious agent's own bundle can only make the comparison harder, so the definition reduces to the usual goods version of EF1, in which one item may be removed from the envied bundle [HS19]. For an all-chore instance, the situation is reversed: removing a chore from the envied bundle can only make that bundle more attractive. Hence, if an item is removed, it may, without loss, be taken from the envious agent's own bundle.

Every instance in this paper is an all-chore instance, and there Definition 3 takes a simpler form: the item removed can always be taken from the envious agent's own bundle. We record that specialisation, since it is the form that the results below establish.

Definition 4 (EF1 for chores). *A complete allocation $\mathcal{A}$ of chores is EF1 iff for all $i, j \in N$, either i does not envy j , that is $v_i(A_i) \geq v_i(A_j)$, or there is a chore $e \in A_i$ with*

$$v_i(A_i \setminus \{e\}) \geq v_i(A_j).$$

Definition 4 is exactly Definition 3 on an all-chore instance. Taking $X = \emptyset$ in the first case and $X = \{e\} \subseteq A_i$ in the second gives one direction; for the converse, a removed item lying in A_j only lowers $c_i(A_j)$'s counterpart on the wrong side of the inequality, so nothing is gained by removing from the envied bundle when every item is a chore.

2.1 Negative Dichotomous Valuations

Definition 5 (Negative dichotomous valuation). *A valuation $v_i : 2^M \rightarrow \mathbb{R}$ with $v_i(\emptyset) = 0$ is negative dichotomous if, for every $S \subseteq M$ and every $g \in M \setminus S$,*

$$\Delta_i^+(S, g) := v_i(S \cup \{g\}) - v_i(S) \in \{-1, 0\}.$$

Equivalently,

$$v_i(S) - v_i(S \cup \{g\}) \in \{0, 1\}.$$

Thus, adding an item weakly decreases an agent's valuation, so every item is a chore and v_i is non-increasing:

$$S \subseteq T \implies v_i(S) \geq v_i(T).$$

Beyond the binary-marginal condition, we impose no further structure; in particular, the valuations need not be additive, submodular, or subadditive. This is the same level of generality considered by Barman et al. [BKNS22] for dichotomous goods.

For negative dichotomous valuations, it is often convenient to work with the corresponding non-negative *cost function*

$$c_i(S) := -v_i(S).$$

Then $c_i(\emptyset) = 0$ and

$$c_i(S \cup \{g\}) - c_i(S) \in \{0, 1\}$$

for every $S \subseteq M$ and every $g \in M \setminus S$.

Definition 6 (Dichotomous cost function). *A set function $c : 2^M \rightarrow \mathbb{R}$ is dichotomous if $c(\emptyset) = 0$ and*

$$c(S \cup \{g\}) - c(S) \in \{0, 1\}$$

for every $S \subseteq M$ and every $g \in M \setminus S$.

Since all marginal costs are non-negative, every dichotomous cost function is automatically non-decreasing. Moreover, the correspondence

$$v_i \longleftrightarrow c_i = -v_i$$

is a bijection between negative dichotomous valuations and dichotomous cost functions. In particular, the cost functions arising in our model belong to the same mathematical class as the dichotomous valuations studied by Barman et al. [BKNS22]; the difference is that here they represent costs rather than utilities.

In cost form, the envy-freeness condition for an allocation $\mathcal{A} = (A_1, \dots, A_n)$ and subsidy vector p becomes

$$c_i(A_i) - p_i \leq c_i(A_j) - p_j \quad \text{for all } i, j \in N.$$

We use the valuation and cost representations interchangeably, according to which is more convenient.

2.2 The Envy Graph and the Halpern–Shah Characterisation

We first fix the directed-graph terminology used below for the envy graph and, in Section 3.1, for the equality graph of the partial-allocation algorithm.

Definition 7 (Strongly connected component; tail component). *Let $G = (V, E)$ be a finite directed graph. Two vertices $u, v \in V$ are strongly connected if G contains a directed path from u to v and a directed path from v to u . This relation is an equivalence relation on V , and its equivalence classes are the strongly connected components of G . A strongly connected component S is a tail component (equivalently, a sink component) if no arc of E leaves it, that is, if there is no arc $(i, j) \in E$ with $i \in S$ and $j \in V \setminus S$.*

Being an equivalence relation, strong connectivity partitions V ; hence every vertex of G lies in exactly one strongly connected component, and G has at least one such component whenever $V \neq \emptyset$. Tail components exist as well.

Observation 1. *Every finite directed graph $G = (V, E)$ with $V \neq \emptyset$ has at least one tail strongly connected component.*

Proof The condensation of G is the directed graph whose vertices are the strongly connected components of G , with an arc from S to S' whenever $S \neq S'$ and some arc of E runs from a vertex of S to a vertex of S' . The condensation is acyclic: a directed cycle through two distinct components would place every vertex on it in a single component, contradicting their distinctness [BJG09, Ch. 1]. Since $V \neq \emptyset$, the condensation has at least one vertex, and a finite acyclic directed graph on a non-empty vertex set has a vertex of out-degree 0 — follow arcs from any vertex, never repeating one, and the walk must terminate. A component with out-degree 0 in the condensation has no arc of E leaving it, so it is a tail component. $\square$

For an allocation $\mathcal{A}$, the *envy graph* $G_{\mathcal{A}}$ is the complete weighted directed graph on vertex set N in which the arc (i, j) carries weight

$$w_{\mathcal{A}}(i, j) := v_i(A_j) - v_i(A_i) = c_i(A_i) - c_i(A_j),$$

the envy that agent i has towards agent j . The weight of a directed path P is $w_{\mathcal{A}}(P) = \sum_{(i,j) \in P} w_{\mathcal{A}}(i, j)$, and $\ell_{\mathcal{A}}(i)$ denotes the maximum weight of any path in $G_{\mathcal{A}}$ starting at i , the empty path of weight 0 included.

The following two results of Halpern and Shah [HS19] underpin much of the literature on subsidies for envy-freeness. They apply to general valuation functions and hence, in particular, to the negative dichotomous valuations considered here. We state them in the notation of our model.

Theorem 1 ([HS19]). *For any allocation $\mathcal{A} = (A_1, \dots, A_n)$, the following are equivalent.*

- (i) $\mathcal{A}$ is envy-freeable.
- (ii) $\mathcal{A}$ maximises utilitarian welfare across all reassignments of its own bundles among the agents: for every permutation σ over N , $\sum_{i \in N} v_i(A_i) \geq \sum_{i \in N} v_i(A_{\sigma(i)})$.
- (iii) The envy graph $G_{\mathcal{A}}$ has no positive-weight directed cycle.

Theorem 2 ([HS19]). *For any envy-freeable allocation $\mathcal{A}$, the subsidy vector p^* given by $p_i^* := \ell_{\mathcal{A}}(i)$ realises an envy-free solution $(\mathcal{A}, p^*)$, and $p_i^* \leq p_i$ for every $i \in N$ and every p such that $(\mathcal{A}, p)$ is an envy-free solution. It can be computed in strongly polynomial time by running an all-pairs longest-path computation on $G_{\mathcal{A}}$.*

Condition (ii) of Theorem 1 implies that, starting from any collection of bundles, one can obtain an envy-freeable allocation by suitably reassigning those bundles among the agents. In particular, such a reassignment can be found by solving a maximum-weight perfect matching problem between agents and bundles. Accordingly, the main issue is not the existence of an envy-freeable allocation, but rather the magnitude of the subsidies required to support one, as characterised by Theorem 2.

For negative dichotomous valuations, all bundle values are integral: indeed, $v_i(S)$ is obtained by telescoping from $v_i(\emptyset) = 0$ along marginals in $\{-1, 0\}$. Consequently, every arc weight $w_{\mathcal{A}}(i, j)$ in the envy graph is an integer, and hence, by Theorem 2, the pointwise-minimal subsidy vector p^* is integral as well. Therefore, showing $p_i^* \leq 1$ for every i is equivalent to showing $p^* \in \{0, 1\}^n$.

Moreover, every pointwise-minimal nonnegative subsidy vector has at least one zero coordinate. Indeed, if $p_i^* > 0$ for every i , then subtracting $\min_i p_i^*$ from every coordinate preserves all envy-freeness inequalities while producing a strictly smaller nonnegative subsidy vector, contradicting pointwise minimality. Thus, if $p^* \in \{0, 1\}^n$, then automatically

$$\sum_{i \in N} p_i^* \leq n - 1.$$

3 Unit Subsidies for Negative Dichotomous Valuations

This section proves the main theorem: for any number of agents n and every negative dichotomous instance there is an allocation admitting an envy-free solution with a subsidy of 0 or 1 per agent, at most $n - 1$ in total, computable in polynomial time.

Our argument starts from an envy-free *partial* allocation produced by the algorithm of Tao, Wu, Yu, and Zhou [TWYZ25], and then completes this partial allocation. We recall in Section 3.1 the relevant terminal-state properties of their algorithm, including Theorem 5.1 and the three update rules used in its construction. These ingredients are due to Tao et al. [TWYZ25]. The completion procedure introduced in Section 3.2, together with the subsequent lemmas and theorems, constitutes our contribution.

We work in cost form throughout, so an envy-free solution (Definition 2) is the requirement $(\mathcal{A}, p)$ such that

$$c_i(A_i) - p_i \leq c_i(A_j) - p_j \quad \text{for all } i, j \in N, \tag{1}$$

and a partial allocation X is envy-free (Definition 1) when

$$c_i(X_i) \leq c_i(X_j) \quad \text{for all } i, j \in N. \tag{2}$$

For $S \subseteq M$ and $e \in M \setminus S$ write $c_i(e \mid S) := c_i(S \cup \{e\}) - c_i(S) \in \{0, 1\}$ for a marginal, and for a partial allocation X write $R(X) := M \setminus \bigcup_{i \in N} X_i$ for its *residue*. Recall that as c_i s are integer valued, for all $S, T \subseteq M$, $c_i(S) < c_i(T) \implies c_i(S) \leq c_i(T) - 1$.

3.1 The terminal state of the partial-allocation algorithm

Theorem 3 ([TWYZ25], Theorem 5.1). *For every n and all dichotomous cost functions $c_1, \dots, c_n$ there is a polynomial-time algorithm returning a partial allocation X that is envy-free in the sense of (2) and satisfies $|R(X)| \leq n - 1$.*

The proof of the results in this paper uses not only the statement of Theorem 3 but also the state in which the algorithm of [TWYZ25] halts, so we record that algorithm. It maintains a partial allocation $X = (X_1, \dots, X_n)$, envy-free throughout, together with the *equality graph*

$$G = (N, E), \quad (i, j) \in E \iff i \neq j \text{ and } c_i(X_i) = c_i(X_j), \quad (3)$$

and repeats the following three rules in the stated order, halting only when none applies.

- (R1) If some $e \in R(X)$ and some $i \in N$ satisfy $c_i(e \mid X_i) = 0$, assign e to i .
- (R2) If some $e \in R(X)$ and some arc $(i, j) \in E$ lying on a directed cycle of G satisfy $c_i(e \mid X_j) = 0$, rotate the bundles along that cycle and assign e to i .
- (R3) Otherwise select a strongly connected component S of G with no arc of E leaving S ; such a component exists by Observation 1. If $|R(X)| \geq |S|$, assign one residual chore to each agent of S ; otherwise halt.

The loop runs while $R(X) \neq \emptyset$, so the only exit with $R(X) \neq \emptyset$ is the final clause of (R3).

The proof of our main result uses not only the statement of Theorem 3, but also structural properties of the terminal state of the algorithm of [TWYZ25]. We therefore recall the relevant part of that algorithm. It maintains an envy-free partial allocation $X = (X_1, \dots, X_n)$ together with the *equality graph*

$$G = (N, E), \quad (i, j) \in E \iff i \neq j \text{ and } c_i(X_i) = c_i(X_j). \quad (4)$$

While $R(X) \neq \emptyset$, the algorithm considers the following rules in the stated order.

- (R1) If there exist $e \in R(X)$ and $i \in N$ such that $c_i(e \mid X_i) = 0$, assign e to i .
- (R2) Otherwise, if there exist $e \in R(X)$ and an arc $(i, j) \in E$ lying on a directed cycle of G such that $c_i(e \mid X_j) = 0$, rotate the bundles along that cycle and assign e to i .
- (R3) Otherwise, select a strongly connected component S of G with no outgoing arc; such a component exists by Observation 1. If $|R(X)| \geq |S|$, assign distinct residual chores to the agents of S , one chore per agent; if $|R(X)| < |S|$, halt.

Consequently, if the algorithm terminates with $R(X) \neq \emptyset$, then neither (R1) nor (R2) is applicable at the terminal state, and the algorithm terminates through the final clause of (R3).

For the remainder of this section fix the following data: let X be the partial allocation at which the algorithm halts, let $R := R(X)$ and $r := |R|$, and, when $r \geq 1$, let S be the tail strongly connected component (Definition 7) selected by (R3) at the halting step, with G and E as in (4) evaluated at that state.

Lemma 1. Suppose $r \geq 1$. Then

- (a) $r < |S|$;
- (b) $c_i(e \mid X_i) = 1$ for every $i \in N$ and every $e \in R$;
- (c) $c_i(e \mid X_j) = 1$ for every arc $(i, j) \in E$ with $i, j \in S$ and every $e \in R$;
- (d) $c_i(X_i) < c_i(X_j)$ for every $i \in S$ and every $j \in N \setminus S$.

Proof Since $r \geq 1$, the algorithm halted at the final clause of (R3); hence neither (R1) nor (R2) applied at the halting state, and $|R| < |S|$, which is (a).

(b) If $c_i(e \mid X_i) = 0$ for some $i \in N$ and some $e \in R$, rule (R1) would apply. Marginals lie in $\{0, 1\}$ by Definition 6, so the value is 1.

(c) Let $(i, j) \in E$ with $i, j \in S$. Since S is strongly connected there is a directed path in G from j to i ; taking one with no repeated vertex and appending the arc (i, j) yields a directed cycle of G through (i, j) . If $c_i(e \mid X_j) = 0$ for some $e \in R$, rule (R2) would apply to that arc and that cycle. Hence the marginal is not 0, so it is 1.

(d) Let $i \in S$ and $j \in N \setminus S$. No arc of E leaves S , so $(i, j) \notin E$, i.e. $c_i(X_i) \neq c_i(X_j)$ by (4). Envy-freeness (2) gives $c_i(X_i) \leq c_i(X_j)$, so the inequality is strict. $\square$

3.2 The completion

Assume $r \geq 1$ and write $R = \{e_1, \dots, e_r\}$. By Lemma 1(a), $r < |S|$, so we may fix r distinct agents

$$T = \{t_1, \dots, t_r\} \subseteq S,$$

chosen arbitrarily. Define $\mathcal{A} = (A_1, \dots, A_n)$ by

$$A_i := \begin{cases} X_{t_k} \cup \{e_k\}, & i = t_k \ (1 \leq k \leq r), \\ X_i, & i \in N \setminus T. \end{cases} \quad (5)$$

The t_k are distinct and each chore of R is used exactly once, so the bundles of $\mathcal{A}$ are pairwise disjoint and cover M : $\mathcal{A}$ is a complete allocation. Members of T are called *recipients*.

Definition 8 (Subsidy set). Let $P \subseteq N$ be the smallest set satisfying

- (P1) $T \subseteq P$; and
- (P2) if $i \in N \setminus S$ and $(i, j) \in E$ for some $j \in P$, then $i \in P$.

Equivalently, P is obtained by initialising $P := T$ and repeatedly adjoining any $i \in N \setminus S$ having an arc of E into the current set, until no further agent can be adjoined; the process terminates because N is finite and P only grows. Define

$$p_i := \begin{cases} 1, & i \in P, \\ 0, & i \in N \setminus P. \end{cases} \quad (6)$$

The complete procedure is summarised in Algorithm 1. The partial-allocation phase in Line 2 is Algorithm 3 of [TWYZ25]; our contribution is the completion and subsidy construction in the subsequent lines.

Algorithm 1 EF1 allocation with at most one unit of subsidy per agent for negative dichotomous valuations

Require: Agents $N = [n]$, chores M , and value-oracle access to negative dichotomous valuations $\{v_i\}_{i \in N}$.

Ensure: A complete allocation $\mathcal{A}$ that is EF1, together with a subsidy vector $p \in \{0, 1\}^n$ such that $(\mathcal{A}, p)$ is envy-free and $\sum_{i \in N} p_i \leq n - 1$.

- 1: Set $c_i = -v_i$ for every $i \in N$.
 - 2: Run Algorithm 3 of [TWYZ25] on $\{c_i\}_{i \in N}$.
 - 3: Let $X = (X_1, \dots, X_n)$ be its terminal partial allocation and let $R = M \setminus \bigcup_{i \in N} X_i$.
 - 4: **if** $R = \emptyset$ **then**
 - 5: **return** $(X, \mathbf{0})$.
 - 6: Let S be the tail strongly connected component (Definition 7) selected at the terminating application of Rule (R3).
 - 7: Write $R = \{e_1, \dots, e_r\}$.
 - 8: Choose arbitrary distinct agents $T = \{t_1, \dots, t_r\} \subseteq S$. $\triangleright r < |S|$ by Lemma 1(a)
 - 9: **for** $k = 1, \dots, r$ **do**
 - 10: $A_{t_k} \leftarrow X_{t_k} \cup \{e_k\}$.
 - 11: **for** $i \in N \setminus T$ **do**
 - 12: $A_i \leftarrow X_i$.
 - 13: Construct the equality graph $G = (N, E)$ on the agents, where
 $(i, j) \in E \iff i \neq j \text{ and } c_i(X_i) = c_i(X_j)$.
 - 14: Initialise $P \leftarrow T$.
 - 15: **while** there exist $i \in N \setminus S$ and $j \in P$ with $(i, j) \in E$ and $i \notin P$ **do**
 - 16: $P \leftarrow P \cup \{i\}$.
 - 17: **for** $i \in N$ **do**
 - 18: $p_i \leftarrow \mathbf{1}[i \in P]$.
 - 19: $\mathcal{A} \leftarrow (A_1, \dots, A_n)$.
 - 20: **return** $(\mathcal{A}, p)$.
-

Lemma 2. $P \setminus T \subseteq N \setminus S$: every agent of P that is not a recipient lies outside S .

Proof Let $P_0 := T$ and let P_{u+1} be P_u together with every $i \in N \setminus S$ having an arc of E into P_u , so that $P = \bigcup_{u \geq 0} P_u$. We show $P_u \setminus T \subseteq N \setminus S$ by induction on u . For $u = 0$ the set $P_0 \setminus T$ is empty. For the step, every agent adjoined in passing from P_u to P_{u+1} lies in $N \setminus S$ by construction. $\square$

Lemma 3. $p \in \{0, 1\}^n$ and $\sum_{i \in N} p_i = |P| \leq n - 1$.

Proof That $p \in \{0, 1\}^n$ with $\sum_i p_i = |P|$ is immediate from (6). By Lemma 2, $P \subseteq T \cup (N \setminus S)$, and this union is disjoint because $T \subseteq S$; hence $|P| \leq |T| + |N \setminus S| = r + n - |S|$. By Lemma 1(a), $r \leq |S| - 1$, so $|P| \leq n - 1$. $\square$

Lemma 4. Let $i \in S$, let $t \in T$ and let $e \in R$. Then

$$c_i(X_t \cup \{e\}) \geq c_i(X_i) + 1.$$

Proof Note that $t \in T \subseteq S$. Envy-freeness (2) gives $c_i(X_i) \leq c_i(X_t)$. If $c_i(X_i) < c_i(X_t)$, then $c_i(X_i) + 1 \leq c_i(X_t)$, and $c_i(X_t \cup \{e\}) \geq c_i(X_t)$ by monotonicity of c_i . Otherwise $c_i(X_i) = c_i(X_t)$. If $i = t$ then $c_i(X_t \cup \{e\}) = c_i(X_i) + c_i(e \mid X_i) = c_i(X_i) + 1$ by Lemma 1(b). If $i \neq t$ then $(i, t) \in E$ by (4), with $i, t \in S$, so $c_i(e \mid X_t) = 1$ by Lemma 1(c) and therefore $c_i(X_t \cup \{e\}) = c_i(X_t) + 1 = c_i(X_i) + 1$. $\square$

3.3 Envy-freeness of the completion

Theorem 4. Let X, R, S be as in Section 3.1 with $r \geq 1$, let $\mathcal{A}$ be as in (5) and let p be as in (6). Then $(\mathcal{A}, p)$ is an envy-free solution.

Proof Fix $i, j \in N$; we verify (1), that is $c_i(A_i) - p_i \leq c_i(A_j) - p_j$. Throughout, $X_j \subseteq A_j$ for every j by (5), so by monotonicity (Observation ??) and (2),

$$c_i(A_j) \geq c_i(X_j) \geq c_i(X_i). \quad (7)$$

Note also that $T \subseteq P$, so $i \notin P$ implies $i \notin T$ and hence $A_i = X_i$; likewise $j \notin P$ implies $A_j = X_j$ and $p_j = 0$. Since N is the disjoint union of $T, P \setminus T$ and $N \setminus P$, the following three cases are exhaustive.

Case 1: $i \in T$. Then $p_i = 1$ and, by Lemma 1(b), $c_i(A_i) = c_i(X_i) + c_i(e \mid X_i) = c_i(X_i) + 1$, so $c_i(A_i) - p_i = c_i(X_i)$. Also $i \in T \subseteq S$.

(1a) $j \in T$. Then $p_j = 1$ and $A_j = X_j \cup \{e\}$ for some $e \in R$, so $c_i(A_j) \geq c_i(X_i) + 1$ by Lemma 4. Hence $c_i(A_j) - p_j \geq c_i(X_i)$.

(1b) $j \in P \setminus T$. Then $p_j = 1$ and $A_j = X_j$. By Lemma 2, $j \notin S$; since $i \in S$, Lemma 1(d) gives $c_i(X_i) < c_i(X_j)$, hence $c_i(X_j) \geq c_i(X_i) + 1$ by Observation ??. Hence $c_i(A_j) - p_j = c_i(X_j) - 1 \geq c_i(X_i)$.

(1c) $j \in N \setminus P$. Then $p_j = 0$ and $A_j = X_j$, so $c_i(A_j) - p_j = c_i(X_j) \geq c_i(X_i)$ by (2).

Case 2: $i \in P \setminus T$. Then $p_i = 1$ and $A_i = X_i$, so $c_i(A_i) - p_i = c_i(X_i) - 1$. For every j we have $p_j \leq 1$ and $c_i(A_j) \geq c_i(X_i)$ by (7), hence $c_i(A_j) - p_j \geq c_i(X_i) - 1 = c_i(A_i) - p_i$.

Case 3: $i \in N \setminus P$. Then $p_i = 0$ and $A_i = X_i$, so $c_i(A_i) - p_i = c_i(X_i)$.

(3a) $j \in N \setminus P$. Then $p_j = 0$ and $A_j = X_j$, so $c_i(A_j) - p_j = c_i(X_j) \geq c_i(X_i)$ by (2).

(3b) $j \in T$. Then $p_j = 1$ and $A_j = X_j \cup \{e\}$ for some $e \in R$; we claim $c_i(A_j) \geq c_i(X_i) + 1$. If $i \in S$ this is Lemma 4. If $i \notin S$, suppose $c_i(X_i) = c_i(X_j)$. Then $i \neq j$, since $j \in T \subseteq P$ while $i \notin P$; so $(i, j) \in E$ by (4), and $j \in P$, so (P2) of Definition 8 places $i \in P$ — contradicting $i \notin P$. Hence $c_i(X_i) < c_i(X_j)$, so $c_i(X_j) \geq c_i(X_i) + 1$ by Observation ??, and $c_i(A_j) \geq c_i(X_j)$ by monotonicity. In either case $c_i(A_j) - p_j \geq c_i(X_i)$.

(3c) $j \in P \setminus T$. Then $p_j = 1$ and $A_j = X_j$; we claim $c_i(X_j) \geq c_i(X_i) + 1$. If $i \notin S$, suppose $c_i(X_i) = c_i(X_j)$. Then $i \neq j$, since $j \in P$ while $i \notin P$; so $(i, j) \in E$, and (P2) places $i \in P$, a contradiction. Hence $c_i(X_i) < c_i(X_j)$. If instead $i \in S$, then $j \notin S$ by Lemma 2, and Lemma 1(d) gives $c_i(X_i) < c_i(X_j)$. In either case Observation ?? yields $c_i(X_j) \geq c_i(X_i) + 1$, so $c_i(A_j) - p_j = c_i(X_j) - 1 \geq c_i(X_i)$.

In every sub-case $c_i(A_j) - p_j \geq c_i(X_i) = c_i(A_i) - p_i$ in Cases 1 and 3, and the inequality was shown directly in Case 2. $\square$

3.4 The EF1 property of the completion

Corollary 1. *The complete allocation $\mathcal{A}$ constructed in Section 3.2 is EF1 in the sense of Definition 4, hence also in the sense of Definition 3. More precisely, every non-recipient $i \in N \setminus T$ is envy-free in $\mathcal{A}$, while for every recipient $t_k \in T$, removing the residual chore e_k assigned to t_k eliminates all of t_k 's envy.*

Proof Recall that the partial allocation $X = (X_1, \dots, X_n)$ is envy-free, and that $X_j \subseteq A_j$ for every $j \in N$. First, let $i \in N \setminus T$. Then $A_i = X_i$. Hence, for every $j \in N$, envy-freeness of X and monotonicity of c_i give

$$c_i(A_i) = c_i(X_i) \leq c_i(X_j) \leq c_i(A_j).$$

Thus i does not envy any agent in $\mathcal{A}$. Now let $i = t_k \in T$. By construction,

$$A_i = X_i \cup \{e_k\},$$

and hence

$$A_i \setminus \{e_k\} = X_i.$$

Therefore, for every $j \in N$,

$$c_i(A_i \setminus \{e_k\}) = c_i(X_i) \leq c_i(X_j) \leq c_i(A_j),$$

where the first inequality follows from envy-freeness of X and the second from monotonicity.

Thus, for every recipient, removing the single newly assigned chore from her own bundle eliminates her envy towards every other agent. Consequently, $\mathcal{A}$ is EF1. $\square$

3.5 The main theorem

Theorem 5. *For every n and every instance $\langle N, M, \{v_i\}_{i \in N} \rangle$ with negative dichotomous valuations there exist a complete allocation $\mathcal{A}$ and a subsidy vector $p \in \{0, 1\}^n$ with $\sum_{i \in N} p_i \leq n - 1$ such that $(\mathcal{A}, p)$ is an envy-free solution. Moreover, the allocation $\mathcal{A}$ is EF1. Such an allocation and subsidy vector can be computed in polynomial time given value-oracle access.*

Proof Pass to cost form, $c_i = -v_i$, which is dichotomous in the sense of Definition 6. Run Algorithm 3 of [TWYZ25] and retain its terminal partial allocation X , the residue R with $r = |R|$. If $r \geq 1$, retain also the tail strongly connected component S selected at the terminating application of rule (R3), all as fixed in Section 3.1. The statement of Theorem 3 alone does not supply S .

If $r = 0$ then X is a complete allocation satisfying (2), so $(X, \mathbf{0})$ is an envy-free solution with $\sum_i p_i = 0 \leq n - 1$. Since every envy-free allocation is EF1, X is also EF1.

If $r \geq 1$, construct T , $\mathcal{A}$ and p as in Section 3.2. Then $\mathcal{A}$ is a complete allocation, Theorem 4 shows $(\mathcal{A}, p)$ is an envy-free solution, and Lemma 3 gives $p \in \{0, 1\}^n$ with $\sum_i p_i \leq n - 1$. The allocation $\mathcal{A}$ is EF1 by Corollary 1.

For the running time, Algorithm 3 of [TWYZ25] is polynomial and yields X , R and the tail component S of its terminating (R3) step if $r \geq 1$. Constructing $\mathcal{A}$ from (5) takes $r \leq n - 1$ assignments. The graph G of (4) has n vertices and is determined by the n^2 values $c_i(X_j)$, one oracle call each. Computing P is a backward reachability search in G from T restricted to vertices outside S , which takes $O(n^2)$ time. Hence the work after Theorem 3 is polynomial, and so is the whole procedure. $\square$

As discussed earlier, the bound of Theorem 5 is tight: there are instances on which every complete allocation requires total subsidy at least $n - 1$.

3.6 Pareto optimality and subsidies

In this section, we show that the unit-subsidy guarantee need not be compatible with Pareto optimality of the underlying allocation. We evaluate Pareto optimality with respect to the allocation of chores, without taking subsidies into account. Formally, a complete allocation $\mathcal{B}$ *Pareto dominates* a complete allocation $\mathcal{A}$ if

$$c_i(\mathcal{B}_i) \leq c_i(\mathcal{A}_i) \quad \text{for every } i \in N,$$

with strict inequality for at least one agent. An allocation is *Pareto optimal* if it is not Pareto dominated by another complete allocation.

Proposition 1. *There exist negative dichotomous instances, even with two agents and identical binary submodular cost functions, for which no Pareto-optimal allocation admits an envy-free subsidy vector in $\{0, 1\}^n$.*

Proof Consider two agents $N = \{1, 2\}$ and three chores $M = \{a, b, c\}$. Both agents have the same cost function

$$c_1(S) = c_2(S) = \min\{|S|, 2\}.$$

Every marginal cost belongs to $\{0, 1\}$, so the cost function is dichotomous. Moreover, it is submodular, since $k \mapsto \min\{k, 2\}$ is a nondecreasing concave function of cardinality. The function is not additive, since

$$c_i(\{a, b, c\}) = 2 < 3 = \sum_{g \in \{a, b, c\}} c_i(\{g\}).$$

Since the agents have identical costs, every complete allocation has, up to symmetry, one of two bundle-size profiles.

First, consider a concentrated allocation with bundle sizes $(3, 0)$. The corresponding cost profile is $(2, 0)$. This allocation is Pareto optimal. Indeed, every allocation in which both agents receive a nonempty bundle has bundle sizes $(2, 1)$ and therefore cost profile $(2, 1)$, which does not improve the loaded agent and strictly worsens the other agent.

However, a concentrated allocation cannot be made envy-free with unit subsidies. If agent 1 receives all three chores, envy-freeness requires

$$2 - p_1 \leq -p_2,$$

and hence

$$p_1 - p_2 \geq 2.$$

Thus, no subsidy vector in $\{0, 1\}^2$ suffices. The case in which agent 2 receives all three chores is symmetric.

Now consider any split allocation, whose bundle sizes are $(2, 1)$ up to symmetry. Its cost profile is $(2, 1)$, so it can be made envy-free with subsidies $(1, 0)$, up to symmetry. Nevertheless, it is not Pareto optimal: if the agent holding two chores is instead given the third chore as well, the cost profile changes from $(2, 1)$ to $(2, 0)$. The first agent's cost remains unchanged, while the second agent's cost strictly decreases.

Hence, every Pareto-optimal allocation requires a subsidy of at least two for one agent, whereas every allocation admitting unit subsidies is Pareto dominated. Therefore, no allocation is simultaneously Pareto optimal and envy-free with subsidies in $\{0, 1\}^2$. $\square$

Thus, unlike the unit-subsidy guarantee of Theorem 5, Pareto optimality cannot be added to the conclusion without weakening the subsidy bound.

References

- [ARSW17] Haris Aziz, Gerhard Rauchecker, Guido Schryen, and Toby Walsh. Algorithms for max-min share fair allocation of indivisible chores. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 31, 2017.
- [BCE⁺16] Felix Brandt, Vincent Conitzer, Ulle Endriss, Jérôme Lang, and Ariel D. Procaccia. *Handbook of Computational Social Choice*. Cambridge University Press, 1st edition, 2016.
- [BDN⁺20] Johannes Brustle, Jack Dippel, Vishnu V. Narayan, Mashbat Suzuki, and Adrian Vetta. One dollar each eliminates envy. In *Proceedings of the 21st ACM Conference on Economics and Computation (EC)*, pages 23–39, 2020.
- [BJG09] Jørgen Bang-Jensen and Gregory Gutin. *Digraphs: Theory, Algorithms and Applications*. Springer Monographs in Mathematics. Springer, 2nd edition, 2009.
- [BKNS22] Siddharth Barman, Anand Krishna, Y. Narahari, and Soumyarup Sadhukhan. Achieving envy-freeness with limited subsidies under dichotomous valuations. In *Proceedings of the 31st International Joint Conference on Artificial Intelligence (IJCAI)*, pages 60–66, 2022. arXiv:2201.07419.
- [BSV20] Umang Bhaskar, AR Sricharan, and Rohit Vaish. On approximate envy-freeness for indivisible chores and mixed resources. *arXiv preprint arXiv:2012.06788*, 2020.

- [BSV21] Umang Bhaskar, A. R. Sricharan, and Rohit Vaish. On approximate envy-freeness for indivisible chores and mixed resources. In *Approximation, Randomization, and Combinatorial Optimization (APPROX/RANDOM)*, LIPIcs, 2021.
- [Bud11] Eric Budish. The combinatorial assignment problem: Approximate competitive equilibrium from equal incomes. *Journal of Political Economy*, 119(6):1061–1103, 2011.
- [CKM⁺19] Ioannis Caragiannis, David Kurokawa, Hervé Moulin, Ariel D. Procaccia, Nisarg Shah, and Junxing Wang. The unreasonable fairness of maximum nash welfare. *ACM Transactions on Economics and Computation*, 7(3):12:1–12:32, 2019.
- [End17] Ulle Endriss. *Trends in Computational Social Choice*. Lulu.com, 2017.
- [Fol67] Duncan Karl Foley. Resource allocation and the public sector. *Yale Economic Essays*, 7(1):45–98, 1967.
- [GIK⁺24] Hiromichi Goko, Ayumi Igarashi, Yasushi Kawase, Kazuhisa Makino, Hanna Sumita, Akihisa Tamura, Yu Yokoi, and Makoto Yokoo. A fair and truthful mechanism with limited subsidy. *Games and Economic Behavior*, 144:49–70, 2024.
- [GSW25] Jugal Garg, Eklavya Sharma, and Xiaowei Wu. Proportional and Pareto-optimal allocation of chores with subsidy, 2025.
- [HS19] Daniel Halpern and Nisarg Shah. Fair division with subsidy. In *Proceedings of the 12th International Symposium on Algorithmic Game Theory (SAGT)*, volume 11801 of *Lecture Notes in Computer Science*, pages 374–389. Springer, 2019.
- [KMS⁺25a] Yasushi Kawase, Kazuhisa Makino, Hanna Sumita, Akihisa Tamura, and Makoto Yokoo. Towards optimal subsidy bounds for envy-freeable allocations. *Artificial Intelligence*, page 104406, 2025.
- [KMS⁺25b] Yasushi Kawase, Kazuhisa Makino, Hanna Sumita, Akihisa Tamura, and Makoto Yokoo. Towards optimal subsidy bounds for envy-freeable allocations. *Artificial Intelligence*, 348:104406, 2025. Preliminary version in AAAI 2024; arXiv:2308.11230.
- [LLTZ26] Zehan Lin, Shengxin Liu, Biaoshuai Tao, and Shengwei Zhou. Non-existence of EFX chore allocations for monotone cost functions with binary marginals, 2026. arXiv:2608.10572.
- [LMMS04] Richard J. Lipton, Evangelos Markakis, Elchanan Mossel, and Amin Saberi. On approximately fair allocations of indivisible goods. In *Proceedings of the 5th ACM Conference on Electronic Commerce (EC)*, pages 125–131, 2004.
- [LMS26] Xinhang Lu, Simon Mackenzie, and Mashbat Suzuki. Optimal subsidy bounds for goods and chores: One dollar each suffices, 2026. arXiv:2607.10089.
- [SCD23] Ankang Sun, Bo Chen, and Xuan Vinh Doan. Fairness criteria for allocating indivisible chores: connections and efficiencies: A. sun et al. *Autonomous Agents and Multi-Agent Systems*, 37(2):39, 2023.

- [TWYZ25] Biaoshuai Tao, Xiaowei Wu, Ziqi Yu, and Shengwei Zhou. On the existence of efx (and pareto-optimal) allocations for binary chores. *Theoretical Computer Science*, 1042:115248, 2025.
- [Var74] Hal Varian. Equity, envy, and efficiency. *Journal of Economic Theory*, 9(1):63–91, 1974.
- [WXZ25] Xiaowei Wu, Quan Xue, and Shengwei Zhou. Revisiting proportional allocation with subsidy: Simplification and improvements. In *IJCAI*, pages 4082–4090, 2025.
- [WZ24] Xiaowei Wu and Shengwei Zhou. Tree splitting based rounding scheme for weighted proportional allocations with subsidy. In *International Conference on Web and Internet Economics*, pages 295–313. Springer, 2024.

A Appendix

Example 1 (Failure of a BKNS-type ([BKNS22]) incremental extension). Consider three agents $N = \{1, 2, 3\}$ and three chores $M = \{a, b, c\}$. We work in cost form, writing

$$c_i = -v_i$$

for agent i 's non-negative cost function. Thus, an allocation $\mathcal{A} = (A_1, \dots, A_n)$ together with a subsidy vector $p = (p_1, \dots, p_n)$ is envy-free iff

$$c_i(A_i) - p_i \leq c_i(A_j) - p_j \quad \text{for all } i, j \in N.$$

The cost functions depend only on bundle size:

$$c_1(S) = \max\{0, |S| - 1\}, \quad c_2(S) = c_3(S) = \min\{|S|, 2\}.$$

Thus,

$ S $	0	1	2	3
$c_1(S)$	0	0	1	2
$c_2(S) = c_3(S)$	0	1	2	2

Every marginal cost belongs to $\{0, 1\}$, so these are dichotomous cost functions. Suppose c is currently unallocated, and the partial allocation is

$$X_1 = \{a, b\}, \quad X_2 = X_3 = \emptyset.$$

Note that this partial allocation is itself reachable from the empty allocation through successive insertions while maintaining the unit-subsidy invariant:

$$(\emptyset, \emptyset, \emptyset) \longrightarrow (\{a\}, \emptyset, \emptyset) \longrightarrow (\{a, b\}, \emptyset, \emptyset),$$

with pointwise-minimal subsidy vectors $(0, 0, 0)$, $(0, 0, 0)$, and $(1, 0, 0)$, respectively. The pointwise-minimal subsidy vector of (X_1, X_2, X_3) is

$$p^* = (1, 0, 0).$$

Indeed, agent 1 has cost 1 for her own bundle and cost 0 for either empty bundle, so she requires one unit of subsidy. Agents 2 and 3 have no envy, since they have cost 0 for their own empty bundles and cost 2 for X_1 .

We claim that the remaining chore c cannot be inserted so that the resulting allocation admits an envy-free subsidy vector in $\{0, 1\}^3$, even if the existing bundles $\{a, b\}, \emptyset, \emptyset$ may first be permuted among the agents.

If c is added to the two-chore bundle, the resulting bundle sizes are $(3, 0, 0)$. Every agent evaluates a three-chore bundle at cost 2 and an empty bundle at cost 0. Hence, the agent receiving the three-chore bundle requires a subsidy at least two units larger than that of an agent receiving an empty bundle. Thus, no subsidy vector in $\{0, 1\}^3$ can make the resulting allocation envy-free.

It remains to consider the case in which c is added to an empty bundle. The resulting bundle sizes are $(2, 1, 0)$, up to permutation. If an agent in $\{2, 3\}$ receives the two-chore bundle, she evaluates it at cost 2 and the empty bundle at cost 0, again requiring a subsidy difference of at least two.

Therefore, the only remaining possibility is that agent 1 receives the two-chore bundle. Let $i \in \{2, 3\}$ receive the singleton and let j receive the empty bundle. Envy-freeness for agent 1 with respect to i requires

$$1 - p_1 \leq 0 - p_i,$$

and hence

$$p_1 \geq p_i + 1.$$

Envy-freeness for agent i with respect to j requires

$$1 - p_i \leq -p_j,$$

and hence

$$p_i \geq p_j + 1.$$

Combining the two inequalities gives

$$p_1 \geq p_j + 2,$$

which is impossible for $p \in \{0, 1\}^3$. Hence, no assignment of the additional chore c , even after an arbitrary permutation of the previously allocated bundles, preserves the unit-subsidy guarantee. Notice that this is an obstruction to the incremental construction, not to the instance itself. The complete allocation

$$(\{a\}, \{b\}, \{c\})$$

is envy-free without subsidies: agent 1 evaluates every singleton at cost 0, while agents 2 and 3 evaluate every singleton at cost 1. $\square$