

Individual Assignment 5

Maisie Prestridge

2026-05-10

Set up

Packages

```
# general use and cleaning
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrap axis labels
library(scales)

# reading .xlsx files
library(readxl)

# visualise missing data
library(naniar)

# arrange plots
library(patchwork)
```

Data

```
# taxonomic information
taxon_list <- read_csv(here("data", "taxon_list.csv"))
```

```
# aquatic invertebrates
aq_ins <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Aquatic Insects")

# site information
sites <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "sites")

# water quality
water_quality <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Water Quality",
  na = c("", "n/a", "N/A", "over", "173+"))
```

```
# creating new clean object from water quality
water_quality_clean <- water_quality |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites in ncos_sites
  semi_join(sites, by = "site") |>
  # create new column to join with later data frames
  unite("date_site", date, site, remove = TRUE) |>
  # group by date_site column
  group_by(date_site) |>
  # calculate median pH, salinity, DO
  summarize(
    med_ph = median(p_h, na.rm = TRUE),
    med_sal = median(salinity_ppt, na.rm = TRUE),
    med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>
  # ungroup data frame
  ungroup() |>
  # filter out salinity outliers
  filter(date_site != "2022-08-12_NVBR")
```

```
# creating new clean object from taxon_list
taxon_list_clean <- taxon_list |>
  # converting taxon name to snake case (no spaces or capital letters,
  # only underscores)
  mutate(verbatim_name = to_any_case(verbatim_name, case = "snake"))
```

```
# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
  # clean column names
  clean_names() |>
```

```

# filtering join: only include sites occurring in ncos_sites
semi_join(sites, by = c("site")) |>
# renaming columns
rename(date = date_on_vial) |>
# select columns of interest
select(date, sample_type, site,
       ostracod,
       copepod,
       hemiptera_corixidae_boatman,
       cladocera,
       nematode,
       diptera_ceratopogonidae,
       annelida_oligochaete,
       diptera_chironomid,
       annelida_polychaete) |>
# filter to only include "filtered beaker" samples
filter(sample_type %in% c("FB 250", "FB250")) |>
# convert the data frame to long format
pivot_longer(cols = ostracod:annelida_polychaete,
             names_to = "taxon",
             values_to = "abundance") |>
# group by date, site, taxon
group_by(date, site, taxon) |>
# calculate average abundance per liter (sum abundance divided by 7.5 liters)
summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
# ungroup data frame
ungroup() |>
# create a new column called `date_site` to join with water quality
unite("date_site", date, site, remove = FALSE) |>
# join with water quality
left_join(water_quality_clean, by = c("date_site")) |>
# join with taxon list
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>
# add full names of sites
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",
  site == "NEC" ~ "South of Venoco Bridge",
  site == "NMC" ~ "Main Channel",
  site == "NPB" ~ "South of Phelps Bridge",
  site == "NPB1" ~ "North of Phelps Bridge",
  site == "NPB2" ~ "Phelps Road",
  site == "NWP" ~ "West Pond",

```

```

    site == "NDC" ~ "Devereux Creek"
  )) |>
  # setting factor levels for site
  mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm = TRUE),
         site_full = fct_rev(site_full))

```

1. Visualizing differences across sites in major taxon abundance

a. Plan your figure

x-axis: site_full y-axis: log_ave_lit geometry to represent average abundance/L: geom_jitter()
 geometry to represent medians: geom_point()

b. Wrangle your data

```

# new object created from clean aquatic inverts
ostracods <- aq_ins_clean |>
  # filter to only include copepods
  filter(taxon == "ostracod") |>
  # log + 1 to transform mean abundance/liter
  mutate(log_ave_lit = log(ave_lit + 1))

slice_sample(ostracods)

```

```

# A tibble: 1 x 24
  date_site      date                site taxon  ave_lit med_ph med_sal med_do
  <chr>          <dtm>                <chr> <chr>    <dbl>  <dbl>   <dbl>  <dbl>
1 2022-02-21_NPB 2022-02-21 00:00:00 NPB   ostrac~    1.33   7.39    34.1   0.34
# i 16 more variables: taxonRank <chr>, scientificName <chr>, kingdom <chr>,
#   Phylum <chr>, Class <chr>, Subclass <chr>, Superorder <chr>, Order <chr>,
#   Infraorder <chr>, Family <chr>, Subfamily <chr>, Tribe <chr>, Genus <chr>,
#   comments <chr>, site_full <fct>, log_ave_lit <dbl>

```


c. Code your figure

```

# base layer
ostracodgraph <- ggplot(data = ostracods,
  # x-axis: site
  # y-axis: mean salinity
  mapping = aes(x = site_full,
    y = log_ave_lit)) +
# salinity measurements at each survey
geom_jitter(height = 0,
  width = 0.1,
  shape = 21) +
# wrapping x-axis text
scale_x_discrete(labels = label_wrap(width = 15)) +
# points to represent means
stat_summary(fun = median,
  color = "orange",
  size = 1) +
# labelling x-axis and y-axis
labs(x = "Site",
  y = "log + 1 transformed average abundance/L",
  title = "Ostracod Abundance Across Sites") +
# change theme from default
theme_minimal()

# displaying object
ostracodgraph

```




d. Create a multipanel figure

```
# salinity
salinity <- ggplot(data = aq_ins_clean,
  # x-axis: site
  # y-axis: mean salinity
  mapping = aes(x = site_full,
    y = med_sal)) +
  # salinity measurements at each survey
  geom_jitter(height = 0,
    width = 0.1,
    shape = 21) +
  # wrapping x-axis text
  scale_x_discrete(labels = label_wrap(width = 15)) +
  # points to represent means
  stat_summary(fun = median,
    color = "red",
    size = 1) +
  # labelling x-axis and y-axis
  labs(x = "Site",
    y = "Median salinity (ppt)",
```

```

    title = "Differences in Median Salinity across sites") +
  # different theme
  theme_minimal()
# displaying object
salinity

```



```

# cleaning copepods before graphing
copepods <- aq_ins_clean |>
  # filter to only include copepods
  filter(taxon == "copepod") |>
  # log + 1 transform mean abundance per liter
  mutate(log_ave_lit = log(ave_lit + 1))

```

8

[1] 8

```


# base layer
copepods <- ggplot(data = copepods,
  # x-axis: site
  # y-axis: mean abundance/L
  mapping = aes(x = site_full,

```

```


                                y = log_ave_lit))) +
# first layer: points to represent observations
geom_jitter(height = 0,
            shape = 21,
            width = 0.1) +
# second layer: points to represent medians
stat_summary(geom = "point",
            fun = "median",
size = 4,
            color = "blue") +
# wrapping x-axis text
scale_x_discrete(labels = label_wrap(15)) +
# relabelling x- and y-axis, adding title
labs(x = "Site",
     y = "log + 1 transformed average abundance/L",
     title = "Copepod Abundance Across Sites") +
# different theme
theme_minimal()
# displaying object
copepods

```




```
combined_plot <-
  (salinity + labs(x = NULL)) /
  plot_spacer() /
  (copepods | ostracodgraph) +
  plot_layout(heights = c(1, 0.4, 1)) &
  theme(
    plot.margin = margin(15, 15, 15, 15),
    axis.text.x = element_text(angle = 45, hjust = 1),
    axis.title.x = element_text(margin = margin(t = 15)),
    axis.title.y = element_text(margin = margin(r = 15))
  )

combined_plot
```



```
ggsave("multipanel_figure.png", combined_plot, width = 14, height = 10, dpi = 300)
```

e. Interpret your figure

- The Copepod and Ostracod abundance both vary across all sites. However, the sites where they are most and least abundant are different. For example, Copepods are most abundant at Venoco Bridge, Devereux Creek, and NA. However, Ostracods are most abundant at West Pond, South of Phelps Bridge, and Devereux Creek.

2. Visualizing total abundance as a function of salinity

- a. Plan your figure**
- b. Wrangle your data**
- c. Code your figure**
- d. Interpret your figure**