

# Finding $E$ from Time of Flight

- Starting at Perigee:
  - $n(t - T) = E - e \sin E$  (from perigee)
  - How do we find  $E$ ?
- No analytic solution, can find by iteration
- Commonly use Newton-Raphson iteration:

$$x_{n+1} = x_n - \frac{f(x)}{f'(x)}$$

# Iteration for $E$ (Starting at Perigee)

$$f(E_k) = n(t - T) - (E_k - e \sin E_k)$$

$$f'(E_k) = -(1 - e \cos E_k)$$

$$n(t - T) = M \quad (\text{fixed})$$

$$E_{k+1} = E_k + \frac{M - (E_k - e \sin E_k)}{1 - e \cos E_k} \quad \text{Continue until } |E_{k+1} - E_k| < \text{TOL}$$

Specify iteration starting value:


$$E_0 = M \quad \text{should usually work}$$

# Alternate Solution (Biondini)

- Let:  
$$M = E - e \sin E$$
$$E = M + e \sin E$$

- Take sine of both sides:

$$\sin E = \sin(M + e \sin E)$$


$$\begin{aligned} x &= \sin E \\ x_{i+1} &= \sin(M + ex_i) \\ E_{i+1} &= \sin^{-1}(M + ex_{i+1}) \end{aligned}$$

- Repeat until

$$\Delta x = |x_{i+1} - x_i| \rightarrow 0$$

# General Form for Time of Flight

- Not starting at perigee, setup for time of flight from  $t_0$  to  $t$
- General Time of Flight Equation

$$t - t_0 = k TP + \frac{1}{n} \underbrace{(E - e \sin E)}_M - \frac{1}{n} \underbrace{(E_0 - e \sin E_0)}_{M_0}$$

–  $k$  = number of perigee crossings between  $E_0$  and  $E$

- General Iteration function

$$f(E_k) = n(t - T) - (E_k - e \sin E_k) + (E_0 - e \sin E_0)$$

- Note: the ( $k TP$ ) term is not included: this version allows  $E_k > 2\pi$

# Newton-Raphson Iteration for E

- As before

$$n(t - T) = n\Delta t = \text{fixed}$$

$$M_0 = (E_0 - e \sin E_0) = \text{fixed}$$

$$f'(E_k) = -(1 - e \cos E_k)$$

$$E_{k+1} = E_k + \frac{n\Delta t + M_0 - (E_k - e \sin E_k)}{1 - e \cos E_k}$$

Specify iteration starting value:

$$E_0 = M_0 \quad \text{should usually work}$$

# Note on the Iteration

- Perigee passages are not included in the iteration,

$$E_{k+1} = E_k + \frac{n\Delta t + M_0 - (E_k - e \sin E_k)}{1 - e \cos E_k}$$

- If  $k > 1$ , result will be a value of  $E > 2\pi$ 
  - Take  $\text{mod}(E, 2\pi)$  to get  $E$  in  $[0, 2\pi]$
  - Can compute  $k$  by

$$k = \text{floor}((E - E_o) / 2\pi)$$


# Getting Right Quadrant for $v$

- Use solved-for value of  $E$  to get  $v$

$$\cos v = \frac{e - \cos E}{e \cos E - 1}$$

$$\sin v = \frac{\sin E \sqrt{1 - e^2}}{1 - e \cos E}$$

$$v = \text{atan2}(\sin v, \cos v)$$



- $\text{atan2}$  returns angle in range  $[-\pi, \pi]$ 
  - If angle  $< 0$ , add  $2\pi$  to get range  $[0, 2\pi]$

# An Aside: Planar Motion

- Things we know:

$$\vec{h} = \vec{r} \times \vec{v}$$

$$\vec{h} \perp \vec{r} \text{ and } \vec{v}$$

$$\vec{h} = \text{constant}$$

- Therefore:
  - Satellite motion is always in the plane containing  $\vec{r}$  and  $\vec{v}$
  - For 2-body problem, can treat prediction problem as a problem in planar motion
    - Don't worry about orientation of orbit in space (yet)




# The Kepler Problem

- Orbit elements are constant functions of  $\vec{r}$  and  $\vec{v}$
- Look at orbit in perifocal frame:
  - Motion in orbit plane

$$\vec{r} = x \hat{P} + y \hat{Q} + 0 \hat{W}$$

$$\dot{\vec{r}} = \dot{x} \hat{P} + \dot{y} \hat{Q} + 0 \hat{W}$$

( $\hat{P}$ ,  $\hat{Q}$ ,  $\hat{W}$  are inertial, so  $d/dt = 0$ )



# Prediction

- Because the orbit elements are constant, it is reasonable to suppose that future states can be found from current states:

$$\vec{r} = f \vec{r}_0 + g \dot{\vec{r}}_0$$

$$\dot{\vec{r}} = \dot{f} \vec{r}_0 + \dot{g} \dot{\vec{r}}_0$$

$$(\vec{r}_0, \dot{\vec{r}}_0 \text{ are fixed, so } d/dt = 0)$$

- Need to find expressions for  $f$ ,  $g$ , etc.

# Find f

- Post-cross multiply both sides by  $\dot{\vec{r}}_0$

$$\vec{r} \times \dot{\vec{r}}_0 = f \vec{r}_0 \times \dot{\vec{r}}_0 + g \cancel{\dot{\vec{r}}_0 \times \dot{\vec{r}}_0} \quad \text{0}$$

$$\vec{r} \times \dot{\vec{r}}_0 = f \vec{r}_0 \times \dot{\vec{r}}_0 = f \vec{h}$$


$$\vec{r} \times \dot{\vec{r}}_0 = \begin{vmatrix} \hat{P} & \hat{Q} & \hat{W} \\ x & y & 0 \\ \dot{x}_0 & \dot{y}_0 & 0 \end{vmatrix} = (x\dot{y}_0 - \dot{x}_0 y) \hat{W} \quad \text{Note : } \vec{h} = h \hat{W}$$

$$(x\dot{y}_0 - \dot{x}_0 y) \cancel{\hat{W}} = f h \cancel{\hat{W}} \implies (x\dot{y}_0 - \dot{x}_0 y) = f h$$

$$f = \frac{(x\dot{y}_0 - \dot{x}_0 y)}{h}$$

# Find g

- Pre-cross multiply both sides by  $\vec{r}_0$

$$\vec{r}_0 \times \vec{r} = f \vec{r}_0 \times \vec{r}_0 + g \vec{r}_0 \times \dot{\vec{r}}_0$$


$$\vec{r}_0 \times \vec{r} = g \vec{r}_0 \times \dot{\vec{r}}_0 = g \vec{h}$$

$$\vec{r}_0 \times \vec{r} = \begin{vmatrix} \hat{P} & \hat{Q} & \hat{W} \\ x_0 & y_0 & 0 \\ x & y & 0 \end{vmatrix} = (x_0 y - x y_0) \hat{W}$$

Similar to above:

$$g = \frac{(x_0 y - x y_0)}{h}$$

# Time derivatives of f and g

$$\dot{f} = \frac{d}{dt} \left( \frac{(x\dot{y}_0 - \dot{x}_0 y)}{h} \right) = \frac{(\dot{x}\dot{y}_0 - \dot{x}_0 \dot{y})}{h}$$

( $x_0$  and  $y_0$  are fixed)

$$\dot{g} = \frac{d}{dt} \left( \frac{(x_0 y - x y_0)}{h} \right) = \frac{(x_0 \dot{y} - \dot{x} y_0)}{h}$$

# Express in terms of $v$

$$x = r \cos v \quad y = r \sin v$$

$$\dot{x} = \dot{r} \cos v - r \sin v \dot{v}$$


$$\dot{y} = \dot{r} \sin v + r \cos v \dot{v}$$

$$v_n = |\vec{v}| \cos \phi = r \dot{v}$$

$$|\vec{h}| = |\vec{r} \times \vec{v}| = rv \sin \gamma = rv \cos \phi$$

$$h = r (r \dot{v}) = r^2 \dot{v}$$

$$r \dot{v} = \frac{h}{r}$$



$$r = \frac{a(1-e^2)}{1+e\cos v} = \frac{p}{1+e\cos v} \quad \dot{r} = -\frac{p}{(1+e\cos v)^2}(-e\sin v)\dot{v}$$

$$\dot{r} = \frac{\cancel{p}}{(1+\cancel{e\cos v})} \frac{(e\sin v)\dot{v}}{(1+e\cos v)} = \frac{re\sin v\dot{v}}{(1+e\cos v)}$$

$$\dot{r}\cos v = \frac{re\cos v\sin v\dot{v}}{(1+e\cos v)}$$

$$\dot{x} = \frac{re\cos v\sin v\dot{v}}{(1+e\cos v)} - r\sin v\dot{v} = \frac{re\cos v\sin v\dot{v}}{(1+e\cos v)} - \frac{r\sin v(1+e\cos v)\dot{v}}{(1+e\cos v)}$$

$$\dot{x} = \frac{\cancel{re\cos v\sin v\dot{v}} - r\sin v\dot{v} - \cancel{re\cos v\sin v\dot{v}}}{(1+e\cos v)} \implies \boxed{\dot{x} = \frac{-r\sin v\dot{v}}{(1+e\cos v)}}$$

# Solving...

$$\dot{x} = \frac{-r \sin v \dot{v}}{(1 + e \cos v)}$$

$$r \dot{v} = \frac{h}{r} \implies \dot{x} = \frac{-h \sin v}{r (1 + e \cos v)}$$

$$p = r (1 + e \cos v) = \frac{h^2}{\mu} \implies h = \sqrt{\mu p}$$

$$\dot{x} = \frac{-\sqrt{\mu p} \sin v}{p} \implies \dot{x} = -\sqrt{\frac{\mu}{p}} \sin v$$



# Similarly for $\dot{y}$

$$\dot{y} = \dot{r} \sin v + r \cos v \dot{v}$$

$$r = p (1 + e \cos v)^{-1}$$

$$\dot{r} = -p (1 + e \cos v)^{-2} (-e \sin v) \dot{v}$$

$$\dot{r} = \frac{p e \sin v}{(1 + e \cos v)^2} \dot{v}$$

$$\dot{r} = \frac{r^2 e \sin v}{p} \dot{v}$$

$$h = r^2 \dot{v}$$

$$\dot{r} = \frac{h}{p} e \sin v$$

$$h = r(r \dot{v}) = \sqrt{\mu p}$$

$$r \dot{v} = \frac{\sqrt{\mu p}}{r}$$

$$r \dot{v} = \frac{\sqrt{\mu p} (1 + e \cos v)}{p}$$

$$r \dot{v} = \sqrt{\frac{\mu}{p}} (1 + e \cos v)$$

$$\dot{r} = \sqrt{\frac{\mu}{p}} e \sin v$$

# Combine

$$\dot{y} = \dot{r} \sin v + r \cos v \dot{v}$$

$$\dot{y} = \sqrt{\frac{\mu}{p}} e \sin^2 v + \sqrt{\frac{\mu}{p}} (1 + e \cos v) \cos v$$

$$\dot{y} = \sqrt{\frac{\mu}{p}} (e [\sin^2 v + \cos^2 v] + \cos v)$$

$$\dot{y} = \sqrt{\frac{\mu}{p}} (e \sin^2 v + \cos v + e \cos^2 v) \implies \boxed{\dot{y} = \sqrt{\frac{\mu}{p}} (e + \cos v)}$$

# Summary so far

$$x = r \cos v$$

$$y = r \sin v$$

$$\dot{x} = -\sqrt{\frac{\mu}{p}} \sin v$$

$$\dot{y} = \sqrt{\frac{\mu}{p}} (e + \cos v)$$

Perifocal Position and  
Velocity Components

$$f = \frac{(x\dot{y}_0 - \dot{x}_0 y)}{h}$$

$$g = \frac{(x_0 y - x y_0)}{h}$$

$$\dot{f} = \frac{(\dot{x}\dot{y}_0 - \dot{x}_0 \dot{y})}{h}$$

$$\dot{g} = \frac{(x_0 \dot{y} - \dot{x} y_0)}{h}$$

$$f\dot{g} - \dot{f}g = 1$$

# Combine to find expressions in $v$

Example: find  $f$

$$f = \frac{(x\dot{y}_0 - \dot{x}_0 y)}{h}$$

$$f = \frac{\left( r \cos v \sqrt{\frac{\mu}{p}} (e + \cos v_0) + \sqrt{\frac{\mu}{p}} \sin v_0 r \sin v \right)}{\sqrt{\mu p}}$$

$$f = \frac{r(\cos v (e + \cos v_0) + \sin v_0 \sin v)}{p}$$

$$f = \frac{r(e \cos v + \cos v \cos v_0 + \sin v_0 \sin v)}{p}$$

Recall the trig identity...

$$\cos A \cos B \pm \sin A \sin B = \cos(A \mp B)$$

$$f = \frac{r(e \cos v + \cos(v-v_0))}{p} = \frac{r(e \cos v + \cos \Delta v)}{p}$$

Add and subtract...

$$\frac{r}{p} - \frac{r}{p} + \frac{r e \cos v}{p} = -\frac{r}{p} + \frac{r}{p}(1 + e \cos v)$$

$$p = r(1 + e \cos v)$$

$$-\frac{r}{p} + \frac{r}{p}(1 + e \cos v) = -\frac{r}{p} + 1$$

$$\frac{r e \cos v}{p} = 1 - \frac{r}{p}$$

$$f = 1 - \frac{r}{p} + \frac{r}{p} \cos \Delta v$$

$$f = 1 - \frac{r}{p}(1 - \cos \Delta v)$$

# Similarly

$$g = \frac{r r_0}{\sqrt{\mu p}} \sin \Delta v$$

$$\dot{g} = 1 - \frac{r_0}{p} (1 - \cos \Delta v)$$

Using  $f\ddot{g} - \dot{f}\dot{g} = 1$ :

$$\dot{f} = \sqrt{\frac{\mu}{p}} \tan \frac{\Delta v}{2} \left( \frac{1 - \cos \Delta v}{p} - \frac{1}{r} - \frac{1}{r_0} \right)$$

# The f&g Functions: So What?

- The f&g functions: undeniably cool, but rather esoteric-seeming
- Does anybody actually use them?
- Answer: yes!
  - Cheap lower-fidelity orbit propagation
    - Can extend to non-two-body formulations
  - Commonly used for initial orbit determination (orbits from observations)
  - Basis of “Lambert Targeting” methods for rendezvous/proximity ops
  - Provides analytic solution for orbit determination State Transition Matrix
  - Universal variable formulation generalizes to all conic sections
  - And more!

# Homework 2

Due Saturday, 31 Jan 2026

# Homework 2 (1)

Given:

Component	Value (meters)
Position X	326151.080726
Position Y	6077471.251787
Position Z	2944583.918767
Velocity X	-7455.178720
Velocity Y	-482.482572
Velocity Z	1910.883434

- Find  $E_o$  (slide 19)
- Find  $v_o$  (Keplerian comps)
- Find  $M_o$
- Find time of flight from perigee to  $v_o$

- Find time of flight from  $v_o$  to  $v = 65^\circ$
- Verify: Starting at  $v_o$ , compute the true anomaly after the time of flight computed above.
- Starting at  $v_o$ , what is the true anomaly after 2700 seconds?
- Starting at  $v_o$ , what is the true anomaly after exactly two orbit periods?
- What is the true anomaly after 15000 seconds?



# Homework 2 (2)

Given:

Component	Value (meters)
Position X	572461.711228
Position Y	-1015437.194396
Position Z	7707337.871302
Velocity X	-6195.262945
Velocity Y	-3575.889650
Velocity Z	-5.423283

- Compute Keplerian elements
- Find perifocal position and velocity,  $\vec{r}_o$  and  $\vec{v}_o$ 
  - See equations on Slide 39
- Find  $f, g, \dot{f}, \dot{g}$  for  $\Delta\nu = 33^\circ$
- Find  $\vec{r}$  and  $\vec{v}$

# Homework 2 (3)

- Derivation 1: Using energy equation and ellipse equation, find an expression for the orbit speed as a function of true anomaly
- Derivation 2: Using the above, show that

$$v_{\text{perigee}} = \sqrt{\frac{\mu}{a} \left( \frac{1+e}{1-e} \right)}$$

$$v_{\text{apogee}} = \sqrt{\frac{\mu}{a} \left( \frac{1-e}{1+e} \right)}$$