

Analysis III Zusammenfassung

Dino Colombo, Michael van Huffel

This summary has been written based on the Lecture of Analysis III by Prof. Dr. Francesca Da Lio (Autumn 20) and the summary of Philip Wolf. There is no guarantee for completeness and/or correctness regarding the content of this summary. Use it at your own discretion

0 Contents			
1	Laplace Transform	2	6.1.3 Solution to problem with $g_1(x)$ (u_B)
1.1	General Formulas	2	6.2 Laplacian in Polar Coordinates
1.2	Heavyside Function or Unit Step Function	2	6.3 Dirichlet Problem for a Disk
1.3	Heavyside Expansions	2	6.4 Dirichlet Problem for a Ring
1.4	Dirac's Delta	2	6.4.1 Cosinus and Sinus variant
1.5	Convolution	2	6.4.2 Complex variant
1.5.1	Properties	2	
1.5.2	Application to Non-Homogeneous ODEs	2	
1.6	ODE/ Initial Value Problems	3	
1.7	Special Linear ODE with Variable Coeff.	3	
1.8	Systems of ODEs	3	
2	Fourier Analysis	3	7 Two-Dimensional Wave Equation
2.1	Periodicity of functions	3	7.1 Rectangular Membrane
2.2	Fourier Series	3	
2.3	Better look: $\cos(nx), \sin(mx)$	4	
2.3.1	Transform properties	4	
2.4	Even and Odd Functions/Expansions	4	
2.5	Complex Fourier Series	4	
2.6	Approximation by Trigonometric Polynomials	4	
2.7	Fourier Integral	5	
2.7.1	Fourier Cosine and Sine Integral	5	
2.8	Fourier Transforms	5	
2.8.1	Transforms of Derivatives	5	
2.8.2	Other properties	5	
3	Partial Differential Equation (PDE)	6	
3.1	Basic Concepts	6	
3.1.1	Simple Approaches	6	
3.2	First order PDE	6	
3.2.1	Method of Characteristics	6	
3.3	Second order PDE	7	
3.3.1	Method of Characteristics	7	
4	Wave Equation	7	
4.1	D'Alembert Solution of the Wave Equation	8	
4.2	Graphic solution (Characteristic lines)	8	
5	Heat Equation	9	
5.1	One-Dimensional Heat Equation	9	
5.2	One - Dimensional Heat Equation with inomogeneous BC	9	
5.3	Inomogeneous One Dimensional Heat Equation	10	
5.4	Heat Equation for Infinite Bars	10	
5.4.1	Using Fourier Transform	10	
5.4.2	Method of Convolution	10	
6	Laplace eq. (Steady 2D Heat Problem)	11	
6.1	Dirichlet problem	11	
6.1.1	Solution to problem with $f_1(x)$ (u_A)	11	
6.1.2	Solution to problem with $f_2(x)$ (u_C)	11	
			8 Tables
			8.1 Laplace Transforms
			8.2 Fourier Transforms
			8.3 Some Integrals
			8.3.1 Basic integrals
			8.3.2 Per Parts Resolution method
			8.4 Some Series
			8.5 Logarithm
			8.6 Differential Calculus
			8.7 Standard ODE
			8.8 Some Limits
			8.9 Limit
			8.9.1 Bernoulli De L'Hôpital
			8.10 Trigonometric Functions
			8.10.1 Periodicity
			8.10.2 Relation between trigonometric functions
			8.10.3 Exact value of trigonometric functions
			8.10.4 Trigonometric function with $t = \tan(\frac{\alpha}{2})$
			8.10.5 Addition Theorems
			8.10.6 Sums and Products
			8.10.7 Inverse Functions
			8.11 Hyperbolic Functions
			8.12 Graph of some common functions

1 Laplace Transform

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt$$

1.1 General Formulas:

$f(t) = \mathcal{L}^{-1}\{F(s)\}$	Inverse
$\mathcal{L}\{af(t) + bg(t)\} = aF(s) + bG(s)$	Linearity
$\mathcal{L}\{e^{+at}f(t)\} = F(s-a)$ $\mathcal{L}^{-1}\{F(s-a)\} = e^{at}f(t)$	s-Shifting
$\mathcal{L}\{f(t-a)u(t-a)\} = e^{-as}F(s)$ $\mathcal{L}^{-1}\{e^{-as}F(s)\} = f(t-a)u(t-a)$	t-Shifting
$\mathcal{L}(f') = s\mathcal{L}(f) - f(0)$ $\mathcal{L}(f'') = s^2\mathcal{L}(f) - sf(0) - f'(0)$ $\mathcal{L}(f^{(n)}) = s^n\mathcal{L}(f) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$	Differentiation of Function
$\mathcal{L}\left(\int_0^t f(\tau) d\tau\right) = \frac{1}{s}\mathcal{L}(f)$	Integration of Function
$\mathcal{L}(tf(t)) = -F'(s)$ $\mathcal{L}(t^n f(t)) = (-1)^n F^{(n)}(s)$	Differentiation of Transform
$\mathcal{L}\left(\frac{f(t)}{t}\right) = \int_s^\infty F(\tilde{s}) d\tilde{s}$	Integration of Transform

Theorem: Partial fraction decomposition

- Single zero: $\frac{A}{x-x_0}$
- Double zeros: $\frac{A}{x-x_0} + \frac{B}{(x-x_0)^2}$
- Complex zeros: $\frac{A \cdot x + B}{\text{i.e.} \rightarrow x^2 + 1}$

Theorem: Existence

If $f(t)$ is defined on piecewise continuous on every finite interval on the semi-axis $t \geq 0$ and satisfies (1) $\forall t \geq 0$ and some constant M, k , the the Laplace transform exists $\forall s > k \rightarrow |f(t)| \leq Me^{kt}$

Theorem: Uniqueness

If two continuous functions have the same transform, they are identical.

1.2 Heavyside Function or Unit Step Function:

$$u(t-a) = \begin{cases} 0 & \text{if } t < a \\ 1 & \text{if } t > a \end{cases} \quad \mathcal{L}\{u(t-a)\} = \frac{e^{-as}}{s}$$

1.3 Heavyside Expansions:

The subsidiary equation usually appears as a quotient $Y(s) = F(s)/G(s)$, so that partial fraction is necessary.

- Repeated real factors** (in $G(s)$)
 $(s-a)^n \rightarrow \sum_{i=1}^n \frac{A_i}{(s-a)^i}$
 $\xrightarrow{\mathcal{L}^{-1}} e^{at}(A_1 + A_2t + \frac{1}{2}A_3t^2 + \dots + \frac{1}{n-1}A_nt^{n-1})$
- Unrepeated complex factors**
 $(s-a)(s-\bar{a}), a = \alpha + i\beta \rightarrow \frac{As+B}{(s-\alpha)^2 + \beta^2}$

1.4 Dirac's Delta:

$$\delta(t-a) = \lim_{k \rightarrow 0} f_k(t-a) = \begin{cases} \infty & \text{if } t = a \\ 0 & \text{otherwise} \end{cases}$$

$$\int_0^\infty g(t)\delta(t-a) dt = g(a) \quad \text{sifting property}$$

$$\int_0^\infty \delta(t-a) dt = 1 \quad \mathcal{L}\{\delta(t-a)\} = e^{-as}$$

1.5 Convolution:

Caution: $\mathcal{L}\{fg\} \neq \mathcal{L}\{f\}\mathcal{L}\{g\}$ (in general)

$$h(t) = (f * g)(t) = \int_0^t f(\tau)g(t-\tau) d\tau$$

$$\mathcal{L}(h) = \mathcal{L}(f)\mathcal{L}(g) \Leftrightarrow H = FG$$

$$\mathcal{L}^{-1}(F(s) \cdot G(s)) = \mathcal{L}^{-1}(F(s)) * \mathcal{L}^{-1}(G(s))$$

Example: Convolution

$$y(t) - \int_0^t y(\tau) \sin(t-\tau) d\tau = t$$

This equation can be written as: $y - y * \sin(t) = t$ (convolution theorem) $\Rightarrow Y - Y \frac{1}{s^2 + 1} = \frac{s^2}{s^2 + 1} Y = \frac{1}{s^2} \Leftrightarrow Y = \frac{s^2 + 1}{s^4}$
 $\Rightarrow Y(s) = \frac{1}{s^2} + \frac{1}{s^4} \Rightarrow y(t) = t + \frac{t^3}{6}$

1.5.1 Properties:

$$\begin{array}{l|l} f * g = g * f & f(g_1 * g_2) = f * g_1 + f * g_2 \\ (f * g) * v = f * (g * v) & f * 0 = 0 * f = 0 \\ f * 1 \neq f & f * f \geq 0 \text{ not necessarily} \end{array}$$

Example: Laplace Transform of a function

$$f(t) = g(t) + \int_0^t f(\tau) \cdot h(t-\tau) d\tau$$
$$\Rightarrow f(t) = \mathcal{L}^{-1}\left(\frac{G(s)}{1-H(s)}\right)$$

1.5.2 Application to Non-Homogeneous ODEs:

$$y(0) = y'(0) = 0 \Rightarrow Y = QR, y(t) = \int_0^t q(t-\tau)r(\tau) d\tau$$

Example: Laplace application to an NHODE

Calculate: $\mathcal{L}^{-1} \ln\left(\frac{s^2 + \omega^2}{s^2}\right)$

Differentiation:
 $F'(s) = \frac{d}{ds} [\ln(s^2 + \omega^2) - \ln(s^2)] = \frac{2s}{s^2 + \omega^2} - \frac{2s}{s^2}$
 $\Rightarrow \mathcal{L}^{-1}\{F'(s)\} = 2 \cos \omega t - 2 = -tf(t)$

Differentiation & Integration:
 $G(s) = F'(s) \Rightarrow g(t) = \mathcal{L}^{-1}\{G\} = 2(\cos(\omega t) - 1)$
 $\mathcal{L}^{-1}\left\{\ln \frac{s^2 + \omega^2}{s^2}\right\} = \mathcal{L}^{-1}\left\{\int_s^\infty G(\tilde{s}) d\tilde{s}\right\} = -\frac{g(t)}{t}$

1.6 ODE/ Initial Value Problems:

$$y'' + ay' + by = r(t), \quad y(0) = K_0, \quad y'(0) = K_1$$

$r(t)$: input $y(t)$: output

1. Setting up subsidiary equation:

$$Y = \mathcal{L}(y), \quad R = \mathcal{L}(r)$$

$$[s^2 Y - sy(0) - y'(0)] + a[sY - y(0)] + bY = R(s)$$

$$(s^2 + as + b)Y = (s + a)y(0) + y'(0) + R(s)$$

2. Solution of subsidiary equation:

$$Q = \frac{1}{s^2 + as + b} = \frac{1}{(s + \frac{1}{2}a)^2 + b - \frac{1}{4}a^2} \quad \text{transfer function}$$

$$\Rightarrow Y(s) = [(s + a)y(0) + y'(0)]Q(s) + R(s)Q(s)$$

$Q(s)$ doesn't depend on $r(t)$ or on the initial conditions

3. Inversion of Y :

Reduce to sum of terms (inverses can be found in tables).

$$y(t) = \mathcal{L}^{-1}(Y)$$

1.7 Special Linear ODE with Variable Coeff.:

$$\mathcal{L}(ty') = -\frac{d}{ds}[sY - y(0)] = -Y - s\frac{dY}{ds}$$

$$\mathcal{L}(ty'') = -\frac{d}{ds}[s^2 - sy(0) - y'(0)] = -2sY - s^2\frac{dY}{ds} + y(0)$$

1.8 Systems of ODEs:

We consider a **first-order** linear system with constant coefficients.

$$\begin{cases} y_1' = a_{11}y_1 + a_{12}y_2 + g_1(t) \\ y_2' = a_{21}y_1 + a_{22}y_2 + g_2(t) \end{cases} \Rightarrow y' = Ay + g \xrightarrow{\mathcal{L}}$$

$$sY - y(0) = AY + G \Leftrightarrow \begin{cases} (a_{11} - s)Y_1 + a_{12}Y_2 = -y_1(0) - G_1(s) \\ a_{21}Y_1 + (a_{22} - s)Y_2 = -y_2(0) - G_2(s) \end{cases}$$

$$\Rightarrow (A - s\mathbb{I})Y = -y(0) - G$$

\rightarrow solve algebraically and take the inverse transform of Y

Theorem: Cramer's Rule

$Ax = b$ with $A \in \mathbb{R}^{n \times n}$, $x, b \in \mathbb{R}^n$, $D := \det A$, and D_k is the determinant obtained by replacing k^{th} column by b

$$\Rightarrow x_k = \frac{D_k}{D}$$

Example: Laplace res of ODE impulse function

$$y'' + 3y' + 2y = r(t),$$

$$r(t) = \begin{cases} 1 & \text{if } 1 < t < 2 \\ 0 & \text{otherwise} \end{cases}, \quad y(0) = y'(0) = 0$$

$$\Rightarrow Q(s) = \frac{1}{s^2 + 3s + 2} = \frac{1}{(s+1)(s+2)} = \frac{1}{s+1} - \frac{1}{s+2}$$

$$\Rightarrow q(t) = e^{-t} - e^{-2t}$$

$$\Rightarrow y(t) = \int q(t-\tau) \cdot 1 d\tau = \int [e^{-(t-\tau)} - e^{-2(t-\tau)}] d\tau$$

$$= e^{-(t-\tau)} - \frac{1}{2}e^{-2(t-\tau)}$$

$r(\tau) = 1$ if $1 < \tau < 2$ only. So if $t < 1$, the integral is zero. If $1 < t < 2$, we have to integrate from $\tau = 1$ to t :

$$y_2(t) = e^{-0} - \frac{1}{2}e^{-0} - (e^{-(t-1)} - \frac{1}{2}e^{-2(t-1)})$$

$$= \frac{1}{2} - e^{-(t-1)} + \frac{1}{2}e^{-2(t-1)}$$

If $t > 2$, we have to integrate from $\tau = 1$ to 2:

$$y_3(t) = e^{-(t-2)} - \frac{1}{2}e^{-2(t-2)} - (e^{-(t-1)} - \frac{1}{2}e^{-2(t-1)})$$

$$\Rightarrow y(t) = \begin{cases} y_1(t) = 0, & t < 1 \\ y_2(t), & 1 < t < 2 \\ y_3(t), & t > 2 \end{cases}$$

2 Fourier Analysis

2.1 Periodicity of functions:

Theorem: Fundamental Theorem of Periodicity

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is periodic of period $P > 0$ if $f(x+P) = f(x) \forall x \in \mathbb{R}$.

A fundamental period is (if it exists) the smallest positive number P for which f is periodic of period P . Ex: for $\sin(mx), \cos(mx)$ $P = \frac{2\pi}{m}$. **Attention:** $\cos(x)^2$ and $\cos(x)^4$ as well as $\sin \dots$ have period of π .

Theorem: Period of the sum of functions

- Let be P_f and P_g the period of 2 periodic functions f and g . The function $f+g$ or $f \cdot g$ is periodic iff $\frac{P_f}{P_g} \in \mathbb{Q}$. The Period of $f+g$ is also the l.c.m(P_f, P_g).
- If f, g are functions periodic of period p and $\alpha, \beta \in \mathbb{R}$, then $\alpha f + \beta g$ is periodic of period p

Theorem: Property of boundedness

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be any function:

- If f is periodic and continuous, then it is bounded.
- If f is differentiable and periodic of period P , then also f' is periodic with the same period.
- If f is periodic and smooth, then it is bounded and all its derivative are bounded as well.

2.2 Fourier Series:

$$F(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos \frac{n\pi}{L}x + b_n \sin \frac{n\pi}{L}x) \quad \text{Fourier series}$$

Beachte: The Fourier series $F(x)$ converges to the function $f(x)$ if the function is continuous.

$$(0) \quad a_0 = \frac{1}{2L} \int_{-L}^L f(x) dx$$

$$(a) \quad a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx$$

$$(b) \quad b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx$$

Theorem: Fourier coefficient

The Fourier coefficients of $f_1 + f_2$ are the sums of the corresponding coefficients of f_1 and f_2 .

The Fourier coefficients of αf are α times the corresponding coefficients of f .

Theorem: Representation of a function by FS/ FI

Let f be the $2L$ -periodic function, piecewise continuous:

- $\lim_{h \rightarrow 0^+} \frac{f(x_0+h) - f(x_0)}{h} = f'_+(x_0),$
- $\lim_{h \rightarrow 0^-} \frac{f(x_0+h) - f(x_0)}{h} = f'_-(x_0)$

Then:

- If x_0 is a point of continuity of $f \Rightarrow$ the Fourier Series (FS/FI) converges to $F(x) = f(x_0)$
- If x_0 is a point of discontinuity of $f \Rightarrow$ the Fourier Series (FS/FI) converges to $F(x) = \frac{f(x_0^+) + f(x_0^-)}{2}$

Example: Fourier coefficients

$$x(\pi - x) \stackrel{!}{=} \sum_{n=1}^{\infty} b_n \sin nx$$

Solution:

$$\begin{aligned} b_n &= \frac{2}{\pi} \int_0^{\pi} x(\pi - x) \sin nx \, dx \\ &= \underbrace{-\frac{2x(\pi - x)}{n\pi} \cos nx \Big|_0^{\pi}}_{=0} + \frac{2}{n\pi} \int_0^{\pi} (\pi - 2x) \cos nx \, dx \\ &= \underbrace{\frac{2(\pi - 2x)}{n^2\pi} \sin nx \Big|_0^{\pi}}_{=0} + \frac{4}{n^2\pi} \int_0^{\pi} \sin nx \, dx = -\frac{4}{n^3\pi} \cos nx \Big|_0^{\pi} \end{aligned}$$

2.3 Better look: $\cos(nx), \sin(mx)$:**Theorem: Orthogonality of Trigonometric System**

The *trigonometric system* is orthogonal on any interval of length 2π ; that is: $n, m \in \mathbb{N}$, $n \neq m$

- (a) $\int_{-\pi}^{\pi} \cos nx \cos mx \, dx = 0$
- (b) $\int_{-\pi}^{\pi} \sin nx \sin mx \, dx = 0$
- (c) $\int_{-\pi}^{\pi} \sin nx \cos mx \, dx = 0$

$$\int \sin(mx) \cdot \cos(nx) \, dx = \begin{cases} -\frac{\cos(mx-nx)}{2 \cdot (m-n)} - \frac{\cos(mx+nx)}{2 \cdot (m+n)}, & m \neq n \\ -\frac{(\cos(mx))^2}{2m}, & m = n \end{cases}$$

$$\int \sin(mx) \cdot \sin(nx) \, dx = \begin{cases} \frac{\sin(mx-nx)}{2 \cdot (m-n)} - \frac{\sin(mx+nx)}{2 \cdot (m+n)}, & m \neq n \\ \frac{x}{2} - \frac{\sin 2mx}{ma}, & m = n \end{cases}$$

$$\int \cos(mx) \cdot \cos(nx) \, dx = \begin{cases} \frac{\sin(mx-nx)}{2 \cdot (m-n)} + \frac{\sin(mx+nx)}{2 \cdot (m+n)}, & m \neq n \\ \frac{x}{2} + \frac{\sin 2mx}{ma}, & m = n \end{cases}$$


2.3.1 Transform properties:Let $n \in \mathbb{N}$:

- $\sin(n\pi) = 0$
- $\cos(n\pi) = (-1)^n$

- $\cos(n\frac{\pi}{2}) = (\frac{1+(-1)^n}{2}) \cdot (-1)^{\frac{n}{2}} = \begin{cases} 0, n = 2j + 1 \\ (-1)^j, n = 2j \end{cases}$
- $\sin(n\frac{\pi}{2}) = (\frac{1+(-1)^n}{2}) \cdot (-1)^{\frac{n+2}{2}} = \begin{cases} 0, n = 2j \\ (-1)^j, n = 2j + 1 \end{cases}$
- $\sin((n \pm 1)\frac{\pi}{2}) = \pm \cos(\frac{n\pi}{2})$
- $\cos((n \pm 1)\frac{\pi}{2}) = \pm \sin(\frac{n\pi}{2})$
- $\cos((n \pm \frac{1}{2})\pi) = 0, \quad \sin((n \pm \frac{1}{2})\pi) = \pm \cos(n\pi)$


2.4 Even and Odd Functions/Expansions:**Theorem: Even and odd functions**Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be any differentiable functions:

- if f is even, f' is odd
- if f is odd, f' is even
- $e_1(x) + e_2(x)$ is even
- $e_1(x) \cdot e_2(x)$ is even
- $o_1(x) + o_2(x)$ is odd
- $o_1(x) \cdot o_2(x)$ is even
- $o_1(x) \cdot e_1(x)$ is odd
- $\int_{-a}^a e_1(x) \, dx = 2 \cdot \int_0^a e_1(x) \, dx$
- $\int_{-a}^a o_1(x) \, dx = 0$

Even Expansion (gerade):If $f(x)$ is an **even $2L$ periodic function** ($f(-x) = f(x)$)

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi}{L} x$$

$$a_0 = \frac{1}{L} \int_0^L f(x) \, dx, \quad a_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi x}{L} \, dx \quad n \in \mathbb{N}$$

Odd Expansion (ungerade):If $f(x)$ is an **odd $2L$ periodic function** ($f(-x) = -f(x)$)

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi}{L} x$$

$$b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi x}{L} \, dx$$

2.5 Complex Fourier Series:

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{\frac{in\pi}{L} x} \quad c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-\frac{in\pi}{L} x} \, dx$$

$$e^{\pm it} = \cos t \pm i \sin t \quad \cos t = \frac{e^{it} + e^{-it}}{2} \quad \sin t = \frac{e^{it} - e^{-it}}{2i}$$

$$\begin{array}{cc} \text{comp} \rightarrow \text{real} & \text{real} \rightarrow \text{comp} \\ a_n = c_n + c_{-n} = 2 \cdot \text{Re}(c_n) & c_n = \frac{1}{2}(a_n - ib_n) \\ b_n = i(c_n - c_{-n}) = -2\text{Re}(c_n) & c_{-n} = \frac{1}{2}(a_n + ib_n) \end{array}$$

- $e^{\pm i\pi n} = \begin{cases} 1, & n \text{ even} \\ -1, & n \text{ odd} \end{cases}$
- $c_n = \bar{c}_{-n}$

2.6 Approximation by Trigonometric Polynomials: $x \in [-\pi, \pi]$, N fixedTrigonometric polynomial of degree N :

$$A_0 + \sum_{n=1}^N (A_n \cos nx + B_n \sin nx)$$

$$E^* = \int_{-L}^L f^2 \, dx - L \left[2a_0^2 + \sum_{n=1}^N (a_n^2 + b_n^2) \right] \quad \text{Square Error}$$

2.7 Fourier Integral:

$$\bullet A(w) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \cos wv \, dv$$

$$\bullet B(w) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \sin wv \, dv$$

$$F(x) = \int_0^{\infty} [A(w) \cos wx + B(w) \sin wx] \, dw$$

Theorem: Existence condition, Fourier integral

The Fourier integral exists if $f(x)$:

- is piecewise continuous in every finite interval
- has a left- and right-hand derivative at every point
- is absolutely integrable
 $\Leftrightarrow \lim_{a \rightarrow -\infty} \int_a^0 |f(x)| \, dx + \lim_{b \rightarrow \infty} \int_0^b |f(x)| \, dx$ exists

2.7.1 Fourier Cosine and Sine Integral:

Fourier cosine integral (f even):

- $f(x) = \int_0^{\infty} A(w) \cos wx \, dw$
- $A(w) = \frac{2}{\pi} \int_0^{\infty} f(v) \cos wv \, dv$

Fourier sine integral (f odd):

- $f(x) = \int_0^{\infty} B(w) \sin wx \, dw$
- $B(w) = \frac{2}{\pi} \int_0^{\infty} f(v) \sin wv \, dv$

Laplace integrals:

- $\int_0^{\infty} \frac{\cos wx}{k^2 + w^2} \, dw = \frac{\pi}{2k} e^{-kx}$
- $\int_0^{\infty} \frac{w \sin wx}{k^2 + w^2} \, dw = \frac{\pi}{2} e^{-kx}$

2.8 Fourier Transforms:

Fourier Transform:

- $\mathcal{F}(f) = \hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-iwx} \, dx$
- $\mathcal{F}^{-1}(\hat{f}) = f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(w) e^{iwx} \, dw$

Fourier cosine transform (f even):

- $\mathcal{F}_c(f) = \hat{f}_c(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \cos wx \, dx$
- $\mathcal{F}_c^{-1}(\hat{f}_c) = f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \hat{f}_c(w) \cos wx \, dw$

Fourier sine transform (f odd):

- $\mathcal{F}_s(f) = \hat{f}_s(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \sin wx \, dx$
- $\mathcal{F}_s^{-1}(\hat{f}_s) = f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \hat{f}_s(w) \sin wx \, dw$

Fourier Transforms are **linear operations**:

$$\mathcal{F}(af + bg) = a\mathcal{F}(f) + b\mathcal{F}(g)$$

Example: Fourier Transform

- (i) $\int_{\mathbb{R}} e^{-ax^2} \, dx$
- (ii) $\int_{\mathbb{R}} x e^{-ax^2} \, dx$
- (iii) $\int_{\mathbb{R}} x^2 e^{-ax^2} \, dx$

$$\text{Hint: } \mathcal{F}[e^{-ax^2}](w) = \frac{1}{\sqrt{2a}} e^{-\frac{w^2}{4a}}$$

$$\mathcal{F}[x^k f(x)] = i^k \frac{d^k}{dw^k} \mathcal{F}[f(x)](w)$$

Solution:

$$(i) \int_{\mathbb{R}} e^{-ax^2} \, dx = \sqrt{2\pi} \mathcal{F}[e^{-ax^2}](0) = \sqrt{\frac{\pi}{a}}$$

$$(ii) \int_{\mathbb{R}} x e^{-ax^2} \, dx = \sqrt{2\pi} i \frac{d}{dw} \mathcal{F}[e^{-ax^2}](0) = 0$$

$$(iii) \int_{\mathbb{R}} x^2 e^{-ax^2} \, dx = \sqrt{2\pi} i^2 \frac{d^2}{dw^2} \mathcal{F}[e^{-ax^2}](0)$$

$$\begin{aligned} &= -\frac{d^2}{dw^2} \sqrt{\frac{\pi}{a}} e^{-\frac{w^2}{4a}} \Big|_{w=0} \\ &= \left(\frac{d}{dw} \frac{w}{2a} \right) \sqrt{\frac{\pi}{a}} e^{-\frac{w^2}{4a}} \Big|_{w=0} \\ &= \sqrt{\frac{\pi}{4a^3}} \end{aligned}$$

2.8.1 Transforms of Derivatives:

- $\mathcal{F}\{f^{(n)}(x)\} = (iw)^n \mathcal{F}\{f(x)\}$
- $\mathcal{F}_c\{f'(x)\} = w \mathcal{F}_s\{f(x)\} - \sqrt{\frac{2}{\pi}} f(0)$
- $\mathcal{F}_c\{f''(x)\} = -w^2 \mathcal{F}_c\{f(x)\} - \sqrt{\frac{2}{\pi}} f'(0)$
- $\mathcal{F}_s\{f'(x)\} = -w \mathcal{F}_c\{f(x)\}$
- $\mathcal{F}_s\{f''(x)\} = -w^2 \mathcal{F}_s\{f(x)\} + \sqrt{\frac{2}{\pi}} w f(0)$

Theorem: Fourier and ramp function

Let $f: \mathbb{R} \rightarrow \mathbb{C}$ differentiable and such that:

- f is absolutely integrable
- xf is absolutely integrable.

So is:

$$(\hat{f})' = -i(x\hat{f})$$

2.8.2 Other properties:

- $\mathcal{F}(f(\lambda x))(\omega) = \begin{cases} \frac{1}{\lambda} \mathcal{F}(f(x))(\frac{\omega}{\lambda}), \lambda > 0 \\ -\frac{1}{\lambda} \mathcal{F}(f(x))(\frac{\omega}{\lambda}), \lambda < 0 \end{cases}$
- $\mathcal{F}(f(x-a))(\omega) = e^{-i\omega a} \mathcal{F}(f(x))(\omega)$
- $\mathcal{F}(x^k \cdot f(x))(w) = i^k \frac{d^k}{dw^k} \mathcal{F}(f(x))(w)$

Theorem: Convolution

Let be f and g piecewise continuous, bounded and absolutely integrable, then $\mathcal{F}(f * g) = \sqrt{2\pi} \mathcal{F}(f) \mathcal{F}(g)$

Theorem: Bounded conditions

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be any function:

- if f is periodic and continuous, then it is bounded.
- if f is differentiable and periodic of period P , then also f' is periodic with the same period

3 Partial Differential Equation (PDE)

3.1 Basic Concepts:

Classification

- A PDE is **linear** if it is of first degree in the unknown function u and its partial derivatives
An equation of the form
 $a_1(x, y)u_{xx} + a_2(x, y)u_{xy} + a_3(x, y)u_{yx} + a_4(x, y)u_{yy} + a_5(x, y)u_x + a_6(x, y)u_y + a_7(x, y)u = f(x, y)$ is linear (a_n must not contain u or its derivatives)
- A *linear* PDE is **homogeneous** if each of its terms contains either u or one of its partial derivatives (See linear, the term $f(x, y)$ must be $= 0$)
- The order of the highest derivative is called the **order** of the PDE

Theorem: Superposition

If u_1, u_2 are solutions of a *homogeneous linear* PDE in some region R , then $u = c_1u_1 + c_2u_2$ with any constant c_1, c_2 is also a solution of that PDE in R .

3.1.1 Simple Approaches:

• Substitution

$$u_{xy} = -u_x \quad \text{setting } u_x = p \\ \Rightarrow p_y = -p \Leftrightarrow p_y/p = -1 \Leftrightarrow \ln |p| = -y + c(x) \Leftrightarrow p = \tilde{c}(x)e^{-y} \Leftrightarrow u(x, y) = f(x)e^{-y} + g(y)$$

• Like an ODE

$$u_{xx} - u = 0 \quad (\text{no } y\text{-derivatives occur}) \\ \Rightarrow u'' - u = 0 \Rightarrow u = A(y)e^x + B(y)e^{-x}$$

Example: PDE

$$\begin{cases} t^2 u_x - u_t = 0 \\ u(x, 0) = 3 \cos x \end{cases}$$

Solution:

- Take Fourier Transform
 $\mathcal{F}(t^2 u_x) = t^2 i w \hat{u}(w)$
 $\mathcal{F}(u_t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u_t e^{-iwx} dx = \frac{1}{\sqrt{2\pi}} \frac{\partial}{\partial t} \int_{-\infty}^{\infty} u e^{-iwx} dx = \frac{\partial \hat{u}}{\partial t}$
- Separation of variables
 $\frac{\partial \hat{u}}{\partial t} = t^2 i w \hat{u} \Leftrightarrow \frac{1}{\hat{u}} \partial \hat{u} = i w t^2 \partial t$
 $\Leftrightarrow \ln \hat{u} = C(w) \frac{1}{3} t^2 i w \Leftrightarrow \hat{u} = C(w) e^{\frac{1}{3} t^2 i w}$
- Initial Condition
 $\hat{u}(w, 0) = C(w) = \hat{f}(w) = \mathcal{F}(3 \cos x)$

Example: PDE

$$\begin{cases} u_t - u_x x = 0 & (x, t) \in \mathbb{R} \times \mathbb{R}_+ \\ u(x, 0) = x^2 & x \in \mathbb{R} \end{cases}$$

$$\Rightarrow u(x, t) = \int_{\mathbb{R}} f(y) K(x - y, t) dy$$

$$K(x, t) = \frac{1}{\sqrt{4\pi t}} e^{-\frac{x^2}{4t}}$$

Solution: Ansatz: $z = x - y$

$$\begin{aligned} u(x, t) &= \int_{\mathbb{R}} (x - z)^2 K(z, t) dz \\ &= \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} (x^2 - 2xz + z^2) e^{-\frac{z^2}{4t}} dz \\ &= \frac{1}{4\pi t} \left(x^2 \sqrt{\frac{\pi}{t}} + 0 + \sqrt{\frac{\pi}{4t^3}} \right) \Big|_{a=\frac{1}{4t}} \\ &= (x^2 + 2t) \end{aligned}$$

3.2 First order PDE:

3.2.1 Method of Characteristics:

A PDE of the form:

$$A(x, y) \frac{\partial u}{\partial x} + B(x, y) \frac{\partial u}{\partial y} + C_1(x, y)u = C_0(x, y)$$

is called a (first order) linear PDE (in two variables). It is called homogeneous if $C_0 = 0$. More generally, a PDE of the form:

$$A(x, y, u) \frac{\partial u}{\partial x} + B(x, y, u) \frac{\partial u}{\partial y} = C(x, y, u)$$

with the initial condition $u(x_0, y_0) = f(x, y) := z_0$, will be called a (first order) quasi-linear PDE (in two variables).

Remark: Every linear PDE is also quasi-linear since we may set: $C(x, y, u) = C_0(x, y) - C_1(x, y)u$

1. **Parametrize the initial curve Γ :** write:

$$\Gamma := \begin{cases} x = x_0(a), \\ y = y_0(a), \\ z = z_0(a) \end{cases}$$

2. **Solve ODE System:** For each a , find the stream line of F that passes through $\Gamma(a)$. That is, solve the system of ODE with initial value problems:

$$\begin{cases} \frac{dx}{ds} = A(x, y, z), x(0) = x_0(a) \\ \frac{dy}{ds} = B(x, y, z), y(0) = y_0(a) \\ \frac{dz}{ds} = C(x, y, z), z(0) = z_0(a) \end{cases}$$

These are the *Characteristic equations of the PDE*. The Solution to the system will be in terms of the parameters a and s :

$$x = X(a, s), y = Y(a, s), z = Z(a, s)$$

This is a parametric expression for the graph of the solution surface $z = u(x, y)$ (in terms of the variables a, s).

3. **Find a, s :** Solve the previous equations for a, s in terms of x, y :

$$a = \Lambda(x, y), s = S(x, y)$$

4. **Substitute the results** of Step 3 into $z = Z(a, s)$ to get the solution to the PDE: $u(x, y) = Z(\Lambda(x, y), S(x, y))$.

Example: First Order PDE

Find the solution to $x\frac{\partial u}{\partial x} - 2y\frac{\partial u}{\partial y} = u^2$ that satisfies $u(x, x) = x^3$.

This is a quasi-linear PDE with:

$$A(x, y, u) = x, B(x, y, u) = -2y, C(x, y, u) = u^2,$$

so we may apply the method of characteristics. The initial curve Γ can be parametrized as:

$$x = a, y = a, z = a^3.$$

Hence the characteristic ODEs are:

$$\begin{cases} \frac{dx}{ds} = x, x(0) = a \\ \frac{dy}{ds} = -2y, y(0) = a \\ \frac{dz}{ds} = z^2, z(0) = a^3 \end{cases}$$

We find immediately that: $x(s) = ae^s$ and $y(s) = ae^{-2s}$ and $z(s) = \frac{a^3}{1-sa^3}$. We now need to solve for a and s . We have $\frac{x}{a} = e^s$ so that:


$$y = a(e^s)^{-2} = a\left(\frac{x}{a}\right)^{-2} = \dots \Rightarrow a = x^{2/3}y^{1/3}$$
$$e^s = \frac{x}{a} = x^{1/3}y^{-1/3} = \dots \Rightarrow s = \frac{1}{3}\ln\left(\frac{x}{y}\right)$$

Substituting is z we get:

$$u(x, y) = z(s) = \frac{x^2y}{1 - \frac{1}{3}x^2y\ln\left(\frac{x}{y}\right)}.$$

3.3 Second order PDE:

3.3.1 Method of Characteristics:


$$A(x, y)u_{xx} + 2B(x, y)u_{xy} + C(x, y)u_{yy} = F(x, y, u, u_x, u_x)$$

Type	Condition	Normal Form
Hyperbolic	$AC - B^2 < 0$	$u_{vw} = F(v, w, u, u_v, u_w)$
Parabolic	$AC - B^2 = 0$	$u_{vv} = F(v, w, u, u_v, u_w)$
Elliptic	$AC - B^2 > 0$	$u_{vv} + u_{ww} = F(\dots)$

Characteristic equation:

$$A(x, y)(y')^2 - 2B(x, y)y' + C(x, y) = 0 \text{ with } y' = \frac{dy}{dx} \xrightarrow{\text{sol'n}}$$
$$y' = \frac{2B \pm \sqrt{(2B)^2 - 4AC}}{2A}$$

$\xi(x, y) = c_1 = \text{const}, \zeta(x, y) = c_2 = \text{const}$
(Solve for $y =$ and then for $c_1 =, c_2 =$)

Type	New Variables	
Hyperbolic	$v = \xi$	$w = \zeta$
Parabolic	$v = x$	$w = \xi = \zeta$
Elliptic	$v = \frac{\xi + \zeta}{2}$	$w = \frac{\xi - \zeta}{2i}$

Theorem: Kettenregel/chain rule

$$\frac{df}{dx} = \frac{df}{dg} \cdot \frac{dg}{dx}$$
$$E(t) = f(g(t)) \rightarrow E'(t) = f'(g(t)) \cdot g'(t) \leftrightarrow \frac{dE}{dt} = \frac{dE}{dv} \cdot \frac{dv}{dt}$$

Example: Normal form of a PDE

Solve: $u_{xx} + 2u_{xy} - 3u_{yy} = e^{x+2y}$

- $u_{xx} + 2u_{xy} - 3u_{yy} = e^{x+2y} \leftrightarrow Au_{xx} + 2Bu_{xy} + Cu_{yy} = F(x, y) \rightarrow A = 1, B = 1, C = -3$
- $AC - B^2 < 0$ PDE is hyperbolic
- Char eq: $A(y')^2 - 2By' + C = 0 \leftrightarrow (y' - 3)(y' + 1) = 0$
- Solutions: $\begin{cases} y' = 3 \leftrightarrow y - 3x = c_1 \\ y' = -1 \leftrightarrow y + x = c_2 \end{cases}$
- $\begin{cases} v = \xi(x, y) = y - 3x \\ w = \zeta(x, y) = y + x \end{cases} \rightarrow \begin{cases} v_x = -3, v_y = 1 \\ w_x = 1, w_y = 1 \end{cases}$
- Partial derivatives:
 - $u_x = u_v v_x + u_w w_x = -3u_v + u_w$
 - $u_y = u_v v_y + u_w w_y = u_v + u_w$
 - $u_{xx} = -3u_{vv} v_x - 3u_{vw} w_x + u_{ww} v_x + u_{ww} w_x = 9u_{vv} - 6u_{vw} + u_{ww}$
 - $u_{xy} = -3u_{vv} v_y - 3u_{vw} w_y + u_{ww} v_y + u_{ww} w_y = -3u_{vv} - 2u_{vw} + u_{ww}$
 - $u_{yy} = u_{vv} v_y + u_{vw} w_y + u_{vw} v_y + u_{ww} w_y = u_{vv} + 2u_{vw} + u_{ww}$
- Transform left side (new coord): $u_{xx} + 2u_{xy} - 3u_{yy} = -16u_{vw}$
- Transform right side (new coord): $x + 2y = \frac{v+7w}{4} \leftrightarrow e^{x+2y} = e^{\frac{v+7w}{4}}$
- Togheter: $\frac{1}{16}e^{\frac{v+7w}{4}} = u_{vw}$
- Integrate and substitute original variables: $u(x, y) = -\frac{1}{7}e^{x+2y} + \varphi(y - 3x) + \psi(y + x)$

4 Wave Equation

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \text{ with } c^2 = \frac{T}{\rho}$$

- Boundary conditions:** $u(0, t) = 0, u(L, t) = 0, \forall t \geq 0$
- Initial conditions:** $u(x, 0) = f(x), u_t(x, 0) = g(x), x \in [0, L]$

1. Separating Variables

Setting $u(x, t) = F(x)G(t)$

$$\Rightarrow \frac{\partial^2 u}{\partial t^2} = F\ddot{G}, \quad \frac{\partial^2 u}{\partial x^2} = F''G \Rightarrow \frac{\ddot{G}}{c^2 G} = \frac{F''}{F} = k = \text{const}$$


$$\Rightarrow F'' - kF = 0, \quad \ddot{G} - c^2 k G = 0$$

2. Boundary Conditions

- $\mathbf{k} = \mathbf{0}$: $F = ax + b \Rightarrow a = b = 0 \Rightarrow F \equiv 0$
- $\mathbf{k} = \mu^2 > \mathbf{0}$: $F = Ae^{\mu x} + Be^{-\mu x} \Rightarrow A = -B = 0 \Rightarrow F \equiv 0$
- $\mathbf{k} = -\mathbf{p}^2 < \mathbf{0}$: $F(x) = A \cos px + B \sin px$
 $\xrightarrow{\text{B.C.}} F(0) = A = 0, \quad F(L) = B \sin pL = 0$
 $\Rightarrow p = \frac{n\pi}{L}, \quad n \in \mathbb{N} \Rightarrow F_n(x) = B_n \sin \frac{n\pi}{L}x$

- $\ddot{G} + \lambda_n^2 G = 0, \quad \lambda_n = cp = \frac{cn\pi}{L}$
 $\Rightarrow G_n(t) = B_n \cos \lambda_n t + B_n^* \sin \lambda_n t$

$$\Rightarrow u_n(x, t) = (B_n \cos \lambda_n t + B_n^* \sin \lambda_n t) \sin \frac{n\pi}{L}x$$

These functions are called **eigenfunctions** and
 $\lambda_n = \frac{cn\pi}{L}$ are called **eigenvalues**. The set $\{\lambda_1, \lambda_2, \dots\}$ is called **spectrum**.


3. Fourier Series

$$u(x, t) = \sum_{n=1}^{\infty} u_n(x, t) =$$

$$= \sum_{n=1}^{\infty} (B_n \cos \lambda_n t + B_n^* \sin \lambda_n t) \sin \left(\frac{n\pi}{L}x \right)$$

• Displacement condition

$$u(x, 0) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi}{L}x = f(x) \quad (\text{Fourier})$$

$$\Rightarrow B_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi x}{L} dx$$

• Velocity condition

$$\frac{\partial u}{\partial t} \Big|_{t=0} = \sum_{n=1}^{\infty} B_n^* \lambda_n \sin \frac{n\pi x}{L} = g(x)$$

$$\Rightarrow B_n^* \lambda_n = \frac{2}{L} \int_0^L g(x) \sin \frac{n\pi x}{L} dx$$

$$\Leftrightarrow B_n^* = \frac{2}{cn\pi} \int_0^L g(x) \sin \frac{n\pi x}{L} dx$$

$$\text{If } g(x) = 0 \Rightarrow u(x, t) = \sum_{n=1}^{\infty} B_n \cos \lambda_n t \sin \frac{n\pi x}{L}$$

$$\Leftrightarrow u(x, t) = \frac{1}{2} \sum_{n=1}^{\infty} B_n \sin \left\{ \frac{n\pi}{L}(x - ct) \right\} + \frac{1}{2} \sum_{n=1}^{\infty} B_n \sin \left\{ \frac{n\pi}{L}(x + ct) \right\}$$

4.1 D'Alembert Solution of the Wave Equation:

• Solve characteristic eqn.

$$u_{tt} = c^2 u_{xx} \Leftrightarrow u_{tt} - c^2 u_{xx} = 0$$

$$\Rightarrow A = -c^2, \quad B = 0, \quad C = 1 \Rightarrow \text{hyperbolic}$$

$$\text{char.eqn.: } -c^2(y')^2 + 1 = 0 \Leftrightarrow 1 \pm cy' = 0$$

$$\Rightarrow v = \xi = x + ct = c_1, \quad w = \zeta = x - ct = c_2$$

• Bring into normal form

$$u_x = u_v v_x + u_w w_x = u_v + u_w$$

$$u_{xx} = (u_{vv} v_x + u_{vw} w_x) + (u_{wv} v_x + u_{ww} w_x)$$

$$= u_{vv} + 2u_{vw} + u_{ww}$$

$$u_t = u_v v_t + u_w w_t = cu_v - cu_w$$

$$u_{tt} = c(u_{vv} v_t + u_{vw} w_t) - c(u_{wv} v_t + u_{ww} w_t)$$

$$= c^2(u_{vv} - 2u_{vw} + u_{ww})$$

$$c^2(u_{vv} - 2u_{vw} + u_{ww}) - c^2(u_{vv} + 2u_{vw} + u_{ww}) = -4c^2 u_{vw} = 0$$

$$\Rightarrow u_{vw} = 0 \Rightarrow u(v, w) = \phi(v) + \psi(w) \Leftrightarrow u(x, t) = \phi(x + ct) + \psi(x - ct)$$

• Solve for initial conditions

$$\text{I.C.: } u(x, 0) = f(x) \quad u_t(x, 0) = g(x), \quad x \in \mathbb{R}, t \geq 0$$

$$f(x) = \phi(x) + \psi(x) \quad g(x) = c\phi'(x) - c\psi'(x)$$

$$\phi(x) + \psi(x) = f(x)$$

$$\phi(x) - \psi(x) = \frac{1}{c} \int_0^x g(s) ds + \underbrace{\phi(0) - \psi(0)}_{k_0}$$

$$\phi(x) = \frac{1}{2} f(x) + \frac{1}{2c} \int_0^x g(s) ds + \frac{1}{2} k_0$$

$$\psi(x) = \frac{1}{2} f(x) - \frac{1}{2c} \int_0^x g(s) ds - \frac{1}{2} k_0$$

by replacing x with $x \pm ct$ and adding the two sol'ns:


$$u(x, t) = \frac{1}{2} [f(x + ct) + f(x - ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds$$

4.2 Graphic solution (Characteristic lines):


- What kind of information has an influence on the solution u at the point (x, t) ?

Through any point (x_0, t_0) with $t_0 > 0$, there are exactly two characteristics, namely:

$$x - ct = x_0 - ct_0 \quad \text{and} \quad x + ct = x_0 + ct_0.$$

These are straight lines whose intersections with the x -axis are respectively the points $(x_0 - ct_0, 0)$ and $(x_0 + ct_0, 0)$. The triangle with vertices $(x_0 - ct_0, 0)$, $(x_0 + ct_0, 0)$ and (x_0, t_0) is called **characteristic triangle**.The interval $[x_0 - ct_0, x_0 + ct_0]$ is called the **domain of dependence** of u at (x_0, t_0) . Changing f or g outside this domain of dependence will not affect the value $u(x_0, t_0)$.

- What region of the (x, t) in the upper half plane is affected by the initial data on an interval $[a, b]$?

The endpoints of the interval define four characteristics: $x \pm ct = a$ and $x \pm ct = b$, whose intersections define six regions indicated in the picture with I, II, III, IV, V and VI.The points that are affected by the initial conditions are exactly the points (x, t) whose domain of dependence $[x - ct, x + ct]$ intersects the interval $[a, b]$ in a non-trivial way. In particular, as shown in the picture, any


point in the regions I and III are such that $u(x, t) = 0$.
For values of $t = \infty$ the point is at rest in a position

$$u(x, t) = \frac{1}{2c} \int_a^b g(s) ds.$$

Example: Region of influence

Let $u(x, t)$ be the solution of the problem ($x \in \mathbb{R}, t > 0$):

$$\begin{cases} u_{tt} = u_{xx} \\ u(x, 0) = f(x) = \begin{cases} 1, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases} \\ u_t(x, 0) = \begin{cases} 1, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases} \end{cases}.$$



- If $(x_0, t_0) \in \text{I}$: $u(x_0, t_0) = 0$
- if $(x_0, t_0) \in \text{II}$: $u(x_0, t_0) = \frac{f(x_0+t_0)+f(x_0-t_0)}{2} + \frac{1}{2} \int_{x_0-t_0}^{x_0+t_0} f(y) dy = 1 + \frac{1}{2}(x_0+t_0 - x_0+t_0) = 1+t_0$
- If $(x_0, t_0) \in \text{III}$: $u(x_0, t_0) = 0$
- If $(x_0, t_0) \in \text{IV}$: $u(x_0, t_0) = \frac{f(x_0+t_0)}{2} + \frac{1}{2} \int_{-1}^{x_0+t_0} g(y) dy = \frac{1}{2} + \frac{x_0+t_0+1}{2} = \frac{x_0+t_0}{2} + 1$
- If $(x_0, t_0) \in \text{V}$: $u(x_0, t_0) = \frac{1}{2} \int_{x_0-t_0}^{x_0+t_0} g(y) dy = \frac{1}{2} \int_{-1}^1 g(y) dy = 1$
- If $(x_0, t_0) \in \text{VI}$: $u(x_0, t_0) = \frac{f(x_0-t_0)}{2} + \frac{1}{2} \int_{x_0-t_0}^1 g(y) dy = \frac{1}{2} + \frac{-x_0+t_0+1}{2} = \frac{-x_0+t_0}{2} + 1$

Observe that the Max value of $u(x, t)$ is obtained at the point $(x_0, t_0) \in \text{II}$ satisfying: $\begin{cases} x_0 - t_0 = -1 \\ x_0 + t_0 = 1 \end{cases} \Rightarrow x_0 = 0 \text{ and } t_0 = 1. \quad \forall x \in \mathbb{R} : \lim_{t \rightarrow \infty} u(x, t) = \frac{1}{2} \int_{-1}^1 g(y) dy = \frac{1}{2} \cdot 2 = 1.$

5 Heat Equation

$$\frac{\partial u}{\partial t} = c^2 \nabla^2 u \quad \text{with } c^2 = \frac{K}{\rho\sigma},$$

$$\text{where } \nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$$

(K = conductivity, ρ = Density, σ = Spec. heat, c = Thermal Diffusivity)

5.1 One-Dimensional Heat Equation:

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$$

- **Assumptions:** laterally insulated, thin, homogeneous metal bar, heat flows in x -direction only
- **Boundary conditions:** $u(0, t) = 0, u(L, t) = 0, \forall t \geq 0$
- **Initial condition:** $u(x, 0) = f(x)$

1. Separating variables

Substituting $u(x, t) = F(x)G(t)$

$$\Rightarrow F\dot{G} = c^2 F''G \Leftrightarrow \frac{\dot{G}}{c^2 G} = \frac{F''}{F} = -p^2 = \text{const}$$

$$\Rightarrow F'' + p^2 F = 0, \quad \dot{G} + c^2 p^2 G = 0$$

Note: Look at the wave equation to see why it is $-p^2$!

2. Boundary Condition

$$\bullet F(x) = A \cos px + B \sin px \xrightarrow{B.C.} A = 0, p = \frac{n\pi}{L}, n \in \mathbb{N}$$

$$\text{Setting } B = 1 \Rightarrow F_n(x) = \sin \frac{n\pi x}{L}$$

$$\bullet \dot{G} + \lambda_n^2 G = 0, \lambda_n = \frac{cn\pi}{L} \Rightarrow G_n(t) = B_n e^{-\lambda_n^2 t}$$

$$\Rightarrow u_n(x, t) = F_n(x)G_n(t) = B_n \sin \frac{n\pi x}{L} e^{-\lambda_n^2 t}$$

3. Fourier Series

$$\bullet u(x, 0) = \sum_{n=1}^{\infty} u_n(x, 0) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi x}{L} = f(x)$$

$$\bullet \Rightarrow B_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n\pi x}{L} dx$$

$$\bullet u(x, t) = \sum_{n=1}^{\infty} u_n(x, t) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi x}{L} e^{-\lambda_n^2 t}$$

Example: Heat equation on a bar

Find $u(x, t)$ in a laterally insulated bar of length 80cm with initial temperature $f(x) = 100 \sin(\pi x/80)^\circ\text{C}$.

$$u(x, 0) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi x}{80} = f(x) = 100 \sin \frac{\pi x}{80} \quad \text{By inspection: } B_1 = 100, B_2 = B_3 = \dots = 0$$

$$\Rightarrow u(x, t) = 100 \sin \frac{\pi x}{80} e^{-\lambda^2 t}$$

Example: Isolated bar

Find $u(x, t)$, bar with insulated ends.

$$u_x(0, t) = 0 \quad u_x(L, t) = 0$$

$$u(x, t) = F(x)G(t) \Rightarrow u_x(0, t) = F'(0)G(t) = 0, u_x(L, t) = F'(L)G(t) = 0$$

$$F' = -Ap \sin px + Bp \cos px \Rightarrow F'(0) = Bp = 0, F'(L) = -Ap \sin pL = 0$$

$$\Rightarrow p_n = \frac{n\pi}{L} \Rightarrow u_n(x, t) = F_n(x)G_n(t) = A_n \cos \frac{n\pi x}{L} e^{-\lambda_n^2 t}$$

$$\Rightarrow u(x, t) = \sum_{n=0}^{\infty} A_n \cos \frac{n\pi x}{L} e^{-\lambda_n^2 t}$$

$$\text{with } A_0 = \frac{1}{L} \int_0^L f(x) dx, \quad A_n = \frac{2}{L} \int_0^L f(x) \cos \frac{n\pi x}{L} dx$$

Note: Here, $\lambda_0 = 0$ is an eigenvalue too.

5.2 One - Dimensional Heat Equation with inhomogeneous BC:

$$\begin{cases} u_t(x, t) = c^2 u_{xx}(x, t) \\ u(0, t) = 2 \\ u(\pi, t) = 3 \\ u(x, 0) = f(x) \end{cases}$$

Solution:

$$1. \text{ Construct a function } w(x) \text{ with } w(0) = 2, w(\pi) = 3, w'' = 0$$

$$\Rightarrow w = \frac{x}{\pi} + 2$$

$$2. \text{ State the boundary value problem for } v(x, t) := u(x, t) - w(x)$$

$$\Rightarrow \begin{cases} v_t(x, t) = c^2 v_{xx}(x, t) \\ v(0, t) = 0 \\ v(\pi, t) = 0 \\ v(x, 0) = f(x) - w(x) \end{cases}$$

3. Solve by separation of variables.

5.3 Inhomogeneous One Dimensional Heat Equation:

$$\begin{cases} u_t = a^2 u_{xx} + B & 0 < x < L, t > 0 \\ u(0, t) = u(L, t) & t \geq 0 \\ u(x, 0) = \sin(\frac{\pi x}{L}) & 0 \leq x \leq L \end{cases}$$

Solution:

- Find the stationary solution:
 $v : x \rightarrow v(x)$ which fulfills the boundary conditions.

$$\begin{cases} a^2 v_{xx} + b = 0 \\ v(0) = v(L) = 0 \end{cases}, x \in \mathbb{R}$$

The unique solution to this PDE is $v(x) = -\frac{b}{2a^2}(x-L)x$ (consider zeros in 0 and L)

- Construct a function $w(x)$ with $w(0) = 2$, $w(\pi) = 3$, $w'' = 0$
 $\Rightarrow w = \frac{x}{\pi} + 2$

- Set $w(x, t) = u(x, t) - v(x)$

$$\Rightarrow \begin{cases} w_t = u_t = a^2(w_{xx} + v_{xx}) + b = a^2 w_{xx} \\ w(0, t) = w(L, t) = 0 \\ w(x, 0) = \sin(\frac{\pi x}{L}) - v(x) \end{cases}$$

- $\dots \Rightarrow w(x, t) = \sum_{n \leq 1} \alpha_n \sin(\frac{n\pi x}{L}) = \sin(\frac{\pi x}{L}) - v(x)$. We extend v to an odd $2L$ periodic function \tilde{v} and determine its Fourier coefficients:

$$B_n := \frac{1}{L} \int_{-L}^L v(\tilde{x}) \sin(\frac{n\pi x}{L}) dx.$$

- By comparing we get $\alpha_n = -B_n$ and for $n = 1 \Rightarrow \alpha_1 = 1 - B_1$.

- $u(x, t) = w(x, t) + v(x)$

5.4 Heat Equation for Infinite Bars:

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$$

1. Separating variables

$$\begin{aligned} u(x, t) = F(x)G(t) &\Rightarrow \begin{cases} F(x) = A \cos px + b \sin px \\ G(t) = e^{-c^2 p^2 t} \end{cases} \\ \Rightarrow u(x, t; p) &= FG = (A \cos px + B \sin px) e^{-c^2 p^2 t} \end{aligned}$$

2. Fourier Integrals

A, B are arbitrary, regard them as $A = A(p), B = B(p)$

$$\begin{aligned} u(x, t) &= \int_0^\infty u(x, t; p) dp \\ &= \int_0^\infty [A(p) \cos px + B(p) \sin px] e^{-c^2 p^2 t} dp \end{aligned}$$

3. $A(p), B(p)$ from Initial Condition

$$\begin{aligned} u(x, 0) &= \int_0^\infty [A(p) \cos px + B(p) \sin px] dp = f(x) \\ A(p) &= \frac{1}{\pi} \int_{-\infty}^\infty f(v) \cos pv dv, B(p) = \frac{1}{\pi} \int_{-\infty}^\infty f(v) \sin pv dv \quad \text{us-} \end{aligned}$$

$$\sin x \sin y = \frac{1}{2} (\cos(x-y) - \cos(x+y))$$

ing:

$$\cos x \cos y = \frac{1}{2} (\cos(x-y) + \cos(x+y))$$

$$\Rightarrow u(x, 0) = \frac{1}{\pi} \int_0^\infty \left[\int_{-\infty}^\infty f(v) \cos(px - pv) dv \right] dp$$

$$\Rightarrow u(x, t) = \frac{1}{\pi} \int_{-\infty}^\infty f(v) \left[\int_0^\infty e^{-c^2 p^2 t} \cos(px - pv) dp \right] dv$$

$$\int_0^\infty e^{-s^2} \cos 2bs ds = \frac{\sqrt{\pi}}{2} e^{-b^2} \quad \text{with } p = \frac{s}{c\sqrt{t}}, \quad b = \frac{x-v}{2c\sqrt{t}}$$

$$\Rightarrow \int_0^\infty e^{-c^2 p^2 t} \cos(px - pv) dp = \frac{\sqrt{\pi}}{2c\sqrt{t}} \exp\left(-\frac{(x-v)^2}{4c^2 t}\right)$$

$$\Rightarrow u(x, t) = \frac{1}{2c\sqrt{\pi t}} \int_{-\infty}^\infty f(v) \exp\left(-\frac{(x-v)^2}{4c^2 t}\right) dv$$

$$\text{with } z = \frac{v-x}{2c\sqrt{t}} \Rightarrow$$

$$u(x, t) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^\infty f(x + 2cz\sqrt{t}) e^{-z^2} dz$$

5.4.1 Using Fourier Transform:

1. Take Fourier Transform

$$\hat{u} = \mathcal{F}(u)$$

$$\bullet \quad c^2 \mathcal{F}(u_{xx}) = c^2 (-w^2) \mathcal{F}(u) = -c^2 w^2 \hat{u}$$

$$\begin{aligned} \mathcal{F}(u_t) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty u_t e^{-iwx} dx \\ \bullet \quad &= \frac{1}{\sqrt{2\pi}} \frac{\partial}{\partial t} \int_{-\infty}^\infty u e^{-iwx} dx = \frac{\partial \hat{u}}{\partial t} \end{aligned}$$

$$\Rightarrow \frac{\partial \hat{u}}{\partial t} = -c^2 w^2 \hat{u} \Rightarrow \hat{u}(w, t) = C(w) e^{-c^2 w^2 t}$$

2. Initial Condition

$$\hat{u}(w, 0) = \hat{f}(w) = C(w) \Rightarrow \hat{u}(w, t) = \hat{f}(w) e^{-c^2 w^2 t}$$

$$\hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty f(v) e^{-i w v} dv$$

$$u(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \hat{f}(w) e^{-c^2 w^2 t} e^{i w x} dw$$

$$\begin{aligned} \Rightarrow u(x, t) &= \frac{1}{2\pi} \int_{-\infty}^\infty f(v) \left[\int_{-\infty}^\infty e^{-c^2 w^2 t} e^{i(w x - w v)} dw \right] dv \\ e^{i(w x - w v)} &= \cos(wx - wv) + i \sin(wx - wv) \\ \rightarrow \text{the imaginary part is odd so its integral is 0} \end{aligned}$$

$$u(x, t) = \frac{1}{\pi} \int_{-\infty}^\infty f(v) \left[\int_{-\infty}^\infty e^{-c^2 w^2 t} \cos(wx - wv) dw \right] dv$$

5.4.2 Method of Convolution:

$$u(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \hat{f}(w) e^{-c^2 w^2 t} e^{i w x} dw$$

$$u(x, t) = (f * g)(x) = \int_{-\infty}^\infty \hat{f}(x) \hat{g}(w) e^{i w x} dw$$

$$\Rightarrow \hat{g}(w) = \frac{1}{\sqrt{2\pi}} e^{-c^2 w^2 t}$$

$$\text{Def. of convolution: } (f * g)(x) = \int_{-\infty}^\infty f(p) g(x-p) dp$$

$$\text{Inverse transform of } \hat{g}: \mathcal{F}\left(e^{-a x^2}\right) = \frac{1}{\sqrt{2a}} e^{-w^2/(4a)}$$

$$\begin{aligned} \text{with } a = 1/(ac^2 t) &\Rightarrow \mathcal{F}\left(e^{-x/(4ac^2 t)}\right) = \sqrt{2c^2 t} e^{-c^2 w^2 t} \\ &= \sqrt{2c^2 t} \sqrt{2\pi} \hat{g}(w) \end{aligned}$$

$$\Rightarrow \mathcal{F}^{-1}(\hat{g}) = \frac{1}{\sqrt{2c^2 t} \sqrt{2\pi}} e^{-x^2/(2c^2 t)}$$

$$u(x, t) = (f * g)(x) = \frac{1}{2c\sqrt{\pi t}} \int_{-\infty}^\infty f(p) \exp\left(-\frac{(x-p)^2}{4c^2 t}\right) dp$$

6 Laplace eq. (Steady 2D Heat Problem)

steady = time-independent $\Rightarrow \frac{\partial u}{\partial t} = 0$

$$\Delta u = \nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \text{ Laplace's equation}$$

Theorem: Liouville's

Suppose $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is harmonic and bounded. Then u is constant.

Theorem: Weak Minimum Principle

Suppose $u_{xx} + u_{yy} = 0$, $B(0, R)$ and u is continuous in $\bar{B}(0, R) = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq R^2\}$. Then the Maximum and the Minimum values of u are obtained on $\partial \bar{B} = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = R^2\}$.

$$\max u(x, y) = \max f(\theta)$$


This principle holds also for a square bound, i.e Section 6.1:

$$\Rightarrow \max u(x, y) = \max f(x)$$

Theorem: Strong Minimum Principle

Let $B \subseteq \mathbb{R}^2$ be a domain (can also not be bounded) and $u : B \rightarrow \mathbb{R}$ be a harmonic function (function that solves the Laplace equation). If u obtains its maximum or minimum in B , then u is constant.

6.1 Dirichlet problem:



Since the equation is linear we can break the problem into simpler problems which do have sufficient homogeneous BC and use superposition to obtain the solution

$$u(x, y) = u_A + u_B + u_C + u_D$$

6.1.1 Solution to problem with $f_1(x)$ (u_A):

1. **Assume** $f_2(x) = g_1(y) = g_2(y) = 0$

2. **Separating variables**

$$\begin{aligned} u_{xx} &= -u_{yy}, \quad u(x, y) = F(x)G(y) \\ \Rightarrow \frac{1}{F} \frac{d^2 F}{dx^2} &= -\frac{1}{G} \frac{d^2 G}{dy^2} = -k^2 \\ \Rightarrow \frac{d^2 F}{dx^2} + k^2 F &= 0, \quad \frac{d^2 G}{dy^2} - k^2 G = 0 \end{aligned}$$

Note: Look at the wave equation to see why it is $-k^2$

3. **Boundary conditions**

- $F(x) = A \sin kx + B \cos kx$
left/right B.C.: $F(0) = 0, F(a) = 0$
 $\Rightarrow A = 1, B = 0, k = \frac{n\pi}{a}$
 $\Rightarrow F(x) = F_n(x) = \sin \frac{n\pi x}{a}$
- $G(y) = C \cosh(ky) + D \sinh(ky)$
upper B.C.: $G_n(b) = C \cosh(kb) + D \sinh(kb) = 0 \Rightarrow C = -D \tanh(ka)$
 $\Rightarrow G(y) = \frac{-D \tanh(kb) \cosh(ky) + D \sinh(ky)}{\cosh(kb)} = \frac{D}{\cosh(kb)} \sinh(k(y-b)) = \tilde{D} \sinh(k(y-b))$
 $\Rightarrow G_n(y) = \tilde{D} \sinh(k(y-b))$

$$u_n(x, y) = F_n(x)G_n(y) = A_n^* \sin\left(\frac{n\pi x}{a}\right) \sinh\left(\frac{n\pi}{a}(y-b)\right)$$

4. **Fourier Series**

$$\begin{aligned} u(x, y) &= \sum_{n=1}^{\infty} u_n(x, y) \\ \text{lower BC } u_{x,0} &= f_1(x) = \sum_{n=1}^{\infty} \left(-A_n^* \sinh \frac{n\pi b}{a} \right) \sin \frac{n\pi x}{a} \\ b_n &= -A_n^* \sinh \frac{n\pi b}{a} = \frac{2}{a} \int_0^a f(x) \sin \frac{n\pi x}{a} dx \\ \Leftrightarrow A_n^* &= -\frac{2}{a \sinh(n\pi b/a)} \int_0^a f_1(x) \sin \frac{n\pi x}{a} dx \end{aligned}$$

$$u_A(x, y) = \sum_{n=1}^{\infty} A_n^* \sin\left(\frac{n\pi x}{a}\right) \sinh\left(\frac{n\pi}{a}(y-b)\right)$$

6.1.2 Solution to problem with $f_2(x)$ (u_C):

1. **Assume** $f_1(x) = g_1(y) = g_2(y) = 0$

2. **Separating variables**

$$\begin{aligned} u_{xx} &= -u_{yy}, \quad u(x, y) = F(x)G(y) \\ \Rightarrow \frac{1}{F} \frac{d^2 F}{dx^2} &= -\frac{1}{G} \frac{d^2 G}{dy^2} = -k^2 \\ \Rightarrow \frac{d^2 F}{dx^2} + k^2 F &= 0, \quad \frac{d^2 G}{dy^2} - k^2 G = 0 \end{aligned}$$

Note: Look at the wave equation to see why it is $-k^2$

3. **Boundary conditions**

- $F(x) = A \sin kx + B \cos kx$
left/right B.C.: $F(0) = 0, F(a) = 0$
 $\Rightarrow A = 1, B = 0, k = \frac{n\pi}{a}$
 $\Rightarrow F(x) = F_n(x) = \sin \frac{n\pi x}{a}$
- $G(y) = A_n \overbrace{e^{n\pi y/a}}^{ky} + B_n \overbrace{e^{-n\pi y/a}}^{-ky}$
lower B.C.: $G_n(0) = A_n + B_n = 0 \Leftrightarrow B_n = -A_n$
 $\Rightarrow G_n(y) = A_n (e^{n\pi y/a} - e^{-n\pi y/a}) = \underbrace{2A_n}_{A_n^*} \sinh \frac{n\pi y}{a}$

$$\Rightarrow u_n(x, y) = F_n(x)G_n(y) = C_n^* \sin \frac{n\pi x}{a} \sinh \frac{n\pi y}{a}$$

4. **Fourier Series**

$$\begin{aligned} u(x, y) &= \sum_{n=1}^{\infty} u_n(x, y) \\ \text{upper B.C } u_{x,b} &= f_2(x) = \sum_{n=1}^{\infty} \left(C_n^* \sinh \frac{n\pi b}{a} \right) \sin \frac{n\pi x}{a} \\ b_n &= C_n^* \sinh \frac{n\pi b}{a} = \frac{2}{a} \int_0^a f(x) \sin \frac{n\pi x}{a} dx \\ \Leftrightarrow C_n^* &= \frac{2}{a \sinh(n\pi b/a)} \int_0^a f_2(x) \sin \frac{n\pi x}{a} dx \end{aligned}$$

$$u_C(x, y) = \sum_{n=1}^{\infty} C_n^* \sin \frac{n\pi x}{a} \sinh \frac{n\pi y}{a}$$

6.1.3 Solution to problem with $g_1(x)$ (u_B):

1. **Assume** $f_1(x) = f_2(x) = g_2(y) = 0$ (for $g_2(y)$ analogous)

2. **Separating variables**

$$u_{xx} = -u_{yy}, \quad u(x, y) = F(x)G(y)$$

$$\Rightarrow \frac{1}{F} \frac{d^2 F}{dx^2} = -\frac{1}{G} \frac{d^2 G}{dy^2} = -k^2$$

$$\Rightarrow \frac{d^2 F}{dx^2} - k^2 F = 0, \quad \frac{d^2 G}{dy^2} + k^2 G = 0$$

Note: Look at the wave equation to see why it is k^2

3. Boundary conditions

- $F(x) = c_1 \cosh(kx) + c_2 \sinh(kx)$
upper/lower B.C.: $F_n(0) = A_n + B_n = 0 \Leftrightarrow B_n = -A_n$
 $\Rightarrow F_n(x) = A_n(e^{n\pi x/b} - e^{-n\pi x/b}) = \underbrace{2A_n}_{A_n^*} \sinh \frac{n\pi x}{b}$

- $G(y) = A \cos ky + B \sin ky$
left B.C.: $G(0) = 0, \quad G(b) = 0$
 $\Rightarrow A = 0, \quad B = 1, \quad k = \frac{n\pi}{b}$
 $\Rightarrow G(y) = G_n(y) = \sin \frac{n\pi y}{b}$

$$\Rightarrow u_n(x, y) = F_n(x)G_n(y) = B_n^* \sin \frac{n\pi y}{b} \sinh \frac{n\pi x}{b}$$

4. Fourier Series

$$u(x, y) = \sum_{n=1}^{\infty} u_n(x, y)$$

right B.C $u_{y,a} = g_1(x) = \sum_{n=1}^{\infty} \left(B_n^* \sinh \frac{n\pi a}{b} \right) \sin \frac{n\pi y}{b}$

$$b_n = B_n^* \sinh \frac{n\pi a}{b} = \frac{2}{b} \int_0^b f(x) \sin \frac{n\pi y}{b} dy$$

$$\Leftrightarrow B_n^* = \frac{2}{b \sinh(n\pi a/b)} \int_0^b g_1(y) \sin \frac{n\pi y}{b} dy$$

$$u_B(x, y) = \sum_{n=1}^{\infty} B_n^* \sin \frac{n\pi y}{b} \sinh \frac{n\pi x}{b}$$

6.2 Laplacian in Polar Coordinates:

Polar Coordinates:

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \arctan \frac{y}{x} \end{cases}$$

$$\begin{cases} u_x = u_r r_x + u_\theta \theta_x \\ u_{xx} = (u_{rr} r_x + u_{r\theta} \theta_x) r_x + u_r r_{xx} + (u_{\theta r} r_x + u_{\theta\theta} \theta_x) \theta_x + u_\theta \theta_{xx} \end{cases}$$

$$\begin{cases} r_x = \frac{x}{\sqrt{x^2 + y^2}} = \frac{x}{r} r_{xx} = \frac{r - x r_x}{r^2} = \frac{1}{r} - \frac{x^2}{r^3} = \frac{y^2}{r^3} \\ \theta_x = \frac{1}{1 + (y/x)^2} \left(-\frac{x}{x^2} \right) = -\frac{y}{r^2} \theta_{xx} = -y \left(-\frac{2}{r^3} \right) r_x = \frac{2xy}{r^4} \end{cases}$$

$$\Rightarrow u_{xx} = \frac{x^2}{r^2} u_{rr} - 2 \frac{xy}{r^3} u_{r\theta} + \frac{y^2}{r^4} u_{\theta\theta} + \frac{y^2}{r^3} u_r + 2 \frac{xy}{r^4} u_\theta$$

Analogously: $u_{yy} = \frac{y^2}{r^2} u_{rr} + 2 \frac{xy}{r^3} u_{r\theta} + \frac{x^2}{r^4} u_{\theta\theta} + \frac{x^2}{r^3} u_r - 2 \frac{xy}{r^4} u_\theta$

$$\Rightarrow \nabla^2 u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2}$$

6.3 Dirichlet Problem for a Disk:

$$\nabla^2 u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

- $u(R, \theta) = f(\theta)$ **Boundary Condition**
- $u(r, 0) = u(r, 2\pi), \quad u_\theta(r, 0) = u_\theta(r, 2\pi)$ **Continuity Conditions**

1. Separating Variables

Setting $u(r, \theta) = F(r)G(\theta)$

$$\Rightarrow F''G + \frac{1}{r} F'G + \frac{1}{r^2} FG'' = 0$$

$$\Leftrightarrow \frac{r^2 F'' + r F'}{F} = -\frac{G''}{G} = k = \text{const}$$

$$r^2 F'' + r F' - k F = 0 \quad G'' + k G = 0$$

2. Continuity Conditions $G(0) = G(2\pi) \quad G'(0) = G'(2\pi)$

- $k < 0$: $G(\theta) = A e^{\sqrt{-k}\theta} + B e^{-\sqrt{-k}\theta}$

$$A + B = A e^{\sqrt{-k}2\pi} + B e^{-\sqrt{-k}2\pi}$$

$$\sqrt{-k}A - \sqrt{-k}B = \sqrt{-k}A e^{k2\pi} - \sqrt{-k}B e^{k2\pi}$$

$$A + B = A e^{\sqrt{-k}2\pi} + B e^{\sqrt{-k}2\pi}$$

$$A - B = A e^{\sqrt{-k}2\pi} - B e^{\sqrt{-k}2\pi}$$

$$\Rightarrow 2A = 2A e^{\sqrt{-k}2\pi} \Rightarrow A = 0 \Rightarrow B = 0$$

- $k = 0$: $G(\theta) = A\theta + B$
 $G(0) = B = G(2\pi) = 2\pi A + B \Rightarrow A = 0$
 $\Rightarrow G(\theta) = B = \text{const}$

- $k > 0$: $G(\theta) = A \cos(\sqrt{k}\theta) + B \sin(\sqrt{k}\theta)$

$$A = A \cos(2\pi\sqrt{k}) + B \sin(2\pi\sqrt{k}) \quad | \cdot B$$

$$\sqrt{k}B = -\sqrt{k}A \sin(2\pi\sqrt{k}) + \sqrt{k}B \cos(2\pi\sqrt{k}) \quad | \cdot A$$

$$\Rightarrow B^2 \sin(2\pi\sqrt{k}) = -A^2 \sin(2\pi\sqrt{k})$$

$$\Rightarrow \sin(2\pi\sqrt{k}) = 0 \Rightarrow \sqrt{k} := n \in \mathbb{N}$$

$$\Rightarrow G_n(\theta) = A_n \cos(n\theta) + B_n \sin(n\theta)$$

$$r^2 F'' + r F' - n^2 F = 0 \Rightarrow F(r) = r^\alpha, \quad \alpha \in \mathbb{Q}$$

$$r^2 \alpha(\alpha - 1) r^{\alpha-2} + r \alpha r^{\alpha-1} - n^2 r^\alpha = 0$$

$$\Leftrightarrow \alpha(\alpha - 1) + \alpha - n^2 = 0$$

$$\Leftrightarrow \alpha^2 - n^2 = 0 \Rightarrow \alpha = \pm n$$

Note: The general solution is $F_n(r) = P_n r^n + Q_n r^{-n}$, which is not bounded for $r = 0$. Therefore the solution to this particular problem is $F_n(r) = P_n r^n$

$$\Rightarrow u(r, \theta) = \sum_{n=0}^{\infty} r^n (A_n \cos(n\theta) + B_n \sin(n\theta))$$

3. Boundary Condition $u(R, \theta) = \sum_{n=0}^{\infty} R^n (A_n \cos(n\theta) + B_n \sin(n\theta)) = f(\theta)$

$$A_n = \frac{1}{R^n \pi} \int_0^{2\pi} f(\phi) \cos(n\phi) d\phi$$

$$B_n = \frac{1}{R^n \pi} \int_0^{2\pi} f(\phi) \sin(n\phi) d\phi$$

4. Poisson Integral Form

$$\begin{aligned} u(r, \theta) &= \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi \\ &+ \frac{1}{\pi} \sum_{n=1}^{\infty} \left(\frac{r}{R} \right)^n \int_0^{2\pi} \underbrace{[\cos n\theta \cos \phi + \sin n\theta \sin \phi]}_{\cos(n(\theta - \phi))} f(\phi) d\phi \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi + \\ &+ \frac{1}{\pi} \sum_{n=1}^{\infty} \left(\frac{r}{R} \right)^n \int_0^{2\pi} \cos(n(\theta - \phi)) f(\phi) d\phi \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[1 + 2 \sum_{n=1}^{\infty} \left(\frac{r}{R} \right)^n \cos(n(\theta - \phi)) \right] f(\phi) d\phi \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{1 - \left(\frac{r}{R} \right)^2}{1 - 2 \frac{r}{R} \cos(\theta - \phi) + \left(\frac{r}{R} \right)^2} f(\phi) d\phi \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2rR \cos(\theta - \phi) + r^2} f(\phi) d\phi \end{aligned}$$

$$\begin{aligned} & * \sum_{n=1}^{\infty} t^n \cos(n\alpha) = \Re \left(\sum_{n=1}^{\infty} t^n e^{in\alpha} \right) \\ & = \Re \left(\frac{te^{i\alpha}}{1 - te^{i\alpha}} \right) = \dots = \frac{t \cos \alpha - t^2}{1 - 2t \cos \alpha + t^2} \end{aligned}$$

Theorem: Poisson Integral Form

$$u(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} K(r, \theta, R, \phi) f(\phi) d\phi,$$

$$\text{where } K(r, \theta, R\phi) := \frac{R^2 - r^2}{R^2 - 2rR \cos(\theta - \phi) + r^2}$$

Theorem: Mean value property

If the disk has center in (x_0, y_0) . Then using the Poisson Integral Form with $K(0, \theta, R, \phi) \Rightarrow u(x_0, y_0) = \frac{1}{2\pi} \int_0^{2\pi} f(\phi) d\phi = \frac{1}{2\pi} \int_0^{2\pi} u(x_0 + R \cos(\tilde{\sigma}), y_0 + R \sin(\tilde{\sigma})) d\tilde{\sigma}$.

Example: Poisson Kessel integral

Prove that $\frac{1}{2\pi} \int_0^{2\pi} \frac{(1-r^2)(\cos^3(\vartheta) \sin(\vartheta) - \sin^3(\vartheta) \cos(\vartheta))}{1-2r \cos(\vartheta-\varphi)+r^2} d\varphi = \frac{r^4}{4} \sin(4\vartheta)$

- Let $D = (r, \vartheta) | 0 \leq r \leq 1, 0 \leq \vartheta \leq 2\pi$, $u(r, \vartheta) = 1$ (constant function)
- $u(r, \vartheta) = \frac{1}{2\pi} \int_0^{2\pi} K(r, \vartheta, 1, \varphi) u(1, \varphi) d\varphi \leftrightarrow 1 = \frac{1}{2\pi} \int_0^{2\pi} \frac{1-r^2}{1-2r \cos(\vartheta-\varphi)+r^2} d\varphi$
- Consider the function $\cos^3(\vartheta) \sin(\vartheta) - \sin^3(\vartheta) \cos(\vartheta)$
- $\begin{cases} \Delta u = 0, \text{ in } D \\ u(1, \vartheta) = f(\vartheta), 0 \leq \vartheta \leq 2\pi \end{cases}$
- Using the boundary condition $\cos^3(\vartheta) \sin(\vartheta) - \sin^3(\vartheta) \cos(\vartheta) = x^3 y + xy^3$
- $\Delta(x^3 y + xy^3) = 6xy - 6xy = 0$ is a well defined harmonic function. It is the solution of the problem
- Convert in polar coordinates: $x^3 y + xy^3 = \dots \frac{r^4}{4} \sin(4\vartheta)$

Example: Property of Laplace on Circular Membrane

$$\begin{aligned} D_1 &= \text{unit disk} \\ \begin{cases} \Delta u = 0 \\ u(x, y) = xy + 3, (x, y) \in \partial D_1 \end{cases} \end{aligned}$$

1. Find $u(x, y)$: $\Delta(xy + 3) = \partial_{xx}(xy + 3) + \partial_{yy}(xy + 3) = 0$. By uniqueness of the solution for the Dirichlet problem the solution must be: $u(x, y) = xy + 3$
2. Find $u(0, 0)$: $u(0, 0) = 0 \cdot 0 + 3 = 3$
3. Find maximum of $u(x, y)$
 - Convert in polar coordinates: $u(r, \vartheta) = r^2 \cos(\vartheta) \sin(\vartheta) + 3 = \frac{r^2}{2} \sin(2\vartheta) + 3$
 - Maximise every components (independent variables)
 - Maximum are $u(1, \frac{\pi}{4}) = u(1, \frac{5}{4}\pi) = \frac{7}{2}$

Example: Circular Membrane of Lagrange Equation

Find sol of $u(r, \vartheta)$ from $\begin{cases} \Delta u = 0, 0 \leq r \leq R, 0 \leq \vartheta \leq 2\pi \\ u(R, \vartheta) = \sin^2(\vartheta), 0 \leq \vartheta \leq 2\pi \end{cases}$

1. Solution: $u(r, \vartheta) = \sum_{n=0}^{\infty} r^n (A_n \cos(n\vartheta) + B_n \sin(n\vartheta))$, coefficients found with: $u(R, \vartheta) = \sum_{n=0}^{\infty} R^n (A_n \cos(n\vartheta) + B_n \sin(n\vartheta)) = \sin^2(\vartheta)$
2. Use $\sin^2(\vartheta) = \frac{1}{2} - \frac{1}{2} \cos(2\vartheta)$ to obtain the coefficient: $\begin{cases} B_n = 0, \forall n \geq 0 \\ A_n = 0, \forall n \geq 0, n \neq 0, 2 \\ A_0 = \frac{1}{2} \\ A_2 = -\frac{1}{2R^2} \end{cases}$
3. The solution is $u(r, \vartheta) = \frac{1}{2} - \frac{1}{2R^2} \cos(2\vartheta)$

Maximum: constant doesn't play a role other function components are independent, maximise individually. $\begin{cases} -\cos(2\vartheta) = 1 \leftrightarrow \vartheta = \frac{\pi}{2}, \frac{3}{2}\pi \\ r = R \leftrightarrow r = R \end{cases}$.
In Cartesian coordinates: $\cos(2\vartheta) = \cos^2(\vartheta) - \sin^2(\vartheta) \rightarrow r^2 \cos(2\vartheta) = x^2 - y^2$. The solution is then $u(x, y) = \frac{1}{2} - \frac{1}{2r^2}(x^2 - y^2)$

6.4 Dirichlet Problem for a Ring:

$$\nabla^2 u = \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$$

- $u(R_1, \theta) = f_1(\theta)$ **Boundary Condition** ($R_1 < R_2$)
- $u(R_2, \theta) = f_2(\theta)$ **Boundary Condition** ($R_2 > R_1$)
- $u(r, 0) = u(r, 2\pi)$, $u_\theta(r, 0) = u_\theta(r, 2\pi)$ **Continuity Conditions**

6.4.1 Cosinus and Sinus variant:**1. Separating variables:**

$$u(r, \varphi) = R(r)\Phi(\varphi) \Rightarrow \begin{cases} r^2 R''(r) + rR'(r) + kR(r) = 0 \\ \Phi(\varphi) = k\Phi(\varphi) \\ \Phi(\varphi + 2\pi) = \Phi(\varphi) \end{cases}$$

2. **Hence Φ is periodic**, k must be negativ: the general solution is so:

$$\Phi_n(\varphi) = a_n \cos(n\varphi) + b_n \sin(n\varphi)$$

For the first we take $R(r) = r^\alpha$. The solution to this is:

$$R(r) = r^n, R(r) = r^{-n}, R(r) = \ln(r)$$

We can now use Superposition:

$$u(r, \varphi) = \sum_{n=1}^{\infty} ((a_n r^n + a_{-n} r^{-n}) \cos(n\varphi) + (b_n r^n + b_{-n} r^{-n}) \sin(n\varphi)) + c_0 + d_0 \ln(r)$$

3. **Boundary Condition:** from now suppose that $f_1(\theta) = 1 - 3 \cos(\varphi)$, $f_2(\theta) = 6 \sin(\varphi) + 60 \sin(\varphi) \cos(\varphi)$ and $R_1 = 1, R_2 = 2$.

We observe, by coefficient comparison for $f_1(\theta)$:

$$u(1, \varphi) = \sum_{n=1}^{\infty} ((a_n + a_{-n}) \cos(n\varphi) + (b_n + b_{-n}) \sin(n\varphi)) + c_0 = 1 - 3 \cos(\varphi)$$

- $a_1 + a_{-1} = -3$
- $a_n + a_{-n} = 0 \forall n \neq 1$
- $b_n + b_{-n} = 0 \forall n \in \mathbb{N}$
- $c_0 = 1$

And by coefficient comparison for $f_2(\theta)$:

$$u(2, \varphi) = \sum_{n=1}^{\infty} ((a_n 2^n + a_{-n} 2^{-n}) \cos(n\varphi) + (b_n 2^n + b_{-n} 2^{-n}) \sin(n\varphi)) + 1 + d_0 \ln(2) = 6 \sin(\varphi) + 30 \sin(2\varphi)$$

- $2^n a_n + 2^{-n} a_{-n} = 0 \quad \forall n \in \mathbb{N}$
- $b_n(2^n - 2^{-n}) = 0$ fuer $n \neq 1, 2$
- $b_1(2 - \frac{1}{2}) = 6$
- $b_2(4 - \frac{1}{4}) = 30$
- $d_0 = -\frac{1}{\ln(2)}$

4. **Find all the coefficient:** $u(r, \varphi) = \dots$

6.4.2 Complex variant:

1. **Separating variables:** as 6.5.1
2. **Hence Φ is periodic,** k must be negativ: the general solution is so:

$$u(r, \varphi) = \sum_{n=1}^{\infty} ((A_n r^n + B_n r^{-n}) e^{in\varphi}) + C_0 + D_0 \ln(r)$$

Attention: The solution cal also be written as:

$$u(r, \varphi) = \sum_{n=-\infty}^{\infty} (A_n e^{in\varphi} r^{-|n|}) + C_0 + D_0 \ln(r)$$

3. **Boundary Condition:** by comparison as 6.5.1. **Remember to transform $\sin(x)$ and $\cos(x)$ of the Boundary function in complex exponential form!**

4. **Find all the coefficient:** $u(r, \varphi) = \dots$


7 Two-Dimensional Wave Equation

Physical Assumptions

- Constant density and perfect flexibility of membrane
- Constant tension T
- small deflection compared to the size of the membrane

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) = c^2 \Delta^2 u \quad c^2 = \frac{T}{\rho}$$

7.1 Rectangular Membrane:



- $u = 0$ on the boundary
- $u(x, y, 0) = f(x, y)$ (initial displacement)
- $u_t(x, y, 0) = g(x, y)$ (initial velocity)

1. ODEs From the Wave Equation

$$u(x, y, t) = F(x, y)G(t) \Rightarrow F\ddot{G} = c^2(F_{xx}G + F_{yy}G)$$

$$\Leftrightarrow \frac{\ddot{G}}{c^2 G} = \frac{1}{F}(F_{xx} + F_{yy}) = -\nu^2 = \text{const}$$

$$\ddot{G} + \lambda^2 G = 0 \quad \text{with } \lambda = c\nu, \quad F_{xx} + F_{yy} + \nu^2 F = 0$$

$$F(x, y) = H(x)Q(y) \Rightarrow \frac{d^2 H}{dx^2} Q = - \left(H \frac{d^2 Q}{dy^2} + \nu^2 H Q \right)$$

$$\Leftrightarrow \frac{1}{H} \frac{d^2 H}{dx^2} = - \frac{1}{Q} \left(\frac{d^2 Q}{dy^2} + \nu^2 Q \right) = -k^2 = \text{const}$$

$$\frac{d^2 Q}{dy^2} + p^2 Q = 0 \quad \text{with } p^2 = \nu^2 - k^2, \quad \frac{d^2 H}{dx^2} + k^2 H = 0$$

2. Boundary Condition

$$H(x) = A \cos kx + B \sin kx \quad Q(y) = C \cos py + D \sin py$$

$$H(0) = A = 0, \quad H(a) = B \sin ka = 0 \Rightarrow k = \frac{m\pi}{a}$$

$$Q(0) = C = 0, \quad Q(b) = D \sin pb \Rightarrow p = \frac{n\pi}{b}$$

$$H_m(x) = \sin \frac{m\pi x}{a} \quad Q_n(y) = \sin \frac{n\pi y}{b}$$

$$\Rightarrow F_{mn}(x, y) = H_m(x)Q_n(y) = \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$$

$$\text{Since } p^2 = \nu^2 - k^2, \quad \lambda = c\nu \Rightarrow \lambda = \lambda_{mn} = c\pi \sqrt{\frac{m^2}{a^2} + \frac{n^2}{b^2}}$$

$$\Rightarrow G_{mn}(t) = B_{mn} \cos \lambda_{mn} t + B_{mn}^* \sin \lambda_{mn} t$$

$$u_{mn}(x, y, t) =$$

$$= (B_{mn} \cos \lambda_{mn} t + B_{mn}^* \sin \lambda_{mn} t) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$$

These functions are called **eigenfunctions** and λ_{mn} are called **eigenvalues**.

3. Double Fourier Series

$$u(x, t) =$$

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} (B_{mn} \cos \lambda_{mn} t + B_{mn}^* \sin \lambda_{mn} t) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b}$$

$$u(x, y, 0) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} B_{mn} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} = f(x, y)$$

$$\begin{aligned} \bullet K_m(y) &:= \sum_{n=1}^{\infty} B_{mn} \sin \frac{n\pi y}{b} \\ \Rightarrow B_{mn} &= \frac{2}{b} \int_0^b K_m(y) \sin \frac{n\pi y}{b} dy \end{aligned}$$

$$\begin{aligned} \bullet f(x, y) &= \sum_{m=1}^{\infty} K_m(y) \sin \frac{m\pi x}{a} \\ \Rightarrow K_m(y) &= \frac{2}{a} \int_0^a f(x, y) \sin \frac{m\pi x}{a} dx \end{aligned}$$

$$B_{mn} = \frac{4}{ab} \int_0^b \int_0^a f(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy$$

$$\frac{\partial u}{\partial t} \Big|_{t=0} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} B_{mn}^* \lambda_{mn} \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} = g(x, y)$$

Analogously:

$$B_{mn}^* = \frac{4}{ab\lambda_{mn}} \int_0^b \int_0^a g(x, y) \sin \frac{m\pi x}{a} \sin \frac{n\pi y}{b} dx dy$$

8 Tables

8.1 Laplace Transforms:

$\mathcal{L}\{f(t)\}$	$f(t)$	$\mathcal{L}\{f(t)\}$	$f(t)$
$1/s$	1	e^{-as}/s	$u(t-a)$
$1/s^2$	t	e^{-as}	$\delta(t-a)$
$1/s^n$	$t^{n-1}/(n-1)!$	$\frac{1}{\sqrt{s}}e^{-\omega/s}$	$\frac{1}{\sqrt{\pi t}}\cos 2\sqrt{\omega t}$
$1/\sqrt{s}$	$1/\sqrt{\pi t}$	$e^{-k\sqrt{s}}$	$\frac{k}{2\sqrt{\pi t^3}}e^{-k^2/4t}$
$1/s^{3/2}$	$2\sqrt{t/\pi}$	$\frac{1}{(s-a)(s-b)}$	$\frac{1}{a-b}(e^{at}-e^{bt})$
$1/s^k$	$t^{k-1}/\Gamma(k)$	$\frac{s}{(s-a)(s-b)}$	$\frac{1}{a-b}(ae^{at}-be^{bt})$
$\frac{1}{s-a}$	e^{at}	$\frac{(s^2+\omega^2)^2}{2\omega s}$	$\sin \omega t - \omega t \cos \omega t$
$\frac{1}{(s-a)^2}$	te^{at}	$\frac{(s^2+\omega^2)^2}{2\omega s^2}$	$t \sin \omega t$
$\frac{1}{(s-a)^n}$	$\frac{1}{(n-1)!}t^{n-1}e^{at}$	$\frac{(s^2+\omega^2)^2}{s^2}$	$\sin \omega t + \omega t \cos \omega t$
$\frac{1}{(s-a)^k}$	$\frac{1}{\Gamma(k)}t^{k-1}e^{at}$	$\frac{1}{(s^2+a^2)(s^2+b^2)}$	$\frac{1}{b^2-a^2}(\cos at - \cos bt)$
$\frac{\omega}{s^2+\omega^2}$	$\sin \omega t$	$\frac{s^4+4\omega^4}{2\omega^2 s}$	$\sin \omega t \cos \omega t - \cos \omega t \sinh \omega t$
$\frac{a}{s^2-a^2}$	$\sinh at$	$\frac{s^4+4\omega^4}{2\omega^3}$	$\sin \omega t \sinh \omega t$
$\frac{\omega}{(s-a)^2+\omega^2}$	$e^{at} \sin \omega t$	$\frac{s^4-\omega^4}{2\omega^2 s}$	$\sinh \omega t - \sin \omega t$
$\frac{\omega}{(s-a)^2-\omega^2}$	$e^{at} \sinh \omega t$	$\frac{s^4-\omega^4}{s^4-\omega^4}$	$\cosh \omega t - \cos \omega t$
$\frac{s}{s^2+\omega^2}$	$\cos \omega t$	$\ln \frac{s-a}{s^2+\omega^2}$	$\frac{1}{t}(e^{bt}-e^{at})$
$\frac{s}{s^2-a^2}$	$\cosh at$	$\ln \frac{s^2+\omega^2}{s^2}$	$\frac{2}{t}(1-\cos \omega t)$
$\frac{s-a}{(s-a)^2+\omega^2}$	$e^{at} \cos \omega t$	$\ln \frac{s^2-\omega^2}{s^2}$	$\frac{2}{r}(1-\cosh \omega t)$
$\frac{s-a}{(s-a)^2-\omega^2}$	$e^{at} \cosh \omega t$		
$\frac{\omega^2}{s(s^2+\omega^2)}$	$1 - \cos \omega t$		
$\frac{\omega^3}{s^2(s^2+\omega^2)}$	$\omega t - \sin \omega t$		

 $k > 0, n \in \mathbb{N}, a \neq b, \gamma \approx 0.5772$

8.2 Fourier Transforms:

 $a > 0$

$f(x)$	$\hat{f}_c(w)$	$f(x)$	$\hat{f}_s(w)$
$\begin{cases} 1 & 0 < x < a \\ 0 & \text{otherwise} \end{cases}$	$\sqrt{\frac{2}{\pi}} \frac{\sin aw}{w}$	$\begin{cases} 1 & 0 < x < a \\ 0 & \text{otherwise} \end{cases}$	$\sqrt{\frac{2}{\pi}} \left[\frac{1 - \cos aw}{w} \right]$
e^{-ax}	$\sqrt{\frac{2}{\pi}} \frac{a}{a^2 + w^2}$	$1/\sqrt{x}$	$1/\sqrt{w}$
$e^{-x^2/2}$	$e^{-w^2/2}$	$1/x^{3/2}$	$2\sqrt{w}$
e^{-ax^2}	$\frac{1}{\sqrt{2a}} e^{-w^2/4a}$	e^{-ax}	$\sqrt{\frac{2}{\pi}} \left(\frac{w}{a^2 + w^2} \right)$
$x^n e^{-ax}$	$\sqrt{\frac{2}{\pi}} \frac{n!}{(a^2 + w^2)^{n+1}} \Re(a + iw)^{n+1}$	$\frac{e^{-ax}}{x}$	$\sqrt{\frac{2}{\pi}} \arctan \frac{w}{a}$
$\begin{cases} \cos x & 0 < x < a \\ 0 & \text{otherwise} \end{cases}$	$\frac{1}{\sqrt{2\pi}} \left[\frac{\sin a(1-w)}{1-w} + \frac{\sin a(1+w)}{1+w} \right]$	$x^n e^{-ax}$	$\sqrt{\frac{2}{\pi}} \frac{n!}{(a^2 + w^2)^{n+1}} \Im(a + iw)^{n+1}$
$\cos(ax^2)$	$\frac{1}{\sqrt{2\pi}} \cos \left(\frac{w^2}{4a - \frac{\pi}{4}} \right)$	$xe^{-x^2/2}$	$we^{w^2/2}$
$\sin(ax^2)$	$\frac{1}{\sqrt{2a}} \cos \left(\frac{w^2}{4a} + \frac{\pi}{4} \right)$	xe^{-ax^2}	$\frac{2}{(2a)^{3/2}} e^{-w^2/4a}$
$\frac{\sin ax}{x}$	$\sqrt{\frac{\pi}{2}} [1 - u(w-a)]$	$\begin{cases} \sin x & 0 < x < a \\ 0 & \text{otherwise} \end{cases}$	$\frac{1}{\sqrt{2\pi}} \left[\frac{\sin a(1-w)}{1-w} - \frac{\sin a(1+w)}{1+w} \right]$
$\frac{e^{-x} \sin x}{x}$	$\frac{1}{\sqrt{2\pi}} \arctan \frac{2}{w^2}$	$\frac{\cos ax}{x}$	$\sqrt{\frac{\pi}{2}} u(w-a)$
$\frac{f(x)}{x}$	$\frac{\hat{f}(w)}{w}$	$\arctan \frac{2a}{x}$	$\sqrt{2\pi} \frac{\sin aw}{w} e^{-aw}$
$\begin{cases} 1 & -b < x < b \\ 0 & \text{otherwise} \end{cases}$	$\sqrt{\frac{2}{\pi}} \frac{\sin bw}{w}$		
$\begin{cases} 1 & b < x < c \\ 0 & \text{otherwise} \end{cases}$	$\frac{e^{-ibw} - e^{-icw}}{iw\sqrt{2\pi}}$		
$\frac{1}{x^2 + a^2}$	$\sqrt{\frac{\pi}{2}} \frac{e^{-a w }}{a}$		
$\begin{cases} x & 0 < x < b \\ 2x-b & b < x < 2b \\ 0 & \text{otherwise} \end{cases}$	$\frac{-1 + 2e^{ibw} - e^{2ibw}}{\sqrt{2\pi}w^2}$		
$\begin{cases} e^{-ax} & x > 0 \\ 0 & \text{otherwise} \end{cases}$	$\frac{1}{\sqrt{2\pi}(a+iw)}$		
$\begin{cases} e^{ax} & b < x < c \\ 0 & \text{otherwise} \end{cases}$	$\frac{e^{(a-iw)c} - e^{(a-iw)b}}{\sqrt{2\pi}(a-iw)}$		
$\begin{cases} e^{iax} & -b < x < b \\ 0 & \text{otherwise} \end{cases}$	$\sqrt{\frac{2}{\pi}} \frac{\sin b(w-a)}{w-a}$		
$\begin{cases} e^{iax} & b < x < c \\ 0 & \text{otherwise} \end{cases}$	$\frac{i}{\sqrt{2\pi}} \frac{e^{ib(a-w)} - e^{ic(a-w)}}{a-w}$		
e^{-ax^2}	$\frac{1}{\sqrt{2a}} e^{-w^2/4a}$		
$\frac{\sin ax}{x}$	$\begin{cases} \sqrt{\pi/2} & w < a \\ 0 & w > a \end{cases}$		

8.3 Some Integrals:

	$\int_0^{\pi/4}$	$\int_0^{\pi/2}$	\int_0^{π}	$\int_0^{2\pi}$	$\int_{-\pi/4}^{\pi/4}$	$\int_{\pi/2}^{\pi}$	$\int_{-\pi}^{\pi}$
\sin	$\frac{\sqrt{2}-1}{\sqrt{2}}$	1	2	0	0	0	0
\sin^2	$\frac{\pi-2}{8}$	$\pi/4$	$\pi/2$	π	$\frac{\pi-2}{4}$	$\pi/2$	π
\sin^3	$\frac{8-5\sqrt{2}}{12}$	2/3	4/3	0	0	0	0
\sin^4	$\frac{3\pi-8}{32}$	$\frac{3\pi}{16}$	$\frac{3\pi}{8}$	$\frac{3\pi}{4}$	$\frac{3\pi-8}{16}$	$\frac{3\pi}{8}$	$\frac{3\pi}{4}$
\cos	$1/\sqrt{2}$	1	0	0	$\sqrt{2}$	2	0
\cos^2	$\frac{2+\pi}{8}$	$\pi/4$	$\pi/2$	π	$\frac{2+\pi}{4}$	$\pi/2$	π
\cos^3	$\frac{5}{6\sqrt{2}}$	2/3	0	0	$\frac{5}{3\sqrt{2}}$	4/3	0
\cos^4	$\frac{8+3\pi}{32}$	$\frac{3\pi}{16}$	$\frac{3\pi}{8}$	$\frac{3\pi}{4}$	$\frac{8+3\pi}{16}$	$\frac{3\pi}{8}$	$\frac{3\pi}{4}$
$\sin \cos$	1/4	1/2	0	0	0	0	0
$\sin^2 \cos$	$\frac{2}{6\sqrt{2}}$	1/3	0	0	$\frac{1}{3\sqrt{2}}$	2/3	0
$\sin \cos^2$	$\frac{4-\sqrt{2}}{12}$	1/3	2/3	0	0	0	0

$$\int_{r\frac{\pi}{2}}^{s\frac{\pi}{2}} \sin^n x \, dx = \frac{n-1}{n} \int_{r\frac{\pi}{2}}^{s\frac{\pi}{2}} \sin^{(n-2)} x \, dx \quad \forall n \geq 2$$

$$\int_{r\frac{\pi}{2}}^{s\frac{\pi}{2}} \cos^n x \, dx = \frac{n-1}{n} \int_{r\frac{\pi}{2}}^{s\frac{\pi}{2}} \cos^{(n-2)} x \, dx \quad r, s \in \mathbb{Z}$$

$$\int (ax+b)^n \, dx = \frac{1}{a(n+1)} (ax+b)^{n+1}$$

$$\int \frac{1}{ax+b} \, dx = \frac{1}{a} \ln |ax+b|$$

$$\int (ax^p+b)^n x^{p-1} \, dx = \frac{(ax^p+b)^{n+1}}{ap(n+1)}$$

$$\int (ax^p+b)^{-1} x^{p-1} \, dx = \frac{1}{ap} \ln |ax^p+b|$$

$$\int \frac{ax+b}{cs+d} \, dx = \frac{ax}{c} - \frac{ad-bc}{c^2} \ln |cx+d|$$

$$\int x(ax+b)^n \, dx = \frac{(ax+b)^{n+2}}{(n+2)a^2} - \frac{b(ax+b)^{n+1}}{(n+1)a^2}$$

$$\int \frac{x}{(ax+b)^n} \, dx = -\frac{1}{(n-2)a^2(ax+b)^{n-2}} + \frac{1}{(n-1)a^2(ax+b)^{n-1}}$$

$$\int \frac{x}{(x^2+a)^n} \, dx = -\frac{1}{2(n-1)(a^2+x^2)^{n-1}}$$

$$\int \frac{x}{(x^2-a^2)^n} \, dx = \frac{1}{2(n-1)(a^2+x^2)^{n-1}}$$

$$\int \frac{x}{x^2+a} \, dx = \frac{1}{2} \ln |x^2+a|$$

$$\int \frac{x}{ax^2+b} \, dx = \frac{1}{2a} \ln |ax^2+b|$$

$$\int \frac{1}{a^2+x^2} \, dx = \frac{1}{a} \arctan \frac{x}{a}$$

$$\int \sqrt{x^2+a^2} \, dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \ln |x+\sqrt{x^2+a^2}|$$

$$\int \sqrt{a^2-x^2} \, dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \arcsin \frac{x}{|a|}$$

$$\int \sqrt{x^2-a^2} \, dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \ln |x+\sqrt{x^2-a^2}|$$

$$\int \frac{1}{\sqrt{x^2+a^2}} \, dx = \ln |x+\sqrt{a^2+x^2}|$$

$$\int \frac{1}{\sqrt{x^2-a^2}} \, dx = \ln |x+\sqrt{x^2-a^2}|$$

$$\int \frac{1}{\sqrt{a^2-x^2}} \, dx = \arcsin \frac{x}{|a|}$$

$$\int \frac{x^2-y^2}{x^2+y^2} \, dx = x - 2 \tan^{-1}\left(\frac{x}{y}\right) \cdot y$$

$$\int \frac{1}{\sqrt{1-x^2}} \, dx = \arcsin x$$

$$\int \frac{1}{1-x^2} \, dx = \operatorname{artanh} x = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|, \quad |x| < 1$$

$$\int \frac{1}{\sqrt{x^2+1}} \, dx = \operatorname{arsinh} x = \ln |x+\sqrt{x^2+1}|$$

$$\int \frac{1}{\sqrt{x^2-1}} \, dx = \operatorname{arcosh} x = \ln |x+\sqrt{x^2-1}|, \quad |x| > 1$$

$$\int \frac{1}{\sin^2 x} \, dx = -\cot x$$

$$\int \sin^2 x \, dx = \frac{1}{2} (x - \sin x \cos x)$$

$$\int \sin^3 x \, dx = \frac{1}{12} (\cos 3x - 9 \cos x)$$

$$\int \sin^4 x \, dx = \frac{1}{32} (12x - 8 \sin 2x + \sin 4x)$$

$$\int \frac{1}{\sin x} \, dx = \ln \left| \tan \frac{x}{2} \right|$$

$$\int \sin^n x \, dx = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} \int \sin^{n-2} x \, dx$$

$$\int \arcsin x \, dx = x \arcsin x + \sqrt{1-x^2}$$

$$\int \cos^2 x \, dx = \frac{1}{2} (x + \sin x \cos x)$$

$$\int \cos^3 x \, dx = \frac{1}{12} (9 \sin x + \sin 3x)$$

$$\int \cos^4 x \, dx = \frac{1}{32} (12x + 8 \sin 2x + \sin 4x)$$

$$\int \frac{1}{\cos x} \, dx = \ln \left| \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right|$$

$$\int \cos^n x \, dx = \frac{1}{n} \sin x \cos^{n-1} x + \frac{n-1}{n} \int \cos^{n-2} x \, dx$$

$$\int \arccos x \, dx = x \arccos x - \sqrt{1-x^2}$$

$$\int \sin 2x \, dx = -\frac{1}{2} \cos 2x$$

$$\int \cos 2x \, dx = \sin x \cos x$$

$$\int \sin x \cos x \, dx = -\frac{1}{2} \cos^2 x$$

$$\int \sin^2 x \cos x \, dx = \frac{1}{3} \sin^3 x$$

$$\int \sin x \cos^2 x \, dx = -\frac{1}{3} \cos^3 x$$

$$\int \sin^2 x \cos^2 x \, dx = \frac{1}{32} (4x - \sin 4x)$$

$$\int \sin^n ax \cos ax \, dx = \frac{\sin^{n+1} ax}{a(n+1)}$$

$$\int \sin ax \cos^n ax \, dx = -\frac{\cos^{n+1} ax}{a(n+1)}$$

$$\int \tan x \, dx = -\ln |\cos x|$$

$$\int \tan^2 x \, dx = \tan x - x$$

$$\int \tan^3 x \, dx = \frac{1}{2 \cdot \cos(x)^2} + \ln |\cos x|$$

$$\int \tan^4 x \, dx = x + \frac{1}{3} \tan x \left(\frac{1}{\cos(x)^2} - 4 \right)$$

$$\int \frac{1}{\tan x} \, dx = \ln |\sin x|$$

$$\int \arctan x \, dx = x \arctan x - \frac{1}{2} \ln |1+x^2|$$

$$\int \coth x \, dx = \ln |\sinh x|$$

$$\int \frac{\cos ax}{\sin^n ax} \, dx = -\frac{1}{a(n-1) \sin^{n-1} ax}$$

$$\int \tanh x \, dx = \ln \cosh x$$

$$\int \operatorname{arsinh} x \, dx = x \operatorname{arsinh} x - \sqrt{x^2+1}$$

$$\int \operatorname{arcosh} x \, dx = x \operatorname{arcosh} x - \sqrt{x^2-1}$$

$$\int \operatorname{artanh} x \, dx = x \operatorname{artanh} x + \frac{1}{2} \ln |1-x^2|$$

$$\int \frac{1}{e^x+a} \, dx = \frac{x - \ln |a+e^x|}{a}$$

$$\int \frac{1}{x^2+x} \, dx = \ln |x| - \ln |x+1|$$

$$\int \frac{1}{ax^2+\sqrt{bx+c}} \, dx = \frac{2 \arctan \left(\frac{2ax+b}{\sqrt{4ac-b^2}} \right)}{\sqrt{4ac-b^2}}$$

$$\int x e^{ax} \, dx = \frac{ax-1}{a^2} e^{ax}$$

$$\int x^2 e^{ax} \, dx = \frac{a^2 x^2 - 2ax + 2}{a^3} e^{ax}$$

$$\int \frac{1}{p+qe^{ax}} \, dx = \frac{x}{p} - \frac{1}{ap} \ln |p+qe^{ax}|$$

$$\int x e^{kx^2} \, dx = \frac{1}{2k} e^{kx^2}$$

$$\begin{aligned}\int \frac{\ln^n |x|}{x} dx &= \frac{\ln^{n+1} |x|}{x+1} \\ \int \sin^2 ax \, dx &= \frac{x}{2} - \frac{\sin 2ax}{4a} \\ \int x \cdot \sin ax \, dx &= \frac{\sin ax}{a^2} - \frac{x \cdot \cos ax}{a} \\ \int \cos^2 ax \, dx &= \frac{x}{2} + \frac{\sin 2ax}{4a} \\ \int x \cdot \cos ax \, dx &= \frac{\cos ax}{a^2} + \frac{x \cdot \sin ax}{a} \\ \int \sin ax \cos ax \, dx &= -\frac{\cos^2 ax}{2a} \\ \int e^x \sin x \, dx &= \frac{e^x}{2} (\sin x - \cos x) \\ \int e^x \cos x \, dx &= \frac{e^x}{2} (\sin x + \cos x) \\ \int x^2 \cdot \sin ax \, dx &= \frac{1}{a^3} (-a^2 x^2 \cos ax + 2 \cos ax + 2ax \sin ax) \\ \int x^2 \cdot \cos ax \, dx &= \frac{1}{a^3} (a^2 x^2 \sin ax - 2 \sin ax + 2ax \cos ax) \\ \int \tan ax \, dx &= -\frac{1}{a} \ln |\cos ax| \\ \int e^{ax} \sin nx \, dx &= \frac{e^{ax} (a \sin nx - n \cos nx)}{a^2 + n^2} \\ \int e^{ax} \cos nx \, dx &= \frac{e^{ax} (a \cos nx + n \sin nx)}{a^2 + n^2} \\ \int \cos(kx) \sin(nx)^2 \, dx &= -\frac{1}{4} \left(\frac{\sin(x(k+2n))}{k+2n} + \frac{\sin(x(k-2n))}{k-2n} \right) + \frac{1}{2k} \sin(kx) \\ \int \sin(kx) \cos(nx)^2 \, dx &= -\frac{1}{4} \left(\frac{\cos(x(k+2n))}{k+2n} + \frac{\cos(x(k-2n))}{k-2n} \right) - \frac{1}{2k} \cos(kx) \\ \int \sin(kx) \sin(nx)^2 \, dx &= \frac{1}{4} \left(\frac{\cos(x(k+2n))}{k+2n} + \frac{\cos(x(k-2n))}{k-2n} \right) - \frac{1}{2k} \cos(kx) \\ \int \cos(kx) \cos(nx)^2 \, dx &= \frac{1}{4} \left(\frac{\sin(x(k+2n))}{k+2n} + \frac{\sin(x(k-2n))}{k-2n} \right) + \frac{1}{2k} \sin(kx)\end{aligned}$$

8.3.1 Basic integrals:

$f(x)$	$F(x)$	$f(x)$	$F(x)$
a	ax	x^n	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $	$\frac{1}{x^n}$	$\frac{-1}{(n-1)x^{n-1}}$
\sqrt{x}	$\frac{2}{3} x \sqrt{x}$	$\frac{1}{\sqrt{x}}$	$2\sqrt{x}$
$\frac{1}{(x-a)(x-b)}$	$\frac{1}{a-b} \ln \left \frac{x-a}{x-b} \right $	$\frac{ax+b}{cx+d}$	$\frac{ax}{c} - \frac{ad-bc}{c^2} \ln cx+d $
$\frac{1}{x^2+a^2}$	$\frac{1}{a} \arctan\left(\frac{x}{a}\right)$	$\frac{1}{2a} \ln \left \frac{x-a}{x+a} \right $	$\frac{1}{2a} \ln \left \frac{x-a}{x+a} \right $
e^x	e^x	$\ln(x)$	$x(\ln(x)-1)$
a^x	$\frac{a^x}{\ln(a)}$	$\log_a(x)$	$x(\log_a(x) - \log_a(e))$
xe^ax	$\frac{a^x}{\ln(a)}$	$x \ln(ax)$	$\frac{x^2}{4} (2 \ln(ax) - 1)$
$\sin(mx)$	$-\frac{\cos(mx)}{m}$	$\arcsin(x)$	$x \arcsin(x) + \sqrt{1-x^2}$
$\cos(mx)$	$\frac{\sin(mx)}{m}$	$\arccos(x)$	$x \arccos(x) - \sqrt{1-x^2}$
$\tan(mx)$	$-\frac{\ln \cos(mx) }{m}$	$\arctan(x)$	$x \arctan(x) - \frac{1}{2} \ln(1+x^2)$
$\cot(mx)$	$\frac{\ln \sin(mx) }{m}$	$\operatorname{arccot}(x)$	$x \operatorname{arccot}(x) + \frac{1}{2} \ln(1+x^2)$
$\frac{1}{1+\sin(x)}$	$-\frac{\cos(x)}{1+\sin(x)}$	$\frac{1}{1-\sin(x)}$	$\frac{\cos(x)}{1-\sin(x)}$
$\frac{1}{1+\cos(x)}$	$\frac{\sin(x)}{1+\cos(x)}$	$\frac{1}{1-\cos(x)}$	$-\frac{\sin(x)}{1-\cos(x)}$
$\sinh(x)$	$\cosh(x)$	$\operatorname{arsinh}(x)$	$x \operatorname{arsinh}(x) - \sqrt{x^2+1}$
$\cosh(x)$	$\sinh(x)$	$\operatorname{arcosh}(x)$	$x \operatorname{arcosh}(x) - \sqrt{x^2-1}$
$\tanh(x)$	$\ln(\cosh(x))$	$\operatorname{artanh}(x)$	$x \operatorname{artanh}(x) + \frac{1}{2} \ln(1-x^2)$
$\coth(x)$	$\ln \sinh(x) $	$\operatorname{arcoth}(x)$	$x \operatorname{arcoth}(x) + \frac{1}{2} \ln(x^2-1)$

8.3.2 Per Parts Resolution method:

$$\int_a^b f(x) \cdot g'(x) dx = f(b) \cdot g(b) - f(a) \cdot g(a) - \int_a^b f'(x) g(x) dx$$

8.4 Some Series:

$$\begin{aligned}\sum_{k=0}^{\infty} aq^k &= \frac{a}{1-q}, \quad 0 < |q| < 1 \\ \sum_{k=0}^{\infty} (k+1)q^k &= \frac{1}{(1-q)^2}, \quad 0 < |q| < 1 \\ \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} &= \frac{\pi}{4} & \sum_{k=1}^{\infty} \frac{1}{k^2} &= \frac{\pi^2}{6} \\ \sum_{k=1}^{\infty} \frac{1}{k^4} &= \frac{\pi^4}{90} & \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} &= \ln 2 \\ \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2} &= \frac{\pi^2}{12} & \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^4} &= \frac{7\pi^4}{720} \\ \sum_{k=0}^{\infty} \frac{1}{k!} &= e & \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} &= \frac{1}{e}\end{aligned}$$

8.5 Logarithm:

$$\begin{aligned}\log(uv) &= \log u + \log v & \log \frac{u}{v} &= \log u - \log v \\ \log \frac{1}{v} &= -\log v & \log u^r &= r \cdot \log u \\ \log_a x &= \frac{\log_b x}{\log_b a} & y = \log_a x &\Leftrightarrow a^y = x\end{aligned}$$

8.6 Differential Calculus:

$$\begin{aligned}f(x) &= u(x) \pm v(x) & f'(x) &= u'(x) \pm v'(x) \\ f(x) &= c \cdot u(x) & f'(x) &= c \cdot u'(x) \\ f(x) &= u(x) \cdot v(x) & f'(x) &= u'(x)v(x) + u(x) \cdot v'(x) \\ f(x) &= \frac{u(x)}{v(x)} & f'(x) &= \frac{u'(x)v(x) - u(x)v'(x)}{v^2(x)} \\ f(x) &= u(v(x)) & f'(x) &= u'(v(x)) \cdot v'(x)\end{aligned}$$
$$\begin{aligned}(\arcsin x)' &= \frac{1}{\sqrt{1-x^2}} & (\arccos x)' &= -\frac{1}{\sqrt{1-x^2}} \\ (\arctan x)' &= \frac{1}{1+x^2} & (\tan x)' &= \frac{1}{\cos^2 x} = 1 + \tan^2 x \\ (\operatorname{arsinh} x)' &= \frac{1}{\sqrt{x^2+1}} & (\operatorname{arcosh} x)' &= \frac{1}{\sqrt{x^2-1}} \\ (\operatorname{artanh} x)' &= \frac{1}{1-x^2} & (\tanh x)' &= \frac{1}{\cosh^2 x} = 1 - \tanh^2 x\end{aligned}$$

8.7 Standard ODE:

$$F''(x) = kF(x)$$

- $k = 0 \rightarrow F(x) = Ax + B$
- $k < 0 \rightarrow F(x) = C \cos(\sqrt{-k}x) + D \sin(\sqrt{-k}x)$
- $k > 0 \rightarrow F(x) = Ee^{\sqrt{k}x} + Ie^{-\sqrt{k}x}$
or $F(x) = E \sinh(\sqrt{k}x) + I \cosh(\sqrt{k}x)$
if $E = -I \rightarrow F(x) = C \sinh(\sqrt{k}x)$
if $E = I \rightarrow F(x) = C \cosh(\sqrt{k}x)$

$$F''(x) = -kF(x)$$

- $k = 0 \rightarrow F(x) = Ax + B$
- $k > 0 \rightarrow F(x) = C \cos(\sqrt{k}x) + D \sin(\sqrt{k}x)$
- $k < 0 \rightarrow F(x) = Ee^{\sqrt{-k}x} + Ie^{-\sqrt{-k}x}$
or $F(x) = E \sinh(\sqrt{-k}x) + I \cosh(\sqrt{-k}x)$
if $E = -I \rightarrow F(x) = C \sinh(\sqrt{-k}x)$
if $E = I \rightarrow F(x) = C \cosh(\sqrt{-k}x)$

$$F'(x) = aF(x) \Leftrightarrow F = Le^{ax}$$

8.8 Some Limits:

$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$	$\lim_{x \rightarrow 0} \frac{e^{x-1}}{x} = 1$	$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \ln a$
$\lim_{x \rightarrow 1} \frac{\ln x}{x-1} = 1$	$\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$	$\lim_{x \rightarrow \infty} \frac{\ln x}{x^a} = 0$
$\lim_{x \rightarrow \infty} \frac{x^m}{e^{ax}} = 0$	$\lim_{x \rightarrow 0} (x^a \ln x) = 0$	$\lim_{x \rightarrow 0} \frac{\ln(a+x)}{x} = \frac{1}{a}$
$\lim_{x \rightarrow \infty} \sqrt[x]{x} = 1$	$\lim_{x \rightarrow \infty} \frac{e^{ax}}{b} = \infty$	$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$
$\lim_{x \rightarrow \infty} \sqrt[x]{a} = 1$	$\lim_{x \rightarrow 0} x^a \ln^b x = 0$	$\lim_{x \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$
$\lim_{x \rightarrow \infty} \frac{x^b}{e^{ax}} = 0$		$\lim_{x \rightarrow \infty} \left(\frac{x-1}{x+1}\right)^x = \frac{1}{e^2}$
$\lim_{x \rightarrow \pm \frac{\pi}{2}} \frac{\tan x}{x} = \mp \infty$	$\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 1$	$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0$
$\lim_{x \rightarrow 0} \frac{x}{\sin ax} = \frac{1}{a}$	$\lim_{x \rightarrow 0} \frac{\sin ax}{\sin x} = a$	$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$
$\lim_{x \rightarrow 0} \frac{\arctan x}{x} = 1$	$\lim_{x \rightarrow 0} \frac{\arcsin x}{x} = 1$	$\lim_{x \rightarrow \pm \infty} \arctan x = \pm \frac{\pi}{2}$

8.9 Limit:

$$\lim_{x \rightarrow a} (f(x) \pm g(x)) = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$$

$$\lim_{x \rightarrow a} c \cdot f(x) = c \cdot \lim_{x \rightarrow a} f(x)$$

$$\lim_{x \rightarrow a} (f(x) \cdot g(x)) = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

$$\lim_{x \rightarrow a} f(x) = u_0, \quad g(u) \text{ stetig bei } u_0$$

$$\Rightarrow \lim_{x \rightarrow a} g(f(x)) = g\left(\lim_{x \rightarrow a} f(x)\right) = g(u_0)$$


8.9.1 Bernoulli De L'Hôpital:

Sind f und g differenzierbar, $g(x) \neq 0$ auf (a, b) ,

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0 \text{ (oder } \pm\infty) \text{ und existiert } \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)},$$

$$\text{so gilt: } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

8.10 Trigonometric Functions:



$$\bullet \cos^2(\alpha) + \sin^2(\alpha) = 1$$

$$\bullet \tan(\alpha) = \frac{\sin(\alpha)}{\cos(\alpha)}$$

$$\bullet \cot(\alpha) = \frac{\cos(\alpha)}{\sin(\alpha)}$$

$$\bullet \cot(\alpha) = \frac{1}{\tan(\alpha)}$$

$$\bullet \frac{1}{\cos^2(\alpha)} = 1 + \tan^2(\alpha)$$

$$\bullet \frac{1}{\sin^2(\alpha)} = 1 + \cot^2(\alpha)$$

8.10.1 Periodicity:

$$\bullet \cos(\alpha + 2\pi) = \cos(\alpha)$$

$$\bullet \sin(\alpha + 2\pi) = \sin(\alpha)$$

$$\bullet \tan(\alpha + 2\pi) = \tan(\alpha)$$

8.10.2 Relation between trigonometric functions:

$\cos(-\alpha) = \cos(\alpha)$	$\sin(-\alpha) = -\sin(\alpha)$	$\tan(-\alpha) = -\tan(\alpha)$
$\cos(\pi - \alpha) = -\cos(\alpha)$	$\sin(\pi - \alpha) = \sin(\alpha)$	$\tan(\pi - \alpha) = -\tan(\alpha)$
$\cos(\pi + \alpha) = -\cos(\alpha)$	$\sin(\pi + \alpha) = -\sin(\alpha)$	$\tan(\pi + \alpha) = \tan(\alpha)$
$\cos(\frac{\pi}{2} - \alpha) = \sin(\alpha)$	$\sin(\frac{\pi}{2} - \alpha) = \cos(\alpha)$	$\tan(\frac{\pi}{2} - \alpha) = \cot(\alpha)$
$\cos(\frac{\pi}{2} + \alpha) = -\sin(\alpha)$	$\sin(\frac{\pi}{2} + \alpha) = \cos(\alpha)$	$\tan(\frac{\pi}{2} + \alpha) = -\cot(\alpha)$

8.10.3 Exact value of trigonometric functions:

α deg	α rad	$\cos(\alpha)$	$\sin(\alpha)$	$\tan(\alpha)$	$\cot(\alpha)$
0°	0	1	0	0	—
30°	$\frac{\pi}{6}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$
45°	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1
60°	$\frac{\pi}{3}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$
90°	$\frac{\pi}{2}$	0	1	—	0

8.10.4 Trigonometric function with $t = \tan(\frac{\alpha}{2})$:

$$\bullet \cos(\alpha) = \frac{1-t^2}{1+t^2}$$

$$\bullet \sin(\alpha) = \frac{2t}{1+t^2}$$

$$\bullet \tan(\alpha) = \frac{2t}{1-t^2}$$

8.10.5 Addition Theorems:

$$\sin x = \frac{1}{2i}(e^{ix} - e^{-ix})$$

$$\cos x = \frac{1}{2}(e^{ix} + e^{-ix})$$

$$\tan x = \frac{e^{ix} - e^{-ix}}{i(e^{ix} + e^{-ix})}$$

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 2 \cos^2 \alpha - 1 = 1 - 2 \sin^2 \alpha$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$$

$$\sin 3\alpha = 3 \sin \alpha - 4 \sin^3 \alpha$$

$$\cos 3\alpha = 4 \cos^3 \alpha - 3 \cos \alpha$$

$$\tan 3\alpha = \frac{3 \tan \alpha - \tan^3 \alpha}{1 - 3 \tan^2 \alpha}$$

$$\sin^2 \alpha = \frac{1 - \cos 2\alpha}{2}$$

$$\sin^3 \alpha = \frac{1}{4}(3 \sin \alpha - \sin 3\alpha)$$

$$\sin^4 \alpha = \frac{1}{8}(\cos 4\alpha - 4 \cos 2\alpha + 3)$$

$$\sin^n \alpha = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos((n-2k)(\alpha - \frac{\pi}{2}))$$

$$\cos^2 \alpha = \frac{1 + \cos 2\alpha}{2}$$

$$\cos^3 \alpha = \frac{1}{4}(3 \cos \alpha + \cos 3\alpha)$$

$$\cos^4 \alpha = \frac{1}{8}(3 + 4 \cos 2\alpha + \cos 4\alpha)$$

$$\cos^n \alpha = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos((n-k)\alpha)$$

$$\tan^2 \frac{\alpha}{2} = \frac{1 - \cos \alpha}{1 + \cos \alpha}$$

$$\tan \frac{\alpha}{2} = \frac{1 - \cos \alpha}{\sin \alpha} = \frac{\sin \alpha}{1 + \cos \alpha}$$

$$\binom{n}{k} = \frac{n \cdot (n-1) \cdots (n-k+1)}{n!}$$

8.10.6 Sums and Products:

$$\sin \alpha + \sin \beta = 2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\sin \alpha - \sin \beta = 2 \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$\cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\cos \alpha - \cos \beta = -2 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$\tan(\alpha) + \tan(\beta) = \frac{\sin(\alpha + \beta)}{\cos(\alpha) \cos(\beta)}$$

$$\tan(\alpha) - \tan(\beta) = \frac{\sin(\alpha - \beta)}{\cos(\alpha) \cos(\beta)}$$

$$a \cos(\alpha) + b \sin(\alpha) = A \cos(\alpha - \varphi), \leftrightarrow$$

$$\leftrightarrow A = \sqrt{a^2 + b^2} \& \cos(\varphi) = \frac{a}{A} \& \sin(\varphi) = \frac{b}{A}$$

$$\sin \alpha \sin \beta = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2} [\cos(\alpha - \beta) + \cos(\alpha + \beta)]$$

$$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha - \beta) + \sin(\alpha + \beta)]$$

8.10.7 Inverse Functions:

$$\cos(\arcsin x) = \sqrt{1 - x^2}$$

$$\sin(\arccos x) = \sqrt{1 - x^2}$$

$$\sin(\arctan x) = \frac{x}{\sqrt{x^2 + 1}}$$

$$\cos(\arctan x) = \frac{1}{\sqrt{x^2 + 1}}$$

$$\tan(\arcsin x) = \frac{x}{(1 - x^2)^{1/4}}$$


$$\tan(\arccos x) = \frac{(1 - x)^{1/4}}{x}$$

8.11 Hyperbolic Functions:

$$\sinh x = \frac{1}{2}(e^x - e^{-x}) \quad \operatorname{arsinh} x = \ln(x + \sqrt{x^2 + 1})$$

$$\cosh x = \frac{1}{2}(e^x + e^{-x}) \quad \operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$$

$$\tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad \operatorname{artanh} x = \frac{1}{2} \ln \frac{1 + x}{1 - x}$$



$$\cosh^2 x - \sinh^2 x = 1$$

$$\sinh(x_1 \pm x_2) = \sinh x_1 \cosh x_2 \pm \cosh x_1 \sinh x_2$$

$$\cosh(x_1 \pm x_2) = \cosh x_1 \cosh x_2 \pm \sinh x_1 \sinh x_2$$

$$\tanh(x_1 \pm x_2) = \frac{\tanh x_1 \pm \tanh x_2}{1 \pm \tanh x_1 \tanh x_2}$$

$$\sinh x_1 + \sinh x_2 = 2 \sinh \frac{x_1 + x_2}{2} \cosh \frac{x_1 - x_2}{2}$$

$$\sinh x_1 - \sinh x_2 = 2 \cosh \frac{x_1 + x_2}{2} \sinh \frac{x_1 - x_2}{2}$$

$$\cosh x_1 + \cosh x_2 = 2 \cosh \frac{x_1 + x_2}{2} \cosh \frac{x_1 - x_2}{2}$$

$$\cosh x_1 - \cosh x_2 = 2 \sinh \frac{x_1 + x_2}{2} \sinh \frac{x_1 - x_2}{2}$$

$$\tanh x_1 \pm \tanh x_2 = \frac{\sinh x_1 \pm \sinh x_2}{\cosh x_1 \cosh x_2}$$

8.12 Graph of some common functions:

