The gap between surfaces, not the body size, switches fusion in a pattern-forming chemical medium

- **Type:** research article (measurement, one correction, one methodological result)
- **Version:** v2.0 — one article for the whole path taken after the published previous article (DOI 10.5281/zenodo.23933440,

whose archive carries aggregates up to X155): the crossing and its geometry, what a daughter keeps and whether a merge comes

apart, the local supply, the bridge, the gap that switches fusion, what the corridor forecasts, the full crossing table with

the dispersion it hides, and the unit of observation. It follows the sequence's shape: last time we found this and decided to

check that; here are the results; here is where we move next. v1.0–v1.2 described only the census line

- **Date:** 2026-11-05
- **Status:** draft, prepared for submission; NOT yet submitted, deposited or published anywhere
- **Authors:** Ilya Vladimirov
- **Figures:** `030-figures/`
- **Data behind every number:** `030-figures/tables.json`
- **Primary artifacts:** `morphogenesis/experiments/E09_pond/l1_field_x1*.json` (X156–X198; the exact file and JSON path for

every added number is in the aggregate named by the corresponding [A12]–[A25] entry),

`morphogenesis/experiments/L1_field/x19*.compact.json`

- **Reports (pre-registrations and results):** `../reports/2026-11-04-x1*.md`

Abstract

A chemical medium that forms patterns by itself looks, at a glance, as if it were full of objects that appear, grow, split

and merge. We ask what actually decides whether two such objects fuse into one, and what survives when they do.

We simulate a thin layer of two reacting and diffusing chemicals on a 96 × 96 grid, with the supply of fresh chemical made

uneven in stripes. Two blobs of pattern are placed at chosen positions and followed for 400 steps; the outcome we measure is

whether the two become one. The positions matter because the stripes have a period of 12 cells, so a blob's local supply

depends on where it sits within the stripe, and only **nine** distinct local supplies exist across all positions.

What the distance means. The design places bodies at a chosen distance between their centres, but the medium does not use

that number: from 132 merging runs, the two **surfaces** touch at a median distance of **1.93 cells** (range 0.94 to 4.87), the

same at every body size, and because the field is logged every ten steps that reading is an upper bound which bounds the true

threshold down to a median of **0.7 cells** — a window about one to five cells wide, not a point. The crossing itself is one

number per size: fusion falls below one half at gaps 10, 10 and 9 for body radii 3, 4.5 and 6, it does not move with the size

ratio (a (3, 6) pair crosses between 8 and 10 cells), and it does not move with the inflow rate over three rates tested. One

earlier reading of the crossing — five of five fusions at "gap 11" — is **withdrawn** as a units artefact and replaced.

What a daughter keeps. Shape is inherited about half: the daughter-on-parent slope is **0.5167** for elongation and

0.4811 for width. The discrete state of a body is **re-drawn at every division**, not inherited. The medium adds no length

of its own: the distance to the k-th neighbour is set by the body's inherited radius. A merge comes apart again — a merged body

that divides gives two bodies whose combined area is 0.867 of the parents' — and it divides after 25 steps against a single

body's 40.

What decides fusion. Holding the local supply fixed, the outcome is decided by the **bridge** of chemical between the

blobs: across 144 positions at one supply, the bridge ratio spans 0.006 to 0.454 and the logistic slope of fusion on that

ratio is 6.09 (z = 5.8); at a smaller body size the same census gives 4.85 (z = 2.7) with the **opposite** sign of the

association between supply and fusion. What switches fusion on and off, however, is the **gap between the surfaces** and not

the body size: ordered by the gap, the fusion share runs 0.00, 0.19, 0.40, 0.83, 0.96 as the surfaces come from 12 cells apart

to 8, at two different body sizes, while ordered by size the same five arms are not monotone. A size effect exists — at a

matched gap of 8 the smaller body fuses 0.96 against 0.83 — but it shifts the outcome rather than switching it. The edge is a

cliff, not a slope: at radius 6, moving the gap by one cell takes the share from a third to zero.

What can be read before it happens. The corridor — the concentration on the line joining the bodies, in units of their own

peak — is a size-independent criterion (0.48, 0.49, 0.47 at three sizes) but it is largely a readout of how close to the

crossing the pair is, and where a single curve made a sharp prediction it failed: an interior cell landed in its registered

corridor interval and missed its registered share interval. The number that organises the rate ladder is the measured inflow at

the **colder** of the two bodies (AUC 0.866), not the nominal rate the experiment set (0.587) and not the pair's midpoint

(0.215, anti-predictive). A full enumeration of the series — 57 two-body cells, 772 runs — shows why cells disagree with

themselves: it is the phase of the medium where the pair sits, and the worst cell is reproducible by a fresh draw (12 of 30

against 20 of 49 before) because the dispersion is between-family noise rather than a second mechanism.

And a warning about method. Pooling many simulation runs into one number assumes the runs are independent replicates. We

measure how many independent worlds a pool of 757 runs actually contains: 633, with 549 of them used once. For an association

measured by rank (AUC 0.652 → 0.660) the difference is negligible; for a likelihood-based deviance the same pooling moves the

effect size from 20.7 to 14.3, from above the test's own bar of 20 to below it. Whether a result depends on the unit of

observation has to be checked **per statistic**, not once per paper.

Keywords: reaction–diffusion, neural cellular automata, pattern formation, coalescence, bridge, stripe heterogeneity,

inheritance, reversibility, pre-registration, unit of observation

The line this article continues

Last time. In the previous article of this series — *How lineages begin and end in a pattern-forming chemical medium*

(Zenodo, DOI 10.5281/zenodo.23933440) — we followed bodies of pattern in the same medium and measured four things: the medium

has a **mixing length**, two bodies merge when their surfaces come within about 9 ± 1 cells whatever their size; a fusion is a

merger, both bodies ending on the same step with one body left under a new name; a lineage usually ends by **losing its

name** rather than its cells; and the habit that maximises generations is not the one in which a lineage survives. That work

treated the medium as uniform and asked how far apart two bodies can be before they merge.

Everything it left open, and what we decided to check. The published article closed with a list of questions, and this

article is the attempt to answer them in one place rather than in fragments. Four groups of checks:

1. **The distance it measured was a design label, not a physical distance.** We decided to re-measure the contact itself, on

a finer log, and to ask whether the crossing moves with the body-size ratio or with the inflow rate (§5.1).

2. **What a lineage actually copies.** The published article measured one quantitative trait. We decided to ask the same

question of **shape**, of a **discrete** state, and of whether the medium supplies a length of its own — and to test whether

a merge can be undone (§5.2).

3. **The medium is not uniform.** Since our supply varies in stripes, "how far apart" is an average over positions, and two

pairs at the same distance may sit in different local supplies. We decided to check whether the outcome at a fixed distance

is decided by the **local supply where the bodies sit** rather than by the nominal one, and then what decides fusion once

the supply is held fixed: the **bridge of chemical** between the bodies, or the geometry of the pair (§5.3–§5.5).

4. **Whether any of it can be read before it happens, and whether the pool of runs is what it looks like.** We decided to

enumerate every two-body cell instead of sampling them, to forecast the outcome at an unrun supply, and to measure how many

independent worlds a pool of runs really contains (§5.6–§5.9).

What follows. Section 2 places the checks against what is known; sections 3 and 4 define the medium, the outcome and the

census; **section 5 gives the results of the checks**, in the order of the four groups above — the crossing and its geometry,

what a daughter keeps, the local supply, the bridge, the gap that switches fusion, what the corridor forecasts, the full

crossing table with the dispersion it hides, and the unit of observation; sections 6 and 7 are the limits and what would

overturn the answers; and **section 8 says where we try to move next**.

1. The question

Two blobs of a self-organised pattern sit a few cells apart. Over time the medium between them either thickens into a neck

that pulls them together, or stays empty and keeps them apart. Which is it, and what decides?

The everyday answer is "distance": things that are further apart do not touch. Our first measurement says the answer is not

distance alone but **distance divided by the pattern's own scale** — the medium's supply varies in stripes, so a blob's

behaviour depends on where it sits inside a stripe, not only on how far its neighbour is. The second measurement says that

at equal distance the outcome still differs from place to place, and the thing that decides it is the amount of chemical in

the narrow space between the two blobs. The third says that once both are held fixed, fusion is switched by the width of that

narrow space.

Six concrete questions, each with a way to lose:

5. **Does the local supply decide fusion?** Answer "yes" means: at the same nominal supply, blobs placed at different

positions within the stripe fuse at different rates, and the rate follows the local supply. The claim dies if the

rates are all the same, or if they follow the nominal supply just as well.

6. **At a fixed supply, does the bridge decide the outcome?** Answer "yes" means: within one supply value, the worlds whose

bridge is thicker fuse more often. The claim dies if the outcome does not follow the bridge within a value, or if the

bridge is simply a restatement of the supply.

7. **What switches fusion — the body size or the gap between surfaces?** The claim "the gap switches it" dies if closing the

gap does not turn fusion on, or if the same gap at different sizes behaves differently by more than a shift.

8. **How close do the surfaces actually come, and does that move?** The mixing length of the earlier work was stated as a

centre-to-centre number. The claim "the medium uses a surface gap" dies if the measured contact distance scales with the

body size, or if the crossing moves with the size ratio or with the inflow rate.

9. **What does a daughter keep?** "Shape is partly inherited" dies if the daughter-on-parent slope is near zero or near one;

"the discrete state is inherited" dies unless it survives its shuffled-parent null; and "the medium supplies a ruler" dies

if the neighbour-order slope is set by the body's own inherited radius instead.

10. **Can the outcome be read before it happens, and is a pool of runs what it looks like?** "The corridor predicts" dies if a

single curve through the measured arms misses a cell placed between them, or if the share does not follow it at an unrun

supply; and "the pool is independent replicates" dies if regrouping the runs into worlds moves a statistic across its own

bar.

Nothing in this article depends on a trained network or a learned rule; the medium is the classical two-species system. What

is new here is the measurement design: we enumerate **all** positions modulo the stripe period instead of sampling them, we

check every closure against the medium's own arithmetic, and we count how many independent replicates a pool of runs really

holds.

2. What is already known, and what this adds

Spontaneous pattern formation in a two-component reaction–diffusion system was described by Turing [4], and the two-species

system we use is the standard Gray–Scott kinetics [5, 6]. Learned local update rules — neural cellular automata [1, 2, 3] — are a

modern addition to the same family and are the reason the line of work this article belongs to exists at all: a local rule

trained on a pattern-forming medium reproduces much of its phenomenology [2, 3]. In all of these, the objects one sees —

spots, blobs, "individuals" — are local maxima of a continuous field, not entities, and their merge is a topological event:

two connected components of a level set become one.

The merger of two such objects has been studied mostly in droplets and bubbles, where coalescence is controlled by the

distance between surfaces, the curvature of the necks and the drainage of the film between them [12, 9, 10]. In

reaction–diffusion, the distance at which two spots merge has been treated as a property of the pattern's wavelength [11] and

as a boundary in the parameters of the reaction, and our own earlier work located the transition at about 10 cells from the

seed side [A1].

What is missing, and what this article supplies, is the **conditional** question. "How far apart can two spots be before

they merge?" averages over where the spots sit. If the medium's supply is uneven, the position is a hidden variable: two

pairs at the same distance can differ in whether the space between them fills with chemical at all. We treat that space as

the measured state — the *bridge* — and ask whether the bridge, not the distance, decides the outcome. The answer is that

the bridge does decide, and that the geometric variable that switches fusion is the surface-to-surface gap, with the body

size acting as a small shift on top of it. Spatially varying media have been analysed before [7, 8]; what is new here is

turning the position into a *measured* state rather than a nuisance.

The methodological part belongs to a different literature. Pooling runs as if each were independent is pseudo-replication

[17]; when observations are clustered, a cluster bootstrap gives the honest interval [16]; and the general lesson that a

statistic's value depends on the unit of observation is standard in ecology and epidemiology [16, 17]. We measure the size of

that dependence in our own pool and find that it is not uniform across statistics. Pre-registration and multiplicity control

are the other half of the method [18, 19].

Two further literatures are touched, and neither is claimed as new. **Inheritance of a shape between division events** is the

question of whether a local rule that copies a scalar trait also copies a form; our system copies it about half, which is

neither of the two textbook answers, and the discrete state is not copied at all. **Reversibility of coalescence** is the

classical question of whether two droplets that merged can be separated again [12, 9, 10]; in our medium the material returns

to two bodies when the merged body divides, but the lineages do not resume — the distinction between the material and the

identity, which the previous article made its central point, is what separates our case from the droplet literature.

3. The medium and the measurement

The medium. Two chemicals, u and v, react and diffuse on a 96 × 96 grid with periodic borders, integrated with an

explicit step (parameters in `morphogenesis/config.py`; the exact configuration of every run is in the run's manifest). Left

alone it fills with an irregular pattern of spots. The supply of u is made uneven in vertical stripes of period 12 cells:

the nominal supply `F` is multiplied by `1 + 0.5·(2G − 1)` where `G = (sin(2πy/12)·cos(2πx/12) + 1)/2`, so the local supply

runs from 0.5·F to 1.5·F depending on position (Figure 1a). This is the *only* inhomogeneity in the medium.

A world. A "world" is one run with a specified geometry and seed. We build a world by seeding two small discs of pattern

— "bodies" — at chosen positions, separated by a chosen distance. The bodies are rings, not filled discs: their mass sits at

a radius of about 3, 4 or 6 cells, and the centre of a ring is empty, which matters for what we measure (Figure 1a).

Outcome. We follow the run for 400 steps, recording the number of connected components of the field every 10 steps. The

outcome is `fused = 1` if at any recorded step the two bodies are exactly one component, and `0` otherwise. Later, secondary

spots may appear from the medium itself; those are not the outcome but they are why the component count at the end of a run

is not a clean readout of the event.

Bridge. At the moment of closest approach (the last recorded step at which exactly two components exist and the field is

strong enough), we measure the field value at the midpoint between the two centres, divided by the strongest field value

within a radius of 10 cells around that midpoint (Figure 1b). The ratio is near 1 when the neck between the bodies is as

strong as the bodies themselves, and near 0 when the space between them is empty but for a trace. We call it the bridge

ratio; a value above about 0.4 means a neck is present.

Everything is one number, and the lattice matters. The measured ratio is a rasterised quantity: the grid is 96 × 96, the

bodies are discs rasterised on it, and a "gap" of 8 cells is 8 cells as the raster sees them, not as a continuum would. On a


discrete lattice the *existence* of a pattern can depend on the discretisation alone [13], the integration step is a

consequential parameter rather than a resolution knob [14], and rounding error in a simulated physical quantity leaks into

whatever statistic is computed from it [15]. Whenever we compare arms, the comparison is between measured radii and measured

gaps.

Figure 1 — what the measurement is



The words of this article, in one place. A *world* is one run with a given geometry and seed; a *phase class* is a pair of

seeded positions counted modulo the stripe period of 12 cells; the *local supply* is the value of the striped supply function at

the seeded positions; the *nominal supply* `F` is its average over the whole field, the number an experimenter sets; the *gap* is

the shortest distance between the two bodies' surfaces, which differs from the distance between their centres by twice the body

radius; *fused* is 1 when the two bodies are exactly one connected component at any recorded step and 0 otherwise; and the

bridge ratio is the field value at the midpoint between the bodies divided by the strongest value within 10 cells of that

midpoint, read at the last step before they touch. Later sections use short experiment names like "X184" or "radius 4, gap 8";

those are identifiers of runs in the project's own records, listed in the references, and nothing in the argument depends on

knowing them in advance.

4. Methods: the census, and the two kinds of replication

Nine supplies, 144 positions. Because the supply term has period 12 in both directions, the local supply of a body pair

takes only nine distinct values. Enumerating the positions of a pair with a fixed centre separation modulo that period gives

144 phase classes (a "phase" is the pair of positions modulo 12). We run **one world per class** — the whole population

of positions, not a sample — at a fixed separation and body size. This is what makes the conditional question answerable: at

each of the nine supply values we have between 4 and 26 worlds, all at nominally identical settings, differing only in the

hidden phase (Figure 2a).

Pre-registration. Every arm of this line was pre-registered before its run, with its positive control, its readout, its

sensitivity arithmetic and its refusal rule written into `reports/*-plan.md` and into the plan sections of

`docs/plan/10d-field-metrics-and-form.md`. A criterion that could not be tested is recorded as untestable, with the count

that shows why; it is not recorded as failed.

Two kinds of replication, and they are not the same. A run is one execution. A **world** is a pair of positions modulo

the stripe period (a phase) together with the arm's geometry — radius, gap, supply, length. Many runs can share a world: we

found that seeds placed at the same position produce identical outcomes (within-site spread exactly 0.0000), and that five

distinct jitters inside one phase produce the same bridge ratio (0.39916 five times, spread 0.0). In other words a run

repeated at one position is a **clone**; only the phase carries new information. The question "how many independent

observations does a pool hold?" therefore has a measurable answer, and we compute it by collapsing the pool into worlds.

Statistics. All tests are named where used: a one-sided Fisher exact test for a 2 × 2 contrast, Spearman's rank

correlation over classes and over the means of the nine supply values, an iteratively reweighted logistic fit of fusion on

the bridge ratio (slope with standard error and z), a binomial deviance for the predictive comparison, the area under the

ROC curve for the association between local supply and fusion, and a bootstrap over runs and over worlds for the intervals.

Where an outcome has no variation in an arm, no contrast or fit is reported: the criterion is marked untestable.

The arithmetic of sensitivity. Before each arm we wrote down what difference the manipulation could make: the supply band

(0.5·F to 1.5·F), the measured spread of the bridge ratio at one supply (0.14 to 0.45 across arms), the number of worlds per

value, and the bar each criterion had to beat. Arm-level decisions (which value counts as "crowded", how many worlds have to

reach contact) were fixed in the pre-registration, not after the numbers.

5. Results

5.1 How close do two surfaces have to come? The crossing and its own geometry ([A12]–[A16])

The previous article measured the **mixing length** — the distance at which two bodies merge — as "about 9 ± 1 cells",

using the distance between body **centres** as the design variable and the surfaces as the reading. The first thing this

article did was to check that reading against the medium itself, and the medium disagreed with it in an instructive way.

The gap the medium uses is not the gap the design names. From 132 fusing runs, the median distance between the two

surfaces at the moment they touch is **1.93 cells**, with a range of 0.94 to 4.87 ([A12]). At the four nominal radii the

per-radius medians are 2.07, 1.89, 1.69 and 1.93 cells, so the contact geometry has the same shape at every size: bodies

merge when the thin dark layer between them collapses, not when their centres reach a particular number. The nominal "gap 10"

of the design is therefore a label for a pair of positions, not a physical distance; it is the reason this article reports

the **surface gap** wherever it can and says so when it cannot.

That reading is an upper bound, and it is a window rather than a point. Only one field snapshot every ten steps is kept,

so a pair that touched and merged between two snapshots shows a gap that is too wide. X159 ([A13]) bounded the effect: the

unlogged growth widens the threshold downward by a median of **1.32 cells** (90th percentile 2.34), which places the true

threshold's lower bound at a median of **0.7 cells** and turns "the contact gap" into a **window** roughly one to five cells

wide rather than a single number. Both values are reported as bounds; neither is treated as the mechanism.

The crossing itself is one number per size, and it is flat in the design's other axis. On the crossing arms ([A14]) the

fusion share falls below one half first at gaps 10, 10 and 9 cells for radii 3, 4.5 and 6; at radius 3 and gap 10 the share is

0.467 (7 of 15) — the half-way point that the previous article's table and this article's §5.5 both lean on. Asymmetric

pairs do not move it: a pair with sizes (3, 6) crosses between 8 and 10 cells ([A15]), with the surface gap at contact


rising from 2.60 cells at gap 8 to 4.38 at gap 10, inside the symmetric 10 ± 1 band, so

the law is about surfaces and not about matched sizes. Three bodies behave like pairs: a three-body row at gap 8 coalesces in

100 % of runs, at gap 10 in 20 %, at gap 12 in 0 % ([A15]) — near the pairwise 0.47 at gap 10, as a many-body extension of the

pairwise law would predict.

Figure 6 — the distance the medium uses, the crossing, and what does not move it



A band, not a threshold, once the observation is finer. X166 ([A16]) repeated the contact probe on a one-step log rather

than a ten-step grid. Per-arm medians then differ by **2.07 cells** and the within-arm scatter is **0.44 cells**, so the

"threshold" is a band of about that width at this resolution, and a single number for it would be false precision.

One reading was withdrawn here, and the correction is part of the result. X168 asked whether the crossing moves with the

inflow rate, and its aggregate still reports five fusions out of five at "gap 11". The report withdraws the whole item as a

units artefact: a surface gap was passed where a centre separation was expected, so those runs were really at a surface

gap of 5. The corrected series ([A17]) runs gap 11 at three rates with five seeds per cell and finds shares of **0.0, 0.2 and

0.4** at F = 0.026, 0.030 and 0.036 — no dependence detected — and its companion at ten seeds per cell ([A17]) gives 0.0, 0.1 and 0.3,

with a Fisher p of **0.21053** on the extreme pair and a power of only **15 %** for a 0.3 difference. The honest reading is

therefore *no trend detected, and this design could not have found a small one*; the old five-of-five number is not quoted in

this article without that sentence.

5.2 What a daughter keeps, and whether a merge can be undone ([A18]–[A21])

If the medium has a mixing length, then a lineage is a chain of bodies that were once one body, and the interesting question

is what survives the copying. The previous article measured one quantitative trait — the daughter-on-parent slope, 0.9837

over its own range — and this article asked the same question of **shape**, of a **discrete** state, and of the **medium's own

length**.

Shape is inherited about half, and the two habitats exchange which aspect survives. X156 ([A18]) regressed each

daughter's profile on its parent's for two traits in two habitats: the mean slope is **0.5167** for elongation and

0.4811 for width, but the habitats do not agree — elongation 0.6074 against 0.4260 and width 0.3384 against 0.6238,

with disjoint intervals. The mean is therefore a summary of two opposite trades rather than a rate-invariant value, and a

slope near one half says the daughter is pulled half-way back towards its population's shape at each division: neither a

copy nor a fresh draw.

The discrete state is not inherited at all. X161 ([A18]) asked whether the *class* of a body — its integer-valued state,

not its shape — passes from parent to daughter. Division order and lifespan sit inside their shuffled-parent nulls, and the

only trait that reproduces the earlier value is peak area, which is continuous. The conclusion is deliberately narrow: the

discrete state is **re-drawn at every division**, and fertility cannot be tested from lineage trees by construction, because

every lineage in the sample was traced backwards from a division.

The medium adds no length of its own. X162b ([A18]) measured how a body's neighbourhood orders with distance, and the

slope of neighbour order rises monotonically (0.205, 0.236, 0.408, 0.568) because the distance to the *k*-th neighbour is set

by the body's own inherited radius. The medium is not supplying a second, hidden ruler: the ruler is the body. A companion

probe of the spacing in a maze habitat had too little spread to read at all, and is recorded as **not reportable** rather than

as a null result.

A merge comes apart, and the daughters look like the parents. X167 ([A19]) followed every merged body that divided again:

each divides into two bodies whose combined area is **0.867** of the parents' area, near the parents' old positions — so the

operation is reversible in the weak sense that the material returns to two bodies, not in the sense that the two original

lineages resume. Its control ([A19]) compares a merged body with a single body: the merged one divides after **25 steps**

against the single one's **40** (1.12 standard deviations apart; both are two to four logged steps on a ten-step grid), a

difference in *when*, not in *whether*. The seam a merge might leave behind was probed by X169 ([A20]); its registered

bimodality index fails its own control and the item is recorded as **not reportable**, with one auxiliary measurement kept

because it is visible in the pictures: the merged body's waist ratio is **0.357** (median over steps 50–120), a dumbbell while

the field lasts.

And the corridor depends on where the pair sits. X170 ([A21]) repeated the corridor measurement at three inflow rates and

found the normalised concentration at contact to be **0.52, 0.48 and 0.39** at F = 0.026, 0.030 and 0.036 — a difference of

0.12 against a within-group standard deviation of **0.04**. The corridor at contact is therefore a property of the pair's

position as well as of the pair, which is exactly the question the next section takes up in the census.

5.3 The supply that matters is the local one, and it decides fusion ([A2], [A3])

In a purpose-built arm where the nominal supply was held fixed and the position varied, the local supply at the seeded

positions separated the outcomes completely: worlds whose local supply was below the arm's median fused 0.00 of the time and

those above it fused 1.00, with a rank association of 0.982 and $z = +4.64$ (one-sided $p = 1.7 \times 10^{-6}$). This is the

construction that licenses calling the local quantity the driver; it was measured on an arm designed for it, not on the

pooled natural data.

Over the natural pool, the same comparison was made with a binomial deviance: predicting the outcome from the nominal supply

gives a deviance of 464.5, from the local supply 436.5, a drop of 28.0 ([A2]). That number is one of the three headline

numbers of this line; §5.9 shows what happens to it when the unit of observation is changed.

5.4 At a fixed supply, the phase decides the bridge — and the sign of the supply's effect depends on size ([A4], [A5])

Two censuses were run, one at a body radius of 6 with a gap of 9, one at a radius of 3 with a gap of 10. Each is 144 worlds:

the full population of positions at that geometry.

The bridge is not a restatement of the supply. At the most crowded supply value in the radius-6 census — 26 of the 144

phases share it — the bridge ratio spans 0.0056 to 0.4535, a spread of **0.448** (Figure 2b). Since the supply is identical

by construction, that spread is the phase's own doing.

The outcome follows the bridge. Among those same 26 worlds, the 8 whose bridge is above the value's median fused

1.00 of the time and the 18 below it fused **0.111**; the difference is 0.889 with a one-sided Fisher $p < 0.001$. Over all

144 worlds the logistic slope of fusion on the bridge ratio is **6.09** (se 1.06, $z = 5.8$) — a steep dependence: a bridge

ratio moving from about 0.1 to about 0.5 takes the fusion probability from nearly 0 to nearly 1 (Figure 3, left).

The same census at a smaller body carries the opposite sign. At radius 3, the crowded value holds 44 classes with a

bridge spread of **0.401**, the above/below-median contrast is **0.5** (Fisher $p = 0.0001$), and the logistic slope is

4.85 (se 1.79, $z = 2.7$). But the *association between supply and fusion* runs the other way: over classes, Spearman's

rank correlation between bridge ratio and local supply is **−0.378** at radius 6 and **+0.413** at radius 3 (−0.833 and

+0.515 over the nine value means). The bridge explains the outcome in both; what the supply does to the bridge flips with the

size of the bodies (Figure 3, right; Figure 2b).

Figure 2 — the population of phases and the bridge at a fixed inflow





Figure 3 — at a fixed inflow, the bridge decides the outcome



5.5 The gap between surfaces switches fusion; the body size shifts it ([A6])

The two censuses above sit at different sizes *and* different separations, so the sign reversal cannot be attributed to one

of them. Two further arms closed that gap — radius 4 gap 12 and radius 5 gap 10, both with a centre separation of 20 cells —

and produced a surprise: **not one of their 288 worlds fused**, although every one kept a measurable bridge (spreads 0.139

and 0.180). Read naively, that is a window of body sizes in which fusion never happens.

The window moves when the gap is varied instead. The next experiment changed the surface-to-surface gap at two sizes: radius

4 with the gap closed from 12 to 10 and to 8, and radius 6 at the very centre separation of 20 where radius 4 fuses nothing

— which for radius 6 is a gap of 8. The result (Figure 4) is a monotone table in the gap and a non-monotone table in the

radius:


radius	gap (cells)	centre separation	worlds	fused	fusion share
4	12	20	144	0	**0.00**
4	10	18	144	34	**0.19**
6	9	21	144	58	**0.40**
6	8	20	144	126	**0.83**
4	8	16	144	138	**0.96**

The share column is the **mean over the nine supply values** of the per-value fusion share (the arms differ in how the 144

classes distribute over those values), not the raw fraction of fused worlds; the fused column gives that raw count on the

same rows (34/144 = 0.24 and 126/144 = 0.88 for the two arms where the two differ most), and the two orderings agree.

Figure 4 — the gap between the surfaces switches fusion; the size only shifts it



Radius 4 fuses nothing at a separation of 20 and 0.96 when the surfaces come to 8 cells apart; radius 6 fuses 0.83 at the

same separation of 20 where radius 4 fuses nothing, because its gap there is 8 rather than 12. Ordered by gap the share

rises monotonically as the surfaces approach, at both sizes. The size is not absent — at a matched gap of 8 the smaller body

fuses 0.96 and the larger 0.83 — but it shifts the outcome by about a tenth while the gap switches it. This agrees with the

earlier seed-side boundary work, which placed the transition at about 10 cells [A1]. All three pre-registered criteria of the

gap experiment passed, and the earlier reading of the fusion-free arms as a *size* window is withdrawn in favour of the gap

reading (see the Changelog).

5.6 The corridor at contact, and what it does and does not forecast ([A22])

Everything above measured fusion **after** it happened. The corridor is an attempt to read the outcome **before** it happens:

the concentration along the line joining the two bodies, divided by the peak concentration of the bodies themselves, at the

last logged step before they touch. If the corridor at contact is a fixed fraction of the peak, then a run's fate can be read

off a single number.

The corridor is a size-independent criterion. X171 ([A22]) measured it at three radii and found medians of **0.479**,

0.4918 and **0.4701** (radius 3, 4.5, 6) with spreads of 0.036, 0.0216 and 0.0261. The largest size-to-size difference is


0.0217, inside the pooled within-group spread of **0.036**, so a merge happens at a roughly constant fraction of the body

peak whatever the body's size — a geometric criterion, not a size-specific one. (A caveat carried in the open: the artifact

stores 0.036 for that pooled spread while the report and the plan prose say 0.040; the stored field is what is quoted here,

and the discrepancy is listed in the verification section rather than smoothed over.)

Figure 7 — what the corridor reads, and the edge where reading stops



But the criterion depends on the initial gap, and the two axes are the same variable. X174 ([A22]) fixed the size and

varied the initial gap: the corridor medians are **0.3666**, **0.4107** and **0.479** at gaps 8, 9 and 10, a difference of

0.1124 against a pooled within-gap spread of **0.0595** — the largest difference in this block, and outside the spread.

The fusion shares on the same three arms are 1.0, 0.8 and 0.467. Across the eight arm-cells the corridor and the share move

together (Pearson **−0.9492**, Spearman **−0.881**), which says the corridor is largely a **readout of how close to the

crossing the arm is**, not an independent predictor. One defect is recorded rather than hidden: two of those 8 entries are

the same arm (radius 3, gap 10, F = 0.03) entered twice, so the correlations rest on 7 distinct cells and the eighth

repeats one of them.

Where a single curve makes a sharp prediction, it fails. X175 ([A22]) took the interior cell that sits between the

measured arms: the corridor landed at **0.4373** inside its registered interval [0.42, 0.46], but the share came out **0.2667

(4 of 15)* *against a registered interval of [0.45, 0.72]. Worse for the single-curve reading, three arms at the* same* share

of 0.27 have corridors of 0.437, 0.492 and 0.516. A monotone curve through the arms cannot pass through those three points, so

the corridor is not a sufficient statistic for the share.

The forecast itself is partial. X172 ([A22]) predicted the outcome at an unrun supply from the fitted relation: the

corridor was predicted at 0.436 ± 0.08 and observed at **0.442**, a hit, but only **8 of 20** pairs fused against a predicted

9–15. The corridor interpolates; the share does not.

And two probes came back empty in a way that confirms the premise. X176 ([A22]) ran the cell one gap past radius 6's

crossing and found **0 of 15** contact events (0 of 25 pooled with the earlier arm), and X177 ([A22]) ran one gap past radius

4.5's crossing and found **0 of 20** (0 of 35 pooled). In both the corridor field is not measurable because nothing touched:

the "marginal but alive" cell the band's upper edge would need does not exist at one cell past the crossing. The edge is a

cliff, not a slope: at radius 6 and gap 9 the share is **10 of 30 = 0.3333** with a Wilson interval of [0.1923, 0.5122],

against 3 of 10 in the earlier arm at the same cell and 0 of 15 at gap 10. A change of one cell in the gap takes the share

from a third to zero.

The best single threshold on the bridge misclassifies a quarter of the runs. X173 ([A22]) asked whether the bridge's

maximum, read before contact, can separate fusing from non-fusing pairs. Its best threshold, 0.4284, misclassifies **7 of 30**

runs. The item's own registration declines to call this decidable, because the world's proliferation closes the window in

which the seeded pair is identifiable: the pair can be followed for only 10–27 of 301 sampled steps, up to step 90–260 of

3000.

Finally, the rate ladder — and which number on it is the real one. The nominal supply F is what the experiment sets; the

census gives every body a measured local supply. X183 ([A22]) put 5 candidate readings of "the rate this run had" against

the outcome over 139 ladder runs. The nominal rate gives an AUC of **0.5874** (indistinguishable from chance within a level,

mean within-level AUC **0.5**, deviance **185.0**); the inflow at the **colder** of the two bodies gives **0.8656** (within

levels 0.8951, deviance **129.3**); the inflow at the warmer body gives 0.6359; the mean of the two sites gives 0.8396; and the

midpoint of the pair gives **0.2149** — anti-predictive, because the midpoint of a pair seeded across a stripe often sits

off the bodies' own phase. X182 ([A22]) confirmed the midpoint reading with a design that chose the phase in advance: the

share falls from 1.0 at local G 0.0863 to 0.0 at local G 0.8943, Cochran–Armitage z = **−4.446**. So the single number that

organises the ladder is the inflow **at the colder body**, not the number the experiment set, and not the pair's midpoint —

the same conclusion §5.3 reached from the other direction.

5.7 The full crossing table, and the dispersion it hides ([A23], [A24])

The arms above are single cells of a larger object, and the last block of this article enumerated it rather than sampling it.

The table. X179 ([A23]) aggregated every projection in the series and kept the cells that seeded exactly two bodies:

57 cells, 772 runs, 18 further cells with one or three components (263 runs) listed separately. Each cell carries its

radius, gap, rate, fused and total runs, share, a Wilson interval, and the spread **between families** at that cell. Three

cells are flagged **heterogeneous**, meaning their families disagree by more than 0.4 in share: (radius 3, gap 10),

(4.7, gap 9) and (5.9, gap 7). The control cells behave as the earlier arms did: (radius 3, gap 8) fuses **48 of 48 = 1.00**,

(radius 6, gap 10) fuses **0 of 24 = 0.00**. Two honest notes belong with the table. First, the file scan behind it is a


compatibility count: the artifact's `files_scanned` is **1119** and its two-body run total **772**, while the report and the

plan document say 985 and 722; the artifact is authoritative and both are recorded, because a reader who checks the report's

sentence will find the other number. Second, the radii the census *measures* are 2.9, 3.0, 4.4, 4.7, 5.3, 5.9 and 6.0, so the

nominally identical "radius 4.5" arms split into two rows — the size axis is slightly finer than the design's labels.

Figure 8 — the whole table, the dispersion, and which number on the ladder is real



Why 57 cells disagree with themselves. X180 ([A24]) asked what varies *within* a cell. In the 17 cells where both outcomes

occur, over **316 runs**, the phase of the **midpoint** between the bodies is anti-predictive of fusion: the mean point-biserial

correlation is **−0.552** (pooled **−0.527** over the 321 non-saturated runs), only **1 of 17** cells has its fusing runs at the

higher midpoint G, and the two-proportion z between the bottom and top midpoint-G quartiles is **−7.807**. Sorted by the

measured local inflow instead, the same runs order cleanly: **0.836** (46 of 55) in the lowest band, 0.643, 0.248, and **0.16**

(13 of 81) in the highest. The dispersion between families in a cell is therefore not a hidden law of the pair — it is the

phase of the medium where the pair happens to sit, and the informative site is the colder body, exactly as §5.6 found.

And the worst cell is reproducible by a fresh draw. X181 ([A24]) re-ran the cell with the largest family disagreement

(radius 3, gap 10, F = 0.03) with thirty fresh runs: **12 of 30 = 0.40**, Wilson [0.2459, 0.5768], inside its registered band

of [0.23, 0.59] and inside the six earlier families' span of 0.20 to 0.80. Pooled before the redraw the cell read **20 of 49 =

0.4082**, after it** 32 of 79 = 0.4051**. A cell whose families ranged from 0.20 to 0.80 has a pooled value stable to two

thousandths under a fresh family, which is the strongest statement in this block that the dispersion is between-family noise

rather than a second mechanism.

5.8 The bridge measured a second way: what held, and what was withdrawn ([A25])

The bridge entered this article twice. In §5.4 it is the mid-line concentration between two bodies at a **fixed** supply. Here

it is measured from the census instead — each body's own local inflow, warm against cold — because §5.3 and §5.6 both say the

local supply is the variable that matters.

The registered reading holds in direction, and refutes its own second half. X189 ([A25]) compared the tightest bridge

ratio between the two bodies in a warm and a cold version of the same pair. At radius 6 the warm bridge sits **−0.08655** below

the cold one, past the 0.05 bar; at radius 3 the difference is **+0.00387**, i.e. nothing. The direction therefore holds at the

larger body and fails at the smaller one, and the number that carries the headline — **−0.20158** at contact — is an

unregistered secondary readout. Two further caveats are recorded rather than argued away: 9 of the twelve runs have their

tightest saved slice at step 40, so the reading is contaminated by the initial condition, and each level is effectively **one

or two distinct lattice sites** (radius 6: two cold, one warm; radius 3: two cold, one warm), so the "warm versus cold" contrast

is site versus site at n = 3.

The nucleation reading was withdrawn outright. X188 ([A25]) asked the same question of a different observable — how much

material accumulates near a body — with a registered window. Its registered value gives warm minus cold of **+0.234** at

radius 6 against a refusal bar of 0.5, i.e. a refusal; but its own free parameters move the same estimate from **−0.781** to

+0.394 and flip its sign, and the readout is positive in all 24 runs regardless of outcome. The claim "warm carries more"

is therefore **withdrawn**, and the item is kept in the record as a design lesson: the observable was not a function of the

thing it was supposed to measure.

5.9 The unit of observation, and a published effect that does not survive it ([A8], [A9])

The line's pools mix runs, and runs are not independent: seeds at one position are clones, and so are jitters inside one

phase [A10]. We counted the independent units in the largest natural pool — the runs accumulated for the deviance comparison of

§5.3, none of them laid out to study a chosen phase. The 757 runs collapse to **633 worlds**: the largest world holds 5 runs,

549 of them hold exactly one run, and only 2 worlds contain both outcomes. Replacing runs by worlds

therefore should barely change a pooled statistic — and for one statistic it does not:

statistic	per run	per world
AUC of the local-supply → fusion association	0.6517	0.6596
bootstrap sd of that AUC	0.0207	0.0219
binomial deviance drop (nominal → local supply)	**20.7**	**14.3**
share of bootstrap resamples with drop above 20	**0.517**	**0.208**

For the rank statistic the intervals differ by 6 % (0.0207 → 0.0219; the pre-registered refusal was a ratio of 1.3). For the

deviance — the statistic behind §5.3's headline drop — the same regrouping moves the number from 20.7 to 14.3, from **above**


the test's own bar of 20 to **below** it, and the phase-clustered bootstrap clears that bar in only 21 % of resamples against

52 % for runs (Figure 5). The pool here is not identical to the one the published 28.0 came from (later arms selected on

supply are excluded), so the like-for-like comparison is the pair in the table above; even so, the conclusion is that the

deviance drop is carried partly by repeated runs.

Figure 5 — the unit of observation has to be checked for each statistic



This is not a contradiction of the local-supply law. That law rests on the designed arm of §5.3 (0.00 → 1.00 by

construction) and on the two windows of the censuses, not on this deviance. What it shows is that a pooled effect can be

inflated by the unit of observation — and that the size of the inflation depends on the statistic.

6. What this does not show

- **No analytics.** Nothing here is derived from the reaction–diffusion equations. The statements are about measured

outcomes in a rasterised medium at one grid, one integration step and one horizon of 400 steps.

- **One world per phase.** The population of phases is complete (144 of 144), but each phase is measured once. The claim

that a phase is a unit rests on the measured zero spread within a site and within a phase, not on replicates of the whole

census.

- **Geometry is moved by moving the bodies**, which changes which cells they occupy; the gap table is therefore a statement

about this band of geometries, not a continuous law of separation. The two sizes are not compared at one matched gap for

every phase, so the size shift is a difference between two samples of phases.

- **The bridge is one scalar.** It is the value at the midpoint over the strongest value within 10 cells, at one recorded

step; other neck measures (width, curvature, flux) were not tried.

- **One nominal supply per arm** (F = 0.03 for the censuses) with logging every 10 steps, so the contact moment is known to

within 10 steps.

- **The deviance comparison is within our own pool**, with the test's own bar of 20 kept for comparability rather than

re-derived.

- **The bridge ratios cannot be re-derived from the committed tree.** An independent reviewer recomputed every per-world

outcome and ratio from the raw field arrays and matched all 1008 of them, but those arrays are not committed (the compact

slices carry only the component list: step, id, area, centre). A reader outside this machine can verify the outcomes and the

tables, not the bridge itself. Fixing that requires the contact-step field (or the contact ratio) to travel with the

aggregate, which the next census will do.

- **The contact distance is an upper bound.** With one snapshot every ten steps, a pair that touched between snapshots shows a

gap that is too wide; §5.1 quotes the bound and the window it produces, not a threshold. A one-step snapshot at contact would

pin the window, and that run has not been done.

- **One reading was withdrawn and one is not reportable.** X168's dependence of the crossing on the inflow rate is a units

artefact and is replaced by X168c/X168d, whose own design could detect only a large difference (power 15 % for 0.3). X169's

seam and X162's maze spacing failed their own controls and are recorded as not reportable rather than as null results.

- **The crossing table's counts come from a scan, not from a re-run.** Its aggregate stores 1119 files scanned and 772

two-body runs, while the report and the plan say 985 and 722; the stored fields are quoted here and the discrepancy is

recorded. Its radii are measured (2.9–6.0), so nominally identical arms appear as separate rows.

- **The dispersion block excludes 84 runs** whose seeds were chosen by phase in other items, and the between-family spread is

bounded from a redraw of one cell (thirty runs bound it to about ±0.18), not from a redraw of every heterogeneous cell.

- **The bridge against the local inflow rests on one or two lattice sites per level** (X189: two cold and one warm at radius 6,

two and one at radius 3), with nine of twelve tightest slices taken at step 40, and its contact-moment number is an

unregistered readout. X188's nucleation observable is withdrawn because its own free parameters flip the sign.

7. What would change the answer

11. **A controlled size contrast at one fixed gap.** Run the census at radii 3, 4, 6 with the gap held at 8 cells for every

phase class. If the size shift is not of the order of a tenth, §5.5's second half is wrong.

12. **The gap boundary on a finer grid over all classes.** Gaps of 9, 10, 11, 12 at one radius, every phase: a step at 10

cells confirms the boundary reading; a smooth curve replaces "boundary" with a rate.

13. **A two-seed census.** Two seeds per phase class prices the class unit directly instead of inferring it from clones; if

within-phase spread is not zero, §5.9's unit count is wrong.

14. **A world random effect.** Refit the deviance with a random effect for world (a mixed model over cells). If the drop

returns above the bar, the inflation in §5.9 is an artefact of collapsing rather than a property of the unit.

15. **A different neck observable.** Replace the midpoint ratio with a flux through the neck. If the supply–fusion sign

reversal at small bodies disappears, the bridge is not the deciding variable.

16. **A one-step snapshot at the contact moment.** If the true surface gap at contact has a median below 0.5 or above 4 cells,

the window of §5.1 is wrong and the "mixing length" of the earlier article needs restating in surface units.

17. **Shape heredity at a third rate and on other traits.** If the slope moves with the inflow rate, or if it is 0.5 only for

elongation and width and not for other forms, §5.2's "about half" is a property of two traits rather than of copying.

18. **The corridor on an unrun size and gap.** Fit the corridor relation without one cell and predict that cell's share; if the

share misses its interval, the corridor is a readout of position and not a predictor, which is what §5.6 already suspects.

19. **A mixed model for the between-family dispersion.** If a cell's families stop disagreeing once the site is treated as a

random effect, §5.7's "between-family noise" is the whole story; if they still disagree, there is a site-quality variable

that this article has not measured.

20. **A nucleation observable that is a function of the outcome.** X188's withdrawal is a statement about the observable, not

about nucleation: an observable that separates fusing from non-fusing runs by construction would settle it.

8. Where we move next, and the article after this one

This article is one step of a sequence, and the sequence is the point: each step is a set of runs, a pre-registered question, a

result written down whether or not it is convenient, and a new question that the result creates. The steps leave a trail in the

project's own records — every arm has a dated plan written before its run — and the whole sequence is mapped in `docs/series.md`,

with the published and un-deposited steps distinguished. This step's trail is §5; the step before it is the article of the

previous section, whose published version is the DOI quoted there.

One question from §7 is running now, and it is the one this step created. The finer gap grid (item 2) runs five gaps at

radius 4 and three at radius 6 over every phase class — 1152 worlds — and turns the five-row table of §5.5 into a curve,

replacing "at some point between 9 and 8 cells" with the pair of adjacent gaps the switch falls between. Its criteria were

registered before it ran: the share must fall monotonically, at least one neighbouring pair must differ by 0.4 with intervals

that miss the bar, the radius-6 curve must not lie to the right of the radius-4 curve and must be at least 0.10 higher somewhere,

the phase spreads must exceed 0.05, and the three arms that repeat an earlier measurement must land within ±0.05 of it. Its

refusal rules are written down in the same file, so a flat curve is a result and not a failure to report.

What the next article will be. The result that reorders the work is the one in §5.7 and §5.8: the outcome at a site is

decided by the *inflow at the colder body*, and everything else about that site — the midpoint phase, the family a run belongs

to — is a way of averaging that number badly. The next step is therefore to make the local reading **predictive rather than

descriptive**: fit the corridor and the colder-body inflow on part of the table and forecast the outcome on cells that were not

used to fit it, at a size and a gap the fit has never seen. If the forecast lands, this line has a measurable local variable

with a fixed relation to the outcome, which is a different kind of claim from everything in this article. Alongside it, the two

questions that need runs rather than refitting — the two-seed census and the one-step contact snapshot (items 3 and 6) — price

the two units the whole article rests on.

9. Reproducibility

Every number in this article is written into `030-figures/tables.json` by `scripts/make_article_030_figures.py`, which reads

only committed artifacts. The figures are rebuilt by the same script:

```
python scripts/make_article_030_figures.py     # writes 030-figures/fig1..fig8 and tables.json
python scripts/check_article_030_numbers.py    # re-derives every number and checks it against the article text
```

The censuses and their analyses are produced by `scripts/x192_phase_census.py`, `scripts/x193_phase_census_r3.py`,

`scripts/x194_phase_clusters.py`, `scripts/x195_sign_sweep.py`, `scripts/x196_gap_or_size.py` (each with a `_analyse`

companion), `scripts/x197_deviance_worlds.py` and `scripts/x198_gap_curve.py`. Each run writes a manifest under `lab/runs/`

and a row in the canonical journal; each arm has a pre-registration under `reports/` dated before its run. Raw per-world

slices are committed as compacted JSON under `morphogenesis/experiments/L1_field/`.

How the numbers added in v2.0 were checked, and the six places where the records disagree. The sections about the

crossing, inheritance, the corridor, the table and the dispersion take their values from the aggregates listed in

[A12]–[A25]. Those values were transcribed in a separate reading pass that recorded, for every number, the file and the exact

JSON path it came from, and recomputed 361 of them from the stored fields rather than from the reports' prose ([A26]). The

pass found six disagreements between a report's sentence and the stored field, all of them recorded here rather than smoothed

over. In each case the **stored field is what this article quotes**, because the aggregates are the artifacts:

21. **X171's pooled within-group spread** is `0.036` in the artifact and "0.040" in the report and in the plan's prose; the

artifact's own verdict string says 0.04 as well. This article quotes **0.036** in §5.6 and says so there.

22. **X179's scan counts** are `files_scanned` = **1119** and `two_body_runs` = **772** in the artifact, against "985

projections" and "722 runs" in the report and in `docs/plan/10d-field-metrics-and-form.md` §10in. §5.7 quotes the artifact

and names both.

23. **X174's correlation** is computed over eight arm-cells, two of which are the same arm (radius 3, gap 10, F = 0.03); the

report's table lists seven rows. §5.6 quotes the artifact's coefficients and states the duplication rather than presenting

eight independent cells.

24. **X156's pair denominators** are `pairs_total` = 2284 (f030) and 2022 (f040) in the artifact, against "2 256" and "1 869"

edges in the report. The slopes this article quotes (0.5167, 0.4811) are computed over the pairs the artifact lists.

25. **X160's clonal fraction** is reported as `clonal_fraction` 0.834/0.809/0.836, which are the artifact's `clonal_mean`; the

artifact's `clonal_median` is about 0.15 higher. X160 enters this article only as a negative result, and no clonal number

from it is quoted.

26. **X169's waist ratio** is `pair_ratio_median` = **0.357** in the artifact, while the median recomputed from the artifact's

own twelve listed values is 0.345, and that list contains two zeros against the report's stated range of 0.25 –0.63. §5.2

quotes the stored median and calls the item not reportable, so the discrepancy cannot propagate into a claim.

Two further record-level facts belong here because a reader will notice them. **X168 is withdrawn**: its aggregate still

reports five fusions of five at "gap 11", while the report withdraws the item as a units artefact; this article quotes the

corrected X168c/X168d instead and never the withdrawn number as evidence (§5.1). And **X159 has no aggregate of its own**: its

values are stored in `l1_field_x157_quantum.json`, as the X159 report states; [A13] names that file.

What the independent review of the added blocks changed. The forty-two numbers added in v2.0 were recomputed item by item in

`lab/reviews/2026-11-05-030-v20-numbers.md` (36 items, with an explicit list of what could not be checked from the stored

aggregates). It found **six places where this article's prose was wrong**, and all six are corrected here rather than argued

with: X168c runs five seeds per cell and only its companion X168d runs ten (§5.1); X173's followable window is 10–27 of 301

sampled steps and reaches step 90–260, not 13–18 and 120–170 (that window exists only in the report's prose — the aggregate

stores no field for it, so this article quotes the reviewer's recomputation from the ten committed compact slices); X156's two

habitats **exchange** which aspect survives (0.6074 against 0.4260 for elongation, 0.3384 against 0.6238 for width, disjoint

intervals), so the mean 0.5167/0.4811 is a summary of two opposite trades and not a rate-invariant value (§5.2); X183 tested

5 candidate readings, not seven (§5.6); the free-parameter sensitivity of the X188 reading runs from −0.781 to **+0.394**

as stored, the aggregate's own note writing 0.395 (§5.8); and the daughters' combined area is **0.867** of the parents', not

equal to it (§5.2, §5.3). One item the review flagged without counting it as an error is kept as it stands: X171's stored

`pooled_sd` 0.036 is exactly the radius-3 spread, so it may not be pooled at all — the article quotes the stored field and

names the report's 0.040 beside it.

References

External literature. Numbering follows the verified list in `030-references.md`, which records for

each entry the command or manifest key by which its existence was checked; entries that could not be verified there are not

cited here. Two entries rest on corpus documents at manifest status `partial` (not read end to end): [3] and [20].

27. A. Mordvintsev, E. Randazzo, E. Niklasson, M. Levin. "Growing Neural Cellular Automata". *Distill*, 2020. DOI: 10.23915/distill.00023.
28. W. Gilpin. "Cellular automata as convolutional neural networks". *Physical Review E*, 2019. DOI: 10.1103/PhysRevE.100.032402.
29. H. Kojima, T. Ikegami. "Implementation of Lenia as a Reaction-Diffusion System". *The 2023 Conference on Artificial Life*, 2023. DOI: 10.1162/isal_a_00638.

30. A. M. Turing. "The chemical basis of morphogenesis". *Philosophical Transactions of the Royal Society B*, 1952. DOI: 10.1098/rstb.1952.0012.
31. P. Gray, S. K. Scott. "Autocatalytic reactions in the isothermal, continuous stirred tank reactor". *Chemical Engineering Science*, 1984. DOI: 10.1016/0009-2509(84)87017-7.
32. J. E. Pearson. "Complex Patterns in a Simple System". *Science*, 1993. DOI: 10.1126/science.261.5118.189.
33. D. L. Benson, P. K. Maini, J. A. Sherratt. "Unravelling the Turing bifurcation using spatially varying diffusion coefficients". *Journal of Mathematical Biology*, 1998. DOI: 10.1007/s002850050135.
34. T. Biancalani, D. Fanelli, F. Di Patti. "Stochastic Turing patterns in the Brusselator model". *Physical Review E*, 2010. DOI: 10.1103/PhysRevE.81.046215.
35. H.-G. Purwins, H. U. Bödeker, Sh. Amiranashvili. "Dissipative solitons". *Advances in Physics*, 2010. DOI: 10.1080/00018732.2010.498228.
36. H. U. Bödeker, A. W. Liehr, T. D. Frank, R. Friedrich, et al. "Measuring the interaction law of dissipative solitons". *New Journal of Physics*, 2004. DOI: 10.1088/1367-2630/6/1/062.
37. W. N. Reynolds, S. Ponce-Dawson, J. E. Pearson. "Self-replicating spots in reaction-diffusion systems". *Physical Review E*, 1997. DOI: 10.1103/PhysRevE.56.185.
38. J. Eggers, J. R. Lister, H. A. Stone. "Coalescence of liquid drops". *Journal of Fluid Mechanics*, 1999. DOI: 10.1017/S002211209900662X.
39. Q. T. Davis. "Discretization-Dependent Dissolution of Gliders in (Dis)Continuous Systems: Non-Platonic Self-Organization in Complex Systems". *Innovations in Machine Intelligence*, 2023. DOI: 10.54854/imi2023.01.
40. Q. T. Davis, J. Bongard. "Step Size is a Consequential Parameter in Continuous Cellular Automata". *The 2022 Conference on Artificial Life*, 2022. DOI: 10.1162/isal_a_00526.
41. M. L. Pilat, R. Suzuki, T. Arita. "Dealing with rounding error problems in evolutionary physical simulation". *Artificial Life and Robotics*, 2012. DOI: 10.1007/s10015-012-0038-0.
42. A. C. Cameron, J. B. Gelbach, D. L. Miller. "Bootstrap-Based Improvements for Inference with Clustered Errors". *Review of Economics and Statistics*, 2008. DOI: 10.1162/rest.90.3.414.
43. S. H. Hurlbert. "Pseudoreplication and the Design of Ecological Field Experiments". *Ecological Monographs*, 1984. DOI: 10.2307/1942661.
44. B. A. Nosek, C. R. Ebersole, A. C. DeHaven, D. T. Mellor, et al. "The preregistration revolution". *Proceedings of the National Academy of Sciences*, 2018. DOI: 10.1073/pnas.1708274114.
45. Y. Benjamini, Y. Hochberg. "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing". *Journal of the Royal Statistical Society B*, 1995. DOI: 10.1111/j.2517-6161.1995.tb02031.x.
46. G. K. Sandve, A. Nekrutenko, J. Taylor, E. Hovig, et al. "Ten Simple Rules for Reproducible Computational Research". *PLoS Computational Biology*, 2013. DOI: 10.1371/journal.pcbi.1003285.

This line's own published articles (the step before this one, and its own versions):

- **[P1]** I. Vladimirov. *How lineages begin and end in a pattern-forming chemical medium*. Zenodo, 2026. DOI:

10.5281/zenodo.23933440 — the published previous step (version 2.0; the repository holds a v2.1 with the X161–X170 additions,

not yet deposited as a new version of the same record). The whole sequence, with what is published and what is not, is mapped

in `../docs/series.md`.

This line's own artifacts (each with its pre-registration and its aggregate; see the reports for the numbers).

- **[A1]** X158 boundary measurement: `reports/2026-11-04-x158-crossing.md` (a gap of about 10 cells at radii 3, 4.5 and 6).
- **[A2]** X184, X186: `morphogenesis/experiments/E09_pond/l1_field_x184_*.json`, `l1_field_x186_*.json`; reports `reports/2026-11-04-x18{4,6}-*.md`.
- **[A3]** X185: `reports/2026-11-04-x185-*.md` (one law, two windows).
- **[A4]** X192: `l1_field_x192_phase_census.json`; report `reports/2026-11-04-x192-phase-census.md`.
- **[A5]** X193: `l1_field_x193_phase_census_r3.json`; report `reports/2026-11-04-x193-phase-census-r3.md`.
- **[A6]** X195, X196: `l1_field_x195_sign_sweep.json`, `l1_field_x196_gap_or_size.json`; reports `reports/2026-11-04-x195-sign-sweep.md`, `reports/2026-11-04-x196-gap-or-size.md`.
- **[A7]** X187 (the arm-held grouping used in §5.9): report `reports/2026-11-04-x187-*.md`.
- **[A8]** X194: `l1_field_x194_phase_clusters.json`; report `reports/2026-11-04-x194-phase-clusters.md`.
- **[A9]** X197: `l1_field_x197_deviance_worlds.json`; report `reports/2026-11-04-x197-deviance-worlds.md`.
- **[A10]** X190, X191 (the clone and phase spreads behind the world unit): reports `reports/2026-11-04-x19{0,1}-*.md`.
- **[A11]** X198 (the finer gap grid of §8, running while this edition was written): pre-registration `reports/2026-11-05-x198-gap-curve-plan.md`; aggregate `l1_field_x198_gap_curve.json`.
- **[A12]** X157 (the contact gap, §5.1): `l1_field_x157_gap_geometry.json`; report `reports/2026-11-04-x157-gap-geometry.md`.
- **[A13]** X159 (the unlogged-growth bound on that gap, §5.1): numbers stored in `l1_field_x157_quantum.json` (the X159 amendment of X157); report `reports/2026-11-04-x159-quantum.md`.
- **[A14]** X158 (the crossing per radius, §5.1 and §5.5): `l1_field_x158_crossing.json`; report `reports/2026-11-04-x158-crossing.md`.
- **[A15]** X164, X165 (asymmetric pairs and three-body rows, §5.1): `l1_field_x164_asymmetric.json`, `l1_field_x165_three_bodies.json`.
- **[A16]** X166 (the one-step contact log, §5.1): `l1_field_x166_threshold.json`.
- **[A17]** X168c, X168d, with X168 withdrawn (the crossing against the inflow rate, §5.1): `l1_field_x168c_inflow.json`; the withdrawal is argued in `reports/2026-11-04-x168-inflow-crossing.md`.
- **[A18]** X156, X161, X162b (shape heredity, the discrete state, the neighbour order, §5.2): `l1_field_x156_shape_heredity.json`, `l1_field_x161_discrete.json`, `l1_field_x162b_neighbour_decay.json`.
- **[A19]** X167, X167b (the reversibility of a merge and its control, §5.2): `l1_field_x167_reversibility.json, l1_field_x167b_reversibility.json`.
- **[A20]** X169 (the seam, not reportable, §5.2): `l1_field_x169_seam.json`.
- **[A21]** X170 (the corridor against the source, §5.2): `l1_field_x170_corridor.json`.
- **[A22]** X171–X178 and X182, X183 (the corridor, its forecast, the edge, the cliff, the ladder, §5.6): `l1_field_x171_corridor_sizes.json`, `l1_field_x172_forecast.json`, `l1_field_x173_bridge.json`, `l1_field_x174_gap_corridor.json`, `l1_field_x175_interior.json`, `l1_field_x176_past_crossing.json`, `l1_field_x177_edge.json`,

`l1_field_x178_cliff.json`, `l1_field_x182_local_inflow.json`, `l1_field_x183_local_collapse.json`.

- **[A23]** X179 (the full crossing table, §5.7): `l1_field_x179_crossing_table.json`; report `reports/2026-11-04-x179-crossing-table.md`.
- **[A24]** X180, X181 (the dispersion and its redraw, §5.7): `l1_field_x180_dispersion.json`, `l1_field_x181_worst_cell.json`.
- **[A25]** X188, X189 (the nucleation and bridge readings against the local inflow, §5.8): `l1_field_x188_nucleation_identity.json`, `l1_field_x189_bridge.json`.
- **[A26]** The number digest behind the added sections: every value added in v2.0 was transcribed from the aggregates by a separate reading pass, with the file and JSON path recorded per number; the pass found six disagreements between the reports' prose and the stored fields (X171's pooled spread 0.036 against 0.040, X179's 1119/772 against 985/722, X174's duplicated arm, X156's pair denominators, X160's mean-versus-median clonal fraction, X169's waist median 0.357 against 0.345 from its own list). The stored fields are quoted in this article and the disagreements are listed in §9 and in `docs/series.md`.

Changelog

- **v1.0** (2026-11-04): first edition. Consolidates the phase-census line X184–X197 for an outside reader. Includes the

correction of the "size window" reading (X195) to a **gap** reading (X196), and the unit-of-observation result (X194,

X197) including the fact that the published deviance drop falls from 20.7 to 14.3 when the world replaces the run.

- **v1.1** (2026-11-05): a glossary of the article's own terms in one place (§3), so that the definitions do not have to be

collected while reading; an explicit next-step section (§8) naming what is already running and what the following article

will be; the author line set from the owner's answer (Ilya Vladimirov); a repeated word fixed. No number changed.

- **v1.2** (2026-11-05): the article now opens with the shape the owner asked for — **last time we found this, so we decided to

check that, here are the results, and here is where we move next**. A new opening section states what the previous article in

the series found and the two questions it left open; the published previous step is now cited by DOI, and the un-deposited

v2.1 of that article is named as part of the same record; the sequence is mapped in `docs/series.md`; §8 is retitled to

"Where we move next". No number changed.

- **v2.0** (2026-11-05): **one article for the whole path after the published previous article.** The owner's decision, taken

after asking whether the continuation could be a single piece: yes, and the step is one article rather than four. v1.x

described only the census line (X158, X184–X197); v2.0 opens with the four groups of questions the previous article left

open and answers them in one place, adding **§5.1** (the crossing and its geometry: the contact gap 1.93 cells and the

window it is bounded to, the crossing per radius, the withdrawn X168 and its corrected replacement), **§5.2** (what a

daughter keeps: shape heredity 0.5167/0.4811, the re-drawn discrete state, the absence of a medium-supplied ruler, and the

reversibility of a merge), **§5.6** (the corridor, its failed sharp prediction, the edge and the cliff, and the rate ladder

whose informative reading is the inflow at the colder body, AUC 0.8656 against the nominal 0.5874), **§5.7** (the full

crossing table — 57 two-body cells, 772 runs — and the between-family dispersion, with the worst cell reproduced by a fresh

draw), and **§5.8** (the bridge measured against the local inflow, what held and what was withdrawn). The abstract, §1,

§2, §6, §7, §8 and §9 are rewritten to match, the references gain [A12]–[A26] for the added artifacts, and §9 now lists the

six places where a report's prose disagrees with the stored field, with the artifact quoted in every case. Section numbering

of the earlier results moved: the local supply is §5.3, the bridge §5.4, the gap §5.5 and the unit of observation §5.9.

After the independent review of the added blocks (`lab/reviews/2026-11-05-030-v20-numbers.md`) six sentences of this article

were corrected — the seed count of X168c, X173's followable window, the two-habitat trade behind X156's mean slopes, the

number of X183's candidate readings, the X188 sensitivity end and the X167 area ratio — and the gate grew to **145** needles.

Verification

- **Numbers.** Every number quoted above is re-derived from committed artifacts by

`python scripts/check_article_030_numbers.py --ci` — **145** needles, each of which must also appear verbatim in this text

(run before submission); the table behind them is [`030-figures/tables.json`].

- **References.** Existence of every reference is checked in [`030-references.md`] via the corpus

manifest or Crossref/DataCite; entries that could not be verified are marked there and are not cited here.

- **Independent review.** `lab/reviews/2026-11-04-030-numbers.md` — an independent recomputation of the headline numbers by

the reviewer role, including the two statistics whose intervals depend on the unit (date of review in that file).

- **Not published.** No part of this article has been submitted, deposited or posted anywhere. Submission requires the

owner's explicit approval, and the author list is not yet fixed.