How lineages begin and end in a pattern-forming chemical medium

- **Type:** research article (measurement and one negative result)
- **Version:** v2.1 — plain-language edition for an outside reader; adds the unequal-pair, three-body, discrete-trait,

seed-count and field-probe results of 04.11

- **Date:** 2026-11-04
- **Status:** v2.1, released as a preprint together with its data package
- **Figures:** 029-figures/
- **Data behind every number:** 029-figures/tables.json

Abstract

A chemical medium that spontaneously forms patterns is often described as if it contained individuals that inherit, compete and die. We test how much of that description survives measurement.

The medium is a thin layer of two reacting and diffusing chemicals, simulated on a 96 × 96 grid. Left alone it fills with an irregular pattern; here the supply of fresh chemical is made uneven, in stripes, which produces three distinct habitats: a lattice of separate spots, a maze, and a single connected mass. In all three, blobs of the pattern appear, grow, split, merge and disappear.

We asked three plain questions. *Does anything pass from one blob to its descendants? How close can two lineages get before the medium fuses them into one? What actually ends a lineage?*

Four answers. (i) The medium has a **mixing length**, and it is two numbers rather than one: two lineages become a single blob when their surfaces are about **two cells apart at contact** (1.9 cells, median over 132 fusing runs; at one-step resolution this is a band from about one to five cells rather than a single value), while whether they ever touch is set by the initial gap, which stops working between **ten and eleven cells** — flat in the body's size, flat in the size *ratio* (two bodies of radius 3 and 6 cross in the same band), additive over contacts (a row of three coalesces at the square of the pairwise share), and flat in the chemical inflow across the range this world allows. (ii) A fusion is a **merger of two lineages, not a birth**: both lineages end on the same step and one blob appears under a new name, which forces a three-way classification of close encounters (mutual fusion, absorption, persistence). (iii) At the edge of the mixing length the **outcome switches abruptly while the timing spreads**: at body radius 6 the share of runs that end as one blob reads 1.0 / 1.0 / 1.0 / 0.5 across surface gaps of 6.5 / 7 / 7.5 / 8 cells, while the share of slow fusions rises 0.0 / 0.0 / 0.2 / 1.0. (iv) Lineages mostly end by **losing their name, not their cells**: about half of all endings are absorptions (55.3 / 47.8 / 48.4 % in the three habitats), the absorber is a close relative in 87.2 % of the cases in the lattice and in 37.7 % in the connected mass, and — the result we did not expect — **the habitat that produces the most generations is not the one in which a lineage is most likely to still exist at the end**.

On inheritance: the blob-to-descendant slope of the tracked trait is 0.9837 [0.9680, 1.0002] and 0.9931 [0.9756, 1.0103], the scatter around that line equals the model's own noise parameter to within one per cent (0.1487 and 0.1495 against σ = 0.15), and with the noise switched off the trait has no variance at all. The trait is therefore transmitted perfectly by the bookkeeping rule and not by the medium. We say this plainly rather than quoting a heritability number, because the between-lineage variance we can also compute (0.082 in the lattice, 0.416 in the connected mass) counts lineages that happen to still be present at the end, not a property of the trait. A second measurement, on a trait computed from the field rather than written by the rule, changes the sentence by one clause: **shape**

is heritable at about half, and the habitat decides whether it is elongation or width that a daughter inherits (§4.8).

Two further negatives and one open question. Adding a third, slowly diffusing chemical — the most natural way to give the pattern a heritable scale — does not change the pattern's wavelength at all. Nothing **discrete** crosses a division: division order and lifespan have parent-daughter slopes of −0.163 and 0.101, both inside their shuffled-parent nulls, while area as a positive control reproduces 0.546, so the medium offers a digital notation and not a ratchet. And whether a merge erases the two bodies' structure is **open**: the field probe built to see a seam failed its own control, though it showed the merged body is a dumbbell (a bridge four to five cells wide) for as long as the pattern lasts, which in this world is about three hundred steps.

What changed in v2.1

v2.0 was the plain-language edition. v2.1 adds the results of one working block and corrects two sentences that block made false. The crossing's band is now measured rather than bounded: gap 10 and gap 11 were run at three inflow rates with ten seeds per cell (X168c, X168d), so the earlier sentence "the arms between 9 and 11 were never run" no longer holds, and the insensitivity to the source is stated with the seed count that can support it. The contact gap is no longer described as "the same at every body size": a one-step log (X166) shows a band from about one to five cells across arms. Three new results are folded into §4.1 and §4.8 — an unequal pair crossing in the same band (X164), a three-body row coalescing at the square of the pairwise share (X165), and no inheritance of discrete traits (X161) — and one question is recorded as open rather than answered, the seam a merge may or may not leave behind (X169). Every number in this version is in `029-figures/tables.json` with the aggregate it came from, and `scripts/check_article_029_numbers.py --ci` requires each of them to appear verbatim.

1. What this paper asks

Pattern-forming chemical systems are a standard testbed for questions about order, and they are routinely described in the vocabulary of biology: "individuals", "reproduction", "inheritance", "selection". Each of those words is a claim. The claim can be checked — not by looking at pictures, but by defining an event, counting it, and seeing which intervention moves the count.

We concentrate on three claims that are usually left implicit.

First, **inheritance**. If a blob splits, does its daughter resemble it because something is transmitted, or because our own analysis assigned the daughter a value? In this medium the honest answer turns out to be the second, and the useful result is the precise form of the transmission: a slope of one with an error equal to the model's noise parameter.

Second, **identity**. Two blobs of a pattern are not like two animals: they can run into each other and become one. Where the boundary of an individual lies is therefore an empirical question about the medium — the mixing length — and not a matter of definition.

Third, **endings**. A structure can stop existing in several ways: it can be eaten by a neighbour, it can dissolve, or it can be absorbed into a larger body. Which of these dominates decides what "a lineage" even means.

Each of the three is measured below, with an intervention where an intervention is possible (the mixing length) and with pairs of closely related lineages where it is not (endings and inheritance).

2. The medium

Two chemicals, u and v, react and diffuse:

$$\partial u/\partial t = D_u \nabla^2 u - u v^2 + F (1 - u)$$
$$\partial v/\partial t = D_v \nabla^2 v + u v^2 - (F + k) v$$

with $D_u = 0.16$, $D_v = 0.08$, $k = 0.06$, integrated explicitly with $dt = 1.0$ on a 96×96 grid with periodic edges. This is the standard Gray–Scott system, the workhorse of pattern-formation studies: for these parameters it sustains a maze of bright ridges on a dark background, and a uniform F produces either spots or a connected labyrinth depending on its value.

Our only modification is that the supply F is **uneven**: it is modulated by a fixed sinusoidal field of amplitude 0.5 and period **12 cells**, which is what makes the pattern's own wavelength 12 cells. Because the modulation is fixed in space, a run does not drift into one global state; instead the same dish contains a whole range of local conditions. By choosing three supply levels we obtain three habitats that differ only in that level, and that we can therefore compare with every other condition held fixed:

habitat	supply F	what it looks like
spots	0.030	a lattice of separate compact blobs on a dark background
mixed	0.035	spots and short ridges; the transition regime
connected	0.040	one percolating labyrinth, most "blobs" are parts of the same mass

Each run lasts 6 000 integration steps. At every tenth step the analysis records, for each connected region of the pattern above a brightness threshold (we call them **bodies**), its area, its position, and an identifier. Identifiers are the core of the bookkeeping and the reason the results are trustworthy: when a body splits, **both** daughters get new identifiers, and when two bodies merge, the merged body gets a new identifier as well. No identifier ever spans a division or a fusion. A **lineage** is the tree of identifiers descended from one body, and a **generation** is one level of that tree.

The one artificial ingredient is the **trait**. Each body is given a number s around 1.0 — a "composition" — which is copied to its daughters with small random changes of standard deviation $\sigma = 0.15$, and which in turn nudges the local supply F. This is a deliberate stand-in for a heritable character: it is the only thing in the model that is passed from parent to offspring, and the medium itself has no notion of it. Everything in §4.6 follows from taking that seriously.

Box 1 — the words we use.
Body — one connected region of the pattern above threshold.
Surface gap — the distance between the two nearest surfaces of two bodies, $g = d - r_1 - r_2$, where d is the distance between their centres and r_1, r_2 their effective radii. This is the quantity that matters for diffusion, as opposed to centre-to-centre distance, which grows with body size.
Generation (depth) — how many levels of the identifier tree a lineage reaches before it ends.
Sister lineages — two lineages born from the same parent at the same time and place; comparing them cancels everything the two share.
World — one run of the model with one random seed. Ten to thirty worlds per condition are used below, and we report how many are needed.

3. How we measure

Intervention where possible. To measure the mixing length we do not watch a busy world and guess; we place exactly **two bodies and nothing else** in an empty dish, at a chosen centre-to-centre distance, and vary the distance. Every other condition — supply field, threshold, time step, integration length — is the production configuration, and a control run confirms that adding the intervention does not change any other result.

Counting where intervention is impossible. Endings and inheritance cannot be randomised, so we use pairs: two lineages born from one parent (sisters) differ in size but share place, time and habitat, which turns a comparison of strangers into a within-family comparison. Intervals are 95 %,

computed by resampling over parent lineages, never over individual steps, so that a long-lived lineage counts once.

Decision rules written down first. Before each series we wrote the readout, the criterion and the rule for declaring the series uninformative. That matters here because three earlier designs of the same question produced numbers that had to be withdrawn: a rate with two defensible definitions differing sixfold, a "rate difference" that turned out to be a sample filter, and a pairing statistic in which 22.2 % and 21.5 % of pairs are exact ties. All three are reported below where they bear on a conclusion; the withdrawn values are cited only to show what they were, and with the tie rule applied the paired fractions read 0.526 and 0.388.

4. Results

4.1 Fusion happens at contact, and the gap that stops it is ten cells at every size

Two numbers describe this, and the first version of this section reported one of them wrongly and the other as a single figure rather than a bound. All three statements below are measured, and the corrections are recorded where they belong.

What the medium sets — the distance at contact. In 132 runs in which two seeded bodies became one, at the last logged step before they were one the gap between their surfaces was **1.9 cells** (median; range 0.9 to 4.9) — about a sixth of the pattern's 12-cell wavelength, a length belonging to the medium rather than to the pair. That figure came from a log that samples every tenth integration step, and re-running five arms at **one-step resolution** (X166, 2 000 steps, five fresh seeds each) shows what the coarse log had flattened: the typical gap is about one to two cells, but it is not the same at every size — 1.27 at radius 3 at a gap of 8, 1.86 at radius 3 at a gap of 9, 1.07 at radius 5.2, 3.14 at radius 4.5 and 3.01 at radius 6 — and within each arm the widest value belongs to the slowest fusion (2.9 at step 85 against 1.7 at step 60; 7.8 at step 108 against 3.0 at step 68). The one-step log also settles the resolution question: the residual window shrinks from 1.3 cells to **0.06 cells**, twenty-three times smaller, while the spread stays at 2.07, so the "about two cells" is a band by right and not because of our sampling. A pair that takes longer to merge does so from further away, which is what one expects if the merge is driven by a field bridge that has time to build. And it is not the pair's own geometry: a body of radius 3 against one of radius 6 fuses 5/5 at gaps 4, 6 and 8 and 2/5 at 10 (X164), so a twofold difference between the two bodies does not move the crossing out of the same band of ten — the length belongs to the medium and not to the pair; the gap at the last two-body step rises with the initial gap, 1.84 / 2.21 / 2.60 / 4.38 cells, because a pair with further to travel merges later and from further apart. Nor is it a property of an isolated pair: a row of three bodies with the same surface gaps coalesces into one component at exactly the share that independent pairwise events predict — 5/5, 1/5 and 0/5 at surface gaps of 8, 10 and 12, against the 22 % that 0.47 squared gives (X165) — so the length applies per contact, and a third body neither helps nor prevents. The crossing is also insensitive to the source: at inflow rates 13 % below and 20 % above the one it was measured at, a pair separates between ten and eleven cells just the same (gap 10 fuses 2/5, 2/5 and 3/5 at F = 0.026, 0.030 and 0.036; gap 11 fuses 0/5, 1/5 and 2/5 — X168c) — a length that does not move when the source changes by a third belongs to the pattern-former, not to the source. With ten seeds per rate the gap-11 shares are 0/10, 1/10, 3/10, and no contrast clears a thirty-point difference at $p < 0.05$ (the extreme rates give $p = 0.21$ — X168d), so the insensitivity is measured rather than power-limited; the hint that it might rise with the source is recorded and not claimed. Nor is anything discrete carried across: read from 2 778 parent-daughter pairs in the lineage trees the runs already contain, the number of children a body produces has a parent-daughter slope of −0.163 (per-run median exactly zero, ranging from −1.01 to +0.86) and lifespan 0.101, both inside their shuffled-parent nulls (−0.240 to 0.136 for division order and −0.007 to 0.199 for lifespan), while peak area reproduces 0.546 (X161). A digital medium offers a discrete notation, not a ratchet.

That 1.9 is an *upper bound*, not the threshold, and the difference matters at this resolution. The log samples every tenth integration step and the bodies grow inside that quantum, so the pair is known to

be two bodies at 1.9 cells and one body ten steps later; the threshold itself lies inside the window the growth opens. Measured from the growth the log shows, the window is **1.3 cells** wide: the threshold lies between **0.7 and 1.9 cells** at the median, its lower bound stays above two cells in only **21 %** of runs, and at radius 4.5 the lower bound crosses zero. So the defensible sentence is "fusion happens when the surfaces are within about two cells", not "at 1.9 cells" — and to turn the bound into a threshold needs a one-step log, not more seeds.

What the design sets — the gap at which fusion stops being the rule. Two bodies were placed at a series of initial separations and followed for six thousand steps; the outcome is the share of runs that ended as one body, with ten fresh seeds (701-710) at every cell that had not been run before. The crossing — the first gap at which that share falls below one half — is at **10 cells** for radii 3 and 4.5 and at **9 cells** for radius 6, i.e. **10 ± 1 cells**, flat in the seeded size to within the one cell that ten seeds per arm can resolve.

body radius	area (cells)	gap 8	gap 9	gap 10	gap 11
3	29	5/5	5/5	7/15	2/15
4.5	64	5/5	4/5	2/10	0/15
6	113	10/10	3/10	0/10	0/10

Two features of that table are as informative as the crossing itself. The transition is **gradual over two cells**, not a step: radius 6 goes 1.00 → 0.30 → 0.00 and radius 3 goes 1.00 → 0.47 → 0.13. And the larger bodies are **not** more tolerant, which is what the withdrawn `2r + 9` rule predicted (it would have put the crossing at 9, 13.5 and 21 cells): a body seeded at radius 6 crosses one cell *earlier*, not later.

Why the boundary does not scale with the body. Because the bodies grow, and grow to the same size. A body seeded at radius 3 is 2.9 cells in effective radius at the first logged step and 5.8 at the step before it meets its partner; a body seeded at 6 goes from 6.0 to 8.5. By the time two bodies touch they are all between 5.8 and 8.5 cells in radius, whatever we started with, so the initial separation and not the initial size decides (Fig. 2).

The three outcomes of a close encounter are worth keeping in view here, because "one body" is not the same as "mutual fusion": at radius 3 and a gap of 10 the runs that end as one body split into M 2/5 and A 2/5, with P 1/5; at a gap of 11 they are M 2/5, P 3/5, and the shares of M and A are what the crossing table counts together. ("M" is mutual fusion, "A" absorption of one body by the other, "P" the two bodies persisting side by side; §4.2 explains why the three must be told apart.)

The correction, twice. An earlier version of this paper put the crossing at "16 ± 1 cells" for radius 3 and the conserved quantity at "9 ± 1 cells, the same for bodies of 29, 64 and 113 cells", with the rule `2r + 9`. An independent review recomputed the arm counts from the raw worlds and was right on both points: the radius-3 row came from an older test that counted a partner as present when its last known position was within two cells, which is why it read "two of five" where the classification reads "four of five"; and the third row (`20 ± 2`) was a bound between two arms, not a measured crossing. That correction produced the band "between 8 and 11 cells", which was honest but weak; the arms that could narrow it were then run (§7), and the band became the crossing above. The second correction is the one in this section's first paragraph: the 1.9 cells of the same review-corrected analysis turned out to be a bound set by our own sampling, and it is now reported as a window. Both errors are of the same kind — a number read off a design rather than off the medium — which is why they are stated here rather than in a footnote.

Two controls say what the length is *not*. It is not a size rule: among runs where both outcomes occur, the bodies that divided are not the larger ones — median areas at fusion 267 / 268 and 297 / 287 cells, and where the pattern reverses, 242 / 290. And it is not simply the clock running out: at radius 6 the median time to a single body is 39 steps at gaps 6.5 to 7.5 and 49 steps at 8, while a lone body of the same size does not divide at all until a **median of 290 steps**; the smaller sizes behave the same way — 180 and 190 steps for radii 3 and 4.5 — so the fusing pair is always faster than the division time, at every size.




Figure 1. What is measured, drawn rather than photographed. (a) Two bodies of radius `r` at centre separation `d` leave a surface gap `g = d − 2r`; the results are stated in the gap because that is the quantity diffusion sees. (b) The three outcomes of a close encounter: mutual fusion (both lineages end, one new body appears), absorption (one lineage ends inside the other) and persistence (both continue). (c) The identifier bookkeeping: every split gives both daughters a new name, and every fusion gives the merged body a new name, so no name spans a division or a fusion.




Figure 2. The mixing length, in the two quantities that are measured. (a) Share of runs that end as a single body against the **initial** surface gap, for three seeded sizes, with 95 % Wilson intervals; the crossing is where the curve falls through one half, at 10 cells for radii 3 and 4.5 and at 9 for radius 6. (b) The same runs at the moment of contact: the surface gap between the two bodies at the last logged step before they are one, per seeded size, with the median reading of 1.9 cells and the window that the ten-step logging quantum opens beneath it — the reading bounds the threshold, and the threshold lies inside the window.

What a merge does to the merged body's insides remains open, and it is worth saying why rather than implying an answer: the field-level probe that would settle it (does a seam survive between the parents' positions?) failed its own control, because a body's cross-section is a ring and any intensity profile has a central dip whether there is a seam or not — the single body reads 0.047 against the 0.9 floor the pre-registration set (X169). What the probe did show is that a merged body is a dumbbell: the bridge between the parents' positions is four to five cells wide against lobes of ten to fourteen, a median waist ratio of 0.36 while the pattern lasts, and this world's field fades within about three hundred steps, so the neck never gets the time to close or to persist. The question therefore needs a long-lived field, not a finer measurement of this one.

4.2 A fusion is a merger, not a birth

Our first plan was to map a band of distances at which two bodies "annihilate". Reading the logs killed the word. At separations 10 and 11 the two identifiers die on the **same** logged step, and on the next step there is **one** body under a **new** identifier. Nothing is born: two lineages end together and a third thing starts. The old vocabulary had looked like it was describing a birth only because the analysis counted a partner as present if its last known position was within two cells.

That is why the results speak of M, A and P rather than of "coalescence". It is also why the mixing length is a property of *lineages*: below the fusion boundary, two identities are extinguished at once. Whatever counts as an individual in this medium, it cannot survive contact.

4.3 At the edge, the outcome switches and the timing spreads

At body radius 6, ten runs per gap, the two things separate cleanly (Fig. 3):

surface gap (cells)	**6.5**	7	**7.5**	**8**
runs ending as one body	1.0	1.0	**1.0**	**0.5**
dissolution steps observed	39 ×10	39 ×10	39 ×8, 49 ×2	49 ×3, 59 ×2
slow, among runs that merged	0.0	0.0	**0.2**	**1.0**
slow, among all ten runs	0.0	0.0	0.2	0.5

The share of runs that merge is a step between 7.5 and 8 cells. The share that merge *slowly* — one logged step later than the rest — has already left zero at 7.5, and both denominators are quoted because at 8 only five of ten runs merge at all: the strict figure is 1.0, the conservative one 0.5, and the conclusion (a spread that begins before the outcome changes) holds under both.

One control failed and we keep it in the record: the half-cell arm at 7.5 also moved the drawn bodies by one cell each (110 cells against 108 and 110), so that arm is a gap change *plus* about 2 % of area. The failure works against the finding, not for it.




Figure 3. At the edge of the mixing length. (a) Ten runs per gap at body radius 6: the share ending as one body is a step between 7.5 and 8 cells, while the share that merge slowly has already left zero at 7.5. (b) The same runs by dissolution time; the arms are staircase-shaped rather than spread out.

4.4 Most lineages end by being absorbed by a relative

Classifying an ending is harder than it sounds, and two earlier attempts had to be withdrawn: the first offered a choice between "lethal" and "homogenising", no registered outcome ever fired, and the second was voided by its own controls. The version that works restricts an absorption to the dying body's own neighbourhood — a survivor within `r_dead + r_survivor + 2` cells — and adds a category the lattice needs: a death with no living neighbour at all.

habitat	endings	died alone	absorbed	faded away	absorber was a relative
spots	524	31.3 %	**55.3 %**	10.5 %	**87.2 %**
mixed	844	29.4 %	47.8 %	21.7 %	39.2 %
connected	1 102	21.2 %	48.4 %	**27.4 %**	**37.7 %**

Absorption is the commonest single fate everywhere — about half of all endings — and the remaining two fates trade places with the habitat: dying alone falls from 31.3 / 29.4 / 21.2 % while fading away rises 10.5 / 21.7 / 27.4 %.

The sharper result is *who* does the absorbing. In the lattice, 87.2 % of absorptions are by a close relative; in the connected mass, only 37.7 %. We tried to control this for geometry and could not: every absorption with a partner happens at contact, so matching on distance is degenerate — and the degeneracy is itself the finding. At the same contact geometry, the single neighbour is a relative 87.2 % of the time in the lattice and a stranger 62 % of the time in the connected mass. The lattice keeps families apart; the connected habitat mixes them.

Sisters rarely meet again: 18.4 / 8.6 / 17.4 % of sister pairs re-merge within a run (82 % are censored, i.e. the run ends first), with medians of 80 / 120 / 60 steps after birth, and the distance at birth decides it by a factor of two.

4.5 More generations does not mean more survival

Size buys generations. Sisters that differ in size differ in how many generations their lineages reach — by **+1.479 [1.289, 1.654]** and **+2.213 [1.962, 2.489]** generations in the two habitats where the comparison is defined — and the readout is not a disguised lifetime: with lifetime held fixed the correlation is 0.459 / 0.340, and depth is *negatively* correlated with lifetime itself (−0.185 / −0.070). The advantage survives a per-step rate, an equal time window and a persistence readout; matching sisters on the number of daughters halves it (0.125 and 0.044 against 0.2496 / 0.1372 unmatched), and a two-term model puts area and daughter count in opposite directions, with age the one confound we cannot remove.

Comparing the three habitats separates three things that the two-habitat design had merged (Fig. 4):

habitat	lineages below the generation floor	generations advantage	generations per 1 000 steps	survival advantage
spots	0.162	1.479	6.06	0.2496
mixed	0.302	1.416	9.01	0.2984
connected	0.466	2.213	18.96	0.1372

The generation clock is a gradient in supply (0.162 / 0.302 / 0.466 of lineages never reach the floor), depth is a threshold that jumps only at percolation (1.479 / 1.416 / 2.213), the rate accelerates (6.06 / 9.01 / 18.96), and the survival advantage **reverses**: 0.2496 / 0.2984 / 0.1372, the middle habitat sitting 5.99 standard errors above the straight line between the outer two. The habitat that produces the most generations is the one in which a lineage is least likely to be there at the end.

That reversal cost us a lesson in sample size. On ten worlds the paired survival difference was −0.077, interval −0.166 … +0.012: unresolved, because the spread between worlds (sd 0.14) equalled the effect. A power calculation asked for at least 26 worlds per habitat; at thirty the difference resolved at

−0.1323 (−0.1819 … −0.0840), with the size advantage at +1.8340, the share of lineages that fade away at +0.1419 and the share absorbed by relatives at −0.4164. The ten-world estimate was not merely imprecise: it was biased low by about 40 %. Contrasts of this kind now cost 30–40 worlds, and we say so before quoting one.




Figure 4. Endings and the reversal. (a) Composition of endings in the three habitats: absorption dominates everywhere, while dying alone and fading away trade places as the habitat becomes more connected. (b) The size advantage in generations (blue, left axis) rises only at percolation, while the advantage a large sister has in still being present at the end (orange, right axis) peaks in the mixed habitat and collapses in the connected one.

4.6 What is inherited is copied by the bookkeeping, and we can show it

The trait `s` is copied from parent to daughter with noise. To measure the transmission we reconstruct parent–daughter pairs from the logs. Because a split gives both daughters new identifiers, a parent and its daughter are never recorded on the same step; the parent is read at its last recorded step before the daughter appears, and the daughter at her first. The displacement is at most one logging step (ten integration steps), and the control that decides whether this reconstruction is trustworthy is the artifact's own published correlation, compared per world rather than pooled: 0.9092 against 0.9092 and 0.8967 against 0.8989, over 2 164 of 2 211 and 1 779 of 1 951 pairs.

Then the transmission itself (Fig. 5):

quantity	spots	connected
daughter against parent, slope (95 % interval)	**0.9837 [0.9680, 1.0002]**	**0.9931 [0.9756, 1.0103]**
the same, with division rate partialled out	0.9837	0.9932
parents shuffled (mean, one-sided p)	0.19, **0.001**	0.15, **0.001**
scatter of daughter minus parent	**0.1487**	**0.1495**
the model's own noise	σ = 0.15	σ = 0.15
share of variance between surviving lineages	0.082	0.416

Three readings. The slope is one: daughters are their parents, up to noise. Partialling out how often a body divides does not move it, so the apparent correlation between the trait and the division rate is one rule seen twice. And the part of a daughter's value that does not come from its parent is exactly the model's noise, to one per cent.

The decisive test is the ablation. Two runs with the noise switched off (with a control run reproducing the reference world step by step, so that the switch is the only difference) give a **constant trait**: s is 1.0 in every body at every step, between-body sd 0.0, and 100 % of pairs exactly equal. With the copying rule's noise removed, the trait has no variance at all — the degenerate limit of perfect transmission. Nothing in the chemistry contributes anything transmissible: what passes from parent to daughter passes because the analysis wrote it there.

Which is why the last row of the table must not be read as a heritability. The share of trait variance that sits between lineages at the end of a run is 0.082 in the lattice and 0.416 in the connected mass, but a world that happens to hold ten lineages gives 0.30 where a world with two gives 0.00–0.02, and the connected habitat's horizon can be a handful of bodies, so one surviving root can carry 0.93. That number counts **surviving lineages**; it is not a property of the trait. Reporting it as a heritability would be the exact error this section exists to prevent.




Figure 5. Transmission and its error. (a) The measured daughter-against-parent slope with its 95 % interval, against the same slope when parents are shuffled; the dotted line is perfect copying. (b) The scatter of daughter minus parent, against the noise the copying rule itself adds (σ = 0.15): what is not inherited is exactly the noise, and with the noise switched off the trait has no variance at all.

4.7 Negative result: a third chemical does not set the pattern's scale

If the pattern's wavelength were inherited by the pattern from something in the medium, the obvious carrier would be a third chemical that diffuses slowly and is advected by the pattern. We added one, with its diffusion rate raised to **8** × that of v, and swept its coupling.

It does not work. The pattern's wavevector stays at **7–9** in all six living cells — the same value as without the third chemical — while what does move is the *orientation* of the pattern, by a measure that runs between 0.636 and 0.837, and that movement does not replicate: 2 of 6 cells show it, in opposite directions. The coupling constant of the added field is 0.30 and the field survives in only 2 of 6 seeds. A third diffusing species, in short, is not a scale channel in this medium, and the wavelength therefore is not inherited from any chemical the pattern carries.

4.8 Shape is inherited, at about half, and the habitat chooses which aspect

Everything above concerns s, a number the tracking rule writes into the daughter. That makes it a fair test of the bookkeeping and a poor test of the medium. A shape measured *from the field* is the opposite kind of object: nobody writes it, and a daughter that resembles her mother's shape is evidence that the medium, not the rule, carried it.

We measured two such traits for every body in twenty fresh worlds (ten in the lattice, ten in the connected mass, 4 111 parent–daughter pairs), from the second moments of each body's pixels: **elongation**, the square root of the ratio of the principal axes (1 is round), and **width**, the minor-axis extent in cells. Both were read at every tenth step by an observer that draws no random numbers — with the observer switched off, the same run is bit-identical, which is how we know it cannot have perturbed the world.

The two controls came out where they should. The assigned trait is transmitted at **0.9959 [0.9790, 1.0126]** and **0.9817 [0.9631, 1.0003]** in the two habitats, reproducing §4.6 in a different set of worlds; and area is transmitted at **0.4514 [0.4264, 0.4771]** and **0.4481 [0.4193, 0.4723]**, within a hundredth of the geometric expectation of one half, because a daughter is a part of her mother and not a copy of her.

Shape sits between the two, and it does not sit in the same place for both habitats:

trait	lattice (F = 0.030)	connected (F = 0.040)
s, the assigned trait (control)	0.9959 [0.9790, 1.0126]	0.9817 [0.9631, 1.0003]
area (control)	0.4514 [0.4264, 0.4771]	0.4481 [0.4193, 0.4723]
log elongation	**0.6074 [0.5399, 0.6785]**	0.4260 [0.3585, 0.4924]
log width	0.3384 [0.2877, 0.3850]	**0.6238 [0.5846, 0.6672]**

Four things follow.

Shape is not free of heredity, and it is not the tracker's heredity either. All four shape slopes lie between 0.29 and 0.68 with shuffled-parent nulls of −0.001 to +0.002 and p = 0.001, and the partial slope given log area stays between 0.26 and 0.65 — so this is not area in disguise. The gap between 0.98 and 0.5 is the gap between a number the bookkeeping copies and a shape the medium has to reproduce.

The habitat decides which aspect is transmitted. In the lattice elongation is inherited better than width (0.6074 against 0.3384, intervals disjoint); in the connected mass width is inherited better than elongation (0.6238 against 0.4260, intervals disjoint), and the same trait differs between habitats by more than either interval allows. In a lattice of round bodies the orientation of a ridge is the thing a division can pass on; in a connected labyrinth it is the local width.

Nothing relaxes on the timescale we can see. Reading the daughter one, two and three logged steps after her first appearance moves each slope by at most 0.05 (log elongation in the lattice 0.607 to 0.579; log width in the connected mass 0.624 to 0.571). The value is set at birth rather than restored afterwards, which is what a copied quantity looks like and not what a relaxing shape looks like.

The level is about half. Averaged over the two habitats, elongation 0.517 and width 0.481. A slope of 1 would make the daughter a scaled copy of her mother; 0.5 makes her, on average, halfway between her mother and the habitat's typical shape — which is also where the geometric control sits.

So the sentence from §4.6 needs one clause added: the only *perfectly* heritable character is the one the bookkeeping writes; shape is heritable at about half, in a direction the habitat chooses. Both halves matter — the first is what a reader of parent–daughter correlations needs to know, since a correlation of 1.0 can be bookkeeping; the second is what makes these entities more than labels.




Figure 6. Shape, measured from the field. (a) The daughter-against-parent slope for four quantities in the same worlds, with 95 % intervals: the assigned trait (near 1), the size a daughter must halve (near 0.5), and the two shape traits, which swap places between the two habitats. (b) The same shape slopes when the daughter is read one, two or three logged steps after birth: the lines are flat, so the shape is set at the division rather than relaxed towards a typical value afterwards.

5. What it means

A pattern-forming medium produces entities that look individual and are not: they merge on contact at a length the medium sets, they die mostly by being absorbed, and their only perfectly heritable character is one our own analysis writes into them, with shape inherited at about half along a direction the habitat chooses. None of that makes the system uninteresting — it makes the description precise. Four consequences for anyone using this kind of medium as a model of individuality:

1. **"Inheritance" needs an ablation.** A parent–daughter correlation is not evidence of transmission, because the

bookkeeping that names the daughter can create the correlation. The test is what happens when the copying rule's own noise is removed: if the trait becomes constant, the medium is not transmitting it.

2. **The mixing length is the boundary of the individual, and it is measured at contact.** Two bodies whose surfaces come

within about two cells become one, whatever their size; whether they get that close is set by the initial gap, which stops working between eight and eleven cells. Any statement about a "population" of such bodies has to say what it does about contact.

3. **Survival and reproduction are different measurements and can point opposite ways.** In this medium the habitat that

maximises generations is the one where a lineage is least likely to be present at the end. A study that reports only one of the two can be internally consistent and still tell a reader the wrong story.

4. **Heredity is a property of what you measure.** In the same worlds the trait the rule writes is copied at 0.98, the

size at 0.45, and shape at 0.34 to 0.62 depending on which aspect and which habitat. A single "heritability" for a pattern-forming medium is therefore not a number but a list, ordered by how much of the trait the medium has to rebuild at each division.

The negative result closes the most plausible route by which the trait could have become a property of the medium: a carrier chemical does not set the scale, so the scale belongs to the interaction of the two original chemicals with the imposed supply pattern.

5.1 What to measure next

1. **A third habitat.** $F = 0.035$, between the lattice and the connected mass, twenty worlds: it decides whether the swap

of §4.8 is a continuum or a switch.

2. **The fusion crossing inside the band.** Arms at gaps 9, 10 and 11 for radius 4.5 and radius 6, ten seeds each — the

only way the "eight to eleven cells" of §4.1 becomes a crossing instead of a bound.

3. **The shape slopes at another threshold.** A body is the set of pixels above $v = 0.2$; at 0.1 and 0.3 the same

analysis says whether the half-inheritance belongs to the field or to the mask.

4. **Selection on shape.** The heritable part of shape matters only if it changes a daughter's fate: the fate classes of

§4.4 are recorded per body, so correlating a daughter's elongation with her ending needs no new simulation.

6. Limits

1. **One system.** A single grid size (96 × 96), one reaction rate $k = 0.06$, one threshold, one supply modulation. The

mixing length is a length in *this* medium, stated in cells and in wavelengths; a different ratio of the diffusion rates could move it.

2. **The central chain is correlational.** Size is never randomised, so the size-to-generations result is a within-family

comparison, not an intervention, and age remains an uncontrolled confound. What is causal here is the mixing length, which comes from placing bodies by hand.

3. **Time resolution is coarse.** The model is written every ten steps, so "slow" is a single bit and the width of the

timing distribution is bracketed, not located.

4. **Lineage bookkeeping has no notion of fusion.** Our variance decomposition treats a merged body as a new root, which

makes the between-lineage share an upper bound.

5. **Two runs per ablation arm.** Both are degenerate in the same way, which is strong evidence that the trait has no

variance, but it is not a distribution; and the criterion for reading a degenerate arm was amended after seeing the data, which we state rather than hide.

6. **World counts.** Thirty per habitat were enough for the contrasts reported; they are the price of admission, not a

surplus. The shape result of §4.8 rests on ten worlds per habitat, twenty in all.

7. **The fusion crossing is bounded, not located.** At radius 3 the last measured arm is a gap of 11 cells, where two runs

of five still ended as one body, and for the two larger radii the arms between 9 and 11 were never run, so §4.1 states a band. The earlier claim of a single number for all three sizes was wrong and is corrected in place rather than quietly replaced: the arms in the old table were the design parameters, not measurements.

8. **Shape is a thresholded silhouette.** The shape traits are computed from the component above $v = 0.2$, and their

primary statistic is a log slope chosen after the tails were seen. A different threshold or transform would move the levels; it is not obvious it would move the swap between elongation and width, which is the part with structure.

9. **Nothing here explains the shape inheritance.** A daughter could resemble her mother because she buds off the same

ridge, or because the supply pattern is locally the same; the experiment that separates these — an imposed supply pattern with a fixed orientation — is not run.

7. Data and code

All data are public. The derived tables behind every figure and every number in this paper are 029-figures/tables.json (each entry names the aggregate it came from), table-edge.csv, table-habitats.csv, table-heredity.csv and table-shape.csv. The aggregates themselves, with the command line of every run, are in morphogenesis/experiments/E09_pond/ and the per-run artifacts in morphogenesis/experiments/L1_field/. The figures are produced by scripts/make_article_029_figures.py; a checker (scripts/check_article_029_numbers.py) verifies that every number printed above appears in the derived tables, and the tables are built from the aggregates, not typed by hand.

Two analyses behind §4.1 and §4.8 are new in this version and are reported separately, with their pre-registrations and controls: the contact geometry of fusion (scripts/x157_gap_geometry.py, reports/2026-11-04-x157-gap-geometry.md) and the shape-heredity series (scripts/x156_series.py, scripts/x156_analyse.py, reports/2026-11-04-x156-shape-heredity.md). The numbers in §4.1 were re-derived after an independent review recomputed them from the raw worlds (lab/reviews/2026-11-04-029-three-numbers.md); that review is part of the record, not a private correspondence. The whole computation is one consumer GPU: the ablation arm is 0.11 GPU-hours, the shape series 0.01, and the entire paper's series under 40.

8. Reproduction

```
python scripts/make_article_029_figures.py --write     # tables and all six figures, from
the committed aggregates
python scripts/check_article_029_numbers.py --ci       # every number printed above, traced
to its aggregate
python scripts/x156_analyse.py --write                 # the shape results of §4.8, from the
twenty shape runs
python scripts/x157_gap_geometry.py --write            # the contact geometry of §4.1, from
171 two-body runs
```

The series behind the results are `scripts/x134_series.py` … `scripts/x154_series.py` (the two-body intervention and its sweeps), `scripts/x108_series.py` (the dense logs), `scripts/x155_ablation.py` (the noise-off arm) and `scripts/x155_analyse.py` (reconstruction and statistics). Each artifact records its own command, its seeds and its pre-registered criterion.