

EXERCISES

CHAPTER 2

SEAN LI ¹

1. Reducted

Definition Some rules for reference.

$$\frac{x : \sigma \in \Gamma}{\Gamma \vdash x : \sigma} \text{(T-Var)} \qquad \frac{\Gamma \vdash M : \sigma \rightarrow \tau \quad \Gamma \vdash N : \sigma}{\Gamma \vdash MN : \tau} \text{(T-App)}$$

$$\frac{\Gamma, x : \sigma \vdash M : \tau}{\Gamma \vdash \lambda x : \sigma. M : \sigma \rightarrow \tau} \text{(T-Abst)}$$

In this document, convention is that all type judgements in a proof tree, unless stated otherwise, is derived from a single context per tree.

Problem

(2.1) Type the following terms

$$xxy \quad xyx \quad xyx \quad x(xy) \quad x(yx)$$

Solution. The first term cannot be typed.

Proof. $xxy = (xx)y$. Therefore, x is a function type, denote it as $\tau \rightarrow \sigma$. By the application rule, a subterm applied to x must be of τ , which means that the application xx is not legally typed. ■

The second one is typable where $x : \tau \rightarrow \tau \rightarrow \sigma$ and $y : \tau$.

1. $x : \tau \rightarrow \tau \rightarrow \sigma \quad \neg \Gamma$

2. $y : \tau \quad \neg \Gamma$

3. $xy : \tau \rightarrow \sigma$ **1,2 T-App**
4. $xyy : \sigma$ **3,2 T-App**

The third term is not typable.

Proof. Assume $xyx = (xy)x$ is typable. Therefore, $x : \tau$ where $\tau = \sigma \rightarrow \tau \rightarrow \alpha$ and $y : \sigma$. One can construct an infinite chain of function type by substituting τ : $\tau = \sigma \rightarrow (\sigma \rightarrow (\sigma \rightarrow \dots \rightarrow \alpha) \rightarrow \alpha) \rightarrow \alpha$. By induction, it can be proven that only lambda abstractions can construct function types, meaning that the term is of form

$$(\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots))y(\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots))$$

meaning that an infinite reduction path is needed. This is impossible in STLC. ■

The fourth type is typable where $x : (\tau \rightarrow \tau)$ and $y : \tau$.

1. $x : \tau \rightarrow \tau$ $\vdash \Gamma$
2. $y : \tau$ $\vdash \Gamma$
3. $xy : \tau$ **1,2 T-App**
4. $x(xy) : \tau$ **1,3 T-App**

The fifth term is typable where $x : (\tau \rightarrow \sigma)$ and $y : (\tau \rightarrow \sigma) \rightarrow \tau$:

1. $x : \tau \rightarrow \sigma$ $\vdash \Gamma$
2. $y : (\tau \rightarrow \sigma) \rightarrow \tau$ $\vdash \Gamma$
3. $yx : \tau$ **2,1 T-App**
4. $x(yx) : \sigma$ **1,3 T-App**

Problem

(2.2) Find types for zero, one, and two

Solution. Term for zero is

$$\text{zero} := \lambda f x. x$$

Here x is only used as a

$$\text{zero} := \lambda f : \alpha. \lambda x : \beta. x$$

Type derivation shown as below:

1. $f : \alpha$ **Bound**
2. $\mid x : \beta$ **Bound**

3. $\frac{}{x : \beta}$ **T-Var**
4. $\frac{}{\lambda x.x : \beta \rightarrow \beta}$ **3 T-Abst**
5. $\lambda f : \alpha.x : \beta.x : \alpha \rightarrow \beta \rightarrow \beta$ **4 T-Abst**

Term for one is

$$\text{one} := \lambda f x.f x$$

Let f be an arbitrary function type that consumes x

$$\text{one} := \lambda f : \alpha \rightarrow \beta.x : \alpha.f x$$

Type derivation shown as below

1. $f : \alpha \rightarrow \beta$ **Bound**
2. $x : \alpha$ **Bound**
3. $f : \alpha \rightarrow \beta$ **T-Var**
4. $x : \alpha$ **T-Var**
5. $f x : \beta$ **3,4 T-App**
6. $\lambda x.f x : \alpha \rightarrow \beta$ **5 T-Abst**
7. $\lambda f : \alpha \rightarrow \beta.x : \alpha.f x : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$ **6 T-Abst**

Same type signatures can be given to two

$$\text{two} := \lambda f : \alpha \rightarrow \beta.\lambda x : \alpha.f f x$$

Type derivation shown as below

1. $f : \alpha \rightarrow \beta$ **Bound**
2. $x : \alpha$ **Bound**
3. $f : \alpha \rightarrow \beta$ **T-Var**
4. $x : \alpha$ **T-Var**
5. $f x : \beta$ **3,4 T-App**
6. $f f x : \beta$ **3,5 T-App**
7. $\lambda x.f f x : \alpha \rightarrow \beta$ **6 T-Abst**
8. $\lambda f : \alpha \rightarrow \beta.x : \alpha.f f x : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$ **7 T-Abst**

Problem

(2.3) Find types for

$$K := \lambda xy.x$$

$$S := \lambda xyz.xz(yz)$$

Solution. There are no occurrences of application in K 's subterms. Therefore all its binding variables could be given a simple base type.

$$K := \lambda x : \alpha. \lambda y : \beta. x$$

Type derivation shown as below

1.	$x : \alpha$	Bound
2.	$y : \beta$	Bound
3.	$x : \alpha$	T-Var
4.	$\lambda y : \beta. x : \beta \rightarrow \alpha$	3 T-Abst
5.	$\lambda x : \alpha. \lambda y : \beta. x : \alpha \rightarrow \beta \rightarrow \alpha$	4 T-Abst

For the S combinator, no term was applied to z . Therefore it can be given a simple base type α . As z was applied to y , it implies that $y : \alpha \rightarrow \beta$ for some output type β . As x takes z and (yz) , it must be of type $\alpha \rightarrow \beta \rightarrow \delta$.

$$S := \lambda x : \alpha \rightarrow \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z. \alpha. xz(yz)$$

Complete type derivation shown as below:

1.	$x : \alpha \rightarrow \beta \rightarrow \delta$	Bound
2.	$y : \alpha \rightarrow \beta$	Bound
3.	$z : \alpha$	Bound
4.	$y : \alpha \rightarrow \beta$	T-Var
5.	$z : \alpha$	T-Var
6.	$yz : \beta$	4,5 T-App
7.	$x : \alpha \rightarrow \beta \rightarrow \delta$	T-Var
8.	$xz : \beta \rightarrow \delta$	7,5 T-App
9.	$xz(yz) : \delta$	8,6 T-App
10.	$\lambda z : \alpha. xz(yz) : \alpha \rightarrow \delta$	9 T-Abstr
11.	$\lambda y : \alpha \rightarrow \beta. \lambda z. \alpha. xz(yz) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$	10 T-Abstr

12.

$$\lambda x : \alpha \rightarrow \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. xz(yz) : (\alpha \rightarrow \beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta \quad \mathbf{11\ T-Abstr}$$

Problem

(2.4) Type the bound variables

$$\lambda xyz. x(yz)$$

$$\lambda xyz. y(xz)z$$

Solution. For the first term, z had nothing applied to it. Therefore it could be given a simple base type α . z was applied to y , therefore $y : \alpha \rightarrow \beta$ to satisfy the application rule. Because the application yielded a type of β , by the application rule $x : \beta \rightarrow \delta$ for some type δ .

$$\lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz)$$

Complete type derivation shown below

1.	$x : \beta \rightarrow \delta$	Bound
2.	$y : \alpha \rightarrow \beta$	Bound
3.	$z : \alpha$	Bound
4.	$y : \alpha \rightarrow \beta$	T-Var
5.	$z : \alpha$	T-Var
6.	$yz : \beta$	4,5 T-App
7.	$x : \beta \rightarrow \delta$	T-Var
8.	$x(yz) : \delta$	7,6 T-App
9.	$\lambda z : \alpha. x(yz) : \alpha \rightarrow \delta$	8 T-Abst
10.	$\lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$	9 T-Abst
11.		

$$\lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz)$$

$$: (\beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta \quad \mathbf{10\ T-Abst}$$

In the second term z could still be given a simple base type $z : \alpha$. Therefore $x : \alpha \rightarrow \beta$ for some type β . y takes $xz : \beta$ and $z : \alpha$, therefore it is of type $y : \beta \rightarrow \alpha \rightarrow \delta$ for some δ .

$$\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z : \alpha. y(xz)z$$

. Complete type derivation shown below

1.	$x : \alpha \rightarrow \beta$	Bound
2.	$y : \beta \rightarrow \alpha \rightarrow \delta$	Bound
3.	$z : \alpha$	Bound
4.	$x : \alpha \rightarrow \beta$	T-Var
5.	$z : \alpha$	T-Var
6.	$xz : \beta$	4,5 T-App
7.	$y : \beta \rightarrow \alpha \rightarrow \delta$	T-Var
8.	$y(xz) : \alpha \rightarrow \delta$	7,6 T-App
9.	$y(xz)z : \delta$	8,5 T-App
10.	$\lambda z : \alpha. y(xz)z : \alpha \rightarrow \delta$	9 T-Abst
11.	$\lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y(xz)z : (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta$	10 T-Abst
12.		

$\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y(xz)z :$

$(\alpha \rightarrow \beta) \rightarrow (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta$

11 T-Abst

Problem

(2.5) Try to type the following terms, and prove if not typable.

$\lambda xy. x(\lambda z. y)y$

$\lambda xy. x(\lambda z. x)y.$

Solution. The first term is trivially typable.

1.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	Bound
2.	$y : \alpha$	Bound
3.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	T-Var
4.	$z : \delta$	Bound
5.	$y : \alpha$	T-Var
6.	$\lambda z : \delta. y : \delta \rightarrow \alpha$	5 T-Abst
7.	$x(\lambda z : \delta. y) : \alpha \rightarrow \beta$	3,6 T-App
8.	$y : \alpha$	T-Var
9.	$x(\lambda z : \delta. y)y : \beta$	7,8 T-App
10.	$\lambda y : \alpha. x(\lambda z : \delta. y)y : \alpha \rightarrow \beta$	9 T-Abst

11.

$$\begin{aligned} \lambda x : ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \lambda y : \alpha. x(\lambda z : \delta. y) y \\ : ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta \end{aligned} \quad \mathbf{10\ T-Abst}$$

The second term is not typable in STLC.

Proof. By induction on the type inference rule that constructed the type judgement for subterm $x(\lambda z. x)$. Because the term is an application, the only rule that applies is the application rule.

We denote the context inside the abstraction as Γ' . Suppose $\mathcal{J} \equiv \Gamma' \vdash x(\lambda z. x) : \tau$. By the inference rule of application, x must be a function type that accepts the type of $(\lambda z. x)$. Let $\Gamma' \vdash z : \alpha$, and type of x as τ_x . Therefore, $\Gamma' \vdash \lambda z : \alpha. x : \alpha \rightarrow \tau_x$. Therefore, $\tau_x = (\alpha \rightarrow \tau_x) \rightarrow \tau$. This is a recursive type, which is not constructable as it requires infinitely nested lambda abstractions that requires infinite reduction paths to reach a normal form. ■

Problem

(2.6) Prove the pretyped term below is legal.

$$\lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x(\lambda z : \alpha. y)$$

Using the tree format and the flag format.

Solution. We suppose a context $\Gamma \vdash y : \beta$ that obviously exists.

Proof.

$$\frac{\frac{}{x : (\alpha \rightarrow \beta) \rightarrow \alpha} \text{(Bound)} \quad \frac{\Gamma, z : \alpha \vdash y : \beta}{\Gamma \vdash (\lambda z : \alpha. y) : \alpha \rightarrow \beta} \text{(T-Abst)}}{\Gamma \vdash (\lambda z : \alpha. y) : \alpha \rightarrow \beta} \text{(T-App)} \quad \frac{\Gamma, x : (\alpha \rightarrow \beta) \rightarrow \alpha \vdash (x(\lambda z : \alpha. y)) : \alpha}{\Gamma \vdash \lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x(\lambda z : \alpha. y) : ((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow \alpha} \text{(T-Abst)}$$

A valid type could be given to the term. Therefore, the term is typable under an existing context. ■

The flag derivation is given below:

1.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	Bound
2.	$z : \alpha$	Bound
3.	$y : \beta$	$\vdash \Gamma$
4.	$(\lambda z : \alpha. y) : \alpha \rightarrow \beta$	3 T-Abst
5.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	T-Var

6. $\frac{x(\lambda z : \alpha.y) : \beta}{\text{5,4 T-App}}$
- 7.
- $\lambda x : ((\alpha \rightarrow \beta) \rightarrow \beta).x(\lambda z : \alpha.y)$
- $: (\alpha \rightarrow \beta) \rightarrow \beta \rightarrow \alpha$ **6 T-Abst**

Problem

(2.7 a) Derive

$$f : A \rightarrow B \wedge g : B \rightarrow C \Rightarrow g \circ f : A \rightarrow C$$

Using the rules

$$\frac{f : A \rightarrow B, x \in A}{f(x) \in B} \text{(F-App)} \quad \frac{\forall x \in A, f(x) \in B}{f : A \rightarrow B} \text{(F-Abst)}$$

Solution.

Proof.

- | | | |
|-----|--|----------------------|
| 1. | $f : A \rightarrow B \wedge g : B \rightarrow C$ | Assumption |
| 2. | $f : A \rightarrow B$ | $1 \wedge E$ |
| 3. | $g : B \rightarrow C$ | $1 \wedge E$ |
| 4. | $a \in A$ | |
| 5. | $f(a) \in B$ | 3, 4 F-App |
| 6. | $g(f(a)) \in C$ | 5, 4 F-App |
| 7. | $(g \circ f)(a) \in C$ | 6 Compose Def |
| 8. | $\forall x \in A, (g \circ f)(x) \in C$ | $7 \forall E$ |
| 9. | $g \circ f : A \rightarrow C$ | 8 F-Abst |
| 10. | $f : A \rightarrow B, g : B \rightarrow C \Rightarrow g \circ f : A \rightarrow C$ | $9 \Rightarrow I$ |

■

Problem

(2.7 b) Give a derivation in natural deduction of the following:

$$(A \Rightarrow B) \Rightarrow ((B \Rightarrow C) \Rightarrow (A \Rightarrow C))$$

Using the rules

$$\frac{A \Rightarrow B \quad A}{B} (\Rightarrow E) \qquad \frac{\begin{array}{l} 1. \quad A \quad \text{Premise} \\ 2. \quad \vdots \\ 3. \quad B \end{array}}{A \Rightarrow B} (\Rightarrow I)$$

Solution.

Proof.

$$\begin{array}{ll} 1. & A \Rightarrow B \quad \text{Premise} \\ 2. & \begin{array}{|l} B \Rightarrow C \end{array} \quad \text{Premise} \\ 3. & \begin{array}{|l|} A \end{array} \quad \text{Premise} \\ 4. & \begin{array}{|l|} B \end{array} \quad 1, 3 \Rightarrow E \\ 5. & \begin{array}{|l|} C \end{array} \quad 2, 4 \Rightarrow E \\ 6. & \begin{array}{|l|} A \Rightarrow C \end{array} \quad 3-5 \Rightarrow I \\ 7. & \begin{array}{|l|} (B \Rightarrow C) \Rightarrow (A \Rightarrow C) \end{array} \quad 2-6 \Rightarrow I \\ 8. & (A \Rightarrow B) \Rightarrow ((B \Rightarrow C) \Rightarrow (A \Rightarrow C)) \quad 1-7 \Rightarrow I \end{array}$$

■

Problem

(2.7 c) Prove the following pre-typed term is legal using flag notation

$$\lambda z : \alpha. y(xz)$$

Solution.

Proof. Let $\Gamma \vdash x : \alpha \rightarrow \beta, y : \beta \rightarrow \delta$ for some type β and δ .

$$\begin{array}{ll} 1. & z : \alpha \quad \text{Bound} \\ 2. & \begin{array}{|l} x : \alpha \rightarrow \beta \end{array} \quad \neg \Gamma \\ 3. & \begin{array}{|l|} z : \alpha \end{array} \quad \text{T-Var} \end{array}$$

- | | | |
|----|---|------------------|
| 4. | $xz : \beta$ | 2,3 T-App |
| 5. | $y : \beta \rightarrow \delta$ | $\vdash \Gamma$ |
| 6. | $y(xz) : \delta$ | 5,4 T-App |
| 7. | $\lambda z : \alpha. y(xz) : \alpha \rightarrow \delta$ | 6 T-Abst |

■

Problem

(2.7 d) State the similarity between Q. 2.7 (a), (b), and (c).

Solution. All of these examples requires proving something about composing two maps together as like this:



Problem

(2.8 a) Pre-type the bounding variables for the following term

$$\lambda xy. y(\lambda z. yx) : (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$$

Solution.

$$\lambda x : (\gamma \rightarrow \beta). y : ((\gamma \rightarrow \beta) \rightarrow \beta). y(\lambda z : \gamma. yx)$$

Problem

(2.8 b) Give a derivation in tree format

Solution.

$$\begin{array}{c}
\text{(i)} \frac{}{x : (\gamma \rightarrow \beta)} \quad \text{(ii)} \frac{}{y : (\gamma \rightarrow \beta) \rightarrow \beta} \\
\text{(iii)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta), z : \gamma \vdash yx : \beta} \text{T-App} \\
\text{(v)} \frac{}{y : ((\gamma \rightarrow \beta) \rightarrow \beta)} \quad \text{(iv)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta) \vdash \lambda z : \gamma. yx : \gamma \rightarrow \beta} \text{T-Abst} \\
\text{(vi)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta) \vdash y(\lambda z : \gamma. yx) : \beta} \text{T-App} \\
\text{(vii)} \frac{}{x : (\gamma \rightarrow \beta) \vdash \lambda y : ((\gamma \rightarrow \beta) \rightarrow \beta). y(\lambda z : \gamma. yx) : ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta} \text{T-Abst} \\
\text{(viii)} \frac{}{(\lambda x : (\gamma \rightarrow \beta). y : ((\gamma \rightarrow \beta) \rightarrow \beta). y(\lambda z : \gamma. yx)) : (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta} \text{T-Abst}
\end{array}$$

Problem

(2.8 c) Sketch a diagram of tree structure of derivation

Solution. Trivial.

Problem

(2.8 d) Transform the derivation into flag notation

Solution.

	1.	$x : \gamma \rightarrow \beta$	Bound
	2.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	Bound
	3.	$z : \gamma$	Bound
(ii)	4.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	T-Var
(i)	5.	$x : \gamma \rightarrow \beta$	T-Var
(iii)	6.	$yx : \beta$	4,5 T-App
(iv)	7.	$\lambda z : \gamma. yx : \gamma \rightarrow \beta$	6 T-Abst
(v)	8.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	T-Var
(vi)	9.	$y(\lambda z : \gamma. yx) : \beta$	8,7 T-App
		$\lambda y : (\gamma \rightarrow \beta) \rightarrow \beta. y(\lambda z : \gamma. yx)$	
(vii)	10.	$: ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$	9 T-Abst
(viii)	11.		
		$\lambda x : (\gamma \rightarrow \beta). \lambda y : (\gamma \rightarrow \beta) \rightarrow \beta. y(\lambda z : \gamma. yx)$	
		$: (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$	10 T-Abst

Problem

(2.9 a) Give derivations of the following judgement

$$x : \delta \rightarrow \delta \rightarrow \alpha, y : \gamma \rightarrow \alpha, z : \alpha \rightarrow \beta \vdash \\ \lambda u : \delta. \lambda v : \gamma. z(yv) : \delta \rightarrow \gamma \rightarrow \beta$$

Solution.

1.	$u : \delta$	Bound
2.	$v : \gamma$	Bound
3.	$y : \gamma \rightarrow \alpha$	T-Var
4.	$v : \gamma$	T-Var
5.	$yv : \alpha$	3,4 T-App
6.	$z : \alpha \rightarrow \beta$	T-Var
7.	$z(yv) : \beta$	6,5 T-App
8.	$\lambda v : \gamma. z(yv) : \gamma \rightarrow \beta$	7 T-Abst
9.	$\lambda u : \delta. \lambda v : \gamma. z(yv) : \delta \rightarrow \gamma \rightarrow \beta$	8 T-Abst

Problem

(2.9 b) Give derivations of the following judgement

$$x : \delta \rightarrow \delta \rightarrow \alpha, y : \gamma \rightarrow \alpha, z : \alpha \rightarrow \beta \vdash \\ \lambda u : \delta. \lambda v : \gamma. z(xuu) : \delta \rightarrow \gamma \rightarrow \beta$$

Solution.

1.	$u : \delta$	Bound
2.	$v : \gamma$	Bound
3.	$x : \delta \rightarrow \delta \rightarrow \alpha$	T-Var
4.	$u : \delta$	T-Var
5.	$xu : \delta \rightarrow \alpha$	3,4 T-App
6.	$xuu : \alpha$	5,4 T-App
7.	$z : \alpha \rightarrow \beta$	T-Var
8.	$z(xuu) : \beta$	7,6 T-App
9.	$\lambda v : \gamma. z(xuu) : \gamma \rightarrow \beta$	8 T-Abst
10.	$\lambda u : \delta. \lambda v : \gamma. z(xuu) : \delta \rightarrow \gamma \rightarrow \beta$	9 T-Abst

Problem

(2.10 a) Give derivation for

$$xz(yz)$$

Solution. Assume an context

$$\Gamma \vdash x : \alpha \rightarrow \beta \rightarrow \gamma$$

$$\Gamma \vdash y : \alpha \rightarrow \beta$$

$$\Gamma \vdash z : \alpha$$

1. $x : \alpha \rightarrow \beta \rightarrow \gamma$ **T-Var**
2. $y : \alpha \rightarrow \beta$ **T-Var**
3. $z : \alpha$ **T-Var**
4. $xz : \beta \rightarrow \gamma$ **1,3 T-App**
5. $yz : \beta$ **2,3 T-App**
6. $xz(yz) : \gamma$ **4,5 T-App**

Problem

(2.10 b) Give derivation for

$$\lambda x : (\alpha \rightarrow \beta) \rightarrow \beta.x(yz)$$

Solution. Assume an context

$$\Gamma \vdash y : \gamma \rightarrow (\alpha \rightarrow \beta)$$

$$\Gamma \vdash z : \gamma$$

1. $x : (\alpha \rightarrow \beta) \rightarrow \beta$ **Bound**
2. $x : (\alpha \rightarrow \beta) \rightarrow \beta$ **T-Var**
3. $y : \gamma \rightarrow \alpha \rightarrow \beta$ **T-Var**
4. $z : \gamma$ **T-Var**
5. $yz : \alpha \rightarrow \beta$ **3,4 T-App**
6. $x(yz) : \beta$ **2,5 T-App**
7. $\lambda x : (\alpha \rightarrow \beta) \rightarrow \beta.x(yz) : ((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$ **6 T-Abst**

Problem

(2.10 c) Give derivation for

$$\lambda y : \alpha. \lambda z : \beta \rightarrow \gamma. z(xyy)$$

Solution. Assume a context

$$\Gamma \vdash x : \alpha \rightarrow \alpha \rightarrow \beta$$

1.	$y : \alpha$	Bound
2.	$z : \beta \rightarrow \gamma$	Bound
3.	$z : \beta \rightarrow \gamma$	T-Var
4.	$x : \alpha \rightarrow \alpha \rightarrow \beta$	T-Var
5.	$y : \alpha$	T-Var
6.	$xy : \alpha \rightarrow \beta$	4,5 T-App
7.	$xyy : \beta$	6,5 T-App
8.	$z(xyy) : \gamma$	3,6 T-App
9.	$\lambda z : \beta \rightarrow \gamma. z(xyy) : (\beta \rightarrow \gamma) \rightarrow \gamma$	8 T-Abst
10.	$\lambda y : \alpha. \lambda z : \beta \rightarrow \gamma. z(xyy) : \alpha \rightarrow (\beta \rightarrow \gamma) \rightarrow \gamma$	8 T-Abst

Problem

(2.10 d) Give derivation for

$$\lambda x : \alpha \rightarrow \beta. y(xz)z$$

Solution. Consider a context

$$\begin{aligned} \Gamma \vdash z : \alpha \\ \Gamma \vdash y : \beta \rightarrow \alpha \rightarrow \gamma \end{aligned}$$

1.	$x : \alpha \rightarrow \beta$	Bound
2.	$x : \alpha \rightarrow \beta$	T-Var
3.	$z : \alpha$	T-Var
4.	$xz : \beta$	2,3 T-App
5.	$y : \beta \rightarrow \alpha \rightarrow \gamma$	T-Var
6.	$y(xz) : \alpha \rightarrow \gamma$	5,4 T-App
7.	$y(xz)z : \gamma$	3,5 T-App

8. $\lambda x : \alpha \rightarrow \beta. y(xz)z : (\alpha \rightarrow \beta) \rightarrow \gamma$ **7 T-Abst**