

# EXERCISES

## CHAPTER 2

SEAN LI <sup>1</sup>

### 1. Reducted

---

**Definition** Some rules for reference.

$$\frac{x : \sigma \in \Gamma}{\Gamma \vdash x : \sigma} \text{(T-Var)} \qquad \frac{\Gamma \vdash M : \sigma \rightarrow \tau \quad \Gamma \vdash N : \sigma}{\Gamma \vdash M N : \tau} \text{(T-App)}$$

$$\frac{\Gamma, x : \sigma \vdash M : \tau}{\Gamma \vdash \lambda x : \sigma. M : \sigma \rightarrow \tau} \text{(T-Abst)}$$

In this document, convention is that all type judgements in a proof tree, unless stated otherwise, is derived from a single and unique context per tree.

---

#### Problem

(2.1) Type the following terms

$$x x y \quad x y y \quad x y x \quad x (x y) \quad x (y x)$$

*Solution.* The first term cannot be typed.

*Proof.*  $x x y = (x x) y$ . Therefore,  $x$  is a function type, denote it as  $\tau \rightarrow \sigma$ . By the application rule, a subterm applied to  $x$  must be of  $\tau$ , which means that the application  $x x$  is not legally typed. ■

The second one is typable where  $x : \tau \rightarrow \tau \rightarrow \sigma$  and  $y : \tau$ .

1.  $x : \tau \rightarrow \tau \rightarrow \sigma \quad \neg \Gamma$

2.  $y : \tau \quad \neg \Gamma$

3.  $x y : \tau \rightarrow \sigma$       **1,2 T-App**
4.  $x y y : \sigma$       **3,2 T-App**

The third term is not typable.

*Proof.* Assume  $x y x = (x y) x$  is typable. Therefore,  $x : \tau$  where  $\tau \equiv \sigma \rightarrow \tau \rightarrow \alpha$  and  $y : \sigma$ . One can construct an infinite chain of function type by substituting  $\tau$ :  
 $\tau \equiv \sigma \rightarrow (\sigma \rightarrow (\sigma \rightarrow \dots \rightarrow \alpha) \rightarrow \alpha) \rightarrow \alpha$ . By induction, it can be proven that only lambda abstractions can construct function types, meaning that the term is of form

$$(\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots)) y (\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots))$$

meaning that an infinite reduction path is needed. This is impossible in STLC. ■

The fourth type is typable where  $x : (\tau \rightarrow \tau)$  and  $y : \tau$ .

1.  $x : \tau \rightarrow \tau$        $\vdash \Gamma$
2.  $y : \tau$        $\vdash \Gamma$
3.  $x y : \tau$       **1,2 T-App**
4.  $x (x y) : \tau$       **1,3 T-App**

The fifth term is typable where  $x : (\tau \rightarrow \sigma)$  and  $y : (\tau \rightarrow \sigma) \rightarrow \tau$ :

1.  $x : \tau \rightarrow \sigma$        $\vdash \Gamma$
2.  $y : (\tau \rightarrow \sigma) \rightarrow \tau$        $\vdash \Gamma$
3.  $y x : \tau$       **2,1 T-App**
4.  $x (y x) : \sigma$       **1,3 T-App**

### Problem

(2.2) Find types for zero, one, and two

*Solution.* Term for zero is

$$\text{zero} := \lambda f x . x$$

Here  $x$  is only used as a

$$\text{zero} := \lambda f : \alpha. \lambda x : \beta. x$$

Type derivation shown as below:

1.  $f : \alpha$       **Bound**
2.  $\left| \begin{array}{l} x : \beta \end{array} \right.$       **Bound**

3.  $\frac{}{x : \beta}$  **T-Var**
4.  $\frac{}{\lambda x . x : \beta \rightarrow \beta}$  **3 T-Abst**
5.  $\lambda f : \alpha . x : \beta . x : \alpha \rightarrow \beta \rightarrow \beta$  **4 T-Abst**

Term for one is

$$\text{one} := \lambda f x . f x$$

Let  $f$  be an arbitrary function type that consumes  $x$

$$\text{one} := \lambda f : \alpha \rightarrow \beta . x : \alpha . f x$$

Type derivation shown as below

1.  $f : \alpha \rightarrow \beta$  **Bound**
2.  $x : \alpha$  **Bound**
3.  $f : \alpha \rightarrow \beta$  **T-Var**
4.  $x : \alpha$  **T-Var**
5.  $f x : \beta$  **3,4 T-App**
6.  $\lambda x . f x : \alpha \rightarrow \beta$  **5 T-Abst**
7.  $\lambda f : \alpha \rightarrow \beta . x : \alpha . f x : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$  **6 T-Abst**

Same type signatures can be given to two

$$\text{two} := \lambda f : \alpha \rightarrow \beta . \lambda x : \alpha . f f x$$

Type derivation shown as below

1.  $f : \alpha \rightarrow \beta$  **Bound**
2.  $x : \alpha$  **Bound**
3.  $f : \alpha \rightarrow \beta$  **T-Var**
4.  $x : \alpha$  **T-Var**
5.  $f x : \beta$  **3,4 T-App**
6.  $f f x : \beta$  **3,5 T-App**
7.  $\lambda x . f f x : \alpha \rightarrow \beta$  **6 T-Abst**
8.  $\lambda f : \alpha \rightarrow \beta . \lambda x : \alpha . f f x : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$  **7 T-Abst**

## Problem

(2.3) Find types for

$$K := \lambda x y . x$$

$$S := \lambda x y z . x z (y z)$$

*Solution.* There are no occurrences of application in  $K$ 's subterms. Therefore all its binding variables could be given a simple base type.

$$K := \lambda x : \alpha . \lambda y : \beta . x$$

Type derivation shown as below

1.	$x : \alpha$	<b>Bound</b>
2.	$y : \beta$	<b>Bound</b>
3.	$x : \alpha$	<b>T-Var</b>
4.	$\lambda y : \beta . x : \beta \rightarrow \alpha$	<b>3 T-Abst</b>
5.	$\lambda x : \alpha . \lambda y : \beta . x : \alpha \rightarrow \beta \rightarrow \alpha$	<b>4 T-Abst</b>

For the  $S$  combinator, no term was applied to  $z$ . Therefore it can be given a simple base type  $\alpha$ . As  $z$  was applied to  $y$ , it implies that  $y : \alpha \rightarrow \beta$  for some output type  $\beta$ . As  $x$  takes  $z$  and  $(y z)$ , it must be of type  $\alpha \rightarrow \beta \rightarrow \delta$ .

$$S := \lambda x : \alpha \rightarrow \beta \rightarrow \delta . \lambda y : \alpha \rightarrow \beta . \lambda z . \alpha . x z (y z)$$

Complete type derivation shown as below:

1.	$x : \alpha \rightarrow \beta \rightarrow \delta$	<b>Bound</b>
2.	$y : \alpha \rightarrow \beta$	<b>Bound</b>
3.	$z : \alpha$	<b>Bound</b>
4.	$y : \alpha \rightarrow \beta$	<b>T-Var</b>
5.	$z : \alpha$	<b>T-Var</b>
6.	$y z : \beta$	<b>4,5 T-App</b>
7.	$x : \alpha \rightarrow \beta \rightarrow \delta$	<b>T-Var</b>
8.	$x z : \beta \rightarrow \delta$	<b>7,5 T-App</b>
9.	$x z (y z) : \delta$	<b>8,6 T-App</b>
10.	$\lambda z : \alpha . x z (y z) : \alpha \rightarrow \delta$	<b>9 T-Abstr</b>
11.	$\lambda y : \alpha \rightarrow \beta . \lambda z . \alpha . x z (y z) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$	<b>10 T-Abstr</b>

12.

$$\lambda x : \alpha \rightarrow \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x z (y z) : (\alpha \rightarrow \beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta \quad \mathbf{11 \text{ T-Abstr}}$$

### Problem

(2.4) Type the bound variables

$$\lambda x y z . x (y z)$$

$$\lambda x y z . y (x z) z$$

*Solution.* For the first term,  $z$  had nothing applied to it. Therefore it could be given a simple base type  $\alpha$ .  $z$  was applied to  $y$ , therefore  $y : \alpha \rightarrow \beta$  to satisfy the application rule. Because the application yielded a type of  $\beta$ , by the application rule  $x : \beta \rightarrow \delta$  for some type  $\delta$ .

$$\lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x (y z)$$

Complete type derivation shown below

1.	$x : \beta \rightarrow \delta$	<b>Bound</b>
2.	$y : \alpha \rightarrow \beta$	<b>Bound</b>
3.	$z : \alpha$	<b>Bound</b>
4.	$y : \alpha \rightarrow \beta$	<b>T-Var</b>
5.	$z : \alpha$	<b>T-Var</b>
6.	$y z : \beta$	<b>4,5 T-App</b>
7.	$x : \beta \rightarrow \delta$	<b>T-Var</b>
8.	$x (y z) : \delta$	<b>7,6 T-App</b>
9.	$\lambda z : \alpha. x (y z) : \alpha \rightarrow \delta$	<b>8 T-Abst</b>
10.	$\lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x (y z) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$	<b>9 T-Abst</b>
11.		

$$\lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x (y z)$$

$$: (\beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta \quad \mathbf{10 \text{ T-Abst}}$$

In the second term  $z$  could still be given a simple base type  $z : \alpha$ . Therefore  $x : \alpha \rightarrow \beta$  for some type  $\beta$ .  $y$  takes  $x z : \beta$  and  $z : \alpha$ , therefore it is of type  $y : \beta \rightarrow \alpha \rightarrow \delta$  for some  $\delta$ .

$$\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z : \alpha. y (x z) z$$

. Complete type derivation shown below

1.	$x : \alpha \rightarrow \beta$	<b>Bound</b>
2.	$y : \beta \rightarrow \alpha \rightarrow \delta$	<b>Bound</b>
3.	$z : \alpha$	<b>Bound</b>
4.	$x : \alpha \rightarrow \beta$	<b>T-Var</b>
5.	$z : \alpha$	<b>T-Var</b>
6.	$x z : \beta$	<b>4,5 T-App</b>
7.	$y : \beta \rightarrow \alpha \rightarrow \delta$	<b>T-Var</b>
8.	$y (x z) : \alpha \rightarrow \delta$	<b>7,6 T-App</b>
9.	$y (x z) z : \delta$	<b>8,5 T-App</b>
10.	$\lambda z : \alpha. y (x z) z : \alpha \rightarrow \delta$	<b>9 T-Abst</b>
11.	$\lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y (x z) z : (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta$	<b>10 T-Abst</b>
12.		

$\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y (x z) z :$   
 $(\alpha \rightarrow \beta) \rightarrow (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta$

**11 T-Abst**

### Problem

(2.5) Try to type the following terms, and prove if not typable.

$\lambda x y. x (\lambda z. y) y$   
 $\lambda x y. x (\lambda z. x) y.$

*Solution.* The first term is trivially typable.

1.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	<b>Bound</b>
2.	$y : \alpha$	<b>Bound</b>
3.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	<b>T-Var</b>
4.	$z : \delta$	<b>Bound</b>
5.	$y : \alpha$	<b>T-Var</b>
6.	$\lambda z : \delta. y : \delta \rightarrow \alpha$	<b>5 T-Abst</b>
7.	$x (\lambda z : \delta. y) : \alpha \rightarrow \beta$	<b>3,6 T-App</b>
8.	$y : \alpha$	<b>T-Var</b>
9.	$x (\lambda z : \delta. y) y : \beta$	<b>7,8 T-App</b>
10.	$\lambda y : \alpha. x (\lambda z : \delta. y) y : \alpha \rightarrow \beta$	<b>9 T-Abst</b>

11.

$$\begin{aligned} \lambda x : ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \lambda y : \alpha. x (\lambda z : \delta. y) y \\ : ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta \end{aligned} \quad \mathbf{10 \ T-Abst}$$

The second term is not typable in STLC.

*Proof.* By induction on the type inference rule that constructed the type judgement for subterm  $x (\lambda z . x)$ . Because the term is an application, the only rule that applies is the application rule.

We denote the context inside the abstraction as  $\Gamma'$ . Suppose  $\mathcal{J} \equiv \Gamma' \vdash x (\lambda z . x) : \tau$ . By the inference rule of application,  $x$  must be a function type that accepts the type of  $(\lambda z . x)$ . Let  $\Gamma' \vdash z : \alpha$ , and type of  $x$  as  $\tau_x$ . Therefore,  $\Gamma' \vdash \lambda z : \alpha. x : \alpha \rightarrow \tau_x$ . Therefore,  $\tau_x \equiv (\alpha \rightarrow \tau_x) \rightarrow \tau$ . This is a recursive type, which is not constructable as it requires infinitely nested lambda abstractions that requires infinite reduction paths to reach a normal form. ■

### Problem

(2.6) Prove the pretyped term below is legal.

$$\lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x (\lambda z : \alpha. y)$$

Using the tree format and the flag format.

*Solution.* We suppose a context  $\Gamma \vdash y : \beta$  that obviously exists.

*Proof.*

$$\frac{\frac{}{x : (\alpha \rightarrow \beta) \rightarrow \alpha} \text{(Bound)} \quad \frac{\Gamma, z : \alpha \vdash y : \beta}{\Gamma \vdash (\lambda z : \alpha. y) : \alpha \rightarrow \beta} \text{(T-Abst)}}{\Gamma \vdash (\lambda z : \alpha. y) : \alpha \rightarrow \beta} \text{(T-App)} \quad \frac{\Gamma, x : (\alpha \rightarrow \beta) \rightarrow \alpha \vdash (x (\lambda z : \alpha. y)) : \alpha}{\Gamma \vdash \lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x (\lambda z : \alpha. y) : ((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow \alpha} \text{(T-Abst)}$$

A valid type could be given to the term. Therefore, the term is typable under an existing context. ■

The flag derivation is given below:

1.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>Bound</b>
2.	$z : \alpha$	<b>Bound</b>
3.	$y : \beta$	$\vdash \Gamma$
4.	$(\lambda z : \alpha. y) : \alpha \rightarrow \beta$	<b>3 T-Abst</b>
5.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>T-Var</b>

6.  $\frac{x (\lambda z : \alpha. y) : \beta}{}$  **5,4 T-App**
- 7.
- $\lambda x : ((\alpha \rightarrow \beta) \rightarrow \beta). x (\lambda z : \alpha. y)$
- $: (\alpha \rightarrow \beta) \rightarrow \beta \rightarrow \alpha$  **6 T-Abst**

### Problem

(2.7 a) Derive

$$f : A \rightarrow B \wedge g : B \rightarrow C \Rightarrow g \circ f : A \rightarrow C$$

Using the rules

$$\frac{f : A \rightarrow B, x \in A}{f(x) \in B} \text{(F-App)} \quad \frac{\forall x \in A, f(x) \in B}{f : A \rightarrow B} \text{(F-Abst)}$$

*Solution.*

*Proof.*

- |     |  |                      |
|-----|--|----------------------|
| 1.  | $f : A \rightarrow B \wedge g : B \rightarrow C$                                   | <b>Assumption</b>    |
| 2.  | $f : A \rightarrow B$  | $1 \wedge E$         |
| 3.  | $g : B \rightarrow C$  | $1 \wedge E$         |
| 4.  | $a \in A$  |                      |
| 5.  | $f(a) \in B$   | <b>3, 4 F-App</b>    |
| 6.  | $g(f(a)) \in C$  | <b>5, 4 F-App</b>    |
| 7.  | $(g \circ f)(a) \in C$   | <b>6 Compose Def</b> |
| 8.  | $\forall x \in A, (g \circ f)(x) \in C$  | $7 \forall E$        |
| 9.  | $g \circ f : A \rightarrow C$  | <b>8 F-Abst</b>      |
| 10. | $f : A \rightarrow B, g : B \rightarrow C \Rightarrow g \circ f : A \rightarrow C$ | $9 \Rightarrow I$    |

■

### Problem

(2.7 b) Give a derivation in natural deduction of the following:

$$(A \Rightarrow B) \Rightarrow ((B \Rightarrow C) \Rightarrow (A \Rightarrow C))$$

Using the rules

$$\frac{A \Rightarrow B \quad A}{B} (\Rightarrow E) \qquad \frac{\begin{array}{l} 1. \quad A \quad \text{Premise} \\ 2. \quad \vdots \\ 3. \quad B \end{array}}{A \Rightarrow B} (\Rightarrow I)$$

*Solution.*

*Proof.*

$$\begin{array}{ll} 1. & A \Rightarrow B \quad \text{Premise} \\ 2. & \begin{array}{l} B \Rightarrow C \\ \begin{array}{l} A \\ \begin{array}{l} B \\ \begin{array}{l} C \\ A \Rightarrow C \end{array} \end{array} \end{array} \quad \text{Premise} \\ 3. & \quad \quad \quad \text{Premise} \\ 4. & \quad \quad \quad 1, 3 \Rightarrow E \\ 5. & \quad \quad \quad 2, 4 \Rightarrow E \\ 6. & \quad \quad \quad 3-5 \Rightarrow I \\ 7. & \quad \quad \quad (B \Rightarrow C) \Rightarrow (A \Rightarrow C) \quad 2-6 \Rightarrow I \\ 8. & (A \Rightarrow B) \Rightarrow ((B \Rightarrow C) \Rightarrow (A \Rightarrow C)) \quad 1-7 \Rightarrow I \end{array}$$

■

### Problem

(2.7 c) Prove the following pre-typed term is legal using flag notation

$$\lambda z : \alpha. y (x z)$$

*Solution.*

*Proof.* Let  $\Gamma \vdash x : \alpha \rightarrow \beta, y : \beta \rightarrow \delta$  for some type  $\beta$  and  $\delta$ .

$$\begin{array}{ll} 1. & z : \alpha \quad \text{Bound} \\ 2. & \begin{array}{l} x : \alpha \rightarrow \beta \\ z : \alpha \end{array} \quad \neg \Gamma \\ 3. & \quad \quad \text{T-Var} \end{array}$$


- |    |   |                  |
|----|---|------------------|
| 4. | $x z : \beta$   | <b>2,3 T-App</b> |
| 5. | $y : \beta \rightarrow \delta$                            | $\neg \Gamma$    |
| 6. | $y (x z) : \delta$  | <b>5,4 T-App</b> |
| 7. | $\lambda z : \alpha. y (x z) : \alpha \rightarrow \delta$ | <b>6 T-Abst</b>  |

■

### Problem

(2.7 d) State the similarity between Q. 2.7 (a), (b), and (c).

*Solution.* All of these examples requires proving something about composing two maps together as like this:



### Problem

(2.8 a) Pre-type the bounding variables for the following term

$$\lambda x y . y (\lambda z . y x) : (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$$

*Solution.*

$$\lambda x : (\gamma \rightarrow \beta). y : ((\gamma \rightarrow \beta) \rightarrow \beta). y (\lambda z : \gamma. y x)$$

### Problem

(2.8 b) Give a derivation in tree format

*Solution.*

$$\begin{array}{c}
\text{(i)} \frac{}{x : (\gamma \rightarrow \beta)} \quad \text{(ii)} \frac{}{y : (\gamma \rightarrow \beta) \rightarrow \beta} \\
\text{(iii)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta), z : \gamma \vdash y x : \beta} \text{T-App} \\
\text{(v)} \frac{}{y : ((\gamma \rightarrow \beta) \rightarrow \beta)} \quad \text{(iv)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta) \vdash \lambda z : \gamma. y x : \gamma \rightarrow \beta} \text{T-Abst} \\
\text{(vi)} \frac{}{x : (\gamma \rightarrow \beta), y : ((\gamma \rightarrow \beta) \rightarrow \beta) \vdash y (\lambda z : \gamma. y x) : \beta} \text{T-App} \\
\text{(vii)} \frac{}{x : (\gamma \rightarrow \beta) \vdash \lambda y : ((\gamma \rightarrow \beta) \rightarrow \beta). y (\lambda z : \gamma. y x) : ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta} \text{T-Abst} \\
\text{(viii)} \frac{}{(\lambda x : (\gamma \rightarrow \beta). y : ((\gamma \rightarrow \beta) \rightarrow \beta). y (\lambda z : \gamma. y x)) : (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta} \text{T-Abst}
\end{array}$$

### Problem

(2.8 c) Sketch a diagram of tree structure of derivation

*Solution.* Trivial. See tree above

### Problem

(2.8 d) Transform the derivation into flag notation

*Solution.*

	1.	$x : \gamma \rightarrow \beta$	<b>Bound</b>
	2.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	<b>Bound</b>
	3.	$z : \gamma$	<b>Bound</b>
(ii)	4.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	<b>T-Var</b>
(i)	5.	$x : \gamma \rightarrow \beta$	<b>T-Var</b>
(iii)	6.	$y x : \beta$	<b>4,5 T-App</b>
(iv)	7.	$\lambda z : \gamma. y x : \gamma \rightarrow \beta$	<b>6 T-Abst</b>
(v)	8.	$y : (\gamma \rightarrow \beta) \rightarrow \beta$	<b>T-Var</b>
(vi)	9.	$y (\lambda z : \gamma. y x) : \beta$	<b>8,7 T-App</b>
		$\lambda y : (\gamma \rightarrow \beta) \rightarrow \beta. y (\lambda z : \gamma. y x)$	
(vii)	10.	$: ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$	<b>9 T-Abst</b>
(viii)	11.		
		$\lambda x : (\gamma \rightarrow \beta). \lambda y : (\gamma \rightarrow \beta) \rightarrow \beta. y (\lambda z : \gamma. y x)$	
		$: (\gamma \rightarrow \beta) \rightarrow ((\gamma \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$	<b>10 T-Abst</b>

### Problem

(2.9 a) Give derivations of the following judgement

$$x : \delta \rightarrow \delta \rightarrow \alpha, y : \gamma \rightarrow \alpha, z : \alpha \rightarrow \beta \vdash \\ \lambda u : \delta. \lambda v : \gamma. z (y v) : \delta \rightarrow \gamma \rightarrow \beta$$

*Solution.*

1.	$u : \delta$	<b>Bound</b>
2.	$v : \gamma$	<b>Bound</b>
3.	$y : \gamma \rightarrow \alpha$	<b>T-Var</b>
4.	$v : \gamma$	<b>T-Var</b>
5.	$y v : \alpha$	<b>3,4 T-App</b>
6.	$z : \alpha \rightarrow \beta$	<b>T-Var</b>
7.	$z (y v) : \beta$	<b>6,5 T-App</b>
8.	$\lambda v : \gamma. z (y v) : \gamma \rightarrow \beta$	<b>7 T-Abst</b>
9.	$\lambda u : \delta. \lambda v : \gamma. z (y v) : \delta \rightarrow \gamma \rightarrow \beta$	<b>8 T-Abst</b>

### Problem

(2.9 b) Give derivations of the following judgement

$$x : \delta \rightarrow \delta \rightarrow \alpha, y : \gamma \rightarrow \alpha, z : \alpha \rightarrow \beta \vdash \\ \lambda u : \delta. \lambda v : \gamma. z (x u u) : \delta \rightarrow \gamma \rightarrow \beta$$

*Solution.*

1.	$u : \delta$	<b>Bound</b>
2.	$v : \gamma$	<b>Bound</b>
3.	$x : \delta \rightarrow \delta \rightarrow \alpha$	<b>T-Var</b>
4.	$u : \delta$	<b>T-Var</b>
5.	$x u : \delta \rightarrow \alpha$	<b>3,4 T-App</b>
6.	$x u u : \alpha$	<b>5,4 T-App</b>
7.	$z : \alpha \rightarrow \beta$	<b>T-Var</b>
8.	$z (x u u) : \beta$	<b>7,6 T-App</b>
9.	$\lambda v : \gamma. z (x u u) : \gamma \rightarrow \beta$	<b>8 T-Abst</b>
10.	$\lambda u : \delta. \lambda v : \gamma. z (x u u) : \delta \rightarrow \gamma \rightarrow \beta$	<b>9 T-Abst</b>

### Problem

(2.10 a) Give derivation for

$$x\ z\ (y\ z)$$

*Solution.* Assume an context

$$\Gamma \vdash x : \alpha \rightarrow \beta \rightarrow \gamma$$

$$\Gamma \vdash y : \alpha \rightarrow \beta$$

$$\Gamma \vdash z : \alpha$$

1.  $x : \alpha \rightarrow \beta \rightarrow \gamma$     **T-Var**
2.  $y : \alpha \rightarrow \beta$     **T-Var**
3.  $z : \alpha$     **T-Var**
4.  $x\ z : \beta \rightarrow \gamma$     **1,3 T-App**
5.  $y\ z : \beta$     **2,3 T-App**
6.  $x\ z\ (y\ z) : \gamma$     **4,5 T-App**

### Problem

(2.10 b) Give derivation for

$$\lambda x : (\alpha \rightarrow \beta) \rightarrow \beta. x\ (y\ z)$$

*Solution.* Assume an context

$$\Gamma \vdash y : \gamma \rightarrow (\alpha \rightarrow \beta)$$

$$\Gamma \vdash z : \gamma$$

1.  $x : (\alpha \rightarrow \beta) \rightarrow \beta$     **Bound**
2.  $x : (\alpha \rightarrow \beta) \rightarrow \beta$     **T-Var**
3.  $y : \gamma \rightarrow \alpha \rightarrow \beta$     **T-Var**
4.  $z : \gamma$     **T-Var**
5.  $y\ z : \alpha \rightarrow \beta$     **3,4 T-App**
6.  $x\ (y\ z) : \beta$     **2,5 T-App**
7.  $\lambda x : (\alpha \rightarrow \beta) \rightarrow \beta. x\ (y\ z) : ((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow \beta$     **6 T-Abst**

### Problem

(2.10 c) Give derivation for

$$\lambda y : \alpha. \lambda z : \beta \rightarrow \gamma. z (x y y)$$

*Solution.* Assume a context

$$\Gamma \vdash x : \alpha \rightarrow \alpha \rightarrow \beta$$

1.	$y : \alpha$	<b>Bound</b>
2.	$z : \beta \rightarrow \gamma$	<b>Bound</b>
3.	$z : \beta \rightarrow \gamma$	<b>T-Var</b>
4.	$x : \alpha \rightarrow \alpha \rightarrow \beta$	<b>T-Var</b>
5.	$y : \alpha$	<b>T-Var</b>
6.	$x y : \alpha \rightarrow \beta$	<b>4,5 T-App</b>
7.	$x y y : \beta$	<b>6,5 T-App</b>
8.	$z (x y y) : \gamma$	<b>3,6 T-App</b>
9.	$\lambda z : \beta \rightarrow \gamma. z (x y y) : (\beta \rightarrow \gamma) \rightarrow \gamma$	<b>8 T-Abst</b>
10.	$\lambda y : \alpha. \lambda z : \beta \rightarrow \gamma. z (x y y) : \alpha \rightarrow (\beta \rightarrow \gamma) \rightarrow \gamma$	<b>8 T-Abst</b>

### Problem

(2.10 d) Give derivation for

$$\lambda x : \alpha \rightarrow \beta. y (x z) z$$

*Solution.* Consider a context

$$\begin{aligned} \Gamma \vdash z : \alpha \\ \Gamma \vdash y : \beta \rightarrow \alpha \rightarrow \gamma \end{aligned}$$

1.	$x : \alpha \rightarrow \beta$	<b>Bound</b>
2.	$x : \alpha \rightarrow \beta$	<b>T-Var</b>
3.	$z : \alpha$	<b>T-Var</b>
4.	$x z : \beta$	<b>2,3 T-App</b>
5.	$y : \beta \rightarrow \alpha \rightarrow \gamma$	<b>T-Var</b>
6.	$y (x z) : \alpha \rightarrow \gamma$	<b>5,4 T-App</b>
7.	$y (x z) z : \gamma$	<b>3,5 T-App</b>

8.  $\lambda x : \alpha \rightarrow \beta. y (x z) z : (\alpha \rightarrow \beta) \rightarrow \gamma$  **7 T-Abst**

### Problem

(2.11 a) Find an inhabitant of type and prove through derivation

$$(\alpha \rightarrow \alpha \rightarrow \gamma) \rightarrow \alpha \rightarrow \beta \rightarrow \gamma$$

*Solution.*

$$\lambda x : (\alpha \rightarrow \alpha \rightarrow \gamma). \lambda y : (\alpha). \lambda z : (\beta). x y y$$

*Proof.*

1.	$x : \alpha \rightarrow \alpha \rightarrow \gamma$	<b>Bound</b>
2.	$y : \alpha$	<b>Bound</b>
3.	$z : \beta$	<b>Bound</b>
4.	$x : \alpha \rightarrow \alpha \rightarrow \gamma$	<b>T-Var</b>
5.	$y : \alpha$	<b>Bound</b>
6.	$x y : \alpha \rightarrow \gamma$	<b>4,5 T-App</b>
7.	$x y y : \gamma$	<b>6,5 T-App</b>
8.	$\lambda z : \beta. x y y : \beta \rightarrow \gamma$	<b>7 T-Abst</b>
9.	$\lambda y : \alpha. \lambda z : \beta. x y y : \alpha \rightarrow \beta \rightarrow \gamma$	<b>8 T-Abst</b>
10.		

$$\lambda x : \alpha \rightarrow \alpha \rightarrow \gamma. \lambda y : \alpha. \lambda z : \beta. x y y$$

$$: (\alpha \rightarrow \alpha \rightarrow \gamma) \rightarrow \alpha \rightarrow \beta \rightarrow \gamma \quad \mathbf{8\ T-Abst}$$

■

### Problem

(2.11 b) Find an inhabitant of type and prove through derivation

$$((\alpha \rightarrow \gamma) \rightarrow \alpha) \rightarrow (\alpha \rightarrow \gamma) \rightarrow \beta \rightarrow \gamma$$

*Solution.*

$$\lambda x : (\alpha \rightarrow \gamma) \rightarrow \alpha. \lambda y : (\alpha \rightarrow \gamma). \lambda z : \beta. y (x y)$$

*Proof.*

1.	$x : (\alpha \rightarrow \gamma) \rightarrow \alpha$	<b>Bound</b>
2.	$y : \alpha \rightarrow \gamma$	<b>Bound</b>
3.	$z : \beta$	<b>Bound</b>
4.	$x : (\alpha \rightarrow \gamma) \rightarrow \alpha$	<b>T-Var</b>
5.	$y : \alpha \rightarrow \gamma$	<b>T-Var</b>
6.	$x y : \alpha$	<b>4,5 T-App</b>
7.	$y (x y) : \gamma$	<b>5,6 T-App</b>
8.	$\lambda z : \beta. x (x y) : \beta \rightarrow \gamma$	<b>7 T-Abst</b>
9.	$\lambda y : \alpha \rightarrow \gamma. \lambda z : \beta. x (x y) : (\alpha \rightarrow \gamma) \rightarrow \beta \rightarrow \gamma$	<b>7 T-Abst</b>
10.		
	$\lambda x : (\alpha \rightarrow \gamma) \rightarrow \alpha. \lambda y : \alpha \rightarrow \gamma. \lambda z : \beta. x (x y)$	
	$: ((\alpha \rightarrow \gamma) \rightarrow \alpha) \rightarrow (\alpha \rightarrow \gamma) \rightarrow \beta \rightarrow \gamma$	<b>7 T-Abst</b>

■

### Problem

(2.12 a) Construct a term of type

$$((\alpha \rightarrow \beta) \rightarrow \alpha) \rightarrow (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha$$

*Solution.*

$$\lambda x : (\alpha \rightarrow \beta) \rightarrow \alpha. \lambda y : \alpha \rightarrow \alpha \rightarrow \beta. x (\lambda z : \alpha. y z z)$$

*Proof.*

1.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>Bound</b>
2.	$y : \alpha \rightarrow \alpha \rightarrow \beta$	<b>Bound</b>
3.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>T-Var</b>
4.	$z : \alpha$	<b>Bound</b>
5.	$y : \alpha \rightarrow \alpha \rightarrow \beta$	<b>T-Var</b>
6.	$z : \alpha$	<b>Bound</b>
7.	$y z : \alpha \rightarrow \beta$	<b>5,6 T-App</b>
8.	$y z z : \beta$	<b>7,6 T-App</b>
9.	$\lambda z : \alpha. y z z : \alpha \rightarrow \beta$	<b>8 T-Abst</b>
10.	$x (\lambda z : \alpha. y z z) : \alpha$	<b>3,9 T-App</b>
11.	$\lambda y : \alpha \rightarrow \alpha \rightarrow \beta. x (\lambda z : \alpha. y z z) : (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha$	<b>3,9 T-App</b>

12.

$$\lambda x : (\alpha \rightarrow \beta) \rightarrow \alpha. \lambda y : \alpha \rightarrow \alpha \rightarrow \beta. x (\lambda z : \alpha. y z z)$$

$$: ((\alpha \rightarrow \beta) \rightarrow \alpha) \rightarrow (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha$$

3,9 T-App

■

### Problem

(2.12 b) Construct a term of type

$$((\alpha \rightarrow \beta) \rightarrow \alpha) \rightarrow (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \beta$$

*Solution.*

$$\lambda x : (\alpha \rightarrow \beta) \rightarrow \alpha. \lambda y : \alpha \rightarrow \alpha \rightarrow \beta. (\lambda z : \alpha. y z z)(x (\lambda z : \alpha. y z z))$$

*Proof.*

1.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>Bound</b>
2.	$y : \alpha \rightarrow \alpha \rightarrow \beta$	<b>Bound</b>
3.	$x : (\alpha \rightarrow \beta) \rightarrow \alpha$	<b>T-Var</b>
4.	$z : \alpha$	<b>Bound</b>
5.	$y : \alpha \rightarrow \alpha \rightarrow \beta$	<b>T-Var</b>
6.	$z : \alpha$	<b>Bound</b>
7.	$y z : \alpha \rightarrow \beta$	<b>5,6 T-App</b>
8.	$y z z : \beta$	<b>7,6 T-App</b>
9.	have $f := \lambda z : \alpha. y z z : \alpha \rightarrow \beta$	<b>8 T-Abst</b>
10.	$x f : \alpha$	<b>3,9 T-App</b>
11.	$f (x f) : \beta$	<b>9,10 T-App</b>
12.	$\lambda y : \alpha \rightarrow \alpha \rightarrow \beta. f (x f) : (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \beta$	<b>9,10 T-App</b>
	$\lambda x : (\alpha \rightarrow \beta) \rightarrow \alpha. \lambda y : \alpha \rightarrow \alpha \rightarrow \beta. f (x f)$	
13.	$: (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow (\alpha \rightarrow \alpha \rightarrow \beta) \rightarrow \beta$	<b>9,10 T-App</b>

■

### Problem

(2.13 a) Find a term of type

$$(\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \gamma$$

in context  $\Gamma$

$$x : \alpha \rightarrow \beta \rightarrow \gamma \in \Gamma$$

*Solution.*

$$\lambda u : \alpha \rightarrow \beta. \lambda v : \alpha. x v (u v)$$

*Proof.*

1.	$u : \alpha \rightarrow \beta$	<b>Bound</b>
2.	$v : \alpha$	<b>Bound</b>
3.	$x : \alpha \rightarrow \beta \rightarrow \gamma$	<b>T-Var</b>
4.	$v : \alpha$	<b>T-Var</b>
5.	$x v : \beta \rightarrow \gamma$	<b>3,4 T-App</b>
6.	$u : \alpha \rightarrow \beta$	<b>T-Var</b>
7.	$u v : \beta$	<b>6,4 T-App</b>
8.	$x v (u v) : \gamma$	<b>5,7 T-App</b>
9.	$\lambda v : \alpha. x v (u v) : \alpha \rightarrow \gamma$	<b>8 T-Abst</b>
10.	$\lambda u : \alpha \rightarrow \beta. \lambda v : \alpha. x v (u v) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \gamma$	<b>9 T-Abst</b>

■

### Problem

(2.13 b) Find a term of type

$$\alpha \rightarrow (\alpha \rightarrow \beta) \rightarrow \gamma$$

in context  $\Gamma$

$$x : \alpha \rightarrow \beta \rightarrow \alpha \rightarrow \gamma \in \Gamma$$

*Solution.*

$$\lambda u : \alpha. \lambda v : \alpha \rightarrow \beta. x u (v u) u$$

*Proof.*

1.	$u : \alpha$	<b>Bound</b>
2.	$v : \alpha \rightarrow \beta$	<b>Bound</b>
3.	$x : \alpha \rightarrow \beta \rightarrow \alpha \rightarrow \gamma$	<b>T-Var</b>
4.	$u : \alpha$	<b>Bound</b>
5.	$v : \alpha \rightarrow \beta$	<b>Bound</b>
6.	$v u : \beta$	<b>5,4 T-App</b>
7.	$x u : \beta \rightarrow \alpha \rightarrow \gamma$	<b>3,4 T-App</b>
8.	$x u (v u) : \alpha \rightarrow \gamma$	<b>7,6 T-App</b>
9.	$x u (v u) u : \gamma$	<b>8,4 T-App</b>
10.	$\lambda v : \alpha \rightarrow \beta. x u (v u) u : (\alpha \rightarrow \beta) \gamma$	<b>9 T-Abst</b>
11.	$\lambda u : \alpha. \lambda v : \alpha \rightarrow \beta. x u (v u) u : \alpha \rightarrow (\alpha \rightarrow \beta) \gamma$	<b>10 T-Abst</b>

■

### Problem

(2.13 c) Find a term of type

$$(\alpha \rightarrow \gamma) \rightarrow (\beta \rightarrow \alpha) \rightarrow \gamma$$

in context  $\Gamma$

$$x : (\beta \rightarrow \gamma) \rightarrow \gamma \in \Gamma$$

*Solution.*

$$\lambda u : \alpha \rightarrow \gamma. \lambda v : \beta \rightarrow \alpha. x (\lambda y : \beta. u (v y))$$

*Proof.*

1.	$u : \alpha \rightarrow \gamma$	<b>Bound</b>
2.	$v : \beta \rightarrow \alpha$	<b>Bound</b>
3.	$x : (\beta \rightarrow \gamma) \rightarrow \gamma$	<b>T-Var</b>
4.	$y : \beta$	<b>Bound</b>
5.	$v : \beta \rightarrow \alpha$	<b>T-Var</b>
6.	$y : \beta$	<b>T-Var</b>
7.	$v y : \alpha$	<b>5,6 T-App</b>
8.	$u : \alpha \rightarrow \gamma$	<b>T-Var</b>
9.	$u (v y) : \gamma$	<b>8,7 T-App</b>
10.	$\lambda y : \beta. u (v y) : \beta \rightarrow \gamma$	<b>9 T-Abst</b>

11.	$\frac{x (\lambda y : \beta. u (v y)) : \gamma}{\lambda v : \beta \rightarrow \alpha. x (\lambda y : \beta. u (v y)) : (\beta \rightarrow \alpha) \rightarrow \gamma}$	<b>10,3 T-App</b>
12.		<b>11 T-Abst</b>
13.		
	$\lambda u : \alpha \rightarrow \gamma. \lambda v : \beta \rightarrow \alpha. x (\lambda y : \beta. u (v y))$	
	$: (\alpha \rightarrow \gamma) \rightarrow (\beta \rightarrow \alpha) \rightarrow \gamma$	<b>12 T-Abst</b>

■

### Problem

(2.14) Find an inhabitant of type  $\alpha \rightarrow \beta \rightarrow \gamma$  in the context  $\Gamma$

$$x : (\gamma \rightarrow \beta) \rightarrow \alpha \rightarrow \gamma \in \Gamma$$

*Solution.*

$$\lambda u : \alpha. \lambda v : \beta. x (\lambda z : \gamma. v) u$$

*Proof.*

1.	$u : \alpha$	<b>Bound</b>
2.	$v : \beta$	<b>Bound</b>
3.	$x : (\gamma \rightarrow \beta) \rightarrow \alpha \rightarrow \gamma$	<b>T-Var</b>
4.	$z : \gamma$	<b>Bound</b>
5.	$v : \beta$	<b>T-Var</b>
6.	$\lambda z : \gamma. v : \gamma \rightarrow \beta$	<b>5 T-Abst</b>
7.	$x (\lambda z : \gamma. v) : \alpha \rightarrow \gamma$	<b>3,6 T-App</b>
8.	$u : \alpha$	<b>T-Var</b>
9.	$x (\lambda z : \gamma. v) u : \gamma$	<b>7,8 T-App</b>
10.	$\lambda v : \beta. x (\lambda z : \gamma. v) u : \beta \rightarrow \gamma$	<b>9 T-Abst</b>
11.	$\lambda u : \alpha. \lambda v : \beta. x (\lambda z : \gamma. v) u : \alpha \rightarrow \beta \rightarrow \gamma$	<b>10 T-Abst</b>

■

### Problem

(2.15) Prove the thinning lemma via induction.

*Solution.*

---

*Lemma 1.*      **Thinning Lemma.** Given two contexts  $\Gamma' \subseteq \Gamma$ ,

$$\Gamma' \vdash M : \sigma \Rightarrow \Gamma \vdash M : \sigma$$


---

We can do structural induction over the structure of  $M$  by the finite syntactic constructors of  $\lambda$  terms.

*Case 1 : Variable.* By the generation lemma  $M : \sigma \in \Gamma'$ . By the definition of subcontextes,  $M : \sigma \in \Gamma$ . It follows from the T-Var rule that  $\Gamma \vdash M : \sigma$ . ■

*Case 2 : Application.* Therefore the judgement must be of form

$$\Gamma' \vdash A B : \sigma$$

Where  $M = A B$ . By the generation lemma, there exists a type  $\tau$  such that  $\Gamma' \vdash A : \tau \rightarrow \sigma$  and  $\Gamma' \vdash B : \tau$ . By the principle of induction the thinning lemma holds for the terms  $A$  and  $B$ . By plugging in  $\Gamma \vdash A : \tau \rightarrow \sigma, B : \tau$ , the T-App rule proves  $\Gamma \vdash M : \sigma$ . ■

*Case 3 : Abstraction.* Therefore  $M$  is of a function type. We denote  $M$  as a abstraction term

$$\lambda x : \alpha. N : \alpha \rightarrow \beta$$

where  $x \notin \text{FR } N$ . By the generation lemma,  $\Gamma', x : \alpha \vdash N : \beta$ .

By the principle of induction, the thinning lemma already holds for  $N : \beta$ . Because  $\Gamma', x : \alpha \subseteq \Gamma, x : \alpha$ , the thinning lemma proves  $\Gamma, x : \alpha \vdash N : \beta$ . By the T-Abst rule,  $\Gamma \vdash \lambda x : \alpha. N : \alpha \rightarrow \beta$ . ■

### Problem

(2.16) Prove the subterm lemma.

*Solution.*

---

*Lemma 2.*      **Subterm Lemma.** If  $M \in \Lambda_{\mathbb{T}}$  is legal, then all subterms of  $M$  are.

---

Let's formalize the lemma. The definition of a legal term is a term  $M$  with the existence of a context  $\Gamma$  and a type  $\tau$  such that  $\Gamma \vdash M : \tau$ , in other words,  $M$  is typable. Because  $M$  is typable iff there's an applicable inference rule and each term appearing in the premise of each inference rule is an immediate subterm of  $M$  that

could construct  $M$ , structural induction could be done down the typing tree as it is also an induction down the term's structure, inducting over each subterm.

*Case 1 : Variable.*  $M$  is the only subterm of  $M$  thus is trivially typable and legal. ■

*Case 2 : Application.* Therefore the derivation must be of form

$$\frac{\Gamma \vdash A : \tau \rightarrow \sigma \quad \Gamma \vdash B : \tau}{\Gamma \vdash A B : \sigma} \text{T-App}$$

Where  $M = A B$ . From the generation lemma, those  $A$  and  $B$  must exist typable, thus legal. By the principle of induction, the subterm lemma holds for  $A$  and  $B$ . Denote the lemma as  $P(\Lambda_{\mathbb{T}})$ :

$$\begin{aligned} & P(A B) \wedge P(A) \wedge P(B) \wedge (\forall a \in \text{Sub } A, P(a)) \wedge (\forall b \in \text{Sub } B, P(b)) \\ &= \forall m \in \{A B, A, B\} \cup \text{Sub } A \cup \text{Sub } B, P(m) \\ &= \forall m \in \{A B\} \cup \text{Sub } A \cup \text{Sub } B, P(m) \\ &= \forall m \in \text{Sub } A B, P(m) = P(A B) = P(M) \end{aligned}$$

■

*Case 3 : Abstraction.* Therefore the derivation must be of form

$$\frac{\Gamma, x : \alpha \vdash N : \beta}{\Gamma \vdash \lambda x : \alpha. N : \alpha \rightarrow \beta} \text{T-Abst}$$

Where  $M = \lambda x : \alpha. N$  and  $x \notin \text{FR } N$ . By the generation lemma,  $N$  is typable under the context  $\Gamma, x : \alpha$ , thus valid. By the inductive hypothesis, the subterm lemma holds for  $N$ . Denote the lemma as  $P(\Lambda_{\mathbb{T}})$

$$\begin{aligned} & P(\lambda x : \alpha. N) \wedge (P(N) \wedge \forall n \in \text{Sub } N, P(n)) \\ &= \forall m \in \{\lambda x : \alpha. N, N\} \cup \text{Sub } N, P(m) \\ &= \forall m \in \{\lambda x : \alpha. N\} \cup \text{Sub } N, P(m) = P(\lambda x : \alpha. N) = P(M) \end{aligned}$$

■

### Problem

(2.17) Prove the uniqueness of types lemma.

*Solution.*

---

**Lemma 3.** Assume  $\Gamma \vdash M : \sigma$  and  $\Gamma \vdash M : \tau$ . Then  $\sigma = \tau$ .

---

Again do induction on the construction of  $M$ .

*Case 1 : Variable.* By the generation lemma,  $M : \sigma \in \Gamma$  and  $M : \tau \in \Gamma$  By the definition of contexts. Because the context could only provide one judgement per variable,  $M$  must have a unique type. ■

*Case 2 : Application.*  $M$  must be of form  $A B$ . By the generation lemma, two judgements could be obtained where  $\alpha, \beta \in \mathbb{T}$ :

$$\Gamma \vdash A : \alpha \rightarrow \sigma, B : \alpha$$

$$\Gamma \vdash A : \beta \rightarrow \tau, B : \beta$$

By the inductive hypothesis, the uniqueness lemma already holds for  $A$  and  $B$ . Therefore  $\alpha \equiv \beta$ . Because  $A$  is a function type, it could be expanded into two forms:

$$\Gamma \vdash (\lambda x : \alpha. N) : \alpha \rightarrow \sigma$$

$$\Gamma \vdash (\lambda x : \beta. N) : \beta \rightarrow \tau$$

By the generation lemma and the equivalence  $\alpha \equiv \beta$ , the above further deduce to

$$\Gamma, x : \alpha \vdash N : \sigma$$

$$\Gamma, x : \alpha \vdash N : \tau$$

By the principle of induction, the uniqueness lemma holds for  $N$ , therefore  $\sigma \equiv \tau$ . By substituting  $\tau$  for  $\sigma$ , the final type for  $M$  should be the same in the conclusion for both judgements. ■

*Case 3 : Abstraction.* Because  $M$  is a function type, two judgements could be obtained for some  $\alpha, \beta \in \mathbb{T}$

$$\Gamma \vdash \lambda x : \gamma. N : \sigma \text{ where } \sigma \equiv \gamma \rightarrow \alpha$$

$$\Gamma \vdash \lambda x : \gamma. N : \tau \text{ where } \tau \equiv \gamma \rightarrow \beta$$

By applying the generation lemma

$$\Gamma, x : \gamma \vdash N : \alpha$$

$$\Gamma, x : \gamma \vdash N : \beta$$

By the principle of induction,  $N$  already conform to the lemma, thus  $\alpha \equiv \beta$ . Therefore,  $\gamma \rightarrow \alpha \equiv \gamma \rightarrow \beta$ . ■

### Problem

(2.18) Prove the *Subject Reduction* lemma.

---

*Lemma 4.* If  $\Gamma \vdash L : \rho$  and  $L \xrightarrow[\beta]{\rightarrow} L'$ , then  $\Gamma \vdash L' : \rho$

---

*Solution.* Proof by generation of  $L'$ : which is by establishing all possibilities of how  $L'$  was derived.

*Basis : Direct Reduction.*

$$L \equiv (\lambda x : \sigma. M) N \text{ and } L' \equiv M [x := N]$$

By the generation lemma and some destructuring we have  $M : \gamma$  under  $\Gamma, x : \sigma$  and  $N : \sigma$  under  $\Gamma$ . By applying the T-App rule,  $L : \gamma$ . By the uniqueness of types lemma,  $\gamma \equiv \rho$ . Now we define  $\Gamma' \equiv \emptyset$  and  $\Gamma'' \equiv \Gamma$ , by the substitution lemma we have  $\Gamma \vdash M [x := N] : \gamma$ . We substitute  $\gamma \equiv \rho$  back and get  $\Gamma \vdash M [x := N] : \rho$ . ■

*Case 1 : Left Term Reduction.*

$$L \equiv M N \text{ and } L' \equiv M' N \text{ where } M \xrightarrow[\beta]{\rightarrow} M'$$

We type the terms  $M : \sigma \rightarrow \rho$  and  $N : \sigma$ . By the inductive hypothesis,  $\Gamma \vdash M' : \sigma \rightarrow \rho$ . By the T-App rule,  $\Gamma \vdash M' N : \rho$ . ■

*Case 2 : Left Term Reduction.*

$$L \equiv M N \text{ and } L' \equiv M N' \text{ where } N \xrightarrow[\beta]{\rightarrow} N'$$

We type the terms  $M : \sigma \rightarrow \rho$  and  $N : \sigma$ . By the inductive hypothesis,  $\Gamma \vdash N' : \sigma$ . By the T-App rule,  $\Gamma \vdash M N' : \rho$ . ■

*Case 3 : Reduction In Body.*

$$L \equiv (\lambda x : \tau. M) \text{ and } L' \equiv (\lambda x : \tau. M') \text{ where } M \xrightarrow[\beta]{\rightarrow} M'$$

Let  $\sigma \in \mathbb{T}$  where  $\rho \equiv \tau \rightarrow \sigma$ . By the generation lemma  $\Gamma, x : \tau \vdash M : \sigma$ . By the inductive hypothesis,  $\Gamma, x : \tau \vdash M' : \sigma$ , and by the T-Abst rule,  $\Gamma \vdash (\lambda x : \tau. M') : \tau \rightarrow \sigma$ . Therefore,  $\Gamma \vdash L' : \rho$  by substituting  $\tau \rightarrow \sigma$  for  $\rho$ . ■

—

Completed Dec 15 2:24 am.