

EXERCISES

CHAPTER 2

SEAN LI ¹

1. Reducted

Problem

Type the following terms

$xyx \quad xyx \quad xyx \quad x(xy) \quad x(yx)$

Solution. The first term cannot be typed.

Proof. $xyx = (xx)y$. Therefore, x is a function type, denote it as $\tau \rightarrow \sigma$. By the application rule, a subterm applied to x must be of τ , which means that the application xx is not legally typed. ■

The second one is typable where $x : \tau \rightarrow \tau \rightarrow \sigma$ and $y : \tau$.

a	1.	$x : \tau \rightarrow \tau \rightarrow \sigma$	ctx
b	2.	$y : \tau$	ctx
1	3.	$xy : \tau \rightarrow \sigma$	T-App (a)
c	4.	$xyx : \sigma$	T-App (1)

The third term is not typable.

Proof. Assume $xyx = (xy)x$ is typable. Therefore, $x : \tau$ where $\tau = \sigma \rightarrow \tau \rightarrow \alpha$ and $y : \sigma$. One can construct an infinite chain of function type by substituting τ : $\tau = \sigma \rightarrow (\sigma \rightarrow (\sigma \rightarrow \dots \rightarrow \alpha) \rightarrow \alpha) \rightarrow \alpha$. By induction, it can be proven that only lambda abstractions can construct function types, meaning that the term is of form

$$(\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots))y(\lambda n : \tau. \lambda m : \tau. \dots (\lambda a : \sigma. \lambda b : \sigma. \dots))$$

meaning that an infinite reduction path is needed. This is impossible in STLC. ■

The fourth type is typable where $x : (\tau \rightarrow \tau)$ and $y : \tau$.

a 1. $x : \tau \rightarrow \tau$ **ctx**
 b 2. $y : \tau$ **ctx**
1 3. $\frac{}{xy : \tau}$ **T-App (b) on (a)**
2 4. $x(xy) : \tau$ **T-App (1) on (a)**

The fifth term is typable where $x : (\tau \rightarrow \sigma)$ and $y : (\tau \rightarrow \sigma) \rightarrow \tau$:

a 1. $x : \tau \rightarrow \sigma$ **ctx**
 b 2. $y : (\tau \rightarrow \sigma) \rightarrow \tau$ **ctx**
1 3. $\frac{}{yx : \tau}$ **T-App (a) on (b)**
2 4. $x(yx) : \sigma$ **T-App (1) on (a)**

Problem

Find types for zero, one, and two

Solution. Term for zero is

$$\text{zero} := \lambda f x. x$$

Here x is only used as a

$$\text{zero} := \lambda f : \alpha. \lambda x : \beta. x$$

Type derivation shown as below:

a 1. $f : \alpha$ **bind**
 b 2. $\frac{}{x : \beta}$ **bind**
1 3. $\frac{}{x : \beta}$ **T-Var (b)**
2 4. $\frac{}{\lambda x. x : \beta \rightarrow \beta}$ **T-Abst (1)**
3 5. $\lambda f : \alpha. x : \beta. x : \alpha \rightarrow \beta \rightarrow \beta$ **T-Abst (2)**

Term for one is

$$\text{one} := \lambda f x. f x$$

Let f be an arbitrary function type that consumes x

$$\text{one} := \lambda f : \alpha \rightarrow \beta. x : \alpha. f x$$

Type derivation shown as below

<i>a</i>	1.	$f : \alpha \rightarrow \beta$	bind
<i>b</i>	2.	$x : \alpha$	bind
1	3.	$fx : \beta$	T-App (b) on a
2	4.	$\lambda x. fx : \alpha \rightarrow \beta$	T-Abst (1)
3	5.	$\lambda f : \alpha \rightarrow \beta. x : \alpha. fx : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$	T-Abst (2)

Same type signatures can be given to two

$$\text{two} := \lambda f : \alpha \rightarrow \beta. \lambda x : \alpha. ffx$$

Type derivation shown as below

<i>a</i>	1.	$f : \alpha \rightarrow \beta$	bind
<i>b</i>	2.	$x : \alpha$	bind
1	3.	$fx : \beta$	T-App (b) on (a)
2	4.	$ffx : \beta$	T-App (2) on (b)
3	5.	$\lambda x. ffx : \alpha \rightarrow \beta$	T-Abst (2)
4	6.	$\lambda f : \alpha \rightarrow \beta. x : \alpha. ffx : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$	T-Abst (3)

Problem

Find types for

$$K := \lambda xy. x$$

$$S := \lambda xyz. xz(yz)$$

Solution. There are no occurrences of application in K 's subterms. Therefore all its binding variables could be given a simple base type.

$$K := \lambda x : \alpha. \lambda y : \beta. x$$

Type derivation shown as below

<i>a</i>	1.	$x : \alpha$	bind
<i>b</i>	2.	$y : \beta$	bind
1	3.	$x : \alpha$	T-Var
2	4.	$\lambda y : \beta. x : \beta \rightarrow \alpha$	T-Abst on (1)
3	5.	$\lambda x : \alpha. \lambda y : \beta. x : \alpha \rightarrow \beta \rightarrow \alpha$	T-Abst on (2)

For the S combinator, no term was applied to z . Therefore it can be given a simple base type α . As z was applied to y , it implies that $y : \alpha \rightarrow \beta$ for some output type β . As x takes z and (yz) , it must be of type $\alpha \rightarrow \beta \rightarrow \delta$.

$$S := \lambda x : \alpha \rightarrow \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z. \alpha. xz(yz)$$

Complete type derivation shown as below:

$$a \quad 1. \quad x : \alpha \rightarrow \beta \rightarrow \delta$$

bind

$$b \quad 2. \quad \left| \begin{array}{l} y : \alpha \rightarrow \beta \end{array} \right.$$

bind

$$c \quad 3. \quad \left| \begin{array}{l} \left| \begin{array}{l} z : \alpha \end{array} \right. \end{array} \right.$$

bind

$$1 \quad 4. \quad \left| \begin{array}{l} \left| \begin{array}{l} \left| \begin{array}{l} yz : \beta \end{array} \right. \end{array} \right. \end{array} \right.$$

T-App (c) on (b)

$$2 \quad 5. \quad \left| \begin{array}{l} \left| \begin{array}{l} \left| \begin{array}{l} xz : \beta \rightarrow \delta \end{array} \right. \end{array} \right. \end{array} \right.$$

T-App (c) on (a)

$$3 \quad 6. \quad \left| \begin{array}{l} \left| \begin{array}{l} \left| \begin{array}{l} \underline{xz(yz) : \delta} \end{array} \right. \end{array} \right. \end{array} \right.$$

T-App (1) on (2)

$$4 \quad 7. \quad \left| \begin{array}{l} \left| \begin{array}{l} \left| \begin{array}{l} \underline{\lambda z : \alpha. xz(yz) : \alpha \rightarrow \delta} \end{array} \right. \end{array} \right. \end{array} \right.$$

T-Abstr on (3)

$$5 \quad 8. \quad \left| \begin{array}{l} \left| \begin{array}{l} \left| \begin{array}{l} \underline{\lambda y : \alpha \rightarrow \beta. \lambda z. \alpha. xz(yz) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta} \end{array} \right. \end{array} \right. \end{array} \right.$$

T-Abstr on (4)

$$6 \quad 9. \quad \lambda x : \alpha \rightarrow \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. xz(yz) : (\alpha \rightarrow \beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$$

T-Abstr on (5)

Problem

Type the bound variables

$$\lambda xyz. x(yz)$$

$$\lambda xyz. y(xz)z$$

Solution. For the first term, z had nothing applied to it. Therefore it could be given a simple base type α . z was applied to y , therefore $y : \alpha \rightarrow \beta$ to satisfy the application rule. Because the application yielded a type of β , by the application rule $x : \beta \rightarrow \delta$ for some type δ .

$$\lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz)$$

Complete type derivation shown below

$$a \quad 1. \quad x : \beta \rightarrow \delta$$

bind

$$b \quad 2. \quad \left| y : \alpha \rightarrow \beta \right.$$

bind

$$c \quad 3. \quad \left| \left| z : \alpha \right. \right.$$

bind

$$1 \quad 4. \quad \left| \left| \left| yz : \beta \right. \right. \right.$$

T-App (c) on (b)

$$2 \quad 5. \quad \left| \left| \left| x(yz) : \delta \right. \right. \right.$$

T-App (1) on (a)

$$3 \quad 6. \quad \left| \left| \lambda z : \alpha. x(yz) : \alpha \rightarrow \delta \right. \right.$$

T-Abst on (2)

$$4 \quad 7. \quad \left| \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz) : (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta \right.$$

T-Abst on (3)

$$5 \quad 8. \quad \lambda x : \beta \rightarrow \delta. \lambda y : \alpha \rightarrow \beta. \lambda z : \alpha. x(yz) : (\beta \rightarrow \delta) \rightarrow (\alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \delta$$

T-Abst on (4)

In the second term z could still be given a simple base type $z : \alpha$. Therefore $x : \alpha \rightarrow \beta$ for some type β . y takes $xz : \beta$ and $z : \alpha$, therefore it is of type $y : \beta \rightarrow \alpha \rightarrow \delta$ for some δ .

$$\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z : \alpha. y(xz)z$$

. Complete type derivation shown below

$$a \quad 1. \quad x : \alpha \rightarrow \beta$$

bind

$$b \quad 2. \quad \left| y : \beta \rightarrow \alpha \rightarrow \delta \right.$$

bind

$$c \quad 3. \quad \left| \left| z : \alpha \right. \right.$$

bind

$$1 \quad 4. \quad \left| \left| \left| xz : \beta \right. \right. \right.$$

T-App (c) on (b)

2 5. $\left| \left| \left| y(xz) : \alpha \rightarrow \delta \right. \right. \right|$

T-App (1) on (b)

3 6. $\left| \left| \left| \underline{y(xz)z : \delta} \right. \right. \right|$

T-App (c) on (2)

4 7. $\left| \left| \underline{\lambda z : \alpha. y(xz)z : \alpha \rightarrow \delta} \right. \right|$

T-Abst on (3)

5 8. $\left| \underline{\lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y(xz)z : (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta} \right|$

T-Abst on (4)

6 9. $\lambda x : \alpha \rightarrow \beta. \lambda y : \beta \rightarrow \alpha \rightarrow \delta. \lambda z. y(xz)z : (\alpha \rightarrow \beta) \rightarrow (\beta \rightarrow \alpha \rightarrow \delta) \rightarrow \alpha \rightarrow \delta$

T-Abst on (5)

Problem

Try to type the following terms, and prove if not typable.

$\lambda xy. x(\lambda z. y)y$

$\lambda xy. x(\lambda z. x)y.$

Solution. The first term is trivially typable.

a	1.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	bind
b	2.	$y : \alpha$	bind
1	3.	$x : (\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta$	T-Var
c	4.	$z : \delta$	bind
2	5.	$\left \underline{y : \alpha} \right $	T-Var
3	6.	$\lambda z : \delta. y : \delta \rightarrow \alpha$	T-Abst on (1)
4	7.	$x(\lambda z : \delta. y) : \alpha \rightarrow \beta$	T-App (3) on (1)
5	8.	$\left \underline{x(\lambda z : \delta. y)y : \beta} \right $	T-App (b) on (4)
6	9.	$\left \underline{\lambda y : \alpha. x(\lambda z : \delta. y)y : \alpha \rightarrow \beta} \right $	T-Abst on (5)
7	10.		

$\lambda x : ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \lambda y : \alpha. x(\lambda z : \delta. y)y$

$: ((\delta \rightarrow \alpha) \rightarrow \alpha \rightarrow \beta) \rightarrow \alpha \rightarrow \beta$

T-Abst on (5)

The second term is not typable in STLC.

Proof. By induction on the type inference rule that constructed the type judgement for subterm $x(\lambda z.x)$. Because the term is an application, the only rule that applies is the application rule.

We denote the context inside the abstraction as Γ' . Suppose $\mathcal{J} \equiv \Gamma' \vdash x(\lambda z.x) : \tau$. By the inference rule of application, x must be a function type that accepts the type of $(\lambda z.x)$. Let $\Gamma' \vdash z : \alpha$, and type of x as τ_x . Therefore, $\Gamma' \vdash \lambda z : \alpha. x : \alpha \rightarrow \tau_x$. Therefore, $\tau_x = (\alpha \rightarrow \tau_x) \rightarrow \tau$. This is a recursive type, which is not constructable as it requires infinitely nested lambda abstractions that requires infinite reduction paths to reach a normal form. ■

Problem

Prove the pretyped term below is legal.

$$\lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x(\lambda z : \alpha. y)$$

Using the tree format and the flag format.

Solution. We suppose a context $\Gamma \vdash y : \beta$ that obviously exists.

Proof.

$$\frac{\frac{x : (\alpha \rightarrow \beta) \rightarrow \alpha}{\Gamma, x : (\alpha \rightarrow \beta) \rightarrow \alpha \vdash (x(\lambda z : \alpha. y)) : \alpha} \text{(Bound)} \quad \frac{\Gamma, z : \alpha \vdash y : \beta}{\Gamma \vdash (\lambda z : \alpha. y) : \alpha \rightarrow \beta} \text{(T-Abst)}}{\Gamma \vdash \lambda x : ((\alpha \rightarrow \beta) \rightarrow \alpha). x(\lambda z : \alpha. y) : ((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow \alpha} \text{(T-App)} \text{(T-Abst)}$$

A valid type could be given to the term. Therefore, the term is typable under an existing context. ■

The flag derivation is given below:

- | | | | |
|-----|----|--|-------------------------|
| (a) | 1. | $x : (\alpha \rightarrow \beta) \rightarrow \alpha$ | Bound |
| (b) | 2. | $z : \alpha$ | Bound |
| (1) | 3. | $y : \beta$ | $\neg \Gamma$ |
| (2) | 4. | $(\lambda z : \alpha. y) : \alpha \rightarrow \beta$ | T-Abst on (1) |
| (3) | 5. | $x(\lambda z : \alpha. y) : \beta$ | T-App (2) on (a) |
| (4) | 6. | | |

$$\lambda x : ((\alpha \rightarrow \beta) \rightarrow \beta). x(\lambda z : \alpha. y)$$

$$: (\alpha \rightarrow \beta) \rightarrow \beta \rightarrow \alpha \quad \textbf{T-Abst on (3)}$$

Problem

Derive

$$f : A \rightarrow B \wedge g : B \rightarrow C \vdash g \circ f : A \rightarrow C$$

Using the rules

$$\frac{f : A \rightarrow B, c \in A}{f(c) \in B} \text{(F-App)}$$

$$\frac{\forall x \in A, f(x) \in B}{f : A \rightarrow B} \text{(F-Abst)}$$