

EXERCISES

CHAPTER 5

SEAN LI ¹

1. Reducted


Definition Some rules for reference.

$$\begin{array}{c} \frac{}{\emptyset \vdash * : \square} \text{Sort} \quad \frac{\Gamma \vdash A : s}{\Gamma, x : A \vdash x : A} \text{Var} \quad \frac{\Gamma \vdash A : B \quad \Gamma \vdash C : s}{\Gamma, x : C \vdash A : B} \text{Weak} \\[10pt] \frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : s}{\Gamma \vdash \Pi x : A . B : s} \text{Form} \quad \frac{\Gamma \vdash M : \Pi x : A . B \quad \Gamma \vdash N : A}{\Gamma \vdash M N : B[x := N]} \text{App} \\[10pt] \frac{\Gamma, x : A \vdash M : B \quad \Gamma \vdash \Pi x : A . B : s}{\Gamma \vdash \lambda x : A . M : \Pi x : A . B} \text{Abst} \\[10pt] \frac{\Gamma \vdash A : B \quad B \stackrel{\beta}{=} B' \quad \Gamma \vdash B' : s}{\Gamma \vdash A : B'} \text{Conv} \end{array}$$

Problem

(5.1) Give a diagram of the tree corresponding to the complete tree derivation of line 18 of Section 5.3 (P 107)

Solution.



Problem

(5.2) Give a complete λP derivation of

$$S : * \vdash S \rightarrow S \rightarrow * : \square$$

In tree format and flag format.

Solution.

Tree Derivation.

$$\begin{array}{c}
(4) \frac{\vdash * : \square \quad \vdash * : \square}{\text{Weak}} \\
(6) \frac{S : * \vdash * : \square \quad (3) S : * \vdash S : *}{\text{Weak}} \\
(7) \frac{(3) S : * \vdash S : *}{S : * \vdash S \rightarrow * : \square}
\end{array}$$

$$\begin{array}{c}
(3) \frac{\vdash * : \square}{S : * \vdash S : *} \text{Var} \quad (9) \frac{(7) S : * \vdash S \rightarrow * : \square \quad (3) S : * \vdash S : *}{S : *, x : S \vdash S \rightarrow * : \square} \text{Weak} \\
\hline
S : * \vdash S \rightarrow S \rightarrow * : \square \text{Form}
\end{array}$$

■

Flag Derivation.

1.	$* : \square$	Sort
2.	$S : *$	
3.	$S : *$	1 Var
4.	$* : \square$	1,1 Weak
5.	$x : S$	
6.	$\boxed{* : \square}$	4,3 Weak
7.	$S \rightarrow * : \square$	3,6 Form
8.	$x : S$	
9.	$\boxed{S \rightarrow * : \square}$	7,3 Weak
10.	$S \rightarrow S \rightarrow * : \square$	3,9 Form

■

Problem

(5.3) Derive

$$S : *, Q : S \rightarrow S \rightarrow * \vdash \Pi x : S . \Pi y : S . Q x y : *$$

Solution.

1.	$* : \square$	Sort
2.	$S : *$	
3.	$S : *$	1 Var
4.	$* : \square$	1,1 Weak
5.	$x : S$	
6.	$\boxed{* : \square}$	4,3 Weak
7.	$S \rightarrow * : \square$	3,6 Form
8.	$x : S$	
9.	$\boxed{S \rightarrow * : \square}$	7,3 Weak
10.	$S \rightarrow S \rightarrow * : \square$	3,9 Form
11.	$Q : S \rightarrow S \rightarrow *$	

12.	$Q : S \rightarrow S \rightarrow *$	10 Var
13.	$S : *$	3,10 Weak
14.	$* : \square$	4,10 Weak
15.	$x : S$	
16.	$* : \square$	14,13 Weak
17.	$S : *$	13,13 Weak
18.	$x : S$	13 Var
19.	$Q : S \rightarrow S \rightarrow *$	12,13 Weak
20.	$y : S$	
21.	$y : S$	17 Var
22.	$Q : S \rightarrow S \rightarrow *$	19,17 Weak
23.	$x : S$	18,17 Weak
24.	$Q x : S \rightarrow *$	22,23 App
25.	$Q x y : *$	24,21 App
26.	$\Pi y : S . Q x y : *$	17,25 Form
27.	$\Pi x : S . \Pi y : S . Q x y : *$	13,26 Form

Problem

(5.4) Prove that $*$ is the only valid kind in λP .

Solution.

Proof. The only possible way to construct a new kind is through the Form rule and the Sort axiom. Because we are trying to construct a kind, s here stands for \square .

$$\frac{\Gamma \vdash A : * \quad \Gamma, x : A \vdash B : \square}{\Gamma \vdash \Pi x : A . B : \square} \text{Form}$$

One could only construct new kinds with kinds, which requires $A : \square$ and $B : \square$. This contradicts with $A : *$. ■