




Hasso
Plattner
Institut

Hyperbolic Random Graphs

Degree Sequence and Clustering

Algorithm Engineering Group


Janosch Ruff



Network Features - Distributions


The most commonly considered network features are distributions over parts of the network.

- Degrees
- Assortativity
- Distances
- Centrality
- Clustering




[The Structure and Function of Complex Networks, M. E. J. Newman, Computer Physics Communications 2003]

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$



Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$

The arithmetic mean of the degree distribution is the **average degree** of the network.


Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

The arithmetic mean of the degree distribution is the **average degree** of the network.

- Homogeneous




Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$



The arithmetic mean of the degree distribution is the **average degree** of the network.

- Homogeneous



Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous




Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous



Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.

For directed networks **in- and out-degree distributions** are often considered separately.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.


$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$


- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$


- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.


$$c(G) = \frac{3 |\Delta|}{|\bullet\bullet|}$$

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.

$$c(G) = \frac{3 |\Delta|}{|\bullet\bullet|}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)\|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.

$$c(G) = \frac{3 |\Delta|}{|\bullet\bullet|}$$



Compared to the average local clustering coefficient, the global one basically computes the ratio of the means rather than the mean of the ratios. It does not weight the contributions of low-degree nodes as much.

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$



Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$



$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient
- ... a logarithmic diameter



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient
- ... a logarithmic diameter
- ... negative assortativity



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Parameters of HRG


Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$

Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Parameters of HRG

Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$




Parameters of HRG

Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".




*For $\alpha = 1$ there is a transition phase where the giant component crumbles.

Parameters of HRG

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".



$$\lim_{\alpha \uparrow 1}$$

*For $\alpha = 1$ there is a transition phase where the giant component crumbles.




Parameters of HRG


Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".



*For $\alpha = 1$ there is a transition phase where the giant component crumbles.




Hyperbolic Distance

$$R = 2 \log n + C$$


Hyperbolic Distance

$$R = 2 \log n + C$$




Hyperbolic Distance

$$R = 2 \log n + C$$



Hyperbolic Distance


$$R = 2 \log n + C$$



Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$




Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$




Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$




Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .




Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .



Hyperbolic Distance


$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, their distance equal R)?



Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r',\theta') | d(r,r',\theta - \theta') \leq x\}$$


Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x,y)$ such that x and y are connected (hence, there distance equal R)?

Distance in a hyperbolic disc:

$$\cosh(dist(x,y)) :=$$

$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$



Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .


Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, there distance equal R)?

Distance in a hyperbolic disc:

$$\cosh(dist(x, y)) :=$$

$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$

Set $dist(x, y) = R$ and solving for the angular distance yields:



Hyperbolic Distance

$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, there distance equal R)?


Distance in a hyperbolic disc:

$$\cosh(dist(x, y)) :=$$

$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$


Set $dist(x, y) = R$ and solving for the angular distance yields:

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$



Maximal angle between connected points


$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$



Maximal angle between connected points

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.



Maximal angle between connected points


$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$



Maximal angle between connected points

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.


Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:

$$\frac{1}{1-x} = 1 + \Theta(x)$$



Maximal angle between connected points

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$


2. Use identity for $x \in o(1)$:

$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$

$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$



Maximal angle between connected points

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:


$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$

$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$

4. Apply: $\sqrt{1+x} = 1 + \frac{x}{2} + \Theta(x^2) \iff |x| \leq 1$



Maximal angle between connected points

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:


$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$

$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$

4. Apply: $\sqrt{1+x} = 1 + \frac{x}{2} + \Theta(x^2) \iff |x| \leq 1$




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.


An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.




An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.


Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.

Proof. By our previous lemma:

$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.


Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.

Proof. By our previous lemma:

$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$


Then holds: If $w \sim u$ Then also $w \sim v$.

Proof. By our previous lemma:

$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.

We can express the difference by φ and get:

$$\theta(r(v), R) \leq \varphi(u, v) + \theta(r(u), R)$$



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.


Proof. By our previous lemma:

$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.

We can express the difference by φ and get:

$$\theta(r(v), R) \leq \varphi(u, v) + \theta(r(u), R)$$

Solving for φ : $\varphi(u, v) \leq \theta(r(v), R) - \theta(r(u), R)$



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.

Proof. By our previous lemma:


$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.

We can express the difference by φ and get:

$$\theta(r(v), R) \leq \varphi(u, v) + \theta(r(u), R)$$

Solving for φ : $\varphi(u, v) \leq \theta(r(v), R) - \theta(r(u), R)$

The lemma follows by applying our previous lemma for θ .



An application: Dominant angle

Lemma: Let $r(v) \leq r(u)$.

Furthermore,

$$\text{let } \varphi(u, v) \leq 2(e^{r(u)/2} - e^{r(v)/2} + \Theta(e^{-3/2r(u)})) - \Theta(e^{-3/2r(v)}).$$

Then holds: If $w \sim u$ Then also $w \sim v$.

Proof. By our previous lemma:


$\theta(r(v), R)$ and $\theta(r(u), R)$ is the maximal angle such that a point on the boundary is connected with v and u respectively.

We can express the difference by φ and get:

$$\theta(r(v), R) \leq \varphi(u, v) + \theta(r(u), R)$$

Solving for φ : $\varphi(u, v) \leq \theta(r(v), R) - \theta(r(u), R)$

The lemma follows by applying our previous lemma for θ . ■




Measure of the inner ball

Let $0 \leq x \leq R$.

Measure of the inner ball


Let $0 \leq x \leq R$.



Measure of the inner ball

Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?




Measure of the inner ball

Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$




Measure of the inner ball

Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .



Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$2\pi \int_0^x f(r)dr = \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr$$



Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \end{aligned}$$



Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\ &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\ &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\ &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}.
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}. \quad \blacksquare
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure of the inner ball


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}. \quad \blacksquare
 \end{aligned}$$




Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.

Measure space of vertices connected to a point

Let x be a point with radius r .

Measure space of vertices connected to a point


Let x be a point with radius r .



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?




Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .



Measure space of vertices connected to a point


Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .


Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\mu(B_r(R) \cap B_o(R)) = \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .


Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:


$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Measure space of vertices connected to a point

Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:


$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_o(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$

For the details check the paper! For refinements of the lemma check out the cheat sheet!



A variant of the previous lemma

Lemma:


$$\mu((B_r(R) \cap B_o(R)) - B_O(x)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

A variant of the previous lemma

Lemma:

$$\mu((B_r(R) \cap B_o(R)) - B_O(x)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

This is our previous lemma with the variation of removing the blue ball from the measure.




A variant of the previous lemma

Lemma:

$$\mu((B_r(R) \cap B_o(R)) - B_O(x)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

This is our previous lemma with the variation of removing the blue ball from the measure.




A variant of the previous lemma

Lemma:

$$\mu((B_r(R) \cap B_o(R)) - B_O(x)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

This is our previous lemma with the variation of removing the blue ball from the measure.




A variant of the previous lemma

Lemma:

$$\mu((B_r(R) \cap B_o(R)) - B_O(x)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

This is our previous lemma with the variation of removing the blue ball from the measure.



Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

Expected degree of a vertex


We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:

Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.


An application of a variant of our lemma:



Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:




Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:

Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_0(R))\}$
are exactly the points connected to v .
So we need to compute $\mathbb{E}[|S|] = \delta(v)$.




Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:

Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_o(R))\}$
are exactly the points connected to v .
So we need to compute $\mathbb{E}[|S|] = \delta(v)$.

For this we compute: $\mathbb{E}[|S|] = n\mu(B_r(R) \cap B_o(R))$.



Expected degree of a vertex


We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:

Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_0(R))\}$
 are exactly the points connected to v .
 So we need to compute $\mathbb{E}[|S|] = \delta(v)$.

For this we compute: $\mathbb{E}[|S|] = n\mu(B_r(R) \cap B_o(R))$.

Let $I(\beta) := \{v \in V(G) | 0 \leq r(v) < \beta R\}$ and
 $O(\beta) := \{v \in V(G) | \beta R \leq r(v) < R\}$.



Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:


Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_0(R))\}$
 are exactly the points connected to v .
 So we need to compute $\mathbb{E}[|S|] = \delta(v)$.

For this we compute: $\mathbb{E}[|S|] = n\mu(B_r(R) \cap B_o(R))$.

Let $I(\beta) := \{v \in V(G) | 0 \leq r(v) < \beta R\}$ and
 $O(\beta) := \{v \in V(G) | \beta R \leq r(v) < R\}$.

By the previous lemma we get

$$\delta(v) \in \mathcal{O}(ne^{-\frac{\beta R}{2}}) = \mathcal{O}(n^{1-\beta}) = c'n^{1-\beta}.$$



Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:


Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_0(R))\}$
 are exactly the points connected to v .
 So we need to compute $\mathbb{E}[|S|] = \delta(v)$.

For this we compute: $\mathbb{E}[|S|] = n\mu(B_r(R) \cap B_o(R))$.

Let $I(\beta) := \{v \in V(G) | 0 \leq r(v) < \beta R\}$ and
 $O(\beta) := \{v \in V(G) | \beta R \leq r(v) < R\}$.

By the previous lemma we get
 $\delta(v) \in \mathcal{O}(ne^{-\frac{\beta R}{2}}) = \mathcal{O}(n^{1-\beta}) = c'n^{1-\beta}$.

1. We computed the expectation of the degree for v !



Expected degree of a vertex

We would like to calculate the Expected degree of a vertex with radius r or greater and show that it is concentrated around its expectation.

An application of a variant of our lemma:

Notice: $S = \{u \in G(V) | u \in (B_{r_v}(R) \cap B_0(R))\}$
 are exactly the points connected to v .
 So we need to compute $\mathbb{E}[|S|] = \delta(v)$.


For this we compute: $\mathbb{E}[|S|] = n\mu(B_r(R) \cap B_o(R))$.

Let $I(\beta) := \{v \in V(G) | 0 \leq r(v) < \beta R\}$ and
 $O(\beta) := \{v \in V(G) | \beta R \leq r(v) < R\}$.

By the previous lemma we get

$$\delta(v) \in \mathcal{O}(ne^{-\frac{\beta R}{2}}) = \mathcal{O}(n^{1-\beta}) = c'n^{1-\beta}.$$

1. We computed the expectation of the degree for v !
2. Is the degree concentrated around its expectation?




Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Concentration around the expected degree


To show the concentration we use $I(\beta)$ and $O(\beta)$.



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.




Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.



Concentration around the expected degree


To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.


Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$

The bound for \mathcal{B} follows by union bound.



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.


Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$

The bound for \mathcal{B} follows by union bound. ■



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.


Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$

The bound for \mathcal{B} follows by union bound. ■

Changing the coordinates of a point increases or decreases the degree of at most $2ec'n^{1-\beta} + 1$ points in $O(\beta)$ unless \mathcal{B} holds.



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.


Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$

The bound for \mathcal{B} follows by union bound. ■

Changing the coordinates of a point increases or decreases the degree of at most $2ec'n^{1-\beta} + 1$ points in $O(\beta)$ unless \mathcal{B} holds.

By an Azuma-Hoeffding-type* large deviation inequality follows the concentration.



Concentration around the expected degree

To show the concentration we use $I(\beta)$ and $O(\beta)$.

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e\underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.


Proof. By a Chernoff bound we get:

$$\mathbb{P}[\delta_O(v) > 2ec'n^{1-\beta}] \leq 2^{-2ec'n^{1-\beta}}$$

The bound for \mathcal{B} follows by union bound. ■

Changing the coordinates of a point increases or decreases the degree of at most $2ec'n^{1-\beta} + 1$ points in $O(\beta)$ unless \mathcal{B} holds.

By an Azuma-Hoeffding-type* large deviation inequality follows the concentration.



$$^*\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)]\mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum_i c_i^2} + \mathbb{P}[\mathcal{B}]$$

Average degree of a vertex


Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.

Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.




Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.

Proof. By our previous lemma the degree for a radius r is given by:

$$\delta(r) = n\mu(B_r(R) \cap B_o(R)) = n \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$




Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.

Proof. By our previous lemma the degree for a radius r is given by:

$$\delta(r) = n\mu(B_r(R) \cap B_o(R)) = n \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$



Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.


Proof. By our previous lemma the degree for a radius r is given by:

$$\delta(r) = n\mu(B_r(R) \cap B_o(R)) = n \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

$$\text{Recall: } p(r) = \frac{\alpha \sinh(\alpha r)}{\cosh(\alpha R) - 1} = \alpha e^{\alpha(r-R)} \cdot (1 \pm \mathcal{O}(e^{-2\alpha r} + e^{-\alpha R})).$$

By definition of expectation we have:

$$\bar{\delta} = \int_0^R p(r)\delta(r)dr$$



Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.

Proof. By our previous lemma the degree for a radius r is given by:

$$\delta(r) = n\mu(B_r(R) \cap B_o(R)) = n \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$


$$\text{Recall: } p(r) = \frac{\alpha \sinh(\alpha r)}{\cosh(\alpha R) - 1} = \alpha e^{\alpha(r-R)} \cdot (1 \pm \mathcal{O}(e^{-2\alpha r} + e^{-\alpha R})).$$

By definition of expectation we have:

$$\bar{\delta} = \int_0^R p(r)\delta(r)dr$$

Plugging into the definition of expectation and integrating yields:

$(1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$ without the error term and $o(1)$ for the error term.



Average degree of a vertex

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$ for $G_{\alpha,C}(n)$.

Then, the average degree is $\bar{\delta} = (1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$.

Proof. By our previous lemma the degree for a radius r is given by:

$$\delta(r) = n\mu(B_r(R) \cap B_o(R)) = n \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$


$$\text{Recall: } p(r) = \frac{\alpha \sinh(\alpha r)}{\cosh(\alpha R) - 1} = \alpha e^{\alpha(r-R)} \cdot (1 \pm \mathcal{O}(e^{-2\alpha r} + e^{-\alpha R})).$$

By definition of expectation we have:

$$\bar{\delta} = \int_0^R p(r)\delta(r)dr$$

Plugging into the definition of expectation and integrating yields:

$(1 + o(1)) \frac{2\alpha^2 e^{-C/2}}{\pi(\alpha - 1/2)^2}$ without the error term and $o(1)$ for the error term. ■



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.


Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.




Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.




Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.




Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.



Maximum degree of a node of a HRG


Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.


Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$

This means there is no vertex with radius $r_0 - \phi(n)$ whp.



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.


Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$

This means there is no vertex with radius $r_0 - \phi(n)$ whp.

On the other hand:

$\mathbb{P}[X_{\leq r_0 + \phi(n)}] \leq (1 - \mu(B_0(r_0 + \phi(n))))^n \leq e^{-e^{\alpha(\phi(n)-C)/2}} = o(1)$



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.


We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$

This means there is no vertex with radius $r_0 - \phi(n)$ whp.

On the other hand:

$\mathbb{P}[X_{\leq r_0 + \phi(n)}] \leq (1 - \mu(B_0(r_0 + \phi(n))))^n \leq e^{-e^{\alpha(\phi(n)-C)/2}} = o(1)$

Hence, there are vertices within the radius $r_0 + \phi(n)$ whp.



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$


This means there is no vertex with radius $r_0 - \phi(n)$ whp.

On the other hand:

$\mathbb{P}[X_{\leq r_0 + \phi(n)}] \leq (1 - \mu(B_0(r_0 + \phi(n))))^n \leq e^{-e^{\alpha(\phi(n)-C)/2}} = o(1)$

Hence, there are vertices within the radius $r_0 + \phi(n)$ whp.

Next: We apply Markov's Inequality to show that the number of vertices is very small.



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$

This means there is no vertex with radius $r_0 - \phi(n)$ whp.


On the other hand:

$\mathbb{P}[X_{\leq r_0 + \phi(n)}] \leq (1 - \mu(B_0(r_0 + \phi(n))))^n \leq e^{-e^{\alpha(\phi(n)-C)/2}} = o(1)$

Hence, there are vertices within the radius $r_0 + \phi(n)$ whp.

Next: We apply Markov's Inequality to show that the number of vertices is very small.

$$\mu(B_0(r_0 + \phi(n))) \leq \frac{2e^{\alpha(\phi(n)-C)}}{n} \rightarrow \mathbb{E}[X_{\leq r_0 + \phi(n)}] \leq 2e^{\alpha(\phi(n)-C)}$$



Maximum degree of a node of a HRG

Theorem: Let $\alpha > 1/2$ and $C \in \mathbb{R}$.

Then the maximum vertex degree of $G_{\alpha,C}(n)$ is with high probability $\Delta = n^{\frac{1}{2\alpha} + o(1)}$.

Proof. Set $r_0 := (2 - \frac{1}{\alpha}) \log n$.

Consider $r \in [r_0 - \phi(n), r_0 + \phi(n)]$ where $\phi(n) = o(\log \log n)$.

We get $\mathbb{E}[X_{\leq r_0 - \phi(n)}] = n\mu(B_0(r_0 - \phi(n))) \leq 2e^{-\alpha(\phi(n)+C)} = o(1)$

This means there is no vertex with radius $r_0 - \phi(n)$ whp.

On the other hand:


$$\mathbb{P}[X_{\leq r_0 + \phi(n)}] \leq (1 - \mu(B_0(r_0 + \phi(n))))^n \leq e^{-e^{\alpha(\phi(n)-C)/2}} = o(1)$$

Hence, there are vertices within the radius $r_0 + \phi(n)$ whp.

Next: We apply Markov's Inequality to show that the number of vertices is very small.


$$\mu(B_0(r_0 + \phi(n))) \leq \frac{2e^{\alpha(\phi(n)-C)}}{n} \rightarrow \mathbb{E}[X_{\leq r_0 + \phi(n)}] \leq 2e^{\alpha(\phi(n)-C)}$$

$$\text{Hence, by Markov and } \phi(n) = o(\log \log n): \mathbb{P}[X_{\leq r_0 + \phi(n)} > \log n] \leq \frac{2e^{\alpha(\phi(n)-C)}}{\log n} = o(1).$$



Bounding the maximal degree

What is left to show is an lower and upper bound.




Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$




Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$



Bounding the maximal degree


What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches our claim.



Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by


$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches our claim.

The lower bound we compute in similar fashion by

$$\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0+\phi(n)}{2}}.$$



Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$


$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches our claim.

The lower bound we compute in similar fashion by

$$\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0+\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) \geq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}}.$$



Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$


By Chernoff and union bound this matches our claim.

The lower bound we compute in similar fashion by

$$\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0+\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) \geq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches again our claim.



Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches our claim.


The lower bound we compute in similar fashion by

$$\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0+\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) \geq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches again our claim.

$$(1 + o(1)) \frac{4e\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}} \leq \Delta \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$



Bounding the maximal degree

What is left to show is an lower and upper bound.

The upper boud we compute by

$$\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0-\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0-\phi(n)}(R)) \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches our claim.


The lower bound we compute in similar fashion by

$$\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) = (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} e^{-\frac{r_0+\phi(n)}{2}}.$$

$$\text{Hence, } n\mu(B_0(R) \cap B_{r_0+\phi(n)}(R)) \geq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}}.$$

By Chernoff and union bound this matches again our claim.

$$(1 + o(1)) \frac{4e\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{-\frac{\phi(n)}{2}} \leq \Delta \leq (1 + o(1)) \frac{2\alpha}{\pi(\alpha-1/2)} n^{\frac{1}{2\alpha}} e^{\frac{\phi(n)}{2}}. \blacksquare$$




Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.


Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.




Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.




Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.



Cluster coefficient of HRG's


Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.



*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.




*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.

Proof. Let $Y := \sum_{v \in O(\beta)} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v) \cap O(\beta)\}|}{\binom{\delta(v)}{2}}$.




*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.

Proof. Let $Y := \sum_{v \in O(\beta)} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v) \cap O(\beta)\}|}{\binom{\delta(v)}{2}}$.

Let $\mathbb{E}[Y_r | \mathcal{E}]$ be the expected value of Y for a vertex v with radius r conditioned on the event \mathcal{E} that the vertex has at least degree 2.



*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's


Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.

Proof. Let $Y := \sum_{v \in O(\beta)} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v) \cap O(\beta)\}|}{\binom{\delta(v)}{2}}$.

Let $\mathbb{E}[Y_r | \mathcal{E}]$ be the expected value of Y for a vertex v with radius r conditioned on the event \mathcal{E} that the vertex has at least degree 2.

Observation:

$$\mathbb{E}[Y_r | \mathcal{E}] = \mathbb{P}[u_1 \sim u_2 | v \sim u_1, v \sim u_2, r(v) = r]$$



*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.

Proof. Let $Y := \sum_{v \in O(\beta)} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v) \cap O(\beta)\}|}{\binom{\delta(v)}{2}}$.


Let $\mathbb{E}[Y_r | \mathcal{E}]$ be the expected value of Y for a vertex v with radius r conditioned on the event \mathcal{E} that the vertex has at least degree 2.

Observation:

$$\mathbb{E}[Y_r | \mathcal{E}] = \mathbb{P}[u_1 \sim u_2 | v \sim u_1, v \sim u_2, r(v) = r]$$

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$



*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.

Cluster coefficient of HRG's

Theorem: Let $\alpha > 1/2$, $C \in \mathbb{R}$ and $\bar{c}^* = (G_{\alpha,C}(n))$.
 Then $\mathbb{E}[\bar{c}] = \Theta(1)$ with high probability.

Proof. Let $Y := \sum_{v \in O(\beta)} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v) \cap O(\beta)\}|}{\binom{\delta(v)}{2}}$.

Let $\mathbb{E}[Y_r | \mathcal{E}]$ be the expected value of Y for a vertex v with radius r conditioned on the event \mathcal{E} that the vertex has at least degree 2.

Observation:


$$\mathbb{E}[Y_r | \mathcal{E}] = \mathbb{P}[u_1 \sim u_2 | v \sim u_1, v \sim u_2, r(v) = r]$$

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$

*Let $X := \sum_{v \in V} \frac{|\{\{u_1, u_2\} \in E | u_1, u_2 \in \Gamma(v)\}|}{\binom{\delta(v)}{2}}$. Then $\bar{c} = \frac{X}{n}$.




Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R) \cap B_r(R))}$



Lower bound for the cluster coefficient


Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R)) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$

This gives us:

$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R)) ^ 2} d\varphi dy$$



Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.


The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$

This gives us:

$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R)) ^ 2} d\varphi dy$$

Recall: $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$ and if
 $\varphi(x, y) \leq 2(e^{r(x)/2} - e^{r(y)/2} + \Theta(e^{-3/2r(x)}) - \Theta(e^{-3/2r(y)}))$
 then Then holds: If $w \sim x$ Then also $w \sim y$.



Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.


The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$

This gives us:

$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R)) ^ 2} d\varphi dy$$

Recall: $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$ and if
 $\varphi(x, y) \leq 2(e^{r(x)/2} - e^{r(y)/2} + \Theta(e^{-3/2r(x)}) - \Theta(e^{-3/2r(y)})$
 then Then holds: If $w \sim x$ Then also $w \sim y$.



Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$


This gives us:

$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R))^2} d\varphi dy$$

Recall: $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$ and if
 $\varphi(x, y) \leq 2(e^{r(x)/2} - e^{r(y)/2} + \Theta(e^{-3/2r(x)}) - \Theta(e^{-3/2r(y)}))$
 then Then holds: If $w \sim x$ Then also $w \sim y$.

We set $\xi := 2(e^{r/2} - e^{-y/2})$ and we get for our integral:

$$\frac{1}{(\mu(B_0(R)) \cap B_r(R))^2} \int_r^R \int_0^\xi f(y) \cdot \mu(B_0(R) \cap B_y(R) \setminus B_0(\beta R)) d\varphi dy$$



Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R) \cap B_r(R))}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R))}$

This gives us:


$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R))^2} d\varphi dy$$

Recall: $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$ and if
 $\varphi(x, y) \leq 2(e^{r(x)/2} - e^{r(y)/2} + \Theta(e^{-3/2r(x)}) - \Theta(e^{-3/2r(y)}))$
 then Then holds: If $w \sim x$ Then also $w \sim y$.

We set $\xi := 2(e^{r/2} - e^{-y/2})$ and we get for our integral:

$$\frac{1}{(\mu(B_0(R)) \cap B_r(R))^2} \int_r^R \int_0^\xi f(y) \cdot \mu(B_0(R) \cap B_y(R) \setminus B_0(\beta R)) d\varphi dy$$

Lower bounding the integral and lower bounding $\mathbb{P}[\mathcal{E}]$ gives us with "some" calculations:



Lower bound for the cluster coefficient

Fixing a point u_1 at $(y, \varphi) \in B_0(R) \cap B_r(R) \setminus B_0(\beta R)$.

The Probability is clearly $\frac{f(y)}{\mu(B_0(R)) \cap B_r(R)}$

Hence, the contribution of u_2 to $\mathbb{E}[Y_r | \mathcal{E}]$, given the coordinates of u_2 is $\frac{\mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{\mu(B_0(R)) \cap B_r(R)}$

This gives us:

$$\mathbb{E}[Y_r | \mathcal{E}] \geq \frac{1}{2} \int_{\beta R}^R \int_{-\theta(r,y)}^{\theta(r,y)} \frac{f(y) \mu(B_r(R) \cap B_y(R) \cap B_0(R) \setminus B_0(\beta R))}{(\mu(B_0(R)) \cap B_r(R))^2} d\varphi dy$$


Recall: $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$ and if
 $\varphi(x, y) \leq 2(e^{r(x)/2} - e^{r(y)/2} + \Theta(e^{-3/2r(x)}) - \Theta(e^{-3/2r(y)}))$
 then Then holds: If $w \sim x$ Then also $w \sim y$.

We set $\xi := 2(e^{r/2} - e^{-y/2})$ and we get for our integral:

$$\frac{1}{(\mu(B_0(R)) \cap B_r(R))^2} \int_r^R \int_0^\xi f(y) \cdot \mu(B_0(R) \cap B_y(R) \setminus B_0(\beta R)) d\varphi dy$$

Lower bounding the integral and lower bounding $\mathbb{P}[\mathcal{E}]$ gives us with "some" calculations:

$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\bar{c}_r | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$



Concentration bound for the cluster coefficient

$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha - 1/2)}}}{600\pi^3(\alpha - 1/2)(\alpha + 1)(\alpha + 1/2)} \in \Theta(n).$$

Concentration bound for the cluster coefficient


$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$

Recall:

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum_i c_i^2} + \mathbb{P}[\mathcal{B}].$$



Concentration bound for the cluster coefficient

$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$


Recall:

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum_i c_i^2} + \mathbb{P}[\mathcal{B}].$$

Set $f := Y$, $t := n^{6/7}$ and \mathcal{B} as stated in the lemma.



Concentration bound for the cluster coefficient

$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$

Recall:


Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum_i c_i^2} + \mathbb{P}[\mathcal{B}].$$

Set $f := Y$, $t := n^{6/7}$ and \mathcal{B} as stated in the lemma.

Each coordinate change can influence f by at most $c_i := cn^{1-\beta} + 1$ as long as $\overline{\mathcal{B}}$ holds.



Concentration bound for the cluster coefficient

$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$

Recall:

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.


Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum_i c_i^2} + \mathbb{P}[\mathcal{B}].$$

Set $f := Y$, $t := n^{6/7}$ and \mathcal{B} as stated in the lemma.

Each coordinate change can influence f by at most $c_i := cn^{1-\beta} + 1$ as long as $\overline{\mathcal{B}}$ holds.

Hence, $\mathbb{P}[X \leq \mathbb{E}[Y] - n^{6/7} - \mathbb{P}[\mathcal{B}]] = o(1)$.



Concentration bound for the cluster coefficient


$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$

Recall:

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum c_i^2} + \mathbb{P}[\mathcal{B}].$$



Set $f := Y$, $t := n^{6/7}$ and \mathcal{B} as stated in the lemma.

Each coordinate change can influence f by at most $c_i := cn^{1-\beta} + 1$ as long as $\overline{\mathcal{B}}$ holds.

Hence, $\mathbb{P}[X \leq \mathbb{E}[Y] - n^{6/7} - \mathbb{P}[\mathcal{B}]] = o(1)$.

This concludes that the clustering coefficient is with high probability at least $\mathbb{E}[Y]/n = \Theta(1)$.

Concentration bound for the cluster coefficient


$$\mathbb{E}[Y] = n \int_{\beta R}^R f(y) \cdot \mathbb{P}[\mathcal{E}] \cdot \mathbb{E}[\overline{c_r} | \mathcal{E}] dr \geq \frac{n \cdot e^{-C} \alpha^2 e^{\frac{2\alpha e^{-C/2}}{\pi(\alpha-1/2)}}}{600\pi^3(\alpha-1/2)(\alpha+1)(\alpha+1/2)} \in \Theta(n).$$

Recall:

Let $\mathcal{B} := \{\exists v \in O(\beta) : \delta(v) > 2e \underbrace{c'n^{1-\beta}}_{\mathbb{E}[\delta(v)]}\}$.

Lemma: $\mathbb{P}[\mathcal{B}] = e^{-\Omega(n^{1-\beta})}$.

$$\mathbb{P}[f \geq \mathbb{E}[f] + t + (M - m)] \mathbb{P}[(\mathcal{B})] \leq e^{2t^2/\sum c_i^2} + \mathbb{P}[\mathcal{B}].$$



Set $f := Y$, $t := n^{6/7}$ and \mathcal{B} as stated in the lemma.

Each coordinate change can influence f by at most $c_i := cn^{1-\beta} + 1$ as long as $\overline{\mathcal{B}}$ holds.

Hence, $\mathbb{P}[X \leq \mathbb{E}[Y] - n^{6/7} - \mathbb{P}[\mathcal{B}]] = o(1)$.

This concludes that the clustering coefficient is with high probability at least $\mathbb{E}[Y]/n = \Theta(1)$. ■