




Hasso
Plattner
Institut

Hyperbolic Random Graphs

Degree Sequence and Clustering

Algorithm Engineering Group


Janosch Ruff



Network Features - Distributions


The most commonly considered network features are distributions over parts of the network.

- Degrees
- Assortativity
- Distances
- Centrality
- Clustering




[The Structure and Function of Complex Networks, M. E. J. Newman, Computer Physics Communications 2003]

Network Features – Degree Distribution




$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$



Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$

The arithmetic mean of the degree distribution is the **average degree** of the network.


Network Features – Degree Distribution





$$\deg(v) = |N(v)|$$

The arithmetic mean of the degree distribution is the **average degree** of the network.

- Homogeneous




Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$



The arithmetic mean of the degree distribution is the **average degree** of the network.

- Homogeneous



Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous




Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous





Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.


Network Features – Degree Distribution



$$\deg(v) = |N(v)|$$


The arithmetic mean of the degree distribution is the **average degree** of the network.

■ Homogeneous



Typically described using a **binomial distribution**. Additionally characterized by the **standard deviation**.

■ Heterogeneous



Typically described using a **power-law distribution**. Additionally characterized by the **power-law exponent**.

For directed networks **in- and out-degree distributions** are often considered separately.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.


$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$


- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$


- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)\|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.


$$c(G) = \frac{3 |\Delta|}{|\bullet\bullet|}$$

Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)\|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.

$$c(G) = \frac{3 |\Delta|}{|\bullet\bullet|}$$




Network Features – Clustering

Intuitively, **clustering** measures how often nodes are part of dense groups (clusters).

The **local clustering coefficient** of a node denotes the probability that two randomly chosen neighbors are adjacent.

$$c(v) = \frac{|\{v_1, v_2\} \in E \mid v_1, v_2 \in N(v)\|}{\deg(v)(\deg(v) - 1)/2}$$



$$c(2) = \frac{1}{3} \quad c(1) = \frac{1}{1}$$

- For directed graphs the normalization factor becomes $\deg(v)(\deg(v) - 1)$.
- The whole graph is characterized by the distribution of the local clustering coefficients.
 - Its arithmetic mean is called the **average local clustering coefficient**.

The **global clustering coefficient** of a graph measures how many paths of length 2 induce a triangle.

$$c(G) = \frac{3 |\Delta|}{|\bullet|}$$



Compared to the average local clustering coefficient, the global one basically computes the ratio of the means rather than the mean of the ratios. It does not weight the contributions of low-degree nodes as much.

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2

- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$




Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$



Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs


Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$

$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$



$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$



$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient
- ... a logarithmic diameter



[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Network Models – Hyperbolic Random Graphs

Hyperbolic space expands exponentially, while Euclidean space expands polynomially.

- Area of a Euclidean circle with radius r : πr^2
- Area of a hyperbolic circle with radius r : $2\pi(\cosh(r) - 1)$


$$= 1/2(e^r + e^{-r}) \approx 1/2e^r$$

A **hyperbolic random graph** is obtained by distributing n nodes uniformly at random in a hyperbolic disk of radius R and connecting any two at distance $\leq R$.

$$R \approx 2 \log(n)$$

Hyperbolic random graphs have...

- ... a power-law degree distribution
- ... a constant average degree
- ... a non-vanishing clustering coefficient
- ... a logarithmic diameter
- ... negative assortativity




[Hyperbolic Geometry of Complex Networks. Krioukov et al. Phys. Rev. E (2010)]

Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$


Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)}$$



Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$



Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".




*For $\alpha = 1$ there is a transition phase where the giant component crumbles.

Density function to sample points:


$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".



$$\lim_{\alpha \uparrow 1}$$


*For $\alpha = 1$ there is a transition phase where the giant component crumbles.




Density function to sample points:

$$f(r) = \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R) - 1)}$$

$\alpha \in (0.5, 1^*)$ "power law degree".




*For $\alpha = 1$ there is a transition phase where the giant component crumbles.




$$R = 2 \log n + C$$


$$R = 2 \log n + C$$



$$R = 2 \log n + C$$




$$R = 2 \log n + C$$



$$R = 2 \log n + C$$


$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$



$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$


$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$



$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$




$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .




$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .




$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, their distance equal R)?



$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$


Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, their distance equal R)?

Distance in a hyperbolic disc:

$$\cosh(dist(x, y)) :=$$

$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$



$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .


Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, there distance equal R)?

Distance in a hyperbolic disc:

$$\cosh(dist(x, y)) :=$$

$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$

Set $dist(x, y) = R$ and solving for the angular distance yields:



$$R = 2 \log n + C$$

$$E(G) = \{E_{i,j} | dist(i,j) \leq R\}$$

$$B_{r,\theta}(x) = \{(r', \theta') | d(r, r', \theta - \theta') \leq x\}$$

Consider two nodes with radial coordinates x, y .

Question: What is the maximal angular distance denoted by $\theta(x, y)$ such that x and y are connected (hence, there distance equal R)?


Distance in a hyperbolic disc:

$$\cosh(dist(x, y)) :=$$


$$\cosh(r_x)\cosh(r_y) - \sinh(r_x)\sinh(r_y)\cos(\Delta\varphi_{x,y})$$

Set $dist(x, y) = R$ and solving for the angular distance yields:

$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$




$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$



$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.




$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$



$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.


Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:

$$\frac{1}{1-x} = 1 + \Theta(x)$$



$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$


2. Use identity for $x \in o(1)$:

$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$

$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$



$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x) - \cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:


$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$

$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$

4. Apply: $\sqrt{1+x} = 1 + \frac{x}{2} + \Theta(x^2) \iff |x| \leq 1$



$$\theta(x, y) = \arccos\left(\frac{\cosh(y)\cosh(x)-\cosh(R)}{\sinh(y)\sinh(x)}\right)$$

Lemma: Let $0 \leq x \leq R$ and $y \geq R - x$.
 Then $\theta(x, y) = 2e^{\frac{R-x-y}{2}}(1 + \Theta(e^{R-x-y}))$.

Proof ingredients:

1. Use trigonometric identity:

$$\cosh(x \pm y) = \cosh(x)\cosh(y) \pm \sinh(x)\sinh(y)$$

2. Use identity for $x \in o(1)$:


$$\frac{1}{1-x} = 1 + \Theta(x)$$

3. Get rid of cosine by using Taylor expansions to derive lower and upper bound:

$$\cos(\theta) \geq 1 - \frac{\theta^2}{2}$$


$$\cos(\theta) \leq 1 - \frac{\theta^2}{2} + \frac{\theta^4}{4!}$$

4. Apply: $\sqrt{1+x} = 1 + \frac{x}{2} + \Theta(x^2) \iff |x| \leq 1$




Let $0 \leq x \leq R$.

Let $0 \leq x \leq R$.



Let $0 \leq x \leq R$.


Question: What is the measure of the grey area?



Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$




Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .




Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$2\pi \int_0^x f(r)dr = \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr$$




Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \end{aligned}$$



Let $0 \leq x \leq R$.


Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \end{aligned}$$

Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.




Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned} 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\ &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\ &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1}
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}.
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .

$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}. \quad \blacksquare
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.


Let $0 \leq x \leq R$.

Question: What is the measure of the grey area?

Lemma: $\mu(B_0(x)) = e^{-\alpha(R-x)}(1 + o(1))$

Proof. Just compute the probability measure for the integral from 0 to x .


$$\begin{aligned}
 2\pi \int_0^x f(r)dr &= \int_0^x \frac{\alpha \sinh(\alpha r)}{2\pi(\cosh(\alpha R)-1)} dr \\
 &= \frac{1}{\cosh(\alpha R)-1} \int_0^x \alpha \sinh(\alpha r) dr \\
 &= \left(\frac{1}{\cosh(\alpha R)-1} \right) (\cosh(\alpha r) - 1) \\
 &= \frac{\cosh(\alpha r) - 1}{\cosh(\alpha R) - 1} \\
 &= (1 + o(1))e^{-\alpha(R-x)}. \quad \blacksquare
 \end{aligned}$$



Remember: $\frac{\Delta}{\Delta x} \cosh(x) = \sinh(x)$.


Let x be a point with radius r .

Let x be a point with radius r .



Let x be a point with radius r .


Question: What is the measure of the orange area?



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .




Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .


Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\mu(B_r(R) \cap B_o(R)) = \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .


Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:


$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:


1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_0(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$



Let x be a point with radius r .

Question: What is the measure of the orange area?

Notice: The orange dashed area is the probability measure that a point sampled by randomly drawn point is connected with x .

Lemma:

$$\mu(B_r(R) \cap B_o(R)) = \frac{2\alpha e^{-r/2}}{\pi(\alpha - \frac{1}{2})} (1 \pm \mathcal{O}(e^{-(\alpha - \frac{1}{2})r} + e^{-r}))$$

Proof ingredients:

1. Compute the integral:

$$\begin{aligned} \mu(B_r(R) \cap B_o(R)) &= \int_0^R \int_{-\theta(r,y)}^{\theta(r,y)} f(y) d\theta dy \\ &= 2 \int_0^R \int_0^{\theta(r,y)} f(y) d\theta dy \end{aligned}$$

2. Just do it with the lemma for $\theta(x, y)$!

Notice:

$$\mu(B_r(R) \cap B_o(R)) = \mu(B_o(R-r)) + 2 \int_{R-r}^R \theta(r, y) f(y) dy$$

For the details check the paper! For refinements of the lemma check out the cheat sheet!

