
failure of early neurological improvement
(a) out-of-fold ROC by learner
sensitivity
1 − specificity
logistic 0.76
elastic net 0.78
random forest 0.72
grad. boosting 0.74
SVM 0.66
(b) incremental discrimination
(repeated CV, screens within folds)
AUC
adjustment set
+ screen-selected
+ selected domains
+ inflammatory
+ all labs
+ screen-selected
(c) gradient-boosting importance
Glucose
mTICI 3
INR
Onset-recan.
Creatinine
GPR
Haemoglobin
NLR
ΔAUC (permutation)
(d) screen-selected score:
ρ = 0.51, p = 0.001
permutations
Spearman ρ
(e) progressive adjustment
prespecified
+sex
+comorbidity
+ICA/tandem
+procedure
+renal
+lipids
+Hb, INR
+inflammatory
all blocks
OR per mmol/l
(f) dose-response by glucose tertile
(bar = mean; survivors)
NIHSS improvement to discharge
ENI (≥ 4)
T1
n = 29
T2
n = 21
T3
n = 19
(g) unsupervised anomaly score:
AUC 0.58
anomaly score (rank mean)
no
yes
(h) laboratory-panel PCA
PC2 (13%)
PC1 (34%)
admission glucose, mg/dl
(i) decision curve
adjustment set +
screen-selected factors
treat all
treat none
net benefit
threshold probability