
(a) adjusted per-factor association with outcome
(filled: q < 0.05 across all 49 factors)
Glucose
GPR
Diabetes
NLR
SII
SIRI
PLR
MLR
WBC
Neutrophils
Lymphocytes
Monocytes
CRP
RDW
MPV
Haemoglobin
Haematocrit
RBC
INR
APTT
Sodium
Potassium
Urea
Creatinine
Total chol.
HDL
LDL
Triglycerides
mTICI 3
Onset-recan.
First pass
IV tPA
Tandem
ICA involvement
Age
NIHSS
Female
Hypertension
AF
poor 90-day outcome
fENI
0.1
0.3
1
2
5
10
30
adjusted odds ratio (per SD or per category), log scale
(b) progressive adjustment
(filled, complete case per model; open, common sample)
prespecified set
+ sex
+ comorbidity (HTN, AF, diabetes)
+ ICA involvement, tandem
+ procedure (first pass, IV thrombolysis)
+ renal / electrolyte (Na, urea, creatinine)
+ lipids (LDL, HDL, triglycerides)
+ haemoglobin, INR
+ inflammatory (NLR, CRP, WBC)
+ all blocks jointly
1
1.5
2
3
5
odds ratio for fENI per mmol/l of admission glucose, log scale