

Notes

Maksim Levental

December 18, 2019

Contents

1	Smooth Manifolds	2		
1.1	Directional Derivative	2	1.11.5	Tangent Space at the identity of a Lie Group
1.2	Derivations	3	1.11.6	Left-Invariant Vector Fields on a Lie Group
1.3	Vector Fields	3	1.11.7	The Lie Algebra of a Lie Group
1.4	Dual Space	3	1.12	Differential Forms
1.5	Differential Forms on \mathbb{R}^n	4	1.12.1	Differential 1-Forms
1.5.1	Differential of a Function	4	1.12.2	Local Expression for a Differential 1-Form
1.5.2	Differential k -forms	5	1.12.3	The Cotangent Bundle
1.5.3	Differential Forms as Multilinear Functions on Vector Fields	6	1.12.4	Pullback of 1-Forms
1.5.4	Exterior Derivative	6	1.12.5	Differential k -forms
1.5.5	Closed and Exact Forms	6	1.12.6	The Bundle Point of View
1.5.6	Applications to Vector Calculus	6	1.12.7	Pullback of k -forms
1.5.7	Convention on Subscripts and Superscripts	7	1.12.8	The Wedge Product
1.6	Manifolds	8	1.13	The Exterior Derivative
1.6.1	Compatible Charts	8	1.13.1	Exterior Derivative on a Coordinate Chart
1.6.2	Smooth Manifolds	8	1.13.2	Existence of an Exterior Derivative on a Manifold
1.6.3	Examples of Smooth Manifolds	9	1.13.3	Exterior Differentiation Under a Pullback
1.6.4	Smooths Maps on a Manifold	9	1.14	The Lie Derivative and Interior Multiplication
1.6.5	Partial Derivatives	10	1.14.1	The Lie Derivative of a Vector Field
1.7	Tangent Space	11	1.14.2	The Lie Derivative of a Differential Form
1.7.1	The Tangent Space at a Point	12	1.14.3	Interior Multiplication
1.7.2	The Differential of a Map	12	1.14.4	Properties of the Lie Derivative
1.7.3	Curves in a Manifold	14	1.15	Orientation
1.8	Tangent Bundle	15	1.15.1	Orientations of a Vector Space
1.8.1	Topology of the Tangent Bundle	15	1.15.2	Orientations and n -covectors
1.8.2	Manifold Structure on the Tangent Bundle	16	1.15.3	Orientations on a Manifold
1.8.3	Vector Bundles	16	1.15.4	Orienations and Differential Forms
1.8.4	Smooth Sections	16	1.15.5	Orientations and Atlases
1.8.5	Smooth Frames	16	1.16	Manifolds with Boundary
1.9	Bump Functions and Partitions of Unity	17	1.16.1	The Boundary of a Manifold with Boundary
1.9.1	C^∞ Bump Functions	17	1.16.2	Outward-Pointing Vector Fields
1.9.2	Partitions of Unity	17	1.16.3	Boundary Orientation
1.10	Vector Fields	18		
1.10.1	Integral Curves	18		
1.10.2	Local Flows	18		
1.10.3	The Lie Bracket	19		
1.10.4	Pushforward of Vector Fields	20		
1.11	Lie Groups and Lie Algebras	20		
1.11.1	Lie Group	21		
1.11.2	Matrix Exponential	21		
1.11.3	Trace of a Matrix	21		
1.11.4	The Differential of \det at the identity	22		

1 Smooth Manifolds

Contents

1.1	Directional Derivative	2	1.12.4	Pullback of 1-Forms	25
1.2	Derivations	3	1.12.5	Differential k -forms	25
1.3	Vector Fields	3	1.12.6	The Bundle Point of View	27
1.4	Dual Space	3	1.12.7	Pullback of k -forms	27
1.5	Differential Forms on \mathbb{R}^n	4	1.12.8	The Wedge Product	27
1.5.1	Differential of a Function	4	1.13	The Exterior Derivative	28
1.5.2	Differential k -forms	5	1.13.1	Exterior Derivative on a Coordinate Chart	28
1.5.3	Differential Forms as Multilinear Functions on Vector Fields	6	1.13.2	Existence of an Exterior Derivative on a Manifold	29
1.5.4	Exterior Derivative	6	1.13.3	Exterior Differentiation Under a Pullback	29
1.5.5	Closed and Exact Forms	6	1.14	The Lie Derivative and Interior Multiplication	29
1.5.6	Applications to Vector Calculus	6	1.14.1	The Lie Derivative of a Vector Field	29
1.5.7	Convention on Subscripts and Superscripts	7	1.14.2	The Lie Derivative of a Differential Form	30
1.6	Manifolds	8	1.14.3	Interior Multiplication	31
1.6.1	Compatible Charts	8	1.14.4	Properties of the Lie Derivative	31
1.6.2	Smooth Manifolds	8	1.15	Orientation	32
1.6.3	Examples of Smooth Manifolds	9	1.15.1	Orientations of a Vector Space	32
1.6.4	Smooths Maps on a Manifold	9	1.15.2	Orientations and n -covectors	32
1.6.5	Partial Derivatives	10	1.15.3	Orientations on a Manifold	33
1.7	Tangent Space	11	1.15.4	Orienations and Differential Forms	33
1.7.1	The Tangent Space at a Point	12	1.15.5	Orientations and Atlases	34
1.7.2	The Differential of a Map	12	1.16	Manifolds with Boundary	34
1.7.3	Curves in a Manifold	14	1.16.1	The Boundary of a Manifold with Boundary	35
1.8	Tangent Bundle	15	1.16.2	Outward-Pointing Vector Fields	35
1.8.1	Topology of the Tangent Bundle	15	1.16.3	Boundary Orientation	35
1.8.2	Manifold Structure on the Tangent Bundle	16			
1.8.3	Vector Bundles	16			
1.8.4	Smooth Sections	16			
1.8.5	Smooth Frames	16			
1.9	Bump Functions and Partitions of Unity	17			
1.9.1	C^∞ Bump Functions	17			
1.9.2	Partitions of Unity	17			
1.10	Vector Fields	18			
1.10.1	Integral Curves	18			
1.10.2	Local Flows	18			
1.10.3	The Lie Bracket	19			
1.10.4	Pushforward of Vector Fields	20			
1.11	Lie Groups and Lie Algebras	20			
1.11.1	Lie Group	21			
1.11.2	Matrix Exponential	21			
1.11.3	Trace of a Matrix	21			
1.11.4	The Differential of det at the identity	22			
1.11.5	Tangent Space at the identity of a Lie Group	23			
1.11.6	Left-Invariant Vector Fields on a Lie Group	23			
1.11.7	The Lie Algebra of a Lie Group	24			
1.12	Differential Forms	24			
1.12.1	Differential 1-Forms	24			
1.12.2	Local Expression for a Differential 1-Form	25			
1.12.3	The Cotangent Bundle	25			

1.1 Directional Derivative

Elements of the *tangent space* $T_p(\mathbb{R}^n)$ anchored at a point $p = (p^1, \dots, p^n) \in \mathbb{R}^n$ can be visualized as arrows emanating from p . These arrows are called *tangent vectors* and represented by column vectors:

$$\mathbf{v} = \begin{bmatrix} v^1 \\ \vdots \\ v^n \end{bmatrix} \quad (1)$$

The line through a point p with direction \mathbf{v} has parameterization

$$c(t) = (p^1 + tv^1, \dots, p^n + tv^n) \quad (2)$$

If $f \in C^\infty$ in a neighborhood of p and \mathbf{v} is a tangent vector at p , the *directional derivative* of f in the direction of \mathbf{v} at p is defined

$$D_{\mathbf{v}}f = \lim_{t \rightarrow 0} \frac{f(c(t)) - f(p)}{t} = \left. \frac{d}{dt} f(c(t)) \right|_{t=0} \quad (3)$$

By the chain rule

$$D_v f = \sum_{i=1}^n \frac{\partial f}{\partial x^i} \bigg|_p \frac{dc^i}{dt} \bigg|_{t=0} \quad (4)$$

$$= \sum_{i=1}^n \frac{dc^i}{dt} \bigg|_{t=0} \frac{\partial f}{\partial x^i} \bigg|_p \quad (5)$$

$$= \sum_{i=1}^n v^i \frac{\partial f}{\partial x^i} \bigg|_p \quad (6)$$

(7)

The directional derivative operator at p is defined

$$D_v = \sum_{i=1}^n v^i \frac{\partial}{\partial x^i} \bigg|_p \quad (8)$$

The association $v \mapsto D_v$ offers a way to *isomorphically* identify tangent vectors with operators on functions. The following makes this rigorous.

1.2 Derivations

For each tangent vector v at a point $p \in \mathbb{R}^n$, the directional derivative at p gives a map of vector spaces

$$D_v: C_p^\infty \rightarrow \mathbb{R}$$

D_v is a linear map that satisfies the *Leibniz rule*

$$D_v(fg) = (D_v f)g(p) + f(p)(D_v g) \quad (9)$$

because the partial derivative satisfy the product rule. In general, any linear map $L: C_p^\infty \rightarrow \mathbb{R}$ that satisfies the Leibniz rule is called a *derivation* at p . Denote the set of all derivations at p by $\mathcal{D}_p(\mathbb{R}^n)$. **This set is also a real vector space.**

So far we know directional derivatives D_v at p are derivations at p . Thus, there is a map

$$\begin{aligned} \phi: T_p(\mathbb{R}^n) &\rightarrow \mathcal{D}_p(\mathbb{R}^n) \\ v &\mapsto D_v \end{aligned}$$

Theorem 1.1

The linear map ϕ is an isomorphism of vector spaces.

The implication is that we may identify tangent vectors at p with derivations at p (by way of directional derivatives against germs). Under this isomorphism $T_p(\mathbb{R}^n) \simeq \mathcal{D}_p(\mathbb{R}^n)$, the standard basis $\{e_1, \dots, e_n\}$ for $T_p(\mathbb{R}^n)$ maps to

$$\left\{ \frac{\partial}{\partial x^1} \bigg|_p, \dots, \frac{\partial}{\partial x^n} \bigg|_p \right\} \quad (10)$$

Therefore from now on we write a tangent vector as

$$v = \sum_{i=1}^n v^i \frac{\partial}{\partial x^i} \bigg|_p \quad (11)$$

The point being that, while not as geometrically intuitive as arrows, $\mathcal{D}_p(\mathbb{R}^n)$ generalizes to manifolds.

1.3 Vector Fields

A *vector field* X on an open $U \subset \mathbb{R}^n$ is function that assigns to $p \in U$ a tangent vector $X_p \in T_p(\mathbb{R}^n)$. Notice, carefully, that the vector field assigns at each point a vector in the tangent space anchored at that point. Using the tangent basis (eqn. (10))

$$X: p \mapsto \sum_i a^i(p) \frac{\partial}{\partial x^i} \bigg|_p \quad (12)$$

Note that both the coefficients **and** the partial derivatives are evaluated at p . Having said that, we often omit p in the specification of a vector field when it clear from context.

Example 1.1. On $\mathbb{R}^n - \{0\}$, let $p = (x, y)$. Then

$$\begin{aligned} X &= \frac{-y}{\sqrt{x^2 + y^2}} \frac{\partial}{\partial x} + \frac{x}{\sqrt{x^2 + y^2}} \frac{\partial}{\partial y} \\ &= \begin{bmatrix} \frac{-y}{\sqrt{x^2 + y^2}} \\ \frac{x}{\sqrt{x^2 + y^2}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{-y}{\sqrt{x^2 + y^2}} & \frac{x}{\sqrt{x^2 + y^2}} \end{bmatrix}^T \end{aligned}$$

See figure ??

In general we can identify vector fields with parameterized column vectors

$$X = \sum_i a^i(p) \frac{\partial}{\partial x^i} \bigg|_p \leftrightarrow \begin{bmatrix} a^1(p) \\ \vdots \\ a^n(p) \end{bmatrix} \quad (13)$$

1.4 Dual Space

The *dual space* V^\wedge of V is the set of all real-valued linear functions on V i.e. all $f: V \rightarrow \mathbb{R}$. Elements of V^\wedge are called *covectors*.

Assume V is finite dimensional and let $\{e_1, \dots, e_n\}$ be a basis V . Recall that $e_i := \partial_{x^i}$. Then $X = \sum a^i \partial_{x^i}$ for all $X \in T_p$. Let $\alpha^i: V \rightarrow \mathbb{R}$ be the linear function that picks out the i th coordinate of a **vector**, i.e. $\alpha^i(X) = a^i(p)$. Note that

$$\alpha^i(\partial_j) = \alpha^i(1 \cdot \partial_j) \quad (14)$$

$$= \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad (15)$$

$$= \delta_j^i \quad (16)$$

Note that position of indices is important – upper indices are for covectors.

Proposition 1.1

$\{\alpha^i\}$ form a basis for V^\wedge .

Proof. We first prove that $\{\alpha^i\}$ span V^\wedge . If $f \in V^\wedge$ and $X = \sum a^i \partial_{x_i} \in V$, then

$$f(X) = \sum a^i f(\partial_{x_i}) \quad (17)$$

$$= \sum \alpha^i(X) f(\partial_{x_i}) \quad (18)$$

$$= \sum f(\partial_{x_i}) \alpha^i(X) \quad (19)$$

which shows that any f can be expanded as a linear sum of α^i . To show linear independence, suppose $\sum c_i \alpha^i = 0$ with at least one c_i non-zero. Applying this to an arbitrary ∂_{x_i} gives

$$0 = \left(\sum_i c_i \alpha^i \right) (\partial_{x_i}) = \sum_i c_i \alpha^i(\partial_{x_i}) = \sum_i c_i \delta_j^i = c_j \quad (20)$$

which is a contradiction. Hence α^i are linearly independent. \square

This basis $\{\alpha^i\}$ for V^\wedge is said to be *dual* to the basis $\{\partial_{x_i}\}$ for V .

Example 1.2. Coordinate functions With respect to a basis $\{\partial_{x_i}\}$ for V , every $X \in V$ can be written uniquely as a linear combination $X = \sum a^i \partial_{x_i}$ with $a^i \in \mathbb{R}$. Let $\{\alpha^i\}$ be the dual basis (i.e. the basis for V^\wedge). Then

$$\begin{aligned} \alpha^i(X) &= \alpha^i \left(\sum_j a^j \partial_{x_j} \right) \\ &= \sum_j a^j \alpha^i(\partial_{x_j}) \\ &= \sum_j a^j \delta_j^i \\ &= a^i \end{aligned}$$

Thus, the dual basis $\{\alpha^i\}$ to $\{\partial_{x_i}\}$ is the set of coordinate functions. The sense here is that since tangent vectors are directional derivatives (i.e. operators on functions) and the dual space is a mapping from those operators to \mathbb{R} , then a mapping from operators to scalars means hitting an operator with a function (or vice-versa). **And** the coordinate functions are constant with respect to each other coordinate (and hence partials wrt them are naturally zero).

1.5 Differential Forms on \mathbb{R}^n

A differential k -form assigns a k -covector from the dual space at each point p . The wedge product (alternation of tensor product) of differential forms is defined pointwise (as the wedge product of multi-covectors). Differential forms exist on an open set (why?) there is a notion of differentiation (called exterior derivative). Exterior derivative is coordinate independent and intrinsic to a manifold; it is the abstraction of gradient, curl, divergence to arbitrary manifolds. Differential forms extend Grassmann's

exterior algebra (graded algebra of multi-covectors) from the tangent space at a point globally, i.e. to the entire manifold (how? bundles?).

1.5.1 Differential of a Function

Definition 1.1: Cotangent Space

The *cotangent space* to \mathbb{R}^n at p , denoted $T_p^*(\mathbb{R}^n)$, is defined to be the dual space $(T_p(\mathbb{R}^n))^\vee$ of the tangent space $T_p(\mathbb{R}^n)$.

Thus, an element of the cotangent space $T_p^*(\mathbb{R}^n)$ is a **covector of linear functional on tangent space**.

Definition 1.2: Differential 1-form

A *covector field* or a *differential 1-form* on an open subset U of \mathbb{R}^n is a function ω that assigns at each point p in U a covector $\omega_p \in T_p^*(\mathbb{R}^n)$

$$\begin{aligned} \omega: U &\rightarrow \bigcup_{p \in U} T_p^*(\mathbb{R}^n) \\ p &\mapsto \omega_p \in T_p^*(\mathbb{R}^n) \end{aligned} \quad (21)$$

We call a differential 1-form a *1-form* for short.

Definition 1.3: Differential

From any C^∞ function $f: U \rightarrow \mathbb{R}$, we can construct the 1-form df , called the *differential* of f , as follows: for $p \in U$ and $X_p \in T_p(U)$

$$(df)_p(X_p) := X_p f \quad (22)$$

In words the differential of f is the application of X_p to f or **the directional derivative of f in the direction of the tangent vector defined by the coefficients of X_p** .

Let x^1, \dots, x^n be the standard coordinates on \mathbb{R} , $\{(dx^1)_p, \dots, (dx^n)_p\}$ their differentials defined

$$(dx^i)_p(X_p) := (X_p)(x^i)$$

and

$$\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$$

be the standard basis for $T_p(\mathbb{R}^n)$.

Proposition 1.2

$\{(dx^1)_p, \dots, (dx^n)_p\}$ is the basis for $T_p^*(\mathbb{R}^n)$ dual to the coordinate basis for $T_p(\mathbb{R}^n)$.

Proof. By definition,

$$(\mathrm{d}x^i)_p \left(\frac{\partial}{\partial x^j} \Big|_p \right) = \frac{\partial x^i}{\partial x^j} \Big|_p = \delta_j^i$$

□

If ω is a 1-form on $U \in \mathbb{R}^n$ then by proposition (1.2), at each point $p \in U$

$$\omega_p = \sum a_i(p)(\mathrm{d}x^i)_p$$

Note the lower index on $a_i(p)$ as opposed to the upper index on $X_p = \sum a^i(p)\partial_{x^i}|_p$. If $x := x^1, y := x^2, z := x^3$, then $\mathrm{d}x, \mathrm{d}y, \mathrm{d}z$.

Proposition 1.3: $\mathrm{d}f$ in terms of coordinates

If $f: U \rightarrow \mathbb{R}$, then

$$\mathrm{d}f = \sum \frac{\partial f}{\partial x^i} \mathrm{d}x^i \quad (23)$$

Proof.

$$(\mathrm{d}f)_p = \sum a_i(p)(\mathrm{d}x^i)_p$$

for some real numbers $a_i(p)$ depending on p . Thus

$$\begin{aligned} \mathrm{d}f \left(\frac{\partial}{\partial x^j} \right) &= \sum_i a_i \mathrm{d}x^i \left(\frac{\partial}{\partial x^j} \right) \\ &= \sum_i a_i \delta_j^i = a_j \end{aligned} \quad (24)$$

On the other hand, by the definition of the differential

$$\mathrm{d}f = \left(\frac{\partial}{\partial x^j} \right) = \frac{\partial f}{\partial x^j} \quad (25)$$

Therefore

$$a_j = \frac{\partial f}{\partial x^j} \quad (26)$$

and hence

$$(\mathrm{d}f)_p = \frac{\partial f}{\partial x^i} \Big|_p (\mathrm{d}x^i)_p$$

□

1.5.2 Differential k -forms

Definition 1.4: Differential k -forms

More generally, a *differential form* ω of degree k is a function that at each point assigns an alternating k -linear function on $T_p(\mathbb{R}^n)$, i.e. $\omega_p \in A^k(T_p(\mathbb{R}^n))$.

A basis for $A^k(T_p(\mathbb{R}^n))$ is

$$(\mathrm{d}x^I)_p := (\mathrm{d}x^{i_1})_p \wedge \cdots \wedge (\mathrm{d}x^{i_k})_p \quad (27)$$

where $1 \leq i_1 < \cdots < i_k \leq n$.

What is the nuance here?

Therefore, at each point $p \in U$, ω_p is a linear combination

$$\begin{aligned} \omega_p &= \sum_I a_I(p)(\mathrm{d}x^I)_p \\ 1 \leq i_1 < \cdots < i_k \leq n \end{aligned} \quad (28)$$

and a k -form ω on open U is a linear combination

$$\omega = \sum_I a_I \mathrm{d}x^I \quad (29)$$

with function coefficients $a_I: U \rightarrow \mathbb{R}$. We say that a k -form ω is C^∞ on U if all of the coefficients a_I are C^∞ functions on U . Denote $\Omega^k(U)$ the vector space of k -forms on U . A 0-form on U assigns to each point p an element of $A^0(T_p(\mathbb{R}^n)) := \mathbb{R}$; thus, a 0-form on U is a constant function. Note there are no nonzero differential forms of degree $> n$ on U since if $\deg \mathrm{d}x^I > n$ then at least two of the component 1-forms of $\mathrm{d}x^I$ must be the same and therefore $\mathrm{d}x^I = 0$.

Definition 1.5: Wedge product of forms

The *wedge product* of a k -form ω and ℓ -form τ is defined pointwise

$$\begin{aligned} (\omega \wedge \tau)_p &:= \omega_p \wedge \tau_p \\ \omega \wedge \tau &= \sum_{I,J} (a_I b_J) \mathrm{d}x^I \wedge \mathrm{d}x^J \end{aligned} \quad (30)$$

where $I \cap J = \emptyset$.

Hence the wedge product is bilinear

$$\wedge: \Omega^k(U) \times \Omega^\ell(U) \rightarrow \Omega^{k+\ell}(U) \quad (31)$$

The wedge product of forms is also anticommutative and associate (owing to the associativity and anticommutativity of the wedge product on multi-covectors) as therefore induces a graded algebra on $\Omega(U) := \bigoplus_k \Omega^k(U)$.

Example 1.3. In the case of

$$\wedge: \Omega^0(U) \times \Omega^\ell(U) \rightarrow \Omega^\ell$$

we have the pointwise multiplication of a C^∞ function and a C^∞ ℓ -form

$$(f \wedge \omega)_p = f(p) \wedge \omega_p = f(p)\omega_p$$

Let x, y, z be the coordinates on \mathbb{R}^3 . Then, the 1-forms are

$$f \mathrm{d}x + g \mathrm{d}y + h \mathrm{d}z$$

the 2-forms are

$$f \mathrm{d}y \wedge \mathrm{d}z + g \mathrm{d}x \wedge \mathrm{d}z + h \mathrm{d}x \wedge \mathrm{d}y$$

and the 3-forms are

$$f \mathrm{d}x \wedge \mathrm{d}y \wedge \mathrm{d}z$$

1.5.3 Differential Forms as Multilinear Functions on Vector Fields

If ω is a 1-form and X is a vector field then

$$\begin{aligned}\omega(X)|_p &:= \omega_p(X_p) \\ \omega &= \sum a_i dx^i \quad X = \sum b^j \frac{\partial}{\partial x^j} \\ \omega(X) &= \left(\sum a_i dx^i \right) \left(\sum b^j \frac{\partial}{\partial x^j} \right) \\ &= \sum a_i b^i\end{aligned}\tag{32}$$

1.5.4 Exterior Derivative

Definition 1.6: Exterior Derivative

The exterior derivative of a function $f \in C^\infty(U)$ is defined to be its differential df

$$df = \sum \frac{\partial f}{\partial x^i} dx^i$$

For $k \geq 1$, if $\omega = \sum_I a_I dx^I$ is a k -form, then

$$\begin{aligned}d\omega &:= \sum_I da_I \wedge dx^I \\ &= \sum_I \left(\sum_j \frac{\partial a_I}{\partial x^j} dx^j \right) \wedge dx^I\end{aligned}\tag{33}$$

Example 1.4. Let ω be the 1-form $f dx + g dy$ on \mathbb{R}^2 . Then

$$d\omega = df \wedge dx + dg \wedge dy\tag{34}$$

$$= \left(\frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy \right) \wedge dx\tag{35}$$

$$\begin{aligned}&+ \left(\frac{\partial g}{\partial x} dx + \frac{\partial g}{\partial y} dy \right) \wedge dy \\ &= \left(\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx \wedge dy\end{aligned}\tag{36}$$

where we use that $dx \wedge dy = -dy \wedge dx$ and $dx \wedge dx = 0$.

Definition 1.7: Antiderivation

Let $A = \bigoplus_k A^k$ be a graded algebra over a field K . An *antiderivation of the graded algebra* A is a k -linear map $D: A \rightarrow A$ such that for $a \in A^k, b \in A^\ell$

$$D(ab) = (Da)b + (-1)^k aDb\tag{37}$$

If there is an integer m such that D sends A^k to A^{k+m} for all k , then the antiderivation is of *degree* m .

Proposition 1.4: Properties of exterior differentiation

1. exterior differentiation is an antiderivation of degree 1:

$$d(\omega \wedge \tau) = (d\omega) \wedge \tau + (-1)^{\deg \omega} \omega \wedge d\tau$$

2. $d^2 = 0$

Proposition 1.5: Characterization of the exterior derivative

The properties of proposition (1.4) completely characterize exterior differentiation.

1.5.5 Closed and Exact Forms

A k -form ω is a *closed form* if $d\omega = 0$. ω is an *exact form* if there is a $k-1$ -form τ such that $\omega = d\tau$. Since $d(d\tau) = 0$, every exact form is closed.

Example 1.5. A closed 1-form on the punctured plane. Define ω on the manifold $\mathbb{R}^2 - \{0\}$ by

$$\omega(x, y) = \frac{-y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy$$

To show that ω is closed we take the exterior derivative using example (36):

$$\begin{aligned}d\omega &= \left(\frac{y^2 - x^2}{(x^2 + y^2)^2} - \frac{y^2 - x^2}{(x^2 + y^2)^2} \right) dx \wedge dy \\ &= 0\end{aligned}\tag{38}$$

Definition 1.8: Differential Complex

A collection of vector spaces $\{V^0, V^1, \dots\}$ with linear maps $d_k: V^k \rightarrow V^{k+1}$ such that $d_{k+1} \circ d_k = 0$ is called a *differential complex* or a *cochain complex*. For any open subset $U \subset \mathbb{R}^3$, the exterior derivative makes the vector space $\Omega^*(U)$ of C^∞ forms on U into a cochain complex, called the *de Rham complex of U* :

$$0 \rightarrow \Omega^0(U) \xrightarrow{d} \Omega^1(U) \xrightarrow{d} \Omega^2(U) \xrightarrow{d} \dots$$

The closed forms are the elements of the kernel of d and the exact forms are the elements of the image of d .

1.5.6 Applications to Vector Calculus

Recall the three operators grad, curl, div ($\nabla, \nabla \times, \nabla \cdot$) on scalar fields and vector fields (i.e. scalar valued functions

and vector valued functions) over \mathbb{R}^3 :

$$\begin{aligned}\nabla f &= \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} \\ \nabla \times \begin{bmatrix} P \\ Q \\ R \end{bmatrix} &= \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{bmatrix} \times \begin{bmatrix} P \\ Q \\ R \end{bmatrix} = \begin{bmatrix} \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \\ \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial z} \\ \frac{\partial P}{\partial x} - \frac{\partial Q}{\partial y} \end{bmatrix} \\ \nabla \cdot \begin{bmatrix} P \\ Q \\ R \end{bmatrix} &= \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{bmatrix} \cdot \begin{bmatrix} P \\ Q \\ R \end{bmatrix} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}\end{aligned}\quad (39)$$

Note we can identify 1-forms with vector fields:

$$P dx + Q dy + R dz \longleftrightarrow \begin{bmatrix} P \\ Q \\ R \end{bmatrix}$$

Similarly 2-forms on \mathbb{R}^3

$$P dx \wedge dy + Q dz \wedge dx + R dx \wedge dy \longleftrightarrow \begin{bmatrix} P \\ Q \\ R \end{bmatrix}$$

and 3-forms on U can be identified with functions U

$$f dx \wedge dy \wedge dz \longleftrightarrow f$$

In terms of these identifications the exterior derivative of 0-form f is the 1-form df or ∇f :

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz \longleftrightarrow \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial z} \end{bmatrix} = \nabla f$$

the exterior derivative of a vector field $[P \ Q \ R]^T$ (i.e. 1-form) is the 2-form $\nabla \times [P \ Q \ R]^T$

$$\begin{aligned}d(P dx + Q dy + R dz) &= \\ &+ \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) dy \wedge dz \\ &- \left(\frac{\partial R}{\partial x} - \frac{\partial P}{\partial z} \right) dz \wedge dx \\ &+ \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx \wedge dy \\ &\longleftrightarrow \\ &\nabla \times \begin{bmatrix} P \\ Q \\ R \end{bmatrix}\end{aligned}\quad (40)$$

and the exterior derivative of a 2-form is

$$\begin{aligned}d(P dx \wedge dy + Q dz \wedge dx + R dx \wedge dy) &= \\ \left(\frac{\partial P}{\partial x} dx + \frac{\partial Q}{\partial y} dy + \frac{\partial R}{\partial z} dz \right) dx \wedge dy \wedge dz \\ &\longleftrightarrow \nabla \cdot \begin{bmatrix} P \\ Q \\ R \end{bmatrix}\end{aligned}\quad (41)$$

Thus, the exterior derivatives on 0-forms (functions) is the grad operator, on 1-forms is the curl operator, and on 2-forms is the divergence operator.

$$\begin{array}{ccccccc} \Omega^0(U) & \xrightarrow{d} & \Omega^1(U) & \xrightarrow{d} & \Omega^2(U) & \xrightarrow{d} & \Omega^3(U) \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ C^\infty(U) & \xrightarrow{\nabla} & \mathfrak{X}(U) & \xrightarrow{\nabla \times} & \mathfrak{X}(U) & \xrightarrow{\nabla \cdot} & C^\infty(U) \end{array}$$

where $\mathfrak{X}(U)$ is the *Lie algebra* of C^∞ vector fields on U .

Proposition 1.6: grad, curl, div properties

1. $\nabla \times \nabla f = 0$
2. $\nabla \cdot (\nabla \times [P \ Q \ R]^T) = 0$
3. On \mathbb{R}^3 , a vector field \mathbf{F} is the gradient of some scalar function iff $\nabla \times \mathbf{F} = 0$

Properties (1,2) express that $d^2 = 0$ Property (3) expresses the fact that a 1-form on \mathbb{R}^3 is exact iff it is closed; it need not be true on a region other than \mathbb{R}^3 . It turns out that whether proposition (3) is true depends on the topology of U . One measure of the failure of a closed k -form to be exact is the quotient vector space

$$H^k(U) := \frac{\{\text{closed } k\text{-forms on } U\}}{\{\text{exact } k\text{-forms on } U\}} \quad (42)$$

called the k th *de Rham cohomology* of U . The generalization of proposition (3) to any differential on \mathbb{R}^n is called the *Poincare lemma*: for $k \geq 1$, every closed k -form on \mathbb{R}^n is exact. This is equivalent to the vanishing of the k th de Rham cohomology $H^k(\mathbb{R}^n)$ for $k \geq 1$.

1.5.7 Convention on Subscripts and Superscripts

Vector fields e_1, e_2, \dots have subscripts and differential forms $\omega^1, \omega^2, \dots$ have superscripts. Coordinate functions x^1, x^2, \dots , being 0-forms, have superscripts. Their differentials dx^i should also. Coordinate vector fields $\frac{\partial}{\partial x^i}$ are considered to have subscripts because the index is in the denominator. Coefficient functions have subscripts or superscripts depending on whether they're coefficients functions for vector fields or forms. This allows for "conservation of indices": if $X = \sum a^i \frac{\partial}{\partial x^i}$ and $\omega = \sum b_j dx^j$ then

$$\omega(X) = \left(\sum b_j dx^j \right) \left(\sum a^i \frac{\partial}{\partial x^i} \right) = \sum b_i a^i$$

1.6 Manifolds

Definition 1.9: Locally Euclidean

A topological space M is *locally Euclidean of dimension n* if for every $p \in M$ there exists a neighborhood U such that there is a *homeomorphism*¹ ϕ from U **onto** an opensubset of \mathbb{R}^n . The pair $(U, \phi: U \rightarrow \mathbb{R}^n)$ is called a *chart*, with U being the *coordinate neighborhood* and ϕ the *coordinate system*.

Definition 1.10: Topological Manifold

A *topological manifold* is a *Hausdorff*², *second countable*³, locally Euclidean space.

1.6.1 Compatible Charts

Suppose $(U, \phi: U \rightarrow \mathbb{R}^n)$ and $(V, \psi: V \rightarrow \mathbb{R}^n)$ are two charts of a topological manifold; since $U \cap V$ is open and ϕ is a homeomorphism onto an open subset, the image $\phi(U \cap V)$ is also an open subset (similarly $\psi(U \cap V)$).

Definition 1.11: Charts

Two charts $(U, \phi: U \rightarrow \mathbb{R}^n)$ and $(V, \psi: V \rightarrow \mathbb{R}^n)$ of a topological manifold are *C^∞ -compatible* if the two composed maps

$$\begin{aligned}\phi \circ \psi^{-1}: \psi(U \cap V) &\rightarrow \phi(U \cap V) \\ \psi \circ \phi^{-1}: \phi(U \cap V) &\rightarrow \psi(U \cap V)\end{aligned}\quad (43)$$

See figure ???. These two maps $\phi \circ \psi^{-1}$ and $\psi \circ \phi^{-1}$ are called the *transition functions* between the charts;

$$\begin{aligned}\phi \circ \psi^{-1}: \mathbb{R}^n &\rightarrow U \cap V \rightarrow \mathbb{R}^n \\ \psi \circ \phi^{-1}: \mathbb{R}^n &\rightarrow U \cap V \rightarrow \mathbb{R}^n\end{aligned}\quad (44)$$

Definition 1.12: Atlas

A *C^∞ atlas* on a topological manifold M is a collection $\mathfrak{A} := \{(U_i, \phi_i)\}$ of pairwise *C^∞ -compatible* charts that *cover* M , i.e. such that $M = \bigcup_i U_i$.

¹A homeomorphism is a continuous function between topological spaces that has a continuous inverse function.

²A Hausdorff space, separated space is a topological space where for any two distinct points there exists a neighbourhood of each which is disjoint from the neighbourhood of the other.

³A topological space T is second-countable if there exists some countable collection $\mathcal{U} = \{U_i\}_{i=1}^\infty$ of open subsets of T such that any open subset of T can be written as a union of elements of some subfamily of \mathcal{U} .

Example 1.6. A *C^∞ atlas* on a circle The unit circle S^1 in the complex plane \mathbb{C} maybe described as

$$\{e^{it} \in \mathbb{C} \mid 0 \leq t \leq 2\pi\}$$

Let U_1, U_2 be

$$\begin{aligned}U_1 &:= \{e^{it} \in \mathbb{C} \mid -\pi < t < \pi\} \\ U_2 &:= \{e^{it} \in \mathbb{C} \mid 0 < t < 2\pi\}\end{aligned}\quad (45)$$

and define

$$\begin{aligned}\phi_1(e^{it}) &= t \quad -\pi < t < \pi \\ \phi_2(e^{it}) &= t \quad 0 < t < 2\pi\end{aligned}\quad (46)$$

See figure ??. Both ϕ_1, ϕ_2 are homemorphisms onto their respective images; thus, U_1, ϕ_1 and U_2, ϕ_2 are charts on S^1 . The intersection $U_1 \cap U_2$ consists of two disjoint connected components

$$\begin{aligned}A &:= \{e^{it} \in \mathbb{C} \mid -\pi < t < 0\} \\ B &:= \{e^{it} \in \mathbb{C} \mid 0 < t < \pi\}\end{aligned}\quad (47)$$

with

$$\begin{aligned}\phi_1(U_1 \cap U_2) &= \phi_1(A \sqcup B) = \phi_1(A) \sqcup \phi_2(B) = (-\pi, 0) \sqcup (0, \pi) \\ \phi_2(U_1 \cap U_2) &= \phi_1(B \sqcup A) = \phi_1(B) \sqcup \phi_2(A) = (0, \pi) \sqcup (\pi, 2\pi)\end{aligned}\quad (48)$$

where \sqcup means disjoint union. The transition function

$$\left(\phi_2 \circ \phi_1^{-1}\right)(t) = \begin{cases} t + 2\pi & \text{for } t \in (-\pi, 0) \\ t & \text{for } t \in (0, \pi) \end{cases}$$

and similarly

$$\left(\phi_1 \circ \phi_2^{-1}\right)(t) = \begin{cases} t & \text{for } t \in (0, \pi) \\ t - 2\pi & \text{for } t \in (\pi, 2\pi) \end{cases}$$

Therefore, since the transition functions are *C^∞* , the charts are *C^∞ compatible* and form an atlas on S^1 .

Although compatibility is reflexive and symmetric it is not transitive because we don't know anything about mutual intersections. On the otherhand

Lemma 1.1

Let $\{(U_i, \phi_i)\}$ be an atlas for a topological manifold. If two other charts (not in the atlas) $(V, \psi), (U, \phi)$ are both compatible with the atlas (i.e. with all charts in the atlas), then they are mutually compatible.

1.6.2 Smooth Manifolds

An atlas \mathfrak{M} on a topological manifold is said to be *maximal* if it is not strictly contained in another atlas.

Definition 1.13: Smooth Manifold

A *smooth C^∞ manifold* is a topological manifold M together with a maximal atlas. The maximal atlas is called a *differentiable structure* on M . M is said to have dimension n if all of its connected components have dimension n . A 1-dimensional manifold is called a *curve*, a 2-dimensional manifold is called a *surface*, and an n -dimensional manifold is called an n -manifold.

In the context of manifolds, we denote the standard coordinates on \mathbb{R}^n by r^1, \dots, r^n . If (U, ϕ) is a chart of a manifold, we let $x^i = r^i \circ \phi$ be the i th component of ϕ and write $\phi = (x^1, \dots, x^n)$. Thus, $x^i(p) := (r^i \circ \phi)(p)$ is a point in \mathbb{R}^n . The functions x^i are called the *local coordinates* on U . The sense here is that x^i tell you where you are on the manifold in a standardized way, given some intrinsic description of where you are on the manifold. Think of how lat/lon tell you where you are on the globe given some intrinsic identification of where you are on the globe. This understanding gives sense to the words chart and atlas. Abusing notation, sometimes (x^1, \dots, x^n) (sans p) stands for both the coordinates on U or for a point in \mathbb{R}^n .

1.6.3 Examples of Smooth Manifolds

Example 1.7. Graph of a Smooth Function For a subset $A \subset \mathbb{R}^n$ and a function $f: A \rightarrow \mathbb{R}^m$ the *graph* of f is defined

$$\Gamma(f) = \{(x, f(x)) \in A \times \mathbb{R}^m\}$$

If U is an open subset \mathbb{R}^n and $f: U \rightarrow \mathbb{R}^n$ is C^∞ , then the two maps

$$\begin{aligned} \phi: \Gamma(f) &\rightarrow U & (x, f(x)) &\mapsto x \\ (1, f): \Gamma(f) &\rightarrow \Gamma(f) & x &\mapsto (x, f(x)) \end{aligned}$$

constitute a homomorphism. Hence, $\Gamma(f)$ has an atlas with a single $(\Gamma(f), \phi)$ and is therefore a C^∞ manifold.

Example 1.8. General Linear group Let $\mathbb{R}^{m \times n}$ be the vector space of all $m \times n$ matrices. Since $\mathbb{R}^{m \times n}$ is isomorphic to \mathbb{R}^{mn} , we give it the topology of \mathbb{R}^{mn} . The *general linear group* $\text{GL}(n, \mathbb{R})$ is defined

$$\begin{aligned} \text{GL}(n, \mathbb{R}) &:= \{A \in \mathbb{R}^{n \times n} \mid \det(A) \neq 0\} \\ &= \det^{-1}(\mathbb{R} - \{0\}) \end{aligned} \quad (49)$$

Since the determinant function $\det: \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is continuous, $\text{GL}(n, \mathbb{R})$ is an open subset of $\mathbb{R}^{n \times n} \cong \mathbb{R}^{n^2}$ is therefore a manifold.

Example 1.9. Unit circle in the xy -plane Take S^1 as the unit circle in the real plane \mathbb{R}^2 defined by $x^2 + y^2 = 1$. We can cover S^1 by four open sets: the upper and lower semicircles U_1, U_2 and the left and right semicircles U_3, U_4 (see figure ??). On U_1, U_2 the coordinate function x is a homeomorphism onto the open interval $(-1, 1)$ on the

x -axis. Thus, $\phi_i(x, y) := x$. Similarly, on U_3, U_4 , y is a homeomorphism onto the open interval $(-1, 1)$ on the y -axis, and so $\phi_i(x, y) := y$. You can check that on $U_i \cap U_j$, the transition function $\phi_j \circ \phi_i^{-1}$ is C^∞ . For example, on $U_1 \cap U_3$

$$(\phi_3 \circ \phi_1^{-1})(x) = \phi_3(x, \sqrt{1-x^2}) = \sqrt{1-x^2}$$

while on $U_2 \cap U_4$

$$(\phi_4 \circ \phi_2^{-1})(x) = \phi_4(x, -\sqrt{1-x^2}) = -\sqrt{1-x^2}$$

Thus, $\{(U_i, \phi_i)\}_{i=1}^4$ is a C^∞ atlas on S^1 .

Proposition 1.7: An atlas for a product manifold

If $\{(U_i, \phi_i)\}$ and $\{(V_j, \psi_j)\}$ are C^∞ atlases for the manifolds M, N of dimensions, respectively, then the collection

$$\{(U_i \times V_j, (\phi_i, \psi_j)): U_i \times V_j \rightarrow \mathbb{R}^m \times \mathbb{R}^n\} \quad (50)$$

is a C^∞ atlas on $M \times N$. Therefore $M \times N$ is a C^∞ manifold of dimension $m+n$.

There n -dimensional torus $S^1 \times \dots \times S^1$ is a manifold.

1.6.4 Smooths Maps on a Manifold**Definition 1.14: Smooth at a point to \mathbb{R}**

Let M be a smooth manifold of dimension n . A function $f: M \rightarrow \mathbb{R}$ is said to be C^∞ or *smooth* at a point p in M if there is a chart (U, ϕ) containing p such that $f \circ \phi^{-1}$, a function defined on the open subset $\phi(U) \subset \mathbb{R}^n$, is C^∞ at $\phi(p)$. To summarize f is C^∞ if

$$f \circ \phi^{-1}: \phi(U) \rightarrow \mathbb{R}$$

See figure ??.

Definition 1.15: Pullback

Let $F: N \rightarrow M$ be a map and h a function M . The *pullback* of h by F , denoted F^*h , is the composite function $h \circ F$.

Thus, a function f on M is C^∞ on a chart (U, ϕ) iff $(\phi^{-1})^*f \equiv f \circ \phi^{-1}$ is C^∞ on $\phi(U)$.

Definition 1.16: Smooth at a point

Let M, N be manifolds of dimension m, n . A continuous map $F: N \rightarrow M$ is C^∞ at a point $p \in N$ if there are charts $(V, \psi), (U, \phi)$ about $F(p) \in M$ and $p \in N$ such that $\psi \circ F \circ \phi^{-1}$ is C^∞ at $\phi(p)$. To summarize F is C^∞ if

$$\psi \circ F \circ \phi^{-1}: \phi(F^{-1}(V) \cap U) \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$$

See figure ??.

Proposition 1.8: Composition of C^∞ maps

If $F: N \rightarrow M$ and $G: M \rightarrow P$ are both C^∞ maps of manifolds, then the composite $G \circ F: N \rightarrow P$ is C^∞ .

Definition 1.17: Diffeomorphism

A *diffeomorphism* of manifolds is a bijective C^∞ map $F: N \rightarrow M$ whose inverse F^{-1} is also C^∞ .

Proposition 1.9

If (U, ϕ) is a chart on a manifold M of dimension n , then the coordinate map $\phi: U \rightarrow \phi(U)$ is a diffeomorphism.

Definition 1.18: Lie group

A *Lie group* is a C^∞ manifold G having a group structure such that the multiplication map

$$\mu: G \times G \rightarrow G$$

and the inverse map

$$\iota: G \rightarrow G \quad \iota(x) := x^{-1}$$

are both C^∞ . Similarly, a *topological group* is a topological space having a group structure such that multiplication and inverse maps are both continuous.

Example 1.10. Recall the definition of $\text{GL}(n, \mathbb{R})$. As an open subset of $\mathbb{R}^{n \times n}$, it is a manifold. Since the (i, j) -entry of the product of two matrices A, B in $\text{GL}(n, \mathbb{R})$

$$(AB)_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

is a polynomial in the coordinates of A and B . Therefore matrix multiplication

$$\mu: \text{GL}(n, \mathbb{R}) \times \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R})$$

is a C^∞ map. Furthermore, by Cramer's rule

$$(A^{-1})_{ij} = \frac{(-1)^{i+j}}{\det A} ((j, i) - \text{minor of } A)$$

which is a C^∞ function of the a_{ij} s provided $\det A \neq 0$. Therefore, $\text{GL}(n, \mathbb{R})$ is a Lie group.

1.6.5 Partial Derivatives

Let (U, ϕ) be a chart and f a C^∞ function on the manifold. As a function into \mathbb{R}^n , ϕ has n components x^1, \dots, x^n . Therefore, if r^1, \dots, r^n are the standard coordinates on \mathbb{R}^n , then

$$x^i = r^i \circ \phi$$

What this means is ϕ maps to some point $(\phi^1(p), \dots, \phi^n(p))$ in \mathbb{R}^n . The projection to the i th standard coordinate on \mathbb{R}^n is $r^i \circ \phi$, i.e. just pick out the component of $(\phi^1(p), \dots, \phi^n(p))$. This direct path from U to the i th standard coordinate of \mathbb{R}^n is called a local coordinate for U (since it's only valid in a small neighborhood of p). Then the collection (x^1, \dots, x^n) are the local coordinates on U (but don't forget that the coordinates map from $U \rightarrow \mathbb{R}^n$).

Definition 1.19: Partial derivative

For $p \in U$, we define the *partial derivative* $\frac{\partial f}{\partial x^i}$ of f with respect to x^i at p

$$\begin{aligned} \frac{\partial}{\partial x^i} \Big|_p f &:= \frac{\partial f}{\partial x^i} \Big|_p \\ &:= \frac{\partial (f \circ \phi^{-1})}{\partial r^i} \Big|_{\phi(p)} \\ &:= \frac{\partial}{\partial r^i} \Big|_{\phi(p)} (f \circ \phi^{-1}) \end{aligned} \quad (51)$$

Thus, as functions on $\phi(U)$ (and in terms of the local coordinates on U)

$$\frac{\partial f}{\partial x^i} \circ \phi^{-1} = \frac{\partial f \circ \phi^{-1}}{\partial r^i} \quad (52)$$

and therefore the partial derivative $\partial f / \partial x^i$ is C^∞ on U because its pullback

$$(\phi^{-1})^* \frac{\partial f}{\partial x^i}$$

is C^∞ .

Proposition 1.10: Partial derivatives are dual

Suppose U, x^1, \dots, x^n (with local coordinates x^1, \dots, x^n) is a chart on a manifold. Then $\partial x^i / \partial x^j = \delta_j^i$.

Proof. At a point $p \in U$, by the above definition of $\partial/\partial x^j|_p$

$$\begin{aligned} \frac{\partial x^i}{\partial x^j} \Big|_p &= \frac{\partial x^i \circ \phi^{-1}}{\partial r^j} \Big|_{\phi(p)} \\ &= \frac{\partial (r^i \circ \phi \circ \phi^{-1})}{\partial r^j} \Big|_{\phi(p)} \\ &= \frac{\partial r^i}{\partial r^j} \Big|_{\phi(p)} = \delta_j^i \end{aligned} \quad (53)$$

□

Definition 1.20: Jacobian

Let $F: N \rightarrow M$ be a smooth map, and let $(U, \phi) \equiv (U, x^1, \dots, x^n)$ and $(V, \psi) \equiv (U, y^1, \dots, y^n)$ be charts on N, M respectively such that $F(U) \subset V$. Denote by

$$(F_i := y^i \circ F \equiv r^i \circ \psi \circ F): U \rightarrow \mathbb{R}$$

the i th component of F in the chart V, ψ . Then the matrix $\left[\frac{\partial F^i}{\partial x^j} \right]$ is called the *Jacobian matrix* of F relative to the charts U, ϕ and (V, ψ) .

A diffeomorphism $F: U \rightarrow F(U) \subset \mathbb{R}^n$ may be thought of as coordinate system on U . We say that a C^∞ map $F: N \rightarrow M$ is *locally invertible* or a *local diffeomorphism* at $p \in N$ if p has a neighborhood U on which F is a diffeomorphism. Given n smooth functions F^1, \dots, F^n in a neighborhood of p one would like to know whether they form a coordinate system. This is equivalent to whether $F = (F^1, \dots, F^n)$ is a local diffeomorphism at p .

Theorem 1.2: Inverse function theorem \mathbb{R}^n

Let $F: W \rightarrow \mathbb{R}^n$ be a C^∞ map defined on an open subset $W \subset \mathbb{R}^n$. For any point $p \in W$, the map F is locally invertible iff the determinant of the Jacobian

$$\det \left[\frac{\partial F^i}{\partial x^j} \Big|_p \right]$$

is not zero.

Theorem 1.3: Inverse function theorem for manifolds

Let $F: N \rightarrow M$ be a C^∞ map between manifolds of the same dimension, and $p \in N$. Suppose for some charts $(U, \phi) = (U, x^1, \dots, x^n)$, $(V, \psi) = (U, y^1, \dots, y^n)$ about p and $F(p)$ respectively and $F(U) \subset V$. Set $F^i = y^i \circ F$. Then F is locally invertible at p iff

$$\det \left[\frac{\partial F^i}{\partial x^j} \Big|_p \right]$$

is not zero.

Corollary 1.1

Let N be a manifold of dimension n . A set of smooth functions F^1, \dots, F^n defined on a coordinate neighborhood U, x^1, \dots, x^n of a point $p \in N$ forms a coordinate system about p iff the Jacobian determinant

$$\det \left[\frac{\partial F^i}{\partial x^j} \Big|_p \right]$$

is not zero.

1.7 Tangent Space

The tangent space to a manifold at a point is the vector space of derivations (germs/directional derivatives) at the point. A smooth map of manifolds induces a linear map, called its differential, of tangent spaces at corresponding points. In local coordinates, the differential is represented by the Jacobian. In this sense, the differential of a map is the generalization of the derivative between Euclidean spaces.

A basic theme in manifold theory is linearization, according to which a manifold can be approximated by its tangent space and a smooth map can be approximated by the differential of that map. The differential further categorizes maps as either immersions or submersions (depending on whether the differential is injective or surjective).

The collection of tangent spaces to a manifold can be given the structure of a *vector bundle*; it is thus called the *tangent bundle*. Vector fields, which manifest themselves in the physical world as velocity, force, electricity, and magnetism, maybe viewed as sections of the tangent bundle.

1.7.1 The Tangent Space at a Point

Definition 1.21: Tangent vector

The germ of a C^∞ function $p \in M$ to be an equivalence class of C^∞ functions defined in a neighborhood of $p \in M$, with equivalence defined as agreement on a (possibly smaller) neighborhood of p . The set of germs of C^∞ real-valued functions is denoted $C_p^\infty(M)$. Addition and multiplication of functions induces a ring structure on $C_p^\infty(M)$; with scalar multiplication by real numbers $C_p^\infty(M)$ becomes an algebra over \mathbb{R} . A derivation at a point in M is a linear map $D: C_p^\infty(M) \rightarrow \mathbb{R}$ such that

$$D(fg) = (Df)g + f(p)Dg$$

A tangent vector at a point p is a derivation at p .

Given a coordinate neighborhood $(U, \phi) = (U, x^1, \dots, x^n)$ and (r^1, \dots, r^n) the standard coordinates on \mathbb{R}^n and

$$x^i = r^i \circ \phi: U \rightarrow \mathbb{R}$$

If f is a smooth function in a neighborhood of p

$$\left. \frac{\partial}{\partial x^i} \right|_p f = \left. \frac{\partial}{\partial r^i} \right|_{\phi(p)} (f \circ \phi^{-1})$$

The partial derivatives satisfy the derivation property and therefore qualify as tangent vectors.

1.7.2 The Differential of a Map

Let $F: N \rightarrow M$ be a C^∞ map between two manifolds. At each point $p \in N$, the map F induces a linear map of tangent spaces, called its *differential at p*

$$F_*: T_p N \rightarrow T_{F(p)} M$$

defined as follows: if $X_p \in T_p N$, then $F_*(X_p)$ is the tangent vector in $T_{F(p)} M$ according to

$$(F_*(X_p))f = X_p(f \circ F)$$

for $f \in C_{F(p)}^\infty(M)$ a germ (or representative of the germ). Vectors *pushforward* through F .

Example 1.11. Differential of a map between Euclidean spaces Suppose $F: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is smooth and $p \in \mathbb{R}^n$. Let x^1, \dots, x^n be coordinates on \mathbb{R}^n and y^1, \dots, y^m be coordinates on \mathbb{R}^m . Then

$$\left\{ \left. \frac{\partial}{\partial x^1} \right|_p, \dots, \left. \frac{\partial}{\partial x^n} \right|_p \right\}$$

form a basis for the tangent space $T_p(\mathbb{R}^n)$ and

$$\left\{ \left. \frac{\partial}{\partial y^1} \right|_{F(p)}, \dots, \left. \frac{\partial}{\partial y^m} \right|_{F(p)} \right\}$$

form a basis for the tangent space $T_{F(p)}(\mathbb{R}^m)$. The linear map

$$F_*: T_p(\mathbb{R}^n) \rightarrow T_{F(p)}(\mathbb{R}^m)$$

(the differential of F) is described by a matrix $[a_j^i]$ relative to these two bases:

$$F_* \left(\left. \frac{\partial}{\partial x^j} \right|_p \right) = \sum_k a_j^k \left. \frac{\partial}{\partial y^k} \right|_{F(p)}$$

Let $F^i = y^i \circ F$ be the i th component of F ; we can find a_j^i by evaluating the right-hand side and left-hand side on y^i

$$\begin{aligned} \text{RHS} &= \sum_k a_j^k \left. \frac{\partial}{\partial y^k} \right|_{F(p)} y^i = \sum_k a_j^k \delta_k^i = a_j^i \\ \text{LHS} &= F_* \left(\left. \frac{\partial}{\partial x^j} \right|_p \right) y^i = \left. \frac{\partial}{\partial x^j} \right|_p (y^i \circ F) = \left. \frac{\partial F^i}{\partial x^j} \right|_p \end{aligned} \quad (54)$$

where we've used the fact that $F^i = y^i \circ F$. Thus, F_* relative to the bases is the Jacobian

$$\left[\left. \frac{\partial F^i}{\partial x^j} \right|_p \right]$$

and hence the differential generalizes the derivative of a map between Euclidean spaces (because in this instance the Jacobian is the derivative and in abstract manifolds we use the same F_*). Note this means

$$F_* \iff \text{Jacobian} \iff \text{differential} \iff \text{pushforward}$$

Example 1.12. Define a diffeomorphism F such that

$$F: (r, \theta) \mapsto (r \cos \theta, r \sin \theta) \quad (55)$$

The Jacobian of F

$$J = \begin{bmatrix} \frac{\partial F_1}{\partial r} & \frac{\partial F_1}{\partial \theta} \\ \frac{\partial F_2}{\partial r} & \frac{\partial F_2}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix} \quad (56)$$

where $(r, \theta) \equiv (x_1, x_2)$ and

$$(x, y) \equiv (y_1, y_2) \equiv (F_1, F_2) \equiv (r \cos \theta, r \sin \theta)$$

Note that $\det J = r$ and so F is a diffeomorphism iff $r \neq 0$. Given a vector field

$$X_p = a(r, \theta) \partial_r + b(r, \theta) \partial_\theta := a(r, \theta) \frac{\partial}{\partial r} + b(r, \theta) \frac{\partial}{\partial \theta} \quad (57)$$

we can compute the pushforward F_* wrt the ∂_x, ∂_y basis

$$F_*(X_p) = \begin{bmatrix} \cos(\theta) & -r \sin(\theta) \\ \sin(\theta) & r \cos(\theta) \end{bmatrix} \cdot \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} a \cos(\theta) - br \sin(\theta) \\ a \sin(\theta) + br \cos(\theta) \end{pmatrix} \quad (58)$$

Hence, explicitly

$$F_*(X_p) = (a \cos(\theta) - br \sin(\theta)) \partial_x + (a \sin(\theta) + br \cos(\theta)) \partial_y \quad (59)$$

Since J is invertible we can investigate which vector fields map to ∂_x

$$F_*X_p = \partial_x \iff X_p = F_*^{-1}\partial_x \quad (60)$$

Let $X_p = a\partial_r + b\partial_\theta$. Then

$$X_p = \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\frac{\sin(\theta)}{r} & \frac{\cos(\theta)}{r} \end{bmatrix} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \cos(\theta) \\ -\frac{\sin(\theta)}{r} \end{pmatrix} \quad (61)$$

However we need to write r, θ in terms of x, y

$$F_*^{-1}\partial_x = \frac{x}{\sqrt{x^2 + y^2}}\partial_r + \frac{y}{x^2 + y^2}\partial_\theta \quad (62)$$

F_*^{-1} is called the pullback F^* of the vector field ∂_x along F .

Let $F: N \rightarrow M$ and $G: M \rightarrow P$ be smooth maps of manifolds, and $p \in N$. The differentials of F at p and G at $F(p)$ are linear maps

$$T_p N \xrightarrow{F_{*,p}} T_{F(p)} M \xrightarrow{G_{*,F(p)}} T_{G(F(p))} P$$

Theorem 1.4: Chain rule

If $F: N \rightarrow M$ and $G: M \rightarrow P$ are smooth maps of and $p \in N$, then

$$(G \circ F)_{*,p} = G_{*,F(p)} \circ F_{*,p}$$

Proof. Let $X_p \in T_p N$ and let f be C^∞ at $G(F(p)) \in P$. Then

$$((G \circ F)_* X_p) f = X_p(f \circ G \circ F)$$

and

$$\begin{aligned} ((G_* \circ F_*) X_p) f &= (G_*(F_* X_p)) f \\ &= (F_* X_p)(f \circ G) \\ &= X_p(f \circ G \circ F) \end{aligned} \quad (63)$$

□

Example 1.13. Chain rule in Calculus notation Suppose $w = G(x, y, z)$ is a C^∞ function $\mathbb{R}^3 \rightarrow \mathbb{R}$ and $(x, y, z) = F(t)$ is a C^∞ function $\mathbb{R} \rightarrow \mathbb{R}^3$. Under composition

$$w = (G \circ F)(t) = G(x(t), y(t), z(t))$$

becomes a C^∞ function $\mathbb{R} \rightarrow \mathbb{R}$. The differentials

$$\begin{aligned} F_* &= \begin{bmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \\ \frac{dz}{dt} \end{bmatrix} \\ G_* &= \begin{bmatrix} \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{bmatrix} \\ (G \circ F)_* &= \frac{dw}{dt} \end{aligned} \quad (64)$$

and since composition of linear maps is matrix multiplication

$$\begin{aligned} (G \circ F)_* &= G_* \circ F_* \\ &= \begin{bmatrix} \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{bmatrix} \cdot \begin{bmatrix} \frac{dx}{dt} \\ \frac{dy}{dt} \\ \frac{dz}{dt} \end{bmatrix} \\ &= \frac{\partial w}{\partial x} \frac{dx}{dt} + \frac{\partial w}{\partial y} \frac{dy}{dt} + \frac{\partial w}{\partial z} \frac{dz}{dt} \end{aligned} \quad (65)$$

Proposition 1.11

If $F: N \rightarrow M$ is a diffeomorphism of manifolds then

$$F_*: T_p N \rightarrow T_{F(p)} M$$

is an isomorphism of vector spaces.

Recall r^i the standard coordinates on \mathbb{R}^n , (U, ϕ) a chart about $p \in M$ (M dimension n), and $x^i = r^i \circ \phi$. Since ϕ is a diffeomorphism onto its image, the differential (push-forward) ϕ_* is a vector space isomorphism and

Proposition 1.12

Let $(U, \phi) = (U, x^1, \dots, x^n)$. Then

$$\phi_* \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \frac{\partial}{\partial r^i} \Big|_{\phi(p)} \quad (66)$$

Proposition 1.13

Let $(U, \phi) = (U, x^1, \dots, x^n)$. Then $T_p M$ has basis

$$\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$$

Proposition 1.14: Transition matrix for co-ordinate vectors

Suppose $(U, \phi) = (U, x^1, \dots, x^n)$ and $(V, \psi) = (V, y^1, \dots, y^n)$. Then

$$\frac{\partial}{\partial x^j} = \sum_{i=1}^n \frac{\partial y^i}{\partial x^j} \frac{\partial}{\partial y^i}$$

on $U \cap V$.

Given $F: N \rightarrow M$ and $G: M \rightarrow P$ are smooth maps of and $p \in N, F(p) \in M$. We will find a local expression for the differential

$$F_{*,p}: T_p N \rightarrow T_{F(p)} M$$

relative to the two charts. Using the local coordinate bases (induced by the charts) $F_* \equiv F_{*,p}$, is completely deter-

mined by a_j^i such that

$$F_* \left(\frac{\partial}{\partial x^j} \Big|_p \right) = \sum_{k=1}^m a_j^k \frac{\partial}{\partial y^k} \Big|_{F(p)} \quad (67)$$

Applying y^i , we find that

$$\begin{aligned} a_j^i &= \left(\sum_{k=1}^m a_j^k \frac{\partial}{\partial y^k} \Big|_{F(p)} \right) y^i \\ &= F_* \left(\frac{\partial}{\partial x^j} \Big|_p \right) y^i \\ &= \frac{\partial}{\partial x^j} \Big|_p (y^i \circ F) \\ &= \frac{\partial F^i}{\partial x^j} \Big|_p \end{aligned} \quad (68)$$

We restate as a proposition

Proposition 1.15

Given $F: N \rightarrow M$ and $G: M \rightarrow P$ are smooth maps of and $p \in N, F(p) \in M$. Relative to bases $\partial/\partial x^j|_p$ for $T_p N$ and $\partial/\partial y^i|_{F(p)}$ for $T_{F(p)} M$ the differential $F_{*,p}: T_p \rightarrow T_{F(p)} M$ is represented by the matrix

$$\begin{bmatrix} \frac{\partial F^1}{\partial x^1} \Big|_p & \cdots & \frac{\partial F^1}{\partial x^n} \Big|_p \\ \vdots & \ddots & \vdots \\ \frac{\partial F^m}{\partial x^1} \Big|_p & \cdots & \frac{\partial F^m}{\partial x^n} \Big|_p \end{bmatrix} \quad (69)$$

Proposition 1.16: Velocity of a curve in local coordinates

Let $c: (a, b) \rightarrow M$ be a smooth curve and let U, x^1, \dots, x^n be a coordinate chart about $c(t)$. Write $c^i = x^i \circ c$ for the i th component of c in the chart. Then $c'(t)$ is given by

$$c'(t) = \sum_{i=1}^n \dot{c}^i(t) \frac{\partial}{\partial x^i} \Big|_{c(t)}$$

where \dot{c} is the scalar derivative of the i th component. Thus, relative to the basis $\{\partial/\partial x^j|_p\}$ for $T_{c(t)} M$, the velocity $c'(t)$ is represented by the column vector

$$\begin{bmatrix} \dot{c}^1(t) \\ \vdots \\ \dot{c}^n(t) \end{bmatrix}$$

Every smooth curve c at p in a manifold M gives rise to a tangent vector $c'(0)$ in $T_p M$. Conversely, one can show that every tangent vector $X_p \in T_p M$ is the velocity vector of some curve at p

Proposition 1.17: Existence of a curve with a given initial vector

For any point p in a manifold M and any tangent vector $X_p \in T_p M$, there are $\varepsilon > 0$ and a smooth curve $(-\varepsilon, \varepsilon) \rightarrow M$ such that $c(0) = p$ and $c'(0) = X_p$.

Proof. Let $(U, \phi) = (U, x^1, \dots, x^n)$ be a chart centered at p i.e. $\phi(p) = \mathbf{0} \in \mathbb{R}^n$. Suppose $X_p = \sum a^i \partial/\partial x^i|_p$ and let r^1, \dots, r^n be the standard coordinates on \mathbb{R}^n , with $x^i = r^i \circ \phi$. To find a curve c at p with $c'(0) = X_p$, start with a curve $\alpha \in \mathbb{R}^n$ with $\alpha(0) = \mathbf{0}$ and $\alpha'(0) = \sum a^i \partial/\partial r^i|_0$. We then map α to M via ϕ^{-1} (see figure ??). Let

$$\alpha(t) := (a^1 t, \dots, a^n t) \quad t \in (-\varepsilon, \varepsilon)$$

with ε small enough such that $\alpha(t) \in \phi(U)$. Then define $c = \phi^{-1} \circ \alpha: (-\varepsilon, \varepsilon) \rightarrow M$. Then $c(0) = \phi^{-1}(\alpha(0)) = \phi^{-1}(\mathbf{0}) = p$ and

1.7.3 Curves in a Manifold

A *smooth curve* is a smooth map $c: (a, b) \rightarrow M$ with $0 \in (a, b)$ and we say that c is a curve starting at p if $c(0) = p$. The *velocity vector* $c'(t_0)$ at time t_0 is

$$c'(t_0) := c_* \left(\frac{d}{dt} \Big|_{t_0} \right) \in T_{c(t_0)} M$$

$$\begin{aligned} c'(0) &= (\phi^{-1})_* \alpha_* \left(\frac{d}{dt} \Big|_{t=0} \right) \\ &= (\phi^{-1})_* \left(\sum_i a^i \frac{\partial}{\partial r^i} \Big|_0 \right) \\ &= \sum_i a^i \frac{\partial}{\partial x^i} \Big|_p = X_p \end{aligned} \quad (70)$$

□

This gives us a geometrical perspective on tangent vectors as directional derivatives

Proposition 1.18: Directional Derivatives

Suppose X_p is a tangent vector at a point $p \in M$ and $f \in C_p^\infty(M)$. If $c \in (-\varepsilon, \varepsilon) \rightarrow M$ is a smooth curve starting at p with $c'(0) = X_p$, then

$$X_p f = \left. \frac{d}{dt} \right|_0 (f \circ c)$$

Proposition 1.19

Let $F: N \rightarrow M$ be a smooth map of manifolds, $p \in N$, $X_p \in T_p N$. If c is a smooth curve starting at $p \in N$ with velocity X_p at p ,

$$F_{*,p}(X_p) = \left. \frac{d}{dt} \right|_0 (F \circ c)(t) \quad (71)$$

Example 1.14. Differential of left multiplication If $g \in \text{GL}(n, \mathbb{R})$, let $\ell_g: \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R})$ be left multiplication by g i.e. $\ell_g(B) = gB$. Since $\text{GL}(n, \mathbb{R}) \subset \mathbb{R}^{n \times n}$ the tangent space $T_g(\text{GL}(n, \mathbb{R}))$ can be identified with $\mathbb{R}^{n \times n}$. We show that with this identification the differential at the identity

$$(\ell_g)_{*,\iota}: T_\iota(\text{GL}(n, \mathbb{R})) \rightarrow T_g(\text{GL}(n, \mathbb{R}))$$

is also left multiplication by g .

Let $X \in T_\iota(\text{GL}(n, \mathbb{R})) = \mathbb{R}^{n \times n}$. To compute $(\ell_g)_{*,\iota}(X)$, choose a curve $c(t) \in \text{GL}(n, \mathbb{R})$ with $c(0) = \iota$ and $c'(0) = X$. Then $\ell_g(c(t)) = g \cdot c(t)$ is simply matrix multiplication. Then

$$\begin{aligned} (\ell_g)_{*,\iota}(X) &= \left. \frac{d}{dt} \right|_{t=0} \ell_g(c(t)) \\ &= \left. \frac{d}{dt} \right|_{t=0} gc(t) \\ &= gc'(0) = gX \end{aligned} \quad (72)$$

1.8 Tangent Bundle

A *smooth vector bundle* over a smooth manifold M is a smooth varying family of vector spaces, parameterized by M , that locally looks like a product (of the manifold (base space) and the tangent spaces (fibers)). The collection of tangent spaces to a manifold has the structure of a vector bundle, called the *tangent bundle*. A smooth map between manifolds induces, via its differential at each point, a bundle map of the corresponding tangent bundles. The tangent bundle is canonically associated to a manifold, hence invariants of the tangent bundle give rise to invariants of the manifold. For example *Chern-Weil theory of characteristic classes* uses differential geometry to construct invariants for vector bundles; applied to the tangent bundle,

characteristic classes lead to numerical diffeomorphism invariants of a manifold called *characteristic numbers* (which generalize the *Euler characteristic*). A *section* of a vector bundle is a map that maps from each point of M into the *fiber* of the bundle over the point; both vector fields and differential forms on a manifold are sections of vector bundles.

1.8.1 Topology of the Tangent Bundle

Let M be a smooth manifold. Recall that at each point $p \in M$, the tangent space $T_p M$ is the vector space of all point-derivations of $C_p^\infty(M)$ (itself the algebra of germs of C^∞ functions at p). The *tangent bundle* of M is the union of all the tangent spaces of M

$$TM := \bigsqcup_{p \in M} T_p M$$

There is a natural map (in the sense that it does not depend on choice of atlas or local coordinates) $\pi: TM \rightarrow M$ given by $\pi(v) = p$ if $v \in T_p M$. As a matter of notation, we sometimes write a tangent vector $v \in T_p M$ as a pair (p, v) .

As defined, TM is a set, with no topology or manifold structure. We make it into a smooth manifold and show that it is a C^∞ vector bundle over M . The first step is the topology. If $(U, \phi) = (U, x^1, \dots, x^n)$ is a coordinate chart on M , let

$$TU := \bigsqcup_{p \in U} T_p U = \bigsqcup_{p \in U} T_p M$$

where we've the fact that the algebra $C_p^\infty(U)$ of germs of C^∞ functions in U at p is the same as $C_p^\infty(M)$ (since germ equivalence classes are determined by agreement in a neighborhood of p) and therefore $T_p U = T_p M$. At a point $p \in U$ a tangent vector $v \in T_p M$

$$v = \sum_{i=1}^n c^i \frac{\partial}{\partial x^i} \Big|_p$$

In this expression, $c^i \equiv c^i(v)$ depend on v and so **therefore end up being functions on TU** . Define $\tilde{\phi}: TU \rightarrow \phi(U) \times \mathbb{R}^n$ by

$$v \mapsto (x^1(p), \dots, x^n(p), c^1(v), \dots, c^n(v)) \quad (73)$$

$$\iff \quad (74)$$

$$\tilde{\phi}(v) = (\bar{x}^1(v), \dots, \bar{x}^n(v), c^1(v), \dots, c^n(v)) \quad (75)$$

Then $\tilde{\phi}$ has an inverse (for a point $(x^1, \dots, x^n) = \phi(p)$)

$$\tilde{\phi}^{-1}(x^1(p), \dots, x^n(p), c^1(v), \dots, c^n(v)) = \sum c^i \frac{\partial}{\partial x^i} \Big|_p$$

is therefore a bijection. Therefore we use topology of $\phi(U) \times \mathbb{R}^n$ to induce a topology on TU : a set $A \subset TU$ is open iff $\tilde{\phi}(A)$ is open in $\phi(U) \times \mathbb{R}^n$ (where $\phi(U) \times \mathbb{R}^n$ has the standard topology as an open subset of \mathbb{R}^{2n}). With this identification TU becomes homeomorphic to $\phi(U) \times \mathbb{R}^n$.

1.8.2 Manifold Structure on the Tangent Bundle

Next we show that if $\{(U_i, \phi_i)\}$ is a C^∞ atlas for M , then $\{(TU_i, \tilde{\phi}_i)\}$ is a C^∞ atlas for the tangent bundle TM , where $\tilde{\phi}_i$ is the map on TU_i induced by ϕ_i as in (75). It's immediately clear that $TM = \bigcup_i TU_i$; remains to check that on $TU_i \cap TU_j$, $\tilde{\phi}_i, \tilde{\phi}_j$ are C^∞ compatible.

Recall that if $(U, x^1, \dots, x^n), (V, y^1, \dots, y^n)$ are two charts on M , then for any $p \in U \cap V$ there are two bases and so any vector $v \in T_p M$ has two representations

$$v = \sum_j a^j \frac{\partial}{\partial x^j} \Big|_p = \sum_i b^i \frac{\partial}{\partial y^i} \Big|_p$$

Applying v to either of x^k, y^k we get that

$$\begin{aligned} a^k &= \sum_i b^i \frac{\partial x^k}{\partial y^i} \\ b^k &= \sum_j a^j \frac{\partial y^k}{\partial x^j} \end{aligned} \quad (76)$$

Returning to the atlas $\{(U_i, \phi_i)\}$ and let $\phi_\alpha = (x^1, \dots, x^n), \phi_\beta = (y^1, \dots, y^n)$. Then

$$\tilde{\phi}_\beta \circ \tilde{\phi}_\alpha^{-1}: \phi_\alpha(U_\alpha \cap U_\beta) \times \mathbb{R}^n \rightarrow \phi_\beta(U_\alpha \cap U_\beta) \times \mathbb{R}^n$$

is given by (with $\phi_\alpha(p) = (x^1(p), \dots, x^n(p))$)

$$\begin{aligned} (\phi_\alpha(p), a^1(v), \dots, a^n(v)) &\mapsto \left(p, \sum_j a^j \frac{\partial}{\partial x^j} \Big|_p \right) \\ &\mapsto ((\phi_\beta \circ \phi_\alpha^{-1})(\phi_\alpha(p)), b^1, \dots, b^n) \end{aligned} \quad (77)$$

where by (76) and the Jacobian matrix of a transition map

$$\begin{aligned} b^i &= \sum_j a^j \frac{\partial y^i}{\partial x^j} \Big|_p \\ &= \sum_j a^j \frac{\partial(\phi_\beta \circ \phi_\alpha^{-1})}{\partial x^j} \Big|_{\phi_\alpha(p)} \end{aligned} \quad (78)$$

By definition $\phi_\beta \circ \phi_\alpha^{-1}$ is C^∞ and therefore $\tilde{\phi}_\beta \circ \tilde{\phi}_\alpha^{-1}$ is C^∞ and therefore TM is a C^∞ manifold, with $\{(TU_\alpha, \tilde{\phi}_\alpha)\}$ as a C^∞ atlas.

1.8.3 Vector Bundles

On the tangent bundle TM of a smooth manifold M , the natural projection map $\pi: TM \rightarrow M$

$$\pi(p, v) = p$$

makes TM into a C^∞ vector bundle over M which we now define. Given any map $\pi: E \rightarrow M$, where E is called

the *total space* and M is called the *base space*, we call the inverse image $\pi^{-1}(p) := \pi^{-1}(\{p\})$ of a point $p \in M$ the *fiber at p* . The fiber at p is often written E_p . For any two maps

$$\begin{aligned} \pi: E &\rightarrow M \\ \pi': E' &\rightarrow M \end{aligned} \quad (79)$$

with the same target space M , a map $\phi: E \rightarrow E'$ is said to be *fiber-preserving* if $\phi(E_p) \subset E'_p$ for all $p \in M$.

Example 1.15. Fiber-preserving maps Given two maps π, π' , the map ϕ is fiber-preserving iff the diagram

$$\begin{array}{ccc} E & \xrightarrow{\phi} & E' \\ \pi \searrow & & \swarrow \pi' \\ & M & \end{array}$$

commutes.

1.8.4 Smooth Sections

A *section* of a vector bundle $\pi: E \rightarrow M$ is a map $s: M \rightarrow E$ such that $\pi \circ s = \mathbf{1}_M$. This just means that s maps p into the fiber E_p above p (see figure ??).

Definition 1.22: Vector field

A *vector field* X on a manifold M is a function that assigns a tangent vector $X_p \in T_p M$ to each point $p \in M$. In terms of the tangent bundle, a vector field on M is simply a section of the tangent bundle $\pi: TM \rightarrow M$ and the vector field is smooth if it is as smooth as a map from M to TM .

Example 1.16. The formula

$$X_{(x,y)} = \frac{1}{\sqrt{x^2 + y^2}} \left(-y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} \right) = \frac{1}{\sqrt{x^2 + y^2}} \begin{bmatrix} -y \\ x \end{bmatrix}$$

See figure ??.

1.8.5 Smooth Frames

A *frame* for a vector bundle $\pi: E \rightarrow M$ over an open set U is a collection of sections s_1, \dots, s_r of E over U such that at each point $p \in U$, the elements $s_1(p), \dots, s_r(p)$ form a basis for the fiber $E_p := \pi^{-1}(p)$. A frame for the tangent bundle $TM \rightarrow M$ over an open set U is simply called a *frame on U* .

Example 1.17. The collection of vector fields $\partial/\partial x, \partial/\partial y, \partial/\partial z$ is a smooth frame on \mathbb{R}^3 .

Example 1.18. Let M be a manifold and e_1, \dots, e_r be the standard basis for \mathbb{R}^r . Define $\bar{e}_i: M \rightarrow M \times \mathbb{R}^r$ by $\bar{e}_i(p) := (p, e_i)$. Then $\bar{e}_1, \dots, \bar{e}_r$ is a C^∞ frame for the product bundle $M \times \mathbb{R}^r \rightarrow M$.

1.9 Bump Functions and Partitions of Unity

A partition of unity on a manifold is a collection of non-negative functions that sum to 1.

Usually one demands that the partition of unity should be *subordinate* to an open cover $\{U_\alpha\}_{\alpha \in A}$ i.e. the partition $\{\rho_\alpha\}_{\alpha \in A}$ is indexed by the same set A and ρ_α vanishes outside of U_α .

The existence of a C^∞ partition of unity is one of the most important technical tools in the theory of C^∞ manifolds. It is the single feature that makes the behavior of C^∞ manifolds so different from that of real-analytic or complex manifolds. A partition of unity is used two ways:

1. to decompose a global object on a manifold into a locally finite sum of local objects on the open sets U_α of an open cover.
2. to patch together local objects on the open sets U_α into a global object on the manifold.

1.9.1 C^∞ Bump Functions

The *support* of a real-valued function f on a manifold M is defined to be the closure in M of the subset on which $f \neq 0$

$$\text{supp } f = \text{closure } \{q \in M \mid f(q) \neq 0\} \quad (80)$$

Let q be a point in M , and U a neighborhood of q . A *bump function at q supported in U* we mean any continuous nonnegative function ρ on M that is 1 in a neighborhood of q with $\text{supp } \rho \subset U$. See figure ??

Define

$$f(t) = \begin{cases} e^{-1/t} & \text{for } t > 0 \\ 0 & \text{for } t \leq 0 \end{cases} \quad (81)$$

$$g(t) = \frac{f(t)}{f(t) + f(1-t)} \quad (82)$$

$$h(x) = g\left(\frac{x-a^2}{b^2-a^2}\right) \quad (83)$$

$$k(x) = h(x^2) \quad (84)$$

$$\rho(x) = 1 - k(x) \quad (85)$$

See figure ??. It is easy to extend the construction $\rho(x)$ to a bump function from \mathbb{R} to \mathbb{R}^n : to get a C^∞ bump function at $\mathbf{0} \in \mathbb{R}^n$ that is 1 on the closed ball $\bar{B}(\mathbf{0}, a)$ and has support in the closed ball $\bar{B}(\mathbf{0}, b)$

$$\sigma(x) := \rho(|x|) = 1 - g\left(|x| \frac{r^2 - a^2}{b^2 - a^2}\right) \quad (86)$$

In general, a C^∞ on an open subset U of a manifold cannot be extended to a C^∞ function on M ; an example is the function $\sec(x)$ on the open interval $(-\pi/2, \pi/2) \subset \mathbb{R}$. However, if we require that the global function on M agree

with the given function only on some neighborhood of a point in U , then a C^∞ extension is possible.

Proposition 1.20: C^∞ extension of a function

Suppose f is a C^∞ function defined on a neighborhood U of a point p in a manifold M . Then there is a C^∞ function \tilde{f} on M that agrees with f in some possibly smaller neighborhood of p .

Proof. Choose a C^∞ bump function $\rho: M \rightarrow \mathbb{R}$ supported in U that is identically 1 in a neighborhood V of p . Define

$$\tilde{f}(q) = \begin{cases} \rho(q)f(q) & \text{for } q \in U \\ 0 & \text{for } q \notin U \end{cases}$$

As the product of two C^∞ functions on U , \tilde{f} is C^∞ on U . If $q \notin U$, then $q \notin \text{supp } \rho$, and so there is an open set containing q on which \tilde{f} is 0, since $\text{supp } \rho$ is closed. Therefore, \tilde{f} is also C^∞ at every point $q \notin U$. Finally, since $\rho = 1$ on V , the function \tilde{f} agrees with f on V . \square

1.9.2 Partitions of Unity

If $\{U_i\}_{i \in I}$ is a finite open cover of M , a C^∞ *partition of unity subordinate to $\{U_i\}_{i \in I}$* is a collection of nonnegative C^∞ functions $\{\rho_i: M \rightarrow \mathbb{R}\}_{i \in I}$ such that $\text{supp } \rho_i \subset U_i$ and

$$\sum_i \rho_i = 1$$

When the index set I is an infinite set, for the sum to make sense, we impose an additional *local finiteness* condition: a collection $\{A_i\}$ of subsets of a topological space S is said to be *locally finite* if every point $q \in S$ has a neighborhood that meets only finitely many of the sets A_i . In particular, every $q \in S$ is contained in only finitely many of the A_i .

Example 1.19. An open cover that is not locally finite. Let $U_{r,n}$ be the open interval $(r - \frac{1}{n}, r + \frac{1}{n})$. The open cover $\{U_{r,n} \mid r \in \mathbb{Q}, n \in \mathbb{Z}^+\}$ of \mathbb{R} is not locally finite.

Definition 1.23: C^∞ partition of unity

A C^∞ *partition of unity on a manifold* is a collection of nonnegative C^∞ functions $\{\rho_i: M \rightarrow \mathbb{R}\}_{i \in I}$ such that

1. the collection of supports is locally finite
2. $\sum \rho_i = 1$

Suppose $\{f_i\}$ is a collection of C^∞ functions on a manifold M such that the collection of its supports is locally finite. Then every point $q \in M$ has a neighborhood W_q that intersects $\text{supp } f_i$ for only finitely many i . Thus, on W_q the sum $\sum_i f_i$ is a finite sum. This shows that $f = \sum_i f_i$ is well defined and C^∞ on the manifold M . Such a sum is called a *locally finite sum*.

Theorem 1.5: Existence of a C^∞ partition of unity

Let $\{U_i\}$ be an open cover of a manifold M . Then

1. there is a C^∞ partition of unity $\{\phi_k\}_{k=1}^\infty$ having compact support such that for each k , $\text{supp } \phi_k \subset U_i$ for some i .
2. if we do not require compact support, then there is a C^∞ partition of unity $\{\rho_i\}$ subordinate to $\{U_i\}$.

1.10 Vector Fields

A vector field X on a manifold M is the assignment of a tangent vector $X_p \in T_p M$ to each point $p \in M$ (a section of the tangent bundle TM of M). Every smooth vector field may be viewed locally as the velocity vector field of a fluid flow; the path traced out by a point under this flow is called an *integral curve* of the vector field. **Integral curves are curves whose velocity vector field is the restriction of the manifold's vector field, to the curve**; finding an equation of an integral curve is equivalent to solving a system of first-order ODEs. The set $\mathfrak{X}(M)$ of all C^∞ vector fields on a manifold M has the structure of a vector space; the *Lie Bracket* $[\cdot, \cdot]$ makes it into a *Lie algebra*.

1.10.1 Integral Curves

Definition 1.24: Integral Curves

Let X be a C^∞ vector field on a manifold M , and $p \in M$. An *integral curve* $c: (a, b) \rightarrow M$ such that $c'(t) = X_{c(t)}$ for all $t \in (a, b)$. To show the dependence of an integral curve on its initial point p , we write $c_t(p)$.

Example 1.20. Recall the vector field $X_{(x,y)} = \langle -y, x \rangle$ on \mathbb{R}^2 . The condition for $c(t) = (x(t), y(t))$ to be an integral curve is $c'(t) = X_{c(t)}$ or

$$\begin{bmatrix} \dot{x}(t) \\ \dot{y}(t) \end{bmatrix} = \begin{bmatrix} -y(t) \\ x(t) \end{bmatrix}$$

Hence we need to solve the system of first-order ODEs

$$\begin{aligned} \dot{x} &= -y \\ \dot{y} &= x \end{aligned} \quad (87) \quad \text{or}$$

with initial condition $(x(0), y(0)) = (1, 0)$. Making substitutions we get

$$\ddot{x} = -x$$

which has the solution

$$x = A \cos t + B \sin t$$

and therefore

$$y = -\dot{x} = A \sin t - B \cos t$$

The initial condition forces $A = 1, B = 0$ and hence $c(t) = (\cos t, \sin t)$, which parameterizes the unit circle. More generally, if the initial point of the integral curve, corresponding to $t = 0$, is $p = (x_0, y_0)$, then

$$A = x_0 \quad B = -y_0$$

and

$$\begin{aligned} x &= x_0 \cos t - y_0 \sin t \\ y &= x_0 \sin t + y_0 \cos t \end{aligned} \quad (88)$$

which can be written in matrix notation

$$\begin{aligned} c(t) &= \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} \\ &= \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix} \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} \\ &= \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix} p \end{aligned} \quad (89)$$

This shows that the integral curve of X starting at point p can be obtained by rotating point p counterclockwise about the origin through an angle t . Note that

$$c_s(c_t(p)) = c_{s+t}(p)$$

and also that for each $t \in \mathbb{R}$, $c_t: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a diffeomorphism with inverse c_{-t} . Let $\text{operatorname{Diff}}(M)$ be the group of diffeomorphisms of a manifold M to itself, with the group operation being composition. A homomorphism $c: \mathbb{R} \rightarrow \text{Diff}(M)$ is called a *one-parameter group of diffeomorphisms* of M . **In this example the integral curves of the vector field $X_{(x,y)} = \langle -y, x \rangle$ on \mathbb{R}^2 give rise to a one-parameter group of diffeomorphisms of \mathbb{R}^2 .**

1.10.2 Local Flows

In general, if X is a smooth vector field on a manifold M , to find an integral curve $c(t)$ of X starting at p , we first choose a coordinate chart $(U, \phi) = (U, x^1, \dots, x^n)$ about p . In terms of the local coordinates

$$X_{c(t)} = \sum a^i(c(t)) \frac{\partial}{\partial x^i} \Big|_{c(t)}$$

$$c'(t) = \sum \dot{c}^i(t) \frac{\partial}{\partial x^i} \Big|_{c(t)}$$

where $\dot{c}^i(t) = x^i \circ c(t)$ is the i th component of $c(t)$ in the chart (U, ϕ) . The condition $c'(t) = X_{c(t)}$ is equivalent to

$$\dot{c}^i(t) = a^i(c(t))$$

This is a system of ODEs; by *Picard-Lindelöf theorem* such a system has a unique solution in the following sense

Theorem 1.6

Let V be an open subset of \mathbb{R}^n , $p_0 \in V$, and $f: V \rightarrow \mathbb{R}^n$ a C^∞ function. Then the differential equation

$$\frac{dy}{dt} = f(y) \quad y(0) = p_0$$

has a unique C^∞ solution $y: (a(p_0), b(p_0)) \rightarrow V$.

Suppose s, t in the interval $(-\varepsilon, \varepsilon)$ are such that both $F_t(F_s(q))$ and $F_{t+s}(q)$ are defined. Then both $F_t(F_s(q))$ and $F_{t+s}(q)$, as functions of t , are integral curves of X with initial point $F_s(q)$. By uniqueness of the integral curve starting at a point

$$F_t(F_s(q)) = F_{t+s}(q)$$

The map F is called a *local flow generated by the vector field X* . For each $q \in U$, the function $F_t(q)$ of t is called a *flow line* of the local flow. See figure ???. Each flow line is an integral curve of X ; if a local flow F defined on $\mathbb{R} \times M$, then it is called a *global flow* (every smooth vector field has a local flow, but not necessarily a global flow). A vector field having a global flow is called a *complete vector field*. If F is a global flow, then for every $t \in \mathbb{R}$

$$F_t \circ F_{-t} = F_{-t} \circ F_t = F_0 = \mathbf{1}_M$$

so $F_t: M \rightarrow M$ is a diffeomorphism. Thus, a global flow on M gives rise to a one-parameter group of diffeomorphisms on M .

Definition 1.25: Local Flow

A *local flow* about a point p in an open set U of a manifold is a C^∞ function

$$F: (-\varepsilon, \varepsilon) \times W \rightarrow U$$

where $\varepsilon > 0$ and W is a neighborhood of p and $W \subset U$, such that writing $F_t(q) := F(t, q)$, we have

1. $F_0(q) = q$ for all $q \in W$
2. $F_t(F_s(q)) = F_{t+s}(q)$ whenever both sides are defined

If $F_t(q)$ is a local flow of the vector field X on U , then

$$\begin{aligned} F(0, q) &= q \\ \left. \frac{\partial F}{\partial t} \right|_{0, q} &= X_{F(0, q)} \\ &= X_q \end{aligned} \quad (90)$$

Thus, one can recover the vector field from its flow.

Example 1.21. The function $F: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$F\left(t, \begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

is the global flow on \mathbb{R}^2 generated by the vector field

$$\begin{aligned} X_{(x, y)} &= \left. \frac{\partial F(t, (x, y))}{\partial t} \right|_{t=0} \\ &= \begin{bmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \Big|_{t=0} \\ &= \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix} \\ &= -y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} \end{aligned} \quad (91)$$

1.10.3 The Lie Bracket

Suppose X, Y are smooth vector fields on an open subset U of a manifold M . We view X, Y as derivations on the set of C^∞ functions. For a C^∞ function f on U , the function Yf is C^∞ on U , and the function $(XY)f := X(Yf)$ is also C^∞ on U . Moreover, because X, Y are both \mathbb{R} -linear maps from $C^\infty(U)$ to $C^\infty(U)$, the map $XY: C^\infty(U) \rightarrow C^\infty(U)$ is \mathbb{R} -linear. However, XY does not satisfy the derivation property: if $f, g \in C^\infty(U)$, then

$$\begin{aligned} XY(fg) &= X((Yf)g + fYg) \\ &= (XYf)g + (Yf)(Xg) + (Xf)(Yg) + f(XYg) \end{aligned} \quad (92)$$

Looking more closely we see that the two extra terms $(Yf)(Xg)$ and $(Xf)(Yg)$ that make XY not a derivation, are symmetric in X, Y . Thus if we compute $YX(fg)$ and subtract it from $XY(fg)$, the extra terms will disappear, and $XY - YX$ will be a derivation of $C^\infty(U)$.

Definition 1.26: Lie Bracket

Given two smooth vector fields X, Y on U and with $p \in U$, we define their *Lie Bracket* $[X, Y]$ at p

$$[X, Y]_p f = (X_p Y - Y_p X)f \quad (93)$$

for any germ f of a C^∞ function at p .

By the same calculation as above, but now evaluated at p , it is easy to check that $[X, Y]_p$ is a derivation of $C_p^\infty(U)$ and is therefore a tangent vector at p . As p varies over U , $[X, Y]$ becomes a vector field on U .

Proposition 1.21

If X, Y are smooth vector fields on M , then the vector field $[X, Y]$ is also smooth on M .

From this, we see that the Lie bracket provides a product operation on the vector space $\mathfrak{X}(M)$ of all smooth vector fields on M . Clearly

$$[Y, X] = -[X, Y]$$

Example 1.22. Jacobi identity The *Jacobi identity* holds for the Lie bracket

$$\sum_{\text{cyclic}} [X, [Y, Z]] = 0$$

where

$$\sum_{\text{cyclic}} [X, [Y, Z]] = [X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]]$$

Definition 1.27: Lie Algebra

Let K be a field. A *Lie algebra* over K is a vector space V over K together with a product

$$[\cdot, \cdot] \rightarrow V \times V \rightarrow V$$

called the *bracket*, satisfying the following properties: for all $a, b \in K$ and $X, Y, Z \in V$

1. **bilinearity:**

$$\begin{aligned} [aX + bY, Z] &= a[X, Z] + b[Y, Z] \\ [Z, aX + bY] &= a[Z, X] + b[Z, Y] \end{aligned} \quad (94)$$

2. **anticommutativity:**

$$[Y, X] = -[X, Y]$$

3. **Jacobi identity:**

$$\sum_{\text{cyclic}} [X, [Y, Z]] = 0$$

Example 1.23. Abelian Lie algebra On any vector space V , define $[X, Y] = 0$ for all $X, Y \in V$. With this bracket, V becomes a Lie algebra, called an *abelian Lie algebra*.

An abelian Lie algebra is trivially associative, but in general the bracket of a Lie algebra need not be associative. So despite its name, a Lie algebra is in general not an algebra.

Example 1.24. If M is a manifold, then the vector space $\mathfrak{X}(M)$ of C^∞ vector fields on M is a real Lie algebra with the Lie bracket $[\cdot, \cdot]$ as the bracket.

Example 1.25. Let $K^{n \times n}$ be the vector space of all $n \times n$ matrices over a field K . Define

$$[X, Y] = XY - YX$$

where XY is the matrix product of X and Y . With this bracket, $K^{n \times n}$ becomes a Lie algebra. More generally, if A is any algebra over a field K , then the product

$$[x, y] = xy - yx$$

for $x, y \in A$ makes A into a Lie algebra over K .

Definition 1.28: derivation

A *derivation* of a Lie algebra V over a field K is a K -linear map $D: V \rightarrow V$ satisfying the product rule

$$D[Y, Z] = [DY, Z] + [Y, DZ]$$

for $Y, Z \in V$.

Example 1.26. Let V be a Lie algebra over a field K . For each $X \in V$, define

$$\text{ad}_X(Y) = [X, Y]$$

1.10.4 Pushforward of Vector Fields

Let $F: N \rightarrow M$ be a smooth map of manifolds and let $F_*: T_p N \rightarrow T_{F(p)} M$ be its differential at a point p in N . If $X_p \in T_p N$, we call $F_*(X_p)$ the *pushforward* of the vector X_p at p . This notion doesn't extend in general to vector fields, since it X is a vector field on N and

$$z = F(p) = F(q)$$

for two distinct points $p, q \in N$, then X_p and X_q are both pushed forward to tangent vectors $X_z \in T_z M$, but there is no reason why $F_*(X_p)$ and $F_*(X_q)$ should be equal.

1.11 Lie Groups and Lie Algebras

A *Lie group* is a manifold that is also a group such that the group operations are smooth. A Lie group is a homogeneous space in the sense that left translation by a group element g is a diffeomorphism of the group onto itself that maps the identity element to itself. Therefore, locally the group looks the same around any point in the group. To study the local structure of a Lie group, it is enough to examine a neighborhood of the identity element. The tangent space at the identity of a Lie group G turns out to have a canonical bracket operation $[\cdot, \cdot]$ that makes it into a Lie algebra. The tangent space $T_e G$ with the bracket is called the *Lie algebra* of the Lie group G . The Lie algebra of a Lie group encodes within it much information about the group.

The matrix exponential gives rise to curves in a matrix group with a given initial vector. It is useful in computing the differential of a map on a matrix group.

1.11.1 Lie Group

Definition 1.29: Lie group

A *Lie group* is a C^∞ manifold G that is also a group such that the two group operations, multiplication

$$\mu: G \times G \rightarrow G \quad \mu(a, b) := ab$$

and inverse

$$\iota: G \rightarrow G \quad \iota(a) = a^{-1}$$

are C^∞ . For $a \in G$, denote by

$$\begin{aligned} \ell_a: G &\rightarrow G \\ \ell_a(x) &= \mu(a, x) = ax \end{aligned} \quad (95)$$

the operation of *left multiplication by a* and similarly by r_a the operation of *right multiplication by a* . We also call these operations *left and right translations*.

Example 1.27. Left multiplication For an element $a \in G$, prove that the left multiplication/translation $\ell_a: G \rightarrow G$ is a diffeomorphism.

Definition 1.30: Lie group homomorphism

A map $F: H \rightarrow G$ between two Lie groups H, G is a *Lie group homomorphism* if it is a C^∞ map and a group homomorphism. The group homomorphism condition means that for all $h, x \in H$

$$F(hx) = F(h)F(x)$$

or rewritten

$$F \circ \ell_h = \ell_{F(h)} \circ F$$

for all $h \in H$. Note that a group homomorphism always maps the identity to the identity.

Definition 1.31: Orthogonal group

Recall that the orthogonal group $O(n)$ is the subgroup of $GL(n, \mathbb{R})$ consisting of all matrices A satisfying $A^T A = I$. Thus, $O(n)$ is the inverse image of I under the map $f(A) = A^T A$.

1.11.2 Matrix Exponential

Recall computing the pushforward using a smooth curve

$$F_{*,p}(X_p) = \left. \frac{d}{dt} \right|_0 (F \circ c)(t)$$

To compute the differential of a map on a subgroup of

$GL(n, \mathbb{R})$, we need a curve of nonsingular matrices. Because the matrix exponential is always nonsingular, it is uniquely suited for this purpose.

Definition 1.32: Norm

A *norm* on a vector V is a real-valued $|\cdot|: V \rightarrow \mathbb{R}$ satisfying the following three properties for all $r \in \mathbb{R}, v, w \in V$

1. **positive-definiteness:** $|v| \geq 0$ with equality iff $v = 0$
2. **positive homogeneity:** $|rv| = |r||v|$
3. **subadditivity:** $|v + w| \leq |v| + |w|$

A vector space V with a norm is called a *normed vector space*. For example, the vector space $\mathbb{R}^{n \times n} \cong \mathbb{R}^{n^2}$ with the norm

$$|X| = \sqrt{\sum_{ij} x_{ij}^2}$$

Definition 1.33: Matrix exponential

The *matrix exponential* e^X of a matrix $X \in \mathbb{R}^{n \times n}$ is defined

$$e^X = I + X + \frac{1}{2!}X^2 + \frac{1}{3!}X^3 + \dots$$

If A, B commute then

$$e^A e^B = e^{A+B}$$

Proposition 1.22

For $X \in \mathbb{R}^{n \times n}$

$$\frac{d}{dt} e^{tX} = X e^{tX} = e^{tX} X$$

1.11.3 Trace of a Matrix

Definition 1.34: Trace of a Matrix

For a $n \times n$ matrix X

$$\text{tr}(X) = \sum_{i=1}^n x_{ii}$$

i.e. the sum of the diagonal elements.

Lemma 1.2

1. For any two matrices $X, Y \in \mathbb{R}^{n \times n}$

$$\text{tr}(XY) = \text{tr}(YX)$$

2. For $X \in \mathbb{R}^{n \times n}$ and $A \in \text{GL}(n, \mathbb{R})$

$$\text{tr}(AXA^{-1}) = \text{tr}(X)$$

Proposition 1.23

The trace of a matrix, real or complex, is equal to the sum of its complex *eigenvalues*

Proof. Suppose X has complex eigenvalues $\lambda_1, \dots, \lambda_n$. Then there exists a nonsingular matrix $A \in \text{GL}(n, \mathbb{C})$ such that

$$AXA^{-1} = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

and then

$$\text{tr}(X) = \text{tr}(AXA^{-1}) = \sum_i \lambda_i$$

□

Proposition 1.24

For any $X \in \mathbb{R}^{n \times n}$

$$\det(e^X) = e^{\text{tr}(X)}$$

Proof. Case 1: Assume that X is upper triangular

$$X = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

Then

$$\begin{aligned} e^X &= \sum \frac{1}{k!} X^k \\ &= \sum \frac{1}{k!} \begin{bmatrix} \lambda_1^k & & * \\ & \ddots & \\ 0 & & \lambda_n^k \end{bmatrix} \\ &= \begin{bmatrix} e^{\lambda_1} & & * \\ & \ddots & \\ 0 & & e^{\lambda_n} \end{bmatrix} \end{aligned} \quad (96)$$

and hence

$$\det e^X = \prod e^{\lambda_i} = e^{\sum \lambda_i} = e^{\text{tr} X}$$

. Case 2: Given a general matrix X , with eigenvalues $\lambda_1, \dots, \lambda_n$, we can find a nonsingular matrix A such that

$$AXA^{-1} = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

an upper triangular matrix. Then

$$\begin{aligned} e^{AXA^{-1}} &= I + AXA^{-1} + \frac{1}{2!}(AXA^{-1})^2 + \frac{1}{3!}(AXA^{-1})^3 + \dots \\ &= I + AXA^{-1} + A \left(\frac{1}{2!} X^2 \right) A^{-1} + A \left(\frac{1}{3!} X^3 \right) A^{-1} + \dots \\ &= Ae^X A^{-1} \end{aligned} \quad (97)$$

and hence

$$\begin{aligned} \det e^X &= \det(Ae^X A^{-1}) \\ &= \det e^{AXA^{-1}} \\ &= e^{\text{tr}(AXA^{-1})} \\ &= e^{\text{tr} X} \end{aligned} \quad (98)$$

□

It follows that the matrix exponential e^X is always nonsingular, because $\det(e^X) = e^{\text{tr} X} \neq 0$. This is one reason why the matrix exponential is so useful, for it allows us to write down explicitly a curve in $\text{GL}(n, \mathbb{R})$ with a given initial point and a given initial velocity. For example $c(t) = e^{tX} : \mathbb{R} \rightarrow \text{GL}(n, \mathbb{R})$ is a curve in $\text{GL}(n, \mathbb{R})$ with initial point I and initial velocity X , since

$$c(0) = e^{0X} = e^0 = I$$

and

$$c'(0) = \left. \frac{d}{dt} e^{tX} \right|_{t=0} = X e^{tX} \Big|_{t=0} = X$$

Similarly, $c(t) = Ae^{tX}$ is a curve in $\text{GL}(n, \mathbb{R})$ with initial point A and initial velocity AX .

1.11.4 The Differential of det at the identity

Let $\det : \text{GL}(n, \mathbb{R}) \rightarrow \mathbb{R}$ be the determinant map. The tangent space $T_I \text{GL}(n, \mathbb{R})$ at the identity matrix I is the vector space $\mathbb{R}^{n \times n}$ and the tangent space $T_1 \mathbb{R}$ to \mathbb{R} at 1 is \mathbb{R} . So

$$\det_{*,I} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$$

Proposition 1.25

For any $X \in \mathbb{R}^{n \times n}$ it is the case that $\det_{*,I}(X) = \text{tr} X$

Proof. We use a curve at I to compute the differential. As a curve $c(t)$ with $c(0) = I$ and $c'(0) = X$, choose the matrix exponential $c(t) = e^{tX}$. Then

$$\begin{aligned} \det_{*,I}(X) &= \left. \frac{d}{dt} \det(e^{tX}) \right|_{t=0} \\ &= \left. \frac{d}{dt} e^{t \operatorname{tr} X} \right|_{t=0} \\ &= (\operatorname{tr} X) e^{t \operatorname{tr} X} \Big|_{t=0} \\ &= \operatorname{tr} X \end{aligned} \quad (99)$$

□

1.11.5 Tangent Space at the identity of a Lie Group

Left translation $\ell_g: G \rightarrow G$ by an element $g \in G$ is a diffeomorphism with inverse $\ell_{g^{-1}}$. This diffeomorphism takes the identity e to g and induces an isomorphism on tangent spaces

$$(\ell_g)_* := (\ell_g)_{*,g}: T_e G \rightarrow T_g G$$

Thus, if we can describe the tangent space at the identity, then pushing forward we can describe the tangent space at any point in G .

Example 1.28. Tangent space to $\operatorname{GL}(n, \mathbb{R})$ at I The tangent space at any point $g \in \operatorname{GL}(n, \mathbb{R})$ is $\mathbb{R}^{n \times n}$. The isomorphism

$$(\ell_g)_*: T_I(\operatorname{GL}(n, \mathbb{R})) \rightarrow T_g(\operatorname{GL}(n, \mathbb{R}))$$

is also identified as left multiplication by g :

$$g: X \mapsto gX$$

for $X \in G$.

Example 1.29. Tangent space to $\operatorname{SL}(n, \mathbb{R})$ We begin by finding a condition that a tangent vector $X \in T_I(\operatorname{SL}(n, \mathbb{R}))$ must satisfy. Using the curve velocity version of the tangent vector there is a curve $c: (-\varepsilon, \varepsilon) \rightarrow \operatorname{SL}(n, \mathbb{R})$ with $c(0) = I$ and $c'(0) = X$. Being in $\operatorname{SL}(n, \mathbb{R})$, this curve satisfies

$$\det c(t) = 1$$

for all t in $(-\varepsilon, \varepsilon)$. Differentiating both sides and evaluating at $t = 0$

$$\begin{aligned} \left. \frac{d}{dt} \det c(t) \right|_{t=0} &= (\det \circ c)_* \left(\left. \frac{d}{dt} \right|_{t=0} \right) \\ &= \det_{*,I} \left(\left. c_* \frac{d}{dt} \right|_{t=0} \right) \\ &= \det_{*,I}(c'(0)) \\ &= \det_{*,I}(X) \\ &= \operatorname{tr} X \end{aligned} \quad (100)$$

Thus,

$$\operatorname{tr} X = \left. \frac{d}{dt} 1 \right|_{t=0} = 0$$

and so the tangent space $T_I(\operatorname{SL}(n, \mathbb{R}))$ is contained in the subspace V of $\mathbb{R}^{n \times n}$ defined by

$$V = \{X \in \mathbb{R}^{n \times n} \mid \operatorname{tr} X = 0\}$$

Since $\dim V = n^2 - 1 = \dim T_I \operatorname{SL}(n, \mathbb{R})$, the two spaces must be equal.

Example 1.30. The tangent space to $\operatorname{O}(n)$ at I Let X be a tangent vector to the orthogonal group $\operatorname{O}(n)$ at the identity I . Choose a curve $c(t) \in \operatorname{O}(n)$ such that $c(0) = I$, $c'(0) = X$. Since $c(t) \in \operatorname{O}(n)$

$$c(t)^T c(t) = I$$

Differentiating both sides

$$c'(t)^T c(t) + c(t)^T c'(t) = 0$$

Evaluating at $t = 0$ we get

$$X^T + X = 0$$

Thus, X is skew-symmetric. Let K_n be the space of all $n \times n$ real skew-symmetric matrices; the diagonal entries of such matrices are all 0 and entries below the diagonal are determined by those above the diagonal. Therefore

$$\dim K_n = \frac{n^2 - \# \text{ diagonal entries}}{2} = \frac{1}{2}(n^2 - n)$$

Therefore

$$T_I(\operatorname{O}(n)) \subset K_n$$

By an earlier computation

$$\dim T_I(\operatorname{O}(n)) = \dim \operatorname{O}(n) = \frac{n^2 - n}{2}$$

and therefore $T_I(\operatorname{O}(n)) = K_n$

1.11.6 Left-Invariant Vector Fields on a Lie Group

We say that the vector field X is *left-invariant* if

$$(\ell_g)_* X = X$$

for every $g \in G$; this means that for any $h \in G$

$$(\ell_g)_*(X_h) = X_{gh}$$

i.e. the vector X_h (anchored at h) pushes forward (under the left-multiplication by g map ℓ_g) to X_{gh} .

A left-invariant vector field X is completely determined by its value X_e at the identity, since

$$X_g = (\ell_g)_*(X_e)$$

Conversely, given a tangent vector $A \in T_e(G)$ we can define a vector field \tilde{A} on G

$$\tilde{A}_g := (\ell_g)_* A$$

So defined \tilde{A} is left-invariant

$$\begin{aligned} (\ell_g)_*(\tilde{A}_h) &= (\ell_g)_*(\ell_h)_* A \\ &= (\ell_g \circ \ell_h)_* A \\ &= (\ell_{gh})_* A \\ &= \tilde{A}_{gh} \end{aligned} \quad (101)$$

\tilde{A} is called the *left-invariant vector field on G generated by $A \in T_e G$* . Let $L(G)$ be the vector space of all left-invariant vector fields on G . Then there is a one-to-one correspondence

$$\begin{aligned} T_e(G) &\leftrightarrow L(G) \\ X_e &\leftrightarrow X \\ A &\mapsto \tilde{A} \end{aligned} \quad (102)$$

Example 1.31. Left-invariant vector fields on $\mathrm{GL}(n, \mathbb{R})$. Since $\mathrm{GL}(n, \mathbb{R})$ is an open subset of $\mathbb{R}^{n \times n}$, at any $g \in \mathrm{GL}(n, \mathbb{R})$ there is a canonical identification of the tangent space $T_g(\mathrm{GL}(n, \mathbb{R}))$ with $\mathbb{R}^{n \times n}$, under which a tangent vector corresponds to an $n \times n$ matrix

$$\sum a_{ij} \frac{\partial}{\partial x_{ij}} \Big|_g \leftrightarrow [a_{ij}]$$

We use the same letter B for the tangent vector and the matrix. Let

$$\sum b_{ij} \frac{\partial}{\partial x_{ij}} \Big|_I \in T_I(\mathrm{GL}(n, \mathbb{R}))$$

and let \tilde{B} be the left-invariant vector field on $\mathrm{GL}(n, \mathbb{R})$ generated by B . Then

$$\tilde{B}_g = (\ell_g)_* B \leftrightarrow gB$$

and

$$\begin{aligned} \tilde{B}_g &= \sum_{i,j} (gB)_{ij} \frac{\partial}{\partial x_{ij}} \Big|_g \\ &= \sum_{i,j} \left(\sum_k g_{ik} b_{kj} \right) \frac{\partial}{\partial x_{ij}} \Big|_g \end{aligned} \quad (103)$$

Proposition 1.26

If X, Y are left-invariant vector fields on G , then so is $[X, Y]$.

1.11.7 The Lie Algebra of a Lie Group

The isomorphism $\phi: T_e G \cong L(G)$ (the set of left-invariant vector spaces on G) enables us to define a Lie bracket on $T_e G$ and to push forward left-invariant vector fields under a Lie group homomorphism. We begin with the Lie bracket on $T_e G$. Given $A, B \in T_e G$, we first map them via ϕ to the left-invariant vector fields \tilde{A}, \tilde{B} , take the Lie bracket

$$[\tilde{A}, \tilde{B}] = \tilde{A}\tilde{B} - \tilde{B}\tilde{A}$$

and then map it back to $T_e G$ via ϕ^{-1} . Thus, the definition of the Lie bracket $[A, B] \in T_e G$ should be

$$[A, B] = [\tilde{A}, \tilde{B}]_e$$

Proposition 1.27

If $A, B \in T_e G$ and \tilde{A}, \tilde{B} are the left-invariant vector fields they generate, then

$$[\tilde{A}, \tilde{B}] = [A, B]^\sim$$

With the Lie bracket $[\cdot, \cdot]$, the tangent space $T_e G$ becomes a Lie algebra, called the *Lie algebra of the Lie group G* . As a Lie algebra, $T_e G$ is usually denoted \mathfrak{g} .

1.12 Differential Forms

Differential forms play a crucial role in manifold theory. First and foremost, they are intrinsic objects associated to any manifold, and so can be used to construct diffeomorphism invariants. Differential forms have a richer algebraic structure; due to the existence of the wedge product, a grading, and the exterior derivative, the set of smooth forms on a manifold is both a graded algebra and a differential complex. Such an algebraic structure is called a *differential graded algebra*. Moreover, the differential complex of smooth forms on a manifold can be pulled back under a smooth map, making the complex *contravariant functor* called the *de Rham complex* of the manifold. We will eventually construct the de Rham cohomology of a manifold from the de Rham complex.

1.12.1 Differential 1-Forms

Let M be a smooth manifold and $p \in M$. The *cotangent space* of M at p , denoted by $T_p^* M$, is defined to be the dual space of the tangent space $T_p M$

$$T_p^* M = (T_p M)^\vee = \mathrm{Hom}(T_p M, \mathbb{R})$$

An element of the cotangent space $T_p^* M$ is called a *covector* at p . Thus, a covector $\omega_p: T_p M \rightarrow \mathbb{R}$. A *covector field*, a *differential 1-form* is a function ω that assigns to each point $p \in M$ a covector ω_p at p .

Definition 1.35: Differential

If f is a C^∞ real-valued function on a manifold M , its *differential* is defined to be the 1-form df on M such that for any $p \in M$ and $X_p \in T_p M$

$$(df)_p(X_p) = X_p f$$

1.12.2 Local Expression for a Differential 1-Form

Let $(U, \phi) = (U, x^1, \dots, x^n)$ be a coordinate chart on M . Then the differentials dx^1, \dots, dx^n are 1-forms on U .

Proposition 1.28

At each point $p \in U$, the covectors dx^1, \dots, dx^n form a basis for the cotangent space $T_p^* M$, dual to the basis

$$\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n}$$

of the tangent space $T_p M$.

1.12.3 The Cotangent Bundle

The *cotangent bundle* T^*M of a manifold M is the union of the cotangent spaces at all points of M

$$T^*M := \bigcup_{p \in M} T_p^* M$$

There is a natural map $\pi: T^*M \rightarrow M$ given by $\pi(\alpha) = p$ if $\alpha \in T_p^* M$. Properly speaking, the *cotangent bundle* of a manifold M is the triple (T^*M, M, ϕ) , while T^*M is the *total space* and M is the *base space* of the cotangent bundle, but by abuse of language it is customary to call T^*M the cotangent bundle of M . In terms of the cotangent bundle, a 1-form on M is simply a section of the cotangent bundle T^*M i.e. it is a map $\omega: M \rightarrow T^*M$ such that $\pi \circ \omega = 1_M$

Example 1.32. Liouville form on the cotangent bundle If a manifold M has dimension n , then the total space T^*M of its cotangent bundle $\pi: T^*M \rightarrow M$ is a manifold of dimension $2n$ (chart and coefficients). Remarkably, on T^*M there is a 1-form λ , called the *Liouville form*, or *Poincare form*, defined independently of charts as follows. A point in T^*M is a covector $\omega_p \in T_p^* M$ at some point $p \in M$. If X_{ω_p} is a tangent vector to T^*M at ω_p , then the pushforward $\pi_*(X_{\omega_p})$ is a tangent vector to M at p . Therefore, one can pair up ω_p and $\pi_*(X_{\omega_p})$ to obtain a real number $\omega_p(\pi_*(X_{\omega_p}))$

$$\lambda_{\omega_p}(X_{\omega_p}) = \omega_p(\pi_*(X_{\omega_p}))$$

The cotangent bundle and the Liouville form on it play an important role in classical mathematics.

1.12.4 Pullback of 1-Forms

If $F: N \rightarrow M$ is a C^∞ map of manifolds, then at each point $p \in N$ the differential

$$F_{*,p}: T_p N \rightarrow T_{F(p)} M$$

is a linear map that pushes forward vectors at p from N to M . The *codifferential* i.e. the dual of the differential

$$F^* := (F_{*,p})^\vee: T_{F(p)}^* M \rightarrow T_p^* N$$

pulls back a covector at $F(p)$ from M to N . By the definition of the dual, if $\omega_{F(p)} \in T_{F(p)}^* M$ is a covector at $F(p)$ and $X_p \in T_p N$ then

$$F^*(\omega_{F(p)})(X_p) = \omega_{F(p)}(F_{*,p} X_p)$$

We call $F^*(\omega_{F(p)})$ is the *pullback* of the covector $\omega_{F(p)}$ by F .

Unlike vector fields, which in general cannot be pushed forward under a C^∞ map, every covector field can be pulled back by a C^∞ map. If ω is a 1-form on M , its *pullback* $F^*\omega$ is the 1-form on N defined pointwise by

$$(F^*\omega)_p := F^*(\omega_{F(p)})$$

This means that

$$(F^*\omega)_p(X_p) = \omega_{F(p)}(F_{*,p}(X_p))$$

i.e. pushforward the vector to M and then apply ω to it.

Proposition 1.29: Pullback rules

et $F: N \rightarrow M$ be a C^∞ map of manifolds. For any $g, h \in C^\infty(M)$, $\omega, \tau \in \Omega^1(M)$

1. $F^*(dh) = d(F^*h)$
2. $F^*(\omega + \tau) = F^*\omega + F^*\tau$
3. $F^*(g\omega) = (F^*g)(F^*\omega)$

Example 1.33. Liouville form on the cotangent bundle Let M be a manifold. In terms of the pullback, the Liouville form λ on the cotangent bundle T^*M can be expressed as

$$\lambda_{\omega_p} = \pi^*(\omega_p)$$

1.12.5 Differential k -forms

We now generalize from 1-forms on a manifold to k -forms. In parallel with the construction of the tangent and cotangent bundles on a manifold, we construct the k th exterior power $\bigwedge^k T^*M$ of the cotangent bundle; a differential k -form is seen as a section of this bundle. The pullback and the wedge product of differential forms are defined pointwise.

Recall that a k -tensor on a vector space V is a k -linear function

$$f: \underbrace{V \times \dots \times V}_k \rightarrow \mathbb{R}$$

f is alternating if for any permutation $\sigma \in S_k$

$$f(v_{\sigma(1)}, \dots, v_{\sigma(k)}) = (\text{sgn}(\sigma))f(v_1, \dots, v_k)$$

An alternating k -tensor on V is also called a k -covector on V .

Definition 1.36: $A_k(V)$

For any vector space V , denote by $A_k(V)$ the vector space of alternating k -tensors on V . Another common notation is $\bigwedge^k(V^\vee)$, where V^\vee is the dual space to V . Thus,

$$\begin{aligned} \bigwedge^0(V^\vee) &\equiv A_0(V) = \mathbb{R} \\ \bigwedge^1(V^\vee) &\equiv A_1(V) = V^\vee \\ \bigwedge^2(V^\vee) &\equiv A_2(V) = V^\vee \end{aligned} \quad (104)$$

where we use the fact that all 1-tensors are alternating and so $A_1(V) \equiv V^\vee$. In fact, there is a purely algebraic construction $\bigwedge^k(V^\vee)$, called the *exterior power* of the vector space V , with the property that $\bigwedge^k(V^\vee) \cong A_k(V)$ but we take this for granted.

$A_k()$ is a *functor*; applying it to $T_p M$ we get the vector space $A_k(T_p M)$, usually denoted $\bigwedge^k(T_p^* M)$, the space of all alternating k -tensors on the tangent space $T_p M$. A k -covector field on M is a function ω that assigns at each point $p \in M$ a k -covector $\omega_p \in \bigwedge^k(T_p^* M)$. A k -covector field is also called a *differential k -form*, *differential form of degree k* , or simply a k -form. A *top form* on a manifold is a differential form whose degree is the dimension of the manifold.

If $\omega \in \bigwedge^k(T_p^* M)$ on a manifold M and X_1, \dots, X_k are vector fields on M , then $\omega(X_1, \dots, X_k)$ is the function on M defined

$$(\omega(X_1, \dots, X_k))(p) := \omega_p((X_1)_p, \dots, (X_k)_p)$$

Proposition 1.30: Multilinearity of a form over functions

Let ω be a k -form on a manifold M . For any vector fields X_1, \dots, X_k and any function h on M

$$\omega(X_1, \dots, hX_i, \dots, X_k) = h\omega(X_1, \dots, X_i, \dots, X_k)$$

Example 1.34. Let (U, x^1, \dots, x^n) be a coordinate chart on a manifold. At each point $p \in U$, a basis for the tangent space $T_p U$ is

$$\left. \frac{\partial}{\partial x^1} \right|_p, \dots, \left. \frac{\partial}{\partial x^n} \right|_p$$

A dual basis for the cotangent space is $T_p^* U$ is

$$(dx^1)_p, \dots, (dx^n)_p$$

and a basis for $\bigwedge^k(T_p^* U)$ is

$$(dx^{i_1})_p \wedge \dots \wedge (dx^{i_k})_p$$

If ω is a k -form on \mathbb{R}^n , then at each point $p \in \mathbb{R}^n$, ω_p is a linear combination

$$\omega_p = \sum a_{i_1 \dots i_k}(p) (dx^{i_1})_p \wedge \dots \wedge (dx^{i_k})_p$$

Omitting the point p we write

$$\omega_p = \sum a_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

where $a_{i_1 \dots i_k}$ are functions on U because they vary with the point p . To simplify the notation, let

$$\mathcal{J}_{k,n} := \{I = (i_1, \dots, i_k) | 1 \leq i_1 < \dots < i_k \leq n\}$$

be the set of all strictly ascending multi-indices between 1 and n of length k . Then

$$\omega \equiv \sum_{I \in \mathcal{J}_{k,n}} a_I dx^I$$

where

$$dx^I := dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

Note that for U and $I, J \in \mathcal{J}_{k,n}$

$$dx^I(\partial_J) = \delta_J^I := \begin{cases} 1 & \text{for } I = J \\ 0 & \text{for } I \neq J \end{cases}$$

where

$$\partial_J := \left(\frac{\partial}{\partial x^{j_1}}, \dots, \frac{\partial}{\partial x^{j_k}} \right)$$

Proposition 1.31: A wedge of differentials in local coordinates

Let U, x^1, \dots, x^n be a chart on a manifold and f^1, \dots, f^k be smooth functions on U . Then

$$df^1 \wedge \dots \wedge df^k = \sum_{I \in \mathcal{J}_{k,n}} \frac{\partial(f^1, \dots, f^k)}{\partial(x^{i_1}, \dots, x^{i_k})} dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

where

$$\frac{\partial(f^1, \dots, f^k)}{\partial(x^{i_1}, \dots, x^{i_k})} := \det \left[\frac{\partial f^i}{\partial x^{i_j}} \right]$$

Corollary 1.2: Transition differential

$$dy^I = \sum_J \frac{\partial(y^{i_1}, \dots, y^{i_k})}{\partial(x^{j_1}, \dots, x^{j_k})} dx^J$$

In particular

1.

$$df = \sum \frac{\partial f}{\partial x^i} dx^i$$

2.

$$df^1 \wedge \dots \wedge df^n = \det \left[\frac{\partial f^j}{\partial x^i} \right] dx^1 \wedge \dots \wedge dx^n$$

1.12.6 The Bundle Point of View

Let M be a manifold of dimension n . Then

$$\bigwedge^k(T^*M) := \bigcup_{p \in U} \bigwedge^k(T_p^*M) = \bigcup_{p \in M} A_k(T_pM)$$

where $A_k(T_pM)$ is the vector space of alternating k -tensors. $\bigwedge^k(T^*M)$ is called the k -th exterior power of the cotangent bundle; there is a projection map $\pi: \bigwedge^k(T^*M) \rightarrow M$ given by

$$\pi(\alpha) = p$$

if $\alpha \in \bigwedge^k(T_p^*M)$.

If U, ϕ is a coordinate chart on M , then there is a **bijection**

$$\begin{aligned} \bigwedge^k(T^*U) &:= \bigcup_{p \in U} \bigwedge^k(T_p^*U) \cong \phi(U) \times \mathbb{R}^{\binom{n}{k}} \\ \alpha &\in \bigwedge^k(T_p^*U) \mapsto (\phi(p), c_I(\alpha)) \end{aligned} \quad (105)$$

where $\alpha = \sum c_I(\alpha) dx^I \in \bigwedge^k(T_p^*U)$. This bijection induces a topology and differentiable structure on $\bigwedge^k(T^*M)$. This structure then turns the projection map

$$\pi: \bigwedge^k(T^*M) \rightarrow M$$

into a C^∞ vector bundle of rank $\binom{n}{k}$ and a differential k -form into, simply, a section of this bundle.

Note that if $E \rightarrow M$ is a C^∞ vector bundle, then the vector space of C^∞ sections of E is denoted $\Gamma(E)$ or $\Gamma(M, E)$. The vector spaces of all C^∞ k -forms on M is usually denoted by $\Omega^k(M)$. Thus,

$$\Omega^k(M) \equiv \Gamma(\bigwedge^k(T^*M)) \equiv \Gamma(M, \bigwedge^k(T^*M))$$

1.12.7 Pullback of k -forms

For a C^∞ 0-form on M under a C^∞ map $F: N \rightarrow M$ (i.e. a C^∞ function $f: M \rightarrow \mathbb{R}$) the pullback F^*f is simply the composition

$$F^*f = f \circ F \in \Omega^0(N)$$

Generalizing to k -forms, we first recall the pullback of k -covectors: a linear map $L: V \rightarrow W$ of vector spaces induces a pullback map $L^*: A_k(W) \rightarrow A_k(V)$ by

$$(L^*\alpha)(v_1, \dots, v_k) = \alpha(L(v_1), \dots, L(v_k))$$

for $\alpha \in A_k(W)$ and $v_1, \dots, v_k \in V$. Now suppose $F: N \rightarrow M$; at each point the differential (pushforward)

$$F_{*,p}: T_pN \rightarrow T_{F(p)}M$$

is a linear map of tangent spaces and so there is a corresponding pullback map on k -forms:

$$(F_{*,p})^*: A_k(T_{F(p)}M) \rightarrow A_k(T_pN)$$

Thus, if $\omega_{F(p)}$ is a k -form at $F(p)$ in M , then its *pullback* $F^*(\omega_{F(p)})$ is the k -form

$$(F^*(\omega_{F(p)}))(v_1, \dots, v_k) := \omega_{F(p)}(F_{*,p}v_1, \dots, F_{*,p}v_k)$$

for $v_i \in T_pN$.

Proposition 1.32: Linearity of the pullback

Let $F: N \rightarrow M$ be a C^∞ map. If ω, τ are k -forms on M and a is a real number, then

$$1. F^*(\omega + \tau) = F^*\omega + F^*\tau$$

$$2. F^*(a\omega) = aF^*\omega$$

1.12.8 The Wedge Product

If α, β are alternating tensors of degree k, ℓ on a vector space V , then their wedge product $\alpha \wedge \beta$ is the alternating $(k + \ell)$ -tensor on V defined

$$\begin{aligned} (\alpha \wedge \beta)(v_1, \dots, v_{k+\ell}) &= \\ \sum \text{sgn}(\sigma) \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \beta(v_{\sigma(k+1)}, \dots, v_{\sigma(k+\ell)}) \end{aligned} \quad (106)$$

where $v_i \in V$ and σ runs over all (k, ℓ) -shuffles of $1, \dots, k + \ell$. For example if α, β are 1-covectors, then

$$(\alpha \wedge \beta)(v_1, v_2) = \alpha(v_1)\beta(v_2) - \alpha(v_2)\beta(v_1)$$

For a k -form ω and an ℓ -form τ on M define their *wedge product* $\omega \wedge \tau$ to be the $k + \ell$ -form on M

$$(\omega \wedge \tau)_p := \omega_p \wedge \tau_p$$

Proposition 1.33: Pullback of a wedge product

If $F: N \rightarrow M$ is a C^∞ map of manifolds and ω, τ are differential forms on M , then

$$F^*(\omega \wedge \tau) = F^*\omega \wedge F^*\tau$$

Definition 1.37: Differential Graded Algebra

Define the vector space $\Omega^*(M)$ of C^∞ differential forms on a manifold M of dimension n to be the direct sum

$$\Omega^*(M) := \bigoplus_{k=0}^n \Omega^k(M)$$

which means that each element in $\Omega^*(M)$ is uniquely a sum $\sum_{k=0}^n \omega_k$, where $\omega_k \in \Omega^k(M)$. With the wedge product on forms, $\Omega^*(M)$ becomes a graded algebra, the grading being the degree of differential forms.

1.13 The Exterior Derivative

Recall that an *antiderivation* on a graded algebra $A = \bigoplus_{k=0}^\infty A^k$ is an \mathbb{R} -linear map $D: A \rightarrow A$ such that

$$D(\omega \cdot \tau) = (D\omega) \cdot \tau + (-1)^k \omega \cdot (D\tau)$$

where $\omega \in A^k$ and $\tau \in A^\ell$. In the graded algebra A is called a *homogeneous element* of degree k if only comes from A^k . An *antiderivation of degree m* is such that

$$\deg D\omega = \deg \omega + m$$

On the graded algebra $\Omega^*(M)$ there is a uniquely and intrinsically defined antiderivation called the *exterior derivative*, the application of which is called *exterior differentiation*

Definition 1.38: Exterior derivative

An *exterior derivative* on a manifold M is an \mathbb{R} -linear map

$$D: \Omega^*(M) \rightarrow \Omega^*(M)$$

such that

1. D is an antiderivation of degree 1
2. $D \circ D = 0$
3. if f is a C^∞ function and X a C^∞ vector field on M , then

$$(Df)(X) = Xf$$

Condition 3 says that on 0-forms an exterior derivative agrees with the differential df .

1.13.1 Exterior Derivative on a Coordinate Chart

Let U, x^1, \dots, x^n be a coordinate chart on a manifold M . Then for any k -form ω on U , the exterior derivative D is

$$\begin{aligned} D\omega &= \sum_I (Da_I) \wedge dx^I + \sum_I a_I (D dx^I) \\ &= \sum_I (Da_I) \wedge dx^I \\ &= \sum_I \sum_j \frac{\partial a_I}{\partial x^j} dx^j \wedge dx^I \end{aligned} \quad (107)$$

Like the derivative of a function on \mathbb{R}^n , an antiderivation D on $\Omega^*(M)$ has the property that for a k -form ω , the value of $D\omega$ at a point p depends only on the values of ω in a neighborhood of p . To make this precise we define *local operators*.

An *endomorphism* of a vector space W is often called an *operator* on W . For example if $W = C^\infty(\mathbb{R})$ is the vector space of C^∞ functions on \mathbb{R} , then the derivative $\frac{d}{dx}$ is an operator on W :

$$\frac{d}{dx} f(x) = f'(x)$$

The derivative has the property that the value of $f'(x)$ at a point p depends only on the values of f in a small neighborhood of p ; i.e. if $f = g$ on an open set $U \subset \mathbb{R}$, then $f' = g'$ on U . In this sense the derivative is a *local operator* on $C^\infty(\mathbb{R})$.

Definition 1.39: Local operator

An operator $D: \Omega^*(M) \rightarrow \Omega^*(M)$ is said to be *local* if for all $k \geq 0$, whenever a k -form $\omega \in \Omega^k(M)$ restricts to 0 on an open set U in M , then $D\omega \equiv 0$ on U . Equivalently D is local if for all $k \geq 0$, whenever two k -forms $\omega, \tau \in \Omega^k(M)$ agree on an open set U , then $D\omega \equiv D\tau$ on U .

Example 1.35. Integral operator Define the integral operator

$$I: C^\infty([a, b]) \rightarrow C^\infty([a, b])$$

by

$$I(f) := \int_a^b f(t) dt$$

I is not a local operator, since the value of $I(f)$ at any point p depends on the values of f over the entire interval $[a, b]$.

Proposition 1.34

Any antiderivation D on $\Omega^*(M)$ is a local operator.

Proof. Suppose $\omega \in \Omega^k(M)$ and $\omega \equiv 0$ on an open subset U . It suffices to prove that $(D\omega)_p = 0$. Choose a C^∞ bump function f at p supported in U . In particular, $f \equiv 1$ in a neighborhood of $p \in U$. Then $f\omega \equiv 0$ on M (since $q \in U$, then $\omega_q \equiv 0$ and if $q \notin U$ then $f(q) = 0$). Applying the antiderivation property of D to $f\omega$

$$0 = D(0) = D(f\omega) = (Df) \wedge \omega + (-1)^0 f \wedge (D\omega)$$

Evaluating the right-hand side at p , noting that $\omega_p = 0$ and $f(p) = 1$, gives $0 = (D\omega)_p$. \square

1.13.2 Existence of an Exterior Derivative on a Manifold

Define on a chart U, x^1, \dots, x^n

$$d_U \omega = \sum da_I \wedge dx^I$$

Then define

$$(d\omega)_p := (d_U \omega)_p$$

and we can prove that $(d\omega)_p$ is independent of the chart (using transition functions). As p varies over all points of M , this defines an operator

$$d: \Omega^*(M) \rightarrow \Omega^*(M)$$

1.13.3 Exterior Differentiation Under a Pullback

The pullback of differential forms commutes with the exterior derivative. This fact, and the preservation of the wedge product by the pullback, is a cornerstone of a calculus involving the pullback.

Proposition 1.35: Commutation of the pullback with d

Let $F: N \rightarrow M$ be a smooth map of manifolds. If $\omega \in \Omega^k(M)$, then $dF^*\omega = F^*d\omega$.

Corollary 1.3

If U is an open subset of a manifold M and $\omega \in \Omega^k(M)$, then

$$d\omega|_U = d(\omega|_U)$$

Example 1.36. Let U be the open set $(0, \infty) \times (0, 2\pi)$ in the (r, θ) -plane \mathbb{R}^2 . Define $F: U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$F(r, \theta) = (r \cos \theta, r \sin \theta)$$

If (x, y) are the standard coordinates on the target \mathbb{R}^2 , compute the pullback $F^*(dx \wedge dy)$:

$$\begin{aligned} F^* &= dF^*x \\ &= d(x \circ F) \\ &= d(r \cos \theta) \\ &= (\cos \theta) dr - r \sin \theta d\theta \end{aligned} \tag{108}$$

and similarly

$$F^* dy = (\sin \theta) dr + r \cos \theta d\theta$$

Then since the pullback commutes with the wedge product

$$\begin{aligned} F^*(dx \wedge dy) &= (F^* dx) \wedge (F^* dy) \\ &= [(\cos \theta) dr - r \sin \theta d\theta] \wedge [(\sin \theta) dr + r \cos \theta d\theta] \\ &= (r \cos^2 \theta + r \sin^2 \theta) dr \wedge d\theta \\ &= r dr \wedge d\theta \end{aligned} \tag{109}$$

Proposition 1.36

If $F: N \rightarrow M$ is a C^∞ map of manifolds and ω is a C^∞ k -form on M , then $F^*\omega$ is a C^∞ k -form on N .

In summary, if $F: N \rightarrow M$ is a C^∞ map of manifolds, then the pullback map $F^*: \Omega^*(M) \rightarrow \Omega^*(N)$ is a *morphism of differential graded algebras*, i.e. a degree-preserving algebra homomorphism that commutes with the differential.

1.14 The Lie Derivative and Interior Multiplication

1.14.1 The Lie Derivative of a Vector Field

The problem with generalizing the derivative

$$f'(p) = \lim_{t \rightarrow 0} \frac{f(p+t) - f(p)}{t}$$

to the derivative of a vector field Y on a manifold M is that at two nearby points p, q the tangent vectors Y_p, Y_q are in different vector spaces $T_p M, T_q M$ (and so it is not possible to compare them by subtracting one from the other). **One way to get around this difficulty is to use the local flow of another vector field X to transport Y_q to the tangent space $T_p M$ at p .** This leads to the definition of the *Lie derivative of a vector field*.

Recall that for any smooth vector field X on M and point $p \in M$, there is a neighborhood U of p on which the vector field has a *local flow* (integrate the ODE in the neighborhood);

$$\phi: (-\varepsilon, \varepsilon) \times U \rightarrow M$$

such that $\phi_t(q) := \phi(t, q)$. Then

$$\begin{aligned} \frac{\partial}{\partial t} \phi_t(q) &= X_{\phi_t(q)} \\ \phi_0(q) &= q \end{aligned} \tag{110}$$

In other words, for each $q \in U$, the curve $\phi_t(q)$ is an integral curve of X with initial point q . The local flow satisfies the property

$$\phi_s \circ \phi_t = \phi_{s+t}$$




Figure 1: Comparing the values of Y at nearby points.

Consequently, for each t the map $\phi_t: U \rightarrow \phi_t(U)$ is a diffeomorphism onto its image, with a C^∞ inverse ϕ_{-t} :

$$\begin{aligned}\phi_{-t} \circ \phi_t &= \phi_0 = 1 \\ \phi_t \circ \phi_{-t} &= \text{id}_U = 1\end{aligned}\quad (111)$$

Let Y be a C^∞ vector field on M . To compare the values of Y at $\phi_t(p)$ and at p , we use the diffeomorphism

$$\phi_{-t}: \phi_t(U) \rightarrow U$$

to push $Y_{\phi_t(p)}$ back to $T_p M$.

Definition 1.40: Lie derivative

For $X, Y \in \mathfrak{X}(M)$ (vector space of vector fields on M) and $p \in M$, let

$$\phi: (-\varepsilon, \varepsilon) \times U \rightarrow M$$

be a local flow of X on a neighborhood U of p . Define the *Lie derivative* $\mathcal{L}_X Y$ of Y with respect to X at p to be the vector

$$\begin{aligned}(\mathcal{L}_X Y)_p &:= \lim_{t \rightarrow 0} \frac{(\phi_{-t})_*(Y_{\phi_t(p)}) - Y_p}{t} \\ &= \lim_{t \rightarrow 0} \frac{((\phi_{-t})^* Y)_p - Y_p}{t} \\ &= \left. \frac{d}{dt} \right|_{t=0} ((\phi_{-t})^* Y)_p\end{aligned}\quad (112)$$

where $(\phi_{-t})_*(Y_{\phi_t(p)})$ is the pushforward of $Y_{\phi_t(p)}$ by ϕ_{-t} i.e. transport $Y_{\phi_t(p)}$ back to p by ϕ_{-t} . See figure 1.

In this definition the limit is taken in the finite dimensional vector space $T_p M$. For the derivative to exist, it suffices that $\{(\phi_{-t})^* Y\}$ be a smooth family of vector fields on M .

Let ϕ_t^i, ϕ^i be the i th components of ϕ_t and ϕ respectively. Then

$$\phi_t^i(p) = \phi^i(t, p) = (x^i \circ \phi)(t, p)$$

Relative to the frame

$$\left\{ \frac{\partial}{\partial x^j} \right\}$$

the differential $\phi_{*,t}$ at p is represented by the Jacobian matrix

$$\left[\frac{\partial \phi_t^i}{\partial x^j} \right]_p = \left[\frac{\partial \phi^i}{\partial x^j} \right]_{t,p}$$

This means that

$$\phi_{t,*} \left(\frac{\partial}{\partial x^j} \right)_p = \sum_i \frac{\partial \phi^i}{\partial x^j} \bigg|_{t,p} \frac{\partial}{\partial x^i} \bigg|_{\phi_t(p)}$$

Thus, if $Y = \sum b^j \partial / \partial x^j$, then

$$\begin{aligned}(\phi_{-t})_*(Y_{\phi_t(p)}) &= \sum_j b^j \phi_t(p) (\phi_{-t})_* \left(\frac{\partial}{\partial x^j} \right)_{\phi_t(p)} \\ &= \sum_{i,j} b^j \phi_t(p) \frac{\partial \phi^i}{\partial x^j} \bigg|_{-t,p} \frac{\partial}{\partial x^i} \bigg|_p\end{aligned}\quad (113)$$

When X, Y are C^∞ vector fields on M , both ϕ^i, b^j are C^∞ functions. The above shows that $\{(\phi_{-t})^* Y\}$ is a smooth family of vector fields on M . It follows that the Lie derivative $\mathcal{L}_X Y$ exists and is given in local coordinates by

$$\begin{aligned}(\mathcal{L}_X Y)_p &= \left. \frac{d}{dt} \right|_{t=0} (\phi_{-t})_*(Y_{\phi_t(p)}) \\ &= \sum_{i,j} \left. \frac{\partial}{\partial t} \right|_{t=0} \left(b^j(\phi_t(p)) \frac{\partial \phi^i}{\partial x^j} \bigg|_{-t,p} \right) \frac{\partial}{\partial x^i} \bigg|_p\end{aligned}\quad (114)$$

but in fact the Lie derivative of a vector field is nothing new:

Proposition 1.37

If X, Y are C^∞ vector fields on a manifold M , then the Lie derivative $\mathcal{L}_X Y$ coincides with the Lie bracket $[X, Y]$.

1.14.2 The Lie Derivative of a Differential Form

Let X be a smooth vector field and ω a smooth k -form on a manifold M . Fix a point $p \in M$ and let $\phi_t: U \rightarrow M$ be a flow generated by X in a neighborhood U of p . The Lie derivative of a differential form is similar to that of the Lie derivative of a vector field. The difference is, instead of pushing a vector at $\phi_t(p)$ to p via $(\phi_{-t})_*$, we now pull $\omega_{\phi_t(p)}$ back to p via ϕ_t^* .

Definition 1.41: Lie derivative of a differential form

For X a smooth vector field and ω a smooth k -form on a manifold M , the *Lie derivative* $\mathcal{L}_X \omega$ at $p \in M$ is

$$\begin{aligned} (\mathcal{L}_X \omega)_p &:= \lim_{t \rightarrow 0} \frac{\phi_t^*(\omega_{\phi_t(p)}) - \omega_p}{t} \\ &= \lim_{t \rightarrow 0} \frac{(\phi_t^* \omega)_p - \omega_p}{t} \\ &= \left. \frac{d}{dt} \right|_{t=0} (\phi_t^* \omega)_p \end{aligned} \quad (115)$$

Proposition 1.38

If f is a C^∞ function and X a C^∞ vector field on M , then $\mathcal{L}_X f = Xf$.

Proposition 1.40

For $v \in V$, let

$$\iota_v: \bigwedge^*(V^\vee) \rightarrow \bigwedge^{*-1}(V^\vee)$$

be interior multiplication by v . Then

1. $\iota_v \circ \iota_v = 0$
2. for $\beta \in \bigwedge^k(V^\vee)$ and $\gamma \in \bigwedge^\ell(V^\vee)$

$$\iota_v(\beta \wedge \gamma) = (\iota_v \beta) \wedge \gamma + (-1)^k \beta \wedge \iota_v \gamma$$

In other words, ι_v is an antiderivation of degree -1 whose square is zero.

Let \mathcal{F} be the ring $C^\infty(M)$ of C^∞ functions on the manifold M . Because $\iota_v \omega$ is a point operator — that is, its value at p depends only on X_p and ω_p — it is \mathcal{F} -linear in either argument. This means that $\iota_X \omega$ is additive in each argument and moreover, for $f \in \mathcal{F}$

1. $\iota_f X \omega = f \iota_X \omega$
2. $\iota_X(f \omega) = f \iota_X \omega$

1.14.3 Interior Multiplication

Definition 1.42: Interior Multiplication/Contraction

If β is a k -covector on a vector space V and $v \in V$, for $k \geq 2$ the *interior multiplication* or *contraction* of β with v is the $k-1$ -covector $\iota_v \beta$ defined by

$$(\iota_v \beta)(v_2, \dots, v_k) = \beta(v, v_2, \dots, v_k)$$

i.e. plug v into the first β slot.

Proposition 1.39

For 1-covectors $\alpha^1, \dots, \alpha^k$ on a vector space V and $v \in V$

$$\begin{aligned} \iota_v(\alpha^1 \wedge \dots \wedge \alpha^k) &= \\ \sum_{i=1}^k (-1)^{i-1} \alpha^i(v) \alpha^1 \wedge \dots \wedge \widehat{\alpha^i} \wedge \dots \wedge \alpha^k \end{aligned} \quad (116)$$

where $\widehat{\alpha^i}$ means that α^i is omitted from the wedge product.

1.14.4 Properties of the Lie Derivative

Theorem 1.7

1. The Lie derivative $\mathcal{L}_X: \Omega^*(M) \rightarrow \Omega^*(M)$ is a derivation.
2. The Lie derivative commutes with the exterior derivative d
3. **Cartan homotopy formula:** $\mathcal{L}_X = d \circ \iota_X + \iota_X \circ d$
4. **“Product” formula:** For $\omega \in \Omega^k(M)$ and $Y_1, \dots, Y_k \in \mathfrak{X}(M)$

$$\begin{aligned} \mathcal{L}_X(\omega(Y_1, \dots, Y_k)) &= (\mathcal{L}_X \omega)(Y_1, \dots, Y_k) \\ &+ \sum_{i=1}^k \omega(Y_1, \dots, \mathcal{L}_X Y_i, \dots, Y_k) \end{aligned} \quad (117)$$

Theorem 1.8: Global formula for the Lie derivative

For a smooth k -form ω and smooth vector fields X, Y_1, \dots, Y_k on a manifold M

$$\begin{aligned} \mathcal{L}_X(\omega(Y_1, \dots, Y_k)) &= X(\omega(Y_1, \dots, Y_k)) \\ &- \sum_{i=1}^k \omega(Y_1, \dots, [X, Y_i], \dots, Y_k) \end{aligned} \quad (118)$$

Proposition 1.41

If ω is a C^∞ 1-form and X, Y are C^∞ vector fields on a manifold M , then

$$(d\omega)(X, Y) = X\omega(Y) - \omega([X, Y])$$

Theorem 1.9: Global formula for the exterior derivative

Assume $k \geq 1$. For a smooth k -form ω and smooth vector fields Y_0, Y_1, \dots, Y_k on a manifold M

$$(d\omega)(Y_0, \dots, Y_k) = \sum_{i=0}^k (-1)^i Y_i \omega(Y_0, \dots, \widehat{Y}_i, \dots, Y_k) \\ + \sum_{0 \leq i < j \leq k} (-1)^{i+j} \omega([Y_i, Y_j], Y_0, \dots, \widehat{Y}_i, \dots, \widehat{Y}_j, \dots, Y_k)$$

1.15 Orientation

On a manifold one integrates not functions but differential forms. For simplicity we will discuss integration of smooth forms over a submanifold. For integration over a manifold to be well defined, the manifold needs to be oriented.

An orientation of a finite-dimensional real vector space is simply an equivalence class of ordered bases, two ordered bases being equivalent if and only if their change of basis matrix has positive determinant. By its alternating nature, a multivector of top degree turns out to represent perfectly an orientation of a vector space.

An orientation on a manifold is a choice of an orientation for each tangent space (satisfying a continuity condition). An orientation on n -manifold can also be given by an equivalence class of C^∞ nowhere-vanishing n -forms, two such forms being equivalent iff one is a multiple of the other by a positive function. Finally, a third way to represent an orientation on a manifold is through an *oriented atlas*, an atlas in which any two overlapping charts are related by a transition function with everywhere positive Jacobian determinant.

1.15.1 Orientations of a Vector Space

On \mathbb{R}^3 is either right-handed or left-handed. For any two ordered bases (u_1, u_2) and (v_1, v_2) for \mathbb{R}^2 , there is a unique nonsingular 2×2 matrix $A = [a_j^i]$ such that

$$u_j = \sum_{i=1}^2 a_j^i v_i$$

called the *change-of-basis matrix* from (v_1, v_2) to (u_1, u_2) . In matrix notation

$$[u_1 u_2] = [v_1 v_2] A$$

Two ordered bases are *equivalent* if the change-of-basis A has positive determinant; this partitions ordered bases into two equivalence classes and each class is called an *orientation*.

1.15.2 Orientations and n -covectors

Instead of using an ordered basis, we can also use an n -covector to specify orientation of an n -dimensional vector space V . This approach is based on the fact that the space $\bigwedge^n(V^\vee)$ of n -covectors on V is one-dimensional.

Lemma 1.3

Let u_1, \dots, u_n and v_1, \dots, v_n be vectors in a vector space V . Suppose

$$u_j = \sum_{i=1}^n v_i a_j^i$$

If β is an n -covector on V , then

$$\beta(u_1, \dots, u_n) = (\det A) \beta(v_1, \dots, v_n)$$

Corollary 1.4

If u_1, \dots, u_n and v_1, \dots, v_n are ordered bases of a vector space V , then

$$\begin{aligned} \operatorname{sgn}(\beta(u_1, \dots, u_n)) &= \operatorname{sgn}(\beta(v_1, \dots, v_n)) \\ &\iff \det A > 0 \\ &\iff u_1, \dots, u_n \text{ and } v_1, \dots, v_n \\ &\text{are equivalent ordered bases} \end{aligned} \quad (119)$$

We say that β determines or specifies the orientation (v_1, \dots, v_n) if $\beta(v_1, \dots, v_n) > 0$. By the corollary this is well-defined (independent of basis equivalence class representative). Moreover, β, β' determine the same orientation iff $\beta = a\beta'$ for some positive real number a . To that end we define the equivalence relation

$$\beta \sim \beta' \iff \beta = a\beta' \quad a > 0$$

Thus, an orientation on V can be given by an equivalence class of **nonzero** n -covectors on V . A linear isomorphism $\bigwedge^n(V^\vee) \cong \mathbb{R}$ identifies the set of nonzero n -covectors on V with $\mathbb{R} - \{0\}$, which has two connected components. Two nonzero n -covectors β, β' on V are in the same component iff $\beta = a\beta'$ for some real number $a > 0$. Thus, each connected component of $\bigwedge^n(V^\vee) - \{0\}$ determines an orientation of V .

Example 1.37. Let

$$\left\{ \frac{\partial}{\partial x} \Big|_p, \dots, \frac{\partial}{\partial y} \Big|_p \right\}$$

be the standard basis for the tangent space $T_p\mathbb{R}^2$ and $(dx)_p, (dy)_p$ its dual basis. Then $(dx \wedge dy)_p$ determines the counterclockwise orientation on $T_p\mathbb{R}^2$.

1.15.3 Orientations on a Manifold

To orient a manifold, we orient the tangent space at each point in a coherent way (so that the orientation doesn't change abruptly anywhere). Recall that a *frame* is n -tuple X_1, \dots, X_n of possibly discontinuous vector fields on U such that at every point in U , the n -tuple $X_{1,p}, \dots, X_{n,p}$ of vectors is an ordered basis for the tangent space T_pM . A *global frame* is a frame defined on the entire manifold, while a *local frame* is defined only on some neighborhood of p . We introduce an equivalence relation on frames on U :

$$\begin{aligned} (X_1, \dots, X_n) &\sim (Y_1, \dots, Y_n) \\ &\iff \\ (X_{1,p}, \dots, X_{n,p}) &\sim (Y_{1,p}, \dots, Y_{n,p}) \end{aligned} \quad (120)$$

i.e. if

$$Y_j = \sum_i a_j^i(p) X_i$$

then the two frames (X_1, \dots, X_n) and (Y_1, \dots, Y_n) are in the same equivalence class iff A has positive determinant at every point in U . A *pointwise orientation* assigns to each point an orientation μ_p of the tangent space T_pM . In terms of frames, a pointwise orientation is an equivalence class of (possibly) discontinuous frames on M . A pointwise orientation is said to be *continuous* at p if p has a neighborhood U on which μ is represented by a continuous frame. μ is *continuous* if it is continuous at each point. Note that a continuous pointwise orientation need not be represented by a continuous global frame; it suffices that it be locally representable by a continuous local frame. A continuous pointwise orientation on M is called an *orientation* on M . A manifold is said to be *orientable* if it has an orientation; a manifold together with an orientation is said to be *oriented*.

Example 1.38. \mathbb{R}^n is orientable with orientation given by continuous global frame

$$\left[\frac{\partial}{\partial r^1}, \dots, \frac{\partial}{\partial r^n} \right]$$

Example 1.39. The open Möbius band Let R be the rectangle

$$R := \{(x, y) \in \mathbb{R}^2 | 0 \leq x \leq 1, -1 < y < 1\}$$

The *open Möbius band* M is the quotient of R by the equivalence relation

$$(0, y) \sim (1, -y)$$

See figure ???. The interior of R is the open rectangle

$$R := \{(x, y) \in \mathbb{R}^2 | 0 < x < 1, -1 < y < 1\}$$

Suppose the Möbius band is orientable; an orientation on M restricts to an orientation on U . First assume the orientation on U to be given by (e_1, e_2) . By continuity the orientations at points $(0, 0)$ and $(1, 0)$ are also (e_1, e_2) . But under the quotient the ordered basis (e_1, e_2) at $(1, 0)$ maps to $(e_1, -e_2)$ at $(0, 0)$. Thus, this contradiction proves that the Möbius band is not orientable.

Proposition 1.42

A connected, orientable manifold M has exactly two orientations.

1.15.4 Orientations and Differential Forms

In practice it is easier to manipulate the nowhere-vanishing top forms that specify a pointwise orientation. We show that the continuity condition on a pointwise orientations translates to a C^∞ condition on nowhere-vanishing top forms.

Lemma 1.4

A pointwise orientation $[(X_1, \dots, X_n)]$ on a manifold M is continuous iff each point p in M has a coordinate neighborhood U, x^1, \dots, x^n on which the function

$$(dx^1 \wedge \dots \wedge dx^n)(X_1, \dots, X_n) > 0$$

Theorem 1.10:

A manifold M of dimension n is orientable iff there exists a C^∞ nowhere-vanishing n -form on M .

If ω, ω' are two nowhere-vanishing C^∞ n -forms on a manifold M of dimension n , then $\omega = f\omega'$ for some nowhere-vanishing function f . Locally, on a chart (U, x^1, \dots, x^n) ,

$$\begin{aligned} \omega &= h(p) dx^1 \wedge \dots \wedge dx^n \\ \omega' &= g(p) dx^1 \wedge \dots \wedge dx^n \end{aligned} \quad (121)$$

where h, g are C^∞ nowhere-vanishing functions on U . Therefore $f = h/g$ is also a C^∞ nowhere-vanishing function on U . Since U is an arbitrary chart, f is C^∞ and nowhere-vanishing on all of M . On a *connected* manifold, such a function is either everywhere positive or everywhere negative (since it never crosses 0). In this the nowhere-vanishing C^∞ n -forms on a connected orientable manifold are partitioned into two equivalence classes by the equivalence relation

$$\omega \sim \omega' \iff \omega = f\omega' \quad f > 0$$

To each orientation $\mu = [(X_1, \dots, X_n)]$ on a connected orientable manifold, we associate the equivalence class of a C^∞ nowhere-vanishing n -form ω such that

$\omega(X_1, \dots, X_n) > 0$. If $\mu \mapsto [\omega]$, then $-\mu \mapsto [-\omega]$. This sets up a 1-1 correspondence between orientations on M and equivalence classes of C^∞ nowhere-vanishing n -forms. If ω is such that $\omega(X_1, \dots, X_n) > 0$, we say that ω determines or specifies the orientation $[(X_1, \dots, X_n)]$ and we call ω an *orientation form* on M . An oriented manifold can be described by a pair $M, [\omega]$, where $[\omega]$ is the equivalence class of an orientation form. For example, unless otherwise specified, \mathbb{R}^n is oriented by $dx^1 \wedge \dots \wedge dx^n$.

Definition 1.43: A

iffeomorphism

$$F: (N, [\omega_N]) \rightarrow (M, [\omega_M])$$

of oriented manifolds is said to be *orientation-preserving* if $[F^*\omega_M] = [\omega_N]$; it is *orientation-reversing* if $[F^*\omega_M] = [-\omega_N]$.

Proposition 1.43

Let U, V be open subsets of \mathbb{R}^n , both with the standard orientation inherited from \mathbb{R}^n . A diffeomorphism $F: U \rightarrow V$ is orientation preserving iff the Jacobian determinant is everywhere positive on U .

1.15.5 Orientations and Atlases

Definition 1.44

An atlas on M is said to be oriented if for any two overlapping charts (U, x^1, \dots, x^n) and (V, y^1, \dots, y^n) of the atlas, the Jacobian determinant is everywhere positive on $U \cap V$.

Theorem 1.11

A manifold M is orientable iff it has an oriented atlas.

Definition 1.45

Two oriented atlases $\{(U_\alpha, \phi_\alpha)\}$ and $\{(V_\beta, \psi_\beta)\}$ on a manifold M are said to be equivalent if the transition functions

$$\phi_\alpha \circ \psi_\beta^{-1}: \psi_\beta(U_\alpha \cap V_\beta) \rightarrow \phi_\alpha(U_\alpha \cap V_\beta)$$

have positive Jacobian determinant for all α, β . This induces an equivalence relation on the set of oriented atlases.

1.16 Manifolds with Boundary

If M is a manifold with boundary, the its boundary ∂M turns out to be a manifold of dimension one less and **with-**

out boundary. Moreover, an orientation on M induces an orientation on ∂M ; the choice of the induced orientation on ∂M is a matter of convention, guided by the desire to make Stokes' theorem sign-free. Of the ways to describe the boundary orientation, two stand out for their simplicity

1. contraction of an orientation form on M with an outward-pointing vector field on ∂M
2. "outward vector first"

Definition 1.46: Closed upper half-space

Define the *closed upper half-space*

$$\mathcal{H} := \{(x^1, \dots, x^n) \in \mathbb{R}^n | x^n \geq 0\}$$

with the subspace topology inherited from \mathbb{R}^n . The points such that $x^n > 0$, denoted $(\mathcal{H}^n)^\circ$, are called *interior points* and the points such that $x^n = 0$, denoted $\partial \mathcal{H}^n$, are called *boundary points*.

In \mathcal{H}^n one may distinguish between open subsets that intersect the boundary and open subsets that do not intersect the boundary. Both are still considered open sets (clopen?).

Definition 1.47

A topological space is *locally \mathcal{H}^n* if every point $p \in M$ has a neighborhood U homeomorphic to an open subset of \mathcal{H}^n .

Definition 1.48: Topological n -manifold with boundary

A topological n -manifold with boundary is second countable⁴, Hausdorff⁵ topological space that is locally \mathcal{H}^n .

Let M be a topological n -manifold with boundary. For $n \geq 2$, a chart U, ϕ encodes

$$\phi: U \rightarrow \phi(U) \subset \mathcal{H}^n$$

A C^∞ *manifold with boundary* is a topological manifold with boundary together with a maximal atlas.

A point $p \in M$ is called an *interior point* if in some chart (U, ϕ) , the point $\phi(p)$ is an interior point of \mathcal{H}^n . Similarly, p is a *boundary point* of M , denoted ∂M if $\phi(p)$ is a boundary point of \mathcal{H}^n .

⁴A topological space T is second-countable if there exists some countable collection $\mathcal{U} = \{U_i\}_{i=1}^\infty$ of open subsets of T such that any open subset of T can be written as a union of elements of some subfamily of \mathcal{U} .

⁵A Hausdorff space, separated space or T_2 space is a topological space where for any two distinct points there exists a neighbourhood of each which is disjoint from the neighbourhood of the other.

Example 1.40. Topological boundary versus manifold boundary Let A be the open unit disk in \mathbb{R}^2

$$A := \{x \in \mathbb{R}^2 \mid \|x\| < 1\}$$

Then its topological boundary $\text{bd}(A)$ is the unit circle, while its manifold boundary ∂A is the empty set. On the otherhand if B is the closed unit disk in \mathbb{R}^2

$$A := \{x \in \mathbb{R}^2 \mid \|x\| \leq 1\}$$

then its topological boundary and its manifold boundary coincide.

Example 1.41. Topological interior versus manifold interior Let S be the upper half-plane \mathcal{H}^2 and let D be the subset

$$D := \{(x, y) \in \mathcal{H}^2 \mid y \leq 1\}$$

Then

$$\text{int}(D) = \{(x, y) \in \mathcal{H}^2 \mid 0 \leq y < 1\}$$

containing the x -axis, while the manifold interior of D is the set

$$D^\circ = \{(x, y) \in \mathcal{H}^2 \mid 0 < y < 1\}$$

To indicate the dependence of the topological interior of a set A on its ambient space S , we might denote it by $\text{int}_S(A)$ instead of $\text{int}(A)$. Then, in the previous example, the topological interior $\text{int}_{\mathcal{H}^2}(D)$ of $D \subset \mathcal{H}^2$ is as above, but the topological interior $\text{int}_{\mathbb{R}^2}$ of $D \subset \mathbb{R}^2$ coincides with D° .

1.16.1 The Boundary of a Manifold with Boundary

Let M be a manifold of dimension n with boundary ∂M . If (U, ϕ) is a chart on M , we denote by

$$\phi' := \phi|_{U \cap \partial M}$$

the restriction of the coordinate map ϕ to the boundary. Since ϕ maps boundary points to boundary points

$$\phi' : U \cap \partial M \rightarrow \partial \mathcal{H}^n = \mathbb{R}^{n-1}$$

Moreover, if (U, ϕ) and (V, ψ) are two charts on M , then

$$\psi' \circ (\phi')^{-1} : \phi'(U \cap V \cap \partial M) \rightarrow \psi'(U \cap V \cap \partial M)$$

is C^∞ . Thus, an atlas $\{(U_\alpha, \phi_\alpha)\}$ for M induces on atlas

$$\{(U_\alpha \cap \partial M, \phi_\alpha|_{U_\alpha \cap \partial M})\}$$

for ∂M , making ∂M into a manifold of dimension $n - 1$ without boundary.

1.16.2 Outward-Pointing Vector Fields

Let M be a manifold with boundary and $p \in \partial M$. We say that a tangent vector $X_p \in T_p(M)$ is *inward-pointing* if $X_p \in T_p(\partial M)$ and there is a positive real number ε and a curve $c : [0, \varepsilon) \rightarrow M$ such that $c(0) = p$, $c((0, \varepsilon)) \subset M^\circ$, and $c'(0) = X_p$. A vector $X_p \in T_p(M)$ is *outward-pointing* if $-X_p$ is inward-pointing. For example, on \mathcal{H}^2 and for p on the x -axis, the vector $\partial/\partial y|_p$ is inward-pointing and the vector $-\partial/\partial y|_p$ is outward-pointing.

A *vector field along ∂M* is a function X that assigns to each point $p \in \partial M$ a vector X_p in the tangent space $T_p M$ (as opposed to $T_p(\partial M)$). In a coordinate neighborhood (U, x^1, \dots, x^n) of $p \in M$, such a vector field X can be written as a linear combination

$$X_q = \sum_i a^i(q) \frac{\partial}{\partial x^i} \Big|_q \quad q \in \partial M$$

In terms of local coordinates, a vector X_p is outward-pointing iff $a^i(p) < 0$.

Proposition 1.44

On a manifold M with boundary ∂M , there is a smooth outward-pointing vector field along ∂M .

1.16.3 Boundary Orientation

Proposition 1.45

Let M be an oriented n -manifold with boundary. If ω is an orientation form on M and X is a smooth outward-pointing vector field on ∂M , then $\iota_X \omega$ is a smooth nowhere-vanishing $(n - 1)$ -form on ∂M . Hence, ∂M is orientable.

We define the *boundary orientation on ∂M* to be the orientation with orientation form $\iota_X \omega$.

Proposition 1.46: “outward vector first”

Suppose M is an oriented n -manifold with boundary. Let $p \in \partial M$ and let X_p be an outward-pointing vector in $T_p M$. An ordered basis (v_1, \dots, v_{n-1}) for $T_p(\partial M)$ represents the boundary orientation at p iff the ordered basis $(X_p, v_1, \dots, v_{n-1})$ for $T_p M$ represents the orientation on M at p .

This rule is summarized as “outward vector first”.

Example 1.42. The boundary orientation on $\partial \mathcal{H}^n$ An orientation form for the standard orientation on \mathcal{H}^n is

$$\omega = dx^1 \wedge \dots \wedge dx^n$$

A smooth outward-pointing vector field on $\partial \mathcal{H}^n$ is $-\partial/\partial x^n$. By definition, an orientation form for the bound-

any orientation on $\partial\mathcal{H}^n$ is given by the contraction

$$\iota_{\partial/\partial x^n}\omega \tag{122}$$

Index

- C^∞ partition of unity on a manifold, 17
- C^∞ partition of unity subordinate to $\{U_i\}_{i \in I}$, 17
- C^∞ -compatible, 8
- k -covector, 26
- k -covector field, 26
- k -form, 26
- k -tensor, 25
- k -th exterior power, 27
- 1-form, 4

- abelian Lie algebra, 20
- antiderivation, 28
- antiderivation of degree m , 28
- antiderivation of the graded algebra, 6
- atlas, 8

- base space, 16, 25
- boundary orientation on ∂M , 35
- boundary point, 34
- boundary points, 34
- bracket, 20
- bump function at q supported in U , 17

- change-of-basis matrix, 32
- characteristic numbers, 15
- chart, 8
- Chern-Weil theory of characteristic classes, 15
- closed form, 6
- closed upper half-space, 34
- cochain complex, 6
- codifferential, 25
- complete vector field, 19
- connected, 33
- continuous, 33
- contraction, 31
- contravariant functor, 24
- coordinate neighborhood, 8
- coordinate system, 8
- cotangent bundle, 25
- cotangent space, 4, 24
- covector, 24
- covector field, 4, 24
- covectors, 3
- curve, 9

- de Rham cohomology, 7
- de Rham complex, 24
- de Rham complex of U , 6
- derivation, 3
- derivation of a Lie algebra V , 20
- diffeomorphism, 10
- differentiable structure, 9
- differential, 4, 25
- differential 1-form, 4, 24
- differential k -form, 26
- differential at p , 12
- differential complex, 6
- differential form ω of degree k , 5
- differential form of degree k , 26
- differential graded algebra, 24
- directional derivative, 2
- dual space, 3

- eigenvalues, 22
- endomorphism, 28
- equivalent, 32
- Euler characteristic, 15
- exact form, 6
- exterior derivative, 28
- exterior differentiation, 28
- exterior power, 26

- fiber, 15
- fiber at p , 16
- fiber-preserving, 16
- flow line, 19
- frame, 16, 33
- frame on U , 16
- functor, 26

- general linear group $GL(n, \mathbb{R})$, 9
- global flow, 19
- global frame, 33
- graph of f , 9

- Hausdorff, 8
- homeomorphism, 8
- homogeneous element, 28

- integral curve, 18
- interior multiplication, 31
- interior point, 34
- interior points, 34
- inward-pointing, 35
- isomorphically, 3

- Jacobi identity, 20
- Jacobian matrix of F , 11

- left and right translations, 21
- left multiplication by a , 21
- left-invariant, 23
- left-invariant vector field on G generated by $A \in T_e G$, 24
- Leibniz rule, 3
- Lie algebra, 7, 18, 20
- Lie algebra of the Lie group G , 24
- Lie Bracket, 18, 19
- Lie derivative $\mathcal{L}_X \omega$, 31
- Lie derivative $\mathcal{L}_X Y$ of Y , 30
- Lie derivative of a vector field, 29
- Lie group, 10, 20, 21

- Lie group homomorphism, 21
- Liouville form, 25
- local coordinates on U , 9
- local diffeomorphism, 11
- local finiteness, 17
- local flow, 19, 29
- local flow generated by the vector field X , 19
- local frame, 33
- local operator, 28
- local operators, 28
- locally \mathcal{H}^n , 34
- locally Euclidean of dimension n , 8
- locally finite, 17
- locally invertible, 11

- manifold with boundary, 34
- matrix exponential, 21
- maximal, 8
- morphism of differential graded algebras, 29

- norm, 21
- normed vector space, 21

- one-parameter group of diffeomorphisms, 18
- open Möbius band, 33
- operator, 28
- orientable, 33
- orientation, 32, 33
- orientation form, 34
- orientation-preserving, 34
- orientation-reversing, 34
- oriented, 33
- oriented atlas, 32
- outward-pointing, 35

- partial derivative $\frac{\partial f}{\partial x^i}$ of f with respect to x^i at p , 10
- Picard–Lindelöf theorem, 18
- Poincare form, 25
- Poincare lemma, 7
- pointwise orientation, 33
- pullback, 9, 25, 27
- pullback $F^*\omega$, 25
- pushforward, 12, 20

- right multiplication by a , 21

- second countable, 8
- section, 15, 16
- smooth C^∞ manifold, 9
- smooth at a point p , 9
- smooth curve, 14
- smooth vector bundle, 15
- subordinate, 17
- support, 17
- surface, 9

- tangent bundle, 11, 15
- tangent space, 2
- tangent vectors, 2

- top form, 26
- topological group, 10
- topological manifold, 8
- total space, 16, 25
- transition functions, 8

- vector bundle, 11, 16
- vector field, 3
- vector field X on a manifold M , 16
- vector field along ∂M , 35
- velocity vector, 14

- wedge product, 27
- wedge product of a k -form ω and ℓ -form τ , 5