

Homework 3 (Deadline: Jun 6)

1. (10 pts) Given a set $T \subset K$. Recall the definition of the covering number: $N(T, d, \varepsilon)$ is the smallest number of points in T which form a ε -covering of T under the metric d . Suppose we are allowed to use the points outside of T to do covering. Then the smallest number of points that are needed to form a ε -covering of T is referred to as the exterior covering number, denoted $N^{ext}(T, d, \varepsilon)$. Show that

$$N^{ext}(T, d, \varepsilon) \leq N(T, d, \varepsilon) \leq N^{ext}(T, d, \varepsilon/2).$$

2. (10 pts) Suppose T is a finite set and $\log \mathbb{E} [e^{\lambda X_t}] \leq \psi(\lambda)$ for all $\lambda \geq 0$ and $t \in T$, where $\psi(\cdot)$ is a convex function and $\psi(0) = \psi'(0) = 0$. Show that

$$\mathbb{E} \left[\max_{t \in T} X_t \right] \leq \psi^{*-1}(\log |T|),$$

where $\psi^*(x) = \sup_{\lambda \geq 0} \{\lambda x - \psi(\lambda)\}$.

3. (10pts) Define

$$T^n(s) = \{x \in \mathbb{R}^n : \|x\|_0 \leq s, \|x\|_2 \leq 1\},$$

where $\|x\|_0$ counts the number of non-zero entries in x . Show that the Gaussian complexity of $T^n(s)$, denoted $\mathcal{G}(T^n(s)) = \mathbb{E} \left[\sup_{x \in T^n(s)} \langle g, x \rangle \right]$, $g \sim \mathcal{N}(0, I_n)$, satisfies

$$\mathcal{G}(T^n(s)) \lesssim \sqrt{s \log \left(\frac{en}{s} \right)}.$$

4. (10 pts) Let $W \in \mathbb{R}^{m \times n}$ be a random matrix with i.i.d $\mathcal{N}(0, 1)$ entries. We have established an upper bound of $\mathbb{E} [\|W\|_2]$ using the one-step discretization bound as well as the Gaussian comparison inequality. In this problem you are required to establish an upper bound of $\mathbb{E} [\|W\|_2]$ using the Dudley integral.
5. (20 pts) Let $B_1^n = \{x : \|x\|_1 \leq 1\}$ be the ℓ_1 -norm unit ball. We have already seen that the Gaussian complexity of B_1^n satisfies

$$\mathcal{G}(B_1^n) = \mathbb{E} \left[\sup_{\|x\|_1 \leq 1} \langle g, x \rangle \right] \lesssim \sqrt{\log n}, \quad g \sim \mathcal{N}(0, I_n)$$

based on the duality between ℓ_1 -norm and ℓ_∞ -norm. In this problem, we attempt to provide a bound for $\mathcal{G}(B_1^n)$ based on the Dudley integral.

- For $\varepsilon > 0$ being sufficiently small, show that the covering of B_1^n under the ℓ_2 -norm satisfies

$$\sqrt{\log \mathcal{N}(B_1^n, \|\cdot\|_2, \varepsilon)} \lesssim \min\{\varepsilon^{-1} \sqrt{\log n}, \sqrt{n} \cdot \log(1/\varepsilon)\}.$$

Hint: Volume argument as presented in Lecture 4 may be useful in providing one bound.

- Using the above result and the Dudley integral to provide a bound for $\mathcal{G}(B_1^n)$.
6. (10 pts) Assume $X, Y \in \mathbb{R}^n$ are finite centered Gaussian Processes. Suppose that there exist a pair of index sets $A, B \subset \{1, \dots, n\}^2$ for which

$$\begin{aligned}\mathbb{E}[X_i X_j] &\leq \mathbb{E}[Y_i Y_j] & \text{for all } (i, j) \in A; \\ \mathbb{E}[X_i X_j] &\geq \mathbb{E}[Y_i Y_j] & \text{for all } (i, j) \in B; \\ \mathbb{E}[X_i X_j] &= \mathbb{E}[Y_i Y_j] & \text{for all } (i, j) \notin A \cup B.\end{aligned}$$

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function whose second derivative satisfies

$$\begin{aligned}\partial_{ij} f &\geq 0 & \text{for all } (i, j) \in A; \\ \partial_{ij} f &\leq 0 & \text{for all } (i, j) \in B.\end{aligned}$$

Show that

$$\mathbb{E}[f(X)] \leq \mathbb{E}[f(Y)].$$

7. (10 pts) Let $\phi_j : \mathbb{R} \rightarrow \mathbb{R}$ ($j = 1, \dots, n$) be 1-Lipschitz (i.e., $|\phi_j(t) - \phi_j(s)| \leq |t - s|$). Let w_j ($j = 1, \dots, n$) be i.i.d $\mathcal{N}(0, 1)$ random variables. For any $T \subset \mathbb{R}^n$, show that

$$\mathbb{E} \left[\sup_{t=(t_1, \dots, t_n) \in T} \sum_{j=1}^n w_j \phi_j(t_j) \right] \leq \mathbb{E} \left[\sup_{t=(t_1, \dots, t_n) \in T} \sum_{j=1}^n w_j t_j \right].$$

What does the above result mean in terms of the Gaussian complexity?