Summary in Graph


Exam Summary (GO Classes DA Test Series 2026 | Linear Algebra | Subject Wise Test 2)

Qs. Attempted:	11 11 + 0	Correct Marks:	6 6 + 0
Correct Attempts:	6 6 + 0	Penalty Marks:	1.33 1.33 + 0
Incorrect Attempts:	5 5 + 0	Resultant Marks:	4.67 4.67 + 0

Total Questions:	30 13 + 17
Total Marks:	47 13 + 34
Exam Duration:	90 Minutes
Time Taken:	90 Minutes

Individual Question Statistics

Select Metric: Marks Obtained



* Average Stats based on first-attempt stats of Top 10% (25) out of 250 users

EXAM RESPONSE | EXAM STATS | FEEDBACK

Technical

Q #1 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Consider the quadratic form

$$Q(x_1, x_2) = 5x_1^2 - 10x_1x_2 + x_2^2$$

Which of the following correctly represents $Q(x_1, x_2)$ as a matrix product?

A. $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 5 & -5 \\ -5 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

B. $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 5 & -10 \\ -10 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

C. $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 5 & -10 \\ -5 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

D. $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 1 & -5 \\ -5 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

Your Answer: A | Correct Answer: A | Correct | Time taken: 07min 25sec | Discuss

Q #2 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

For which value(s) of k is the set $\left\{ \begin{bmatrix} 1 \\ 3 \\ -4 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -1 \\ k \end{bmatrix} \right\}$ a basis for $\mathbb{R}^3$?

A. $k = 48$

B. $k \neq 48$

C. $k = 24$

D. $k \neq 24$

Your Answer: A | Correct Answer: B | Incorrect | Time taken: 13min 57sec | Discuss

Q #3 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Suppose $A = \begin{bmatrix} a & b \\ b & d \end{bmatrix}$, with $b \neq 0$. If one of the eigenvectors of A is $\begin{bmatrix} 2 \\ 3 \end{bmatrix}$, then another eigenvector of A is

A. $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$

B. $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

C. $\begin{bmatrix} -3 \\ 2 \end{bmatrix}$

D. $\begin{bmatrix} -2 \\ 3 \end{bmatrix}$

Your Answer: A | Correct Answer: C | Incorrect | Time taken: 24min 12sec | Discuss

Q #4 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

If $A = \begin{bmatrix} 1 & 0 & a & 2 \\ -2 & -1 & b & 3 \\ 3 & 1 & c & 1 \end{bmatrix}$ and the reduced row-echelon form of A is $R = \begin{bmatrix} 1 & 0 & 3 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$, then $(a, b, c) =$

A. $(3, -2, 0)$

B. $(3, -4, 7)$

C. $(0, 0, 1)$

D. $(3, -2, 7)$

Your Answer: A | Correct Answer: B | Incorrect | Time taken: 19min 06sec | Discuss

Q #5 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Suppose A is a 6×6 diagonalizable matrix with characteristic polynomial

$$C_A(x) = x^2(x-3)^2(x-4)(x-5).$$

What is the rank of A ?

A. 2

B. 3

C. 4

D. 5

Your Answer: D | Correct Answer: C | Incorrect | Time taken: 01min 19sec | Discuss

Q #6 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Let

$$\vec{d} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \quad \vec{v} = \begin{bmatrix} 6 \\ 6 \\ 6 \end{bmatrix}$$

Then, the projection of $\vec{v}$ onto $\vec{d}$, denoted as $\text{proj}_{\vec{d}}(\vec{v})$, is:

A. $\begin{bmatrix} 6 \\ 6 \\ 6 \end{bmatrix}$

B. $\begin{bmatrix} 3 \\ -3 \\ 3 \end{bmatrix}$

C. $\begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix}$

D. $\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

Your Answer: | Correct Answer: C | Not Attempted | Time taken: 00min 15sec | Discuss

Q #7 | Numerical Type | Award: 1 | Penalty: 0 | Linear Algebra

Let A be a 4×4 matrix such that $\det(A) = -2$. What is the value of $\det(\text{adj}(A))$?

Your Answer: 1 | Correct Answer: -8 | Incorrect | Time taken: 01min 03sec | Discuss

Q #8 | Multiple Select Type | Award: 1 | Penalty: 0 | Linear Algebra

For which of the following values of a are the vectors

$$\mathbf{u} = \begin{bmatrix} 1 \\ a \\ 4 \end{bmatrix} \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 3a \\ -1 \\ -6 \end{bmatrix}$$

orthogonal?

A. 6

B. 8

C. 12

D. 24

Your Answer: C | Correct Answer: C | Correct | Time taken: 10min 41sec | Discuss

Q #9 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

The values of a for which the matrix $\begin{bmatrix} 6 & 0 & a \\ 0 & 2 & 0 \\ a & 0 & 6 \end{bmatrix}$ is not invertible, are

A. $a = \pm 3$
B. $a = \pm 2$
C. $a = \pm 6$
D. $a = \pm 9$

Your Answer: C | Correct Answer: C | Correct | Time taken: 00min 33sec | Discuss

Q #10 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

If A and B are 3×3 matrices such that $\det(A) = 2$ and $\det(B) = -3$, then the value of $\det\left(A^2 B^{-1} A^T B^3\right)$ is

A. -72
B. 72
C. -108
D. 108

Your Answer: B | Correct Answer: B | Correct | Time taken: 01min 23sec | Discuss

Q #11 | Numerical Type | Award: 1 | Penalty: 0 | Linear Algebra

Let A be a 5×7 real matrix whose rows are linearly independent. What is the dimension of the null space of A ?

Your Answer: 2 | Correct Answer: 2 | Correct | Time taken: 01min 33sec | Discuss

Q #12 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Which of the following statements are correct?
A. If U is a subspace of $\mathbb{R}^6$ and $\dim(U) = 4$, then $\dim\left(U^{\perp}\right) = 2$.
B. If A is an invertible matrix, then $\lambda = 0$ is an eigenvalue of A.
C. If $\{X, Y, Z\}$ is a linearly independent set of vectors, then $\{X, Y\}$ is also linearly independent.

A. Only (A) and (C) are correct
B. Only (A) is correct
C. All (A), (B), and (C) are correct
D. Only (C) is correct

Your Answer: A | Correct Answer: A | Correct | Time taken: 00min 57sec | Discuss

Q #13 | Multiple Choice Type | Award: 1 | Penalty: 0.33 | Linear Algebra

Let U be a subspace of $\mathbb{R}^n$, and let $\{\vec{v}_1, \vec{v}_2, \ldots, \vec{v}_k\}$ be an orthonormal basis for U. Let

$$A = [\vec{v}_1 \quad \vec{v}_2 \quad \cdots \quad \vec{v}_k]$$

be the $n \times k$ matrix whose columns are the basis vectors.
Then for any $\vec{x} \in \mathbb{R}^n$, what is $\text{proj}_U(\vec{x})$ equal to?

A. $A^T A\vec{x}$
B. $AA^T\vec{x}$
C. $A\vec{x}$
D. $AA\vec{x}$

Your Answer: | Correct Answer: B | Not Attempted | Time taken: 08min 24sec | Discuss

Q #14 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Linear Algebra

Which of the following matrices are diagonalizable?
Let:

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} -9 & 13 \\ -2 & 6 \end{bmatrix}$$

$$C = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

A. Only B

B. B and C

C. All three: A, B, and C

D. Only C

Your Answer: | Correct Answer: B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #15 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let V be a vector space over $\mathbb{R}$, and let $S \subseteq V$ be defined in each case below. Determine which of the sets S are subspaces of V.

A. $V = \mathbb{R}^3$,
$S = \left\{ \begin{bmatrix} x \\ 12 \\ 3x \end{bmatrix} : x \in \mathbb{R} \right\}$

B. $V = \mathbb{R}^2$,
$S = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 : 2x - 5y = 11 \right\}$

C. $V = \mathbb{R}^n$,
$S = \{\vec{x} \in \mathbb{R}^n : A\vec{x} = 2\vec{x}\}$, where A is a fixed $n \times n$ matrix

D. $V = \mathbb{R}^3$,
$S = \left\{ (x, y, z) \in \mathbb{R}^3 : x + 2y = 0, 2x + 3z = 0 \right\}$

Your Answer: | Correct Answer: C;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #16 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let A be an $n \times n$ matrix. Define the eigenspace V_λ associated with eigenvalue $\lambda \in \mathbb{R}$ as:

$$V_\lambda = \{\vec{v} \in \mathbb{R}^n : A\vec{v} = \lambda\vec{v}\}$$

Which of the following statements is/are correct?

A. $\vec{0} \notin V_\lambda$ because eigenvectors must be nonzero.

B. If $\vec{v} \in V_\lambda \cap V_\tau$ for $\lambda \neq \tau$, then $\vec{v} \neq \vec{0}$.

C. The zero vector always belongs to every eigenspace, but no nonzero vector can belong to two distinct eigenspaces.

D. The dimension of V_λ is always equal to the algebraic multiplicity of λ.

Your Answer: | Correct Answer: C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #17 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Linear Algebra

Find $\vec{w}$, the projection of the vector $\vec{v} = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$ onto the plane with equation $2x + y + 2z = 0$.

A. $\vec{w} = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$

B. $\vec{w} = \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$

C. $\vec{w} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$

D. $\vec{w} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$

Your Answer: | Correct Answer: C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #18 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Suppose A is a 3×5 matrix such that the three vectors

$$\begin{bmatrix} 1 \\ 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 7 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ -2 \\ -7 \\ 5 \\ 0 \end{bmatrix}$$

span $\text{null}(A)$. Select all of the following statements that are True.

A. The rows of A are linearly independent.
B. The columns of A are linearly independent.
C. The system $A\vec{x} = \vec{b}$ is consistent for every $\vec{b}$ in $\mathbb{R}^3$.
D. The rank of A is 3 .

Your Answer: | Correct Answer: A;C;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #19 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let $u = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3$, and define the matrix $A = uu^T$.

Which of the following statements is/are correct?

A. $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix}^T$ is an eigenvector of A

B. A has only one nonzero eigenvalue, which is 14

C. $\text{Rank}(A) = 1$

D. Matrix A has three equal eigenvalues, all equal to 14

Your Answer: | Correct Answer: A;B;C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #20 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Linear Algebra

Let $\mathbf{c} \neq \mathbf{0}$ be a column vector in $\mathbb{R}^m$ and $\mathbf{r} \neq \mathbf{0}$ be a row vector in $\mathbb{R}^n$. Define the matrix:

$$A = \mathbf{cr}$$

Match the following:

(P)	An eigenvector of A^TA	(i)	$\mathbf{r}^T$
(Q)	An eigenvalue of A^TA	(ii)	$\|\mathbf{c}\|^2\|\mathbf{r}\|^2$
(R)	An eigenvector of AA^T	(iii)	$\mathbf{c}$
(S)	An eigenvalue of AA^T	(iv)	$\|\mathbf{c}\|^2\|\mathbf{r}\|^2$

A. $\text{P} \leftrightarrow \text{i}, \text{Q} \leftrightarrow \text{ii}, \text{R} \leftrightarrow \text{iv}, \text{S} \leftrightarrow \text{iii}$

B. $\text{P} \leftrightarrow \text{iii}, \text{Q} \leftrightarrow \text{i}, \text{R} \leftrightarrow \text{ii}, \text{S} \leftrightarrow \text{iv}$

C. $\text{P} \leftrightarrow \text{i}, \text{Q} \leftrightarrow \text{ii}, \text{R} \leftrightarrow \text{iii}, \text{S} \leftrightarrow \text{iv}$

D. $\text{P} \leftrightarrow \text{iv}, \text{Q} \leftrightarrow \text{i}, \text{R} \leftrightarrow \text{ii}, \text{S} \leftrightarrow \text{iii}$

Your Answer: | Correct Answer: C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #21 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let $\mathbf{v}$ be a unit column vector in $\mathbb{R}^n$, and define the matrix

$$A = I - 2\mathbf{v}\mathbf{v}^T$$

where I is the $n \times n$ identity matrix.

Which of the following statements is/are correct?

A. $\mathbf{v}$ is an eigenvector of A corresponding to eigenvalue -1

B. A is symmetric and orthogonal

C. $\mathbf{v}^T\mathbf{v} = 1$ and $\left(\mathbf{v}\mathbf{v}^T\right)\mathbf{v} = \mathbf{v}$

D. $A^{-1} = A^T = A$

Your Answer: | Correct Answer: A;B;C;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #22 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let

$$\vec{u} = \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}, \quad \vec{v} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \quad \vec{w} = \begin{bmatrix} 2 \\ b \\ 5 \end{bmatrix}$$

Given that $\vec{u}$ and $\vec{v}$ form a basis for the solution space of a system of homogeneous linear equations, for which value(s) of b is $\vec{w}$ also a solution to the same system?

A. $b = -3$
B. $b = -7$
C. $b = 1$
D. $b = 0$

Your Answer: | Correct Answer: B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #23 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let $S : \mathbb{R}^3 \to \mathbb{R}^4$ and $T : \mathbb{R}^4 \to \mathbb{R}^3$ be linear transformations defined by:

$$S\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}\right) = \begin{bmatrix} x_1 + 2x_2 \\ x_2 - 2x_3 \\ x_1 + x_3 \\ x_1 - x_2 \end{bmatrix}, \quad T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}\right) = \begin{bmatrix} x_1 - x_2 \\ x_2 - x_3 \\ x_3 - x_4 \end{bmatrix}$$

Which of the following statements are correct?

A. S is one-to-one
B. T is not onto
C. S is not one-to-one
D. T is onto

Your Answer: | Correct Answer: A;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #24 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Linear Algebra

Let $T : \mathbb{R}^2 \to \mathbb{R}^2$ be a linear transformation such that:

$$T\left(\begin{bmatrix} 1 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} -2 \\ -6 \end{bmatrix}, \quad T\left(\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} -1 \\ -9 \end{bmatrix}$$

Let A be the 2×2 matrix such that $T(\vec{x}) = A\vec{x}$ for all $\vec{x} \in \mathbb{R}^2$.
Which of the following is the correct matrix A ?

A. $\begin{bmatrix} -5 & 1 \\ 3 & -3 \end{bmatrix}$

B. $\begin{bmatrix} -2 & -1 \\ -6 & -9 \end{bmatrix}$

C. $\begin{bmatrix} 4 & -3 \\ 1 & 1 \end{bmatrix}$

D. $\begin{bmatrix} -1 & 4 \\ -3 & 3 \end{bmatrix}$

Your Answer: | Correct Answer: A | Not Attempted | Time taken: 00min 00sec | Discuss

Q #25 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let A be an $m \times k$ matrix and B be a $k \times n$ matrix. Consider the product AB, which is an $m \times n$ matrix. Which of the following statements are always true?

A. $\text{nullity}(B) \leq \text{nullity}(AB)$
B. $\text{rank}(AB) \leq \text{rank}(A)$
C. $\text{nullity}(AB) \leq \text{nullity}(B)$
D. $\text{rank}(A) \leq \text{rank}(AB)$

Your Answer: | Correct Answer: A;B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #26 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let $A \in \mathbb{R}^{n \times n}$ be a matrix such that $A^2 = \mathbf{0}$.
Suppose any vector in the column space of A can be expressed as $\vec{v} = A\vec{x}$ for some $\vec{x} \in \mathbb{R}^n$.
Which of the following statements is/are necessarily true?

A. If $\vec{v} \in \text{Col}(A)$, then $\vec{v} \in \text{Null}(A)$
B. $\text{Col}(A) \subseteq \text{Null}(A)$
C. $\text{rank}(A) \leq \frac{n}{2}$
D. $\text{Col}(A) = \text{Null}(A)$

Your Answer: | Correct Answer: A;B;C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #27 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let

$$U = \left\{ \vec{x} \in \mathbb{R}^3 \mid \vec{x} \cdot \vec{u}_1 = 0 \text{ and } \vec{x} \cdot \vec{u}_2 = 0 \right\}$$

where

$$\vec{u}_1 = \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}, \quad \vec{u}_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Which of the following statements is/are correct?

A. $\dim(U) = 2$
B. U is the orthogonal complement of $\text{span}\{\vec{u}_1, \vec{u}_2\}$
C. A basis for U is $\begin{bmatrix} 3 \\ -4 \\ 1 \end{bmatrix}$
D. $U = \text{span}\{\vec{u}_1, \vec{u}_2\}$.

Your Answer: | Correct Answer: B;C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #28 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let

$$A = \begin{bmatrix} 1 & 0 & 0 \\ -4 & -7 & 2 \\ -12 & -24 & 7 \end{bmatrix}, \quad B = \begin{bmatrix} -1 & -3 & 1 \\ -4 & -7 & 2 \\ -16 & -30 & 9 \end{bmatrix}$$

Which of the following statements is/are correct?

A. Matrices A and B have the same characteristic polynomial.
B. A and B have the same rank.
C. A and B are similar matrices.
D. Exactly one of A, B is diagonalizable.

Your Answer: | Correct Answer: A;B;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #29 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let

$$U = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} \in \mathbb{R}^3 \,\middle|\, \det \begin{bmatrix} 2 & -1 & a \\ 3 & 1 & b \\ 1 & -1 & c \end{bmatrix} = 0 \right\}$$

Which of the following statements is/are correct?

A. The set U forms a subspace of $\mathbb{R}^3$.
B. The set U is equal to the null space of the row vector $\begin{bmatrix} -4 & 1 & 5 \end{bmatrix}$.
C. The set U is equal to the span of the vectors $\begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \\ -1 \end{bmatrix}$.
D. The set U contains only finitely many vectors.

Your Answer: | Correct Answer: A;B;C | Not Attempted | Time taken: 00min 00sec | Discuss

Q #30 | Multiple Select Type | Award: 2 | Penalty: 0 | Linear Algebra

Let $\mathbf{u} = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}$, and define the transformation $T : \mathbb{R}^3 \to \mathbb{R}^3$ by $T(\mathbf{x}) = \text{proj}_{\mathbf{u}} \mathbf{x}$.
Let A be the matrix representation of T, i.e., $T(\mathbf{x}) = A\mathbf{x}$.
Which of the following statements is/are correct?

A. The matrix A is given by

$$A = \frac{1}{\|\mathbf{u}\|^2} \mathbf{u}\mathbf{u}^T$$

B. The matrix A is

$$A = \frac{1}{14} \begin{bmatrix} \mathbf{u} & -2\mathbf{u} & 3\mathbf{u} \end{bmatrix}$$

C. The matrix A satisfies both $A^T = A$ and $A^2 = A$
D. The transformation $T(\mathbf{x})$ always projects $\mathbf{x}$ onto the subspace orthogonal to $\mathbf{u}$

Your Answer: | Correct Answer: A;B;C | Not Attempted | Time taken: 00min 00sec | Discuss

Copyright & Stuff