Summary in Graph


Exam Summary (GO Classes CS/DA | Fundamental Course | Weekly Quiz 1)

Qs. Attempted:	**0** (0 + 0)	**Correct Marks:**	**0** (0 + 0)
Correct Attempts:	**0** (0 + 0)	**Penalty Marks:**	**0** (0 + 0)
Incorrect Attempts:	**0** (0 + 0)	**Resultant Marks:**	**0** (0 + 0)

Total Questions:	**10** (4 + 6)
Total Marks:	**16** (4 + 12)
Exam Duration:	**30 Minutes**
Time Taken:	**0 Minutes**

Individual Question Statistics

Select Metric: Marks Obtained



* Average Stats based on first-attempt stats of Top 10% (36) out of 357 users

EXAM RESPONSE | EXAM STATS | FEEDBACK

Aptitude

Q #1 | Multiple Select Type | Award: 1 | Penalty: 0 | Quantitative Aptitude

Let $\mathrm{S} = \sum_{i=1}^{n} f(x_i), \mathrm{A} = \sum_{i=1}^{a} f(x_i)$ and $\mathrm{B} = \sum_{i=a}^{n} f(x_i)$ then

Which of the following is/are ALWAYS true?

(Consider f as some arbitary function)

A. $\mathrm{S} = \mathrm{A} + \mathrm{B}$

B. $\mathrm{S} = \sum_{i=1}^{a-1} f(x_i) + f(x_a) + \mathrm{B}$

C. $\mathrm{S} = \sum_{i=1}^{a-1} f(x_i) + \mathrm{B}$

D. $\mathrm{S} = \mathrm{A} + \mathrm{B} - f(x_a)$

Your Answer: | Correct Answer: C;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #2 Multiple Choice Type | Award: 1 | Penalty: 0.33 | Quantitative Aptitude

Evaluate: $1 + (1+b)r + (1+b+b^2)r^2 + \ldots$ to infinite terms for $|br| < 1$.

A. $\frac{1}{(1-br)(1-r)}$

B. $\frac{1}{(1-r)(1-b)}$

C. $\frac{1}{(1-b)(b-r)}$

D. None of these

Your Answer: | Correct Answer: A | Not Attempted | Time taken: 00min 00sec | Discuss

Q #3 Multiple Choice Type | Award: 1 | Penalty: 0.33 | Quantitative Aptitude

Evaluate $1 + 2x + 3x^2 + 4x^3 + \ldots$ upto infinity, where $|x| < 1$.

A. $\frac{1}{(1-x)^2}$

B. $\frac{x}{(1-x)^2}$

C. $\frac{x^2}{(1-x)^2}$

D. $1 - \frac{1}{(1-x)^2}$

Your Answer: | Correct Answer: A | Not Attempted | Time taken: 00min 00sec | Discuss

Q #4 Numerical Type | Award: 1 | Penalty: 0 | Quantitative Aptitude

Find the sum of first 24 terms of the $\mathrm{A.P.}$ $a_1,\ a_2,\ a_3,\ \ldots$ if it is known that

$$a_1 + a_5 + a_{10} + a_{15} + a_{20} + a_{24} = 225$$

Your Answer: | Correct Answer: 900 | Not Attempted | Time taken: 00min 00sec | Discuss

Q #5 Multiple Select Type | Award: 2 | Penalty: 0 | Quantitative Aptitude

Let k be some integer and it is given that $a \equiv b (\bmod n)$ then which of the following(s) is/are ALWAYS true? More than one option can be true.

A. $a \equiv b - 3n (\bmod n)$

B. $a \equiv b + k (\bmod n)$

C. $a + k \equiv b + k (\bmod n)$

D. $a + 5n \equiv b - 3n (\bmod n)$

Your Answer: | Correct Answer: A;C;D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #6 | Multiple Select Type | Award: 2 | Penalty: 0 | Quantitative Aptitude

Which of the following(s) ALWAYS hold given that $a \equiv b(\bmod n)$ is true for some integers a, b and n. More than one option can be true.

A. $a \bmod n = b \bmod n$
B. $n \mid (a - b)$
C. $n \mid a$
D. $n \mid b$

Your Answer: | Correct Answer: A;B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #7 | Multiple Select Type | Award: 2 | Penalty: 0 | Quantitative Aptitude

Which of the following options is/are TRUE?

A. $43 \equiv -2(\bmod 5)$
B. $5 \equiv -7(\bmod 3)$
C. $-10 \equiv -25(\bmod 7)$
D. $8 \equiv -8(\bmod 5)$

Your Answer: | Correct Answer: A;B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #8 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Quantitative Aptitude

If S is the sum to infinity of a $\mathrm{G.P.}$ whose first term is $'a'$, then the sum of the first n terms is

A. $\mathrm{S}\left(1 - \frac{a}{\mathrm{S}}\right)^n$
B. $\mathrm{S}\left[1 - \left(1 - \frac{a}{\mathrm{S}}\right)^n\right]$
C. $a\left[1 - \left(1 - \frac{a}{\mathrm{S}}\right)^n\right]$
D. $\mathrm{S}\left[1 - \left(1 - \frac{\mathrm{S}}{a}\right)^n\right]$

Your Answer: | Correct Answer: B | Not Attempted | Time taken: 00min 00sec | Discuss

Q #9 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Quantitative Aptitude

If the sum of p terms of an $\mathrm{A.P.}$ is q and the sum of q terms is p, then the sum of $p + q$ terms is _____.

A. 0
B. $p - q$
C. $p + q$
D. $-(p + q)$

Your Answer: | Correct Answer: D | Not Attempted | Time taken: 00min 00sec | Discuss

Q #10 | Multiple Choice Type | Award: 2 | Penalty: 0.67 | Quantitative Aptitude

If the sum of an infinitely decreasing GP is 3, and the sum of the squares of its terms is $9/2$, the sum of the cubes of the terms is _____

A. $\frac{105}{13}$
B. $\frac{108}{13}$
C. $\frac{729}{8}$
D. None of these

Your Answer: | Correct Answer: B | Not Attempted | Time taken: 00min 02sec | Discuss

Copyright & Stuff