

MATH101 ASSIGNMENT 11

MARK VILLAR

- (1) To show that the system is inconsistent if $\gamma \neq 2\alpha - 3\beta$, we can write the equivalent statement: the system is consistent if and only if $\gamma = 2\alpha - 3\beta$. We now use Gauss-Jordan elimination to solve for the three unknowns x, y and z . Assume $\gamma = 2\alpha - 3\beta$. The augmented matrix then becomes

$$\begin{array}{ccc|c} 2 & -1 & 3 & \alpha \\ 3 & 1 & -5 & \beta \\ -5 & -5 & 21 & 2\alpha - 3\beta \end{array}$$

Next we convert the matrix to its reduced row echelon form.

$$\begin{array}{l} R_1 + R_2 \\ \hline \begin{array}{ccc|c} 5 & 0 & -2 & \alpha + \beta \\ 3 & 1 & -5 & \beta \\ -5 & -5 & 21 & 2\alpha - 3\beta \end{array} \end{array}$$

$$\begin{array}{l} R_2 + R_1 \\ \hline \begin{array}{ccc|c} 5 & 0 & -2 & \alpha + \beta \\ 8 & 1 & -7 & \alpha + 2\beta \\ -5 & -5 & 21 & 2\alpha - 3\beta \end{array} \end{array}$$

$$\begin{array}{l} R_3 + R_1 \\ \hline \begin{array}{ccc|c} 5 & 0 & -2 & \alpha + \beta \\ 8 & 1 & -7 & \alpha + 2\beta \\ 0 & -5 & 19 & 3\alpha - 2\beta \end{array} \end{array}$$

$$\begin{array}{l} \frac{1}{5}R_1 \\ \hline \begin{array}{ccc|c} 1 & 0 & -\frac{2}{5} & \frac{\alpha+\beta}{5} \\ 8 & 1 & -7 & \alpha + 2\beta \\ 0 & -5 & 19 & 3\alpha - 2\beta \end{array} \end{array}$$

$$\begin{array}{l} R_2 - 8R_1 \\ \hline \begin{array}{ccc|c} 1 & 0 & -\frac{2}{5} & \frac{\alpha+\beta}{5} \\ 0 & 1 & -\frac{19}{5} & \frac{2\beta-3\alpha}{5} \\ 0 & -5 & 19 & 3\alpha - 2\beta \end{array} \end{array}$$

$$\begin{array}{l} -\frac{1}{5}R_3 \\ \hline \begin{array}{ccc|c} 1 & 0 & -\frac{2}{5} & \frac{\alpha+\beta}{5} \\ 0 & 1 & -\frac{19}{5} & \frac{2\beta-3\alpha}{5} \\ 0 & 1 & -\frac{19}{5} & \frac{2\beta-3\alpha}{5} \end{array} \end{array}$$

$$\begin{array}{l} R_3 - R_2 \\ \hline \begin{array}{ccc|c} 1 & 0 & -\frac{2}{5} & \frac{\alpha+\beta}{5} \\ 0 & 1 & -\frac{19}{5} & \frac{2\beta-3\alpha}{5} \\ 0 & 0 & 0 & 0 \end{array} \end{array}$$

The last row implies that the equations are not independent and that the planes are intersecting in a single straight line. From the first two rows we can write

$$\begin{aligned}x - \frac{2}{5}z &= \frac{\alpha + \beta}{5} \Rightarrow x = \frac{2z + \alpha + \beta}{5} \\y - \frac{19}{5}z &= \frac{2\beta - 3\alpha}{5} \Rightarrow y = \frac{19z + 2\beta - 3\alpha}{5}\end{aligned}$$

Thus we have shown that the system requires $\gamma = 2\alpha - 3\beta$ to solve for two of the unknowns, x and y , in terms of the third, z . Consequently, the system is consistent if and only if $\gamma = 2\alpha - 3\beta$.

- (2) Let x , y and z be the number of shares the investor owns in BHP-Billiton, AMP and WMC respectively. Given the changes in both daily share prices and the value of total shares, the following system summarises the data available.

$$\begin{aligned}-x - 1.5y + 0.5z &= -350 \\1.5x - 0.5y + z &= 600\end{aligned}$$

Thus we have a system of two linear equations in three unknowns. To solve this 2×3 system we use Gauss-Jordan elimination.

$$\begin{aligned}&\begin{array}{ccc|c} -1 & -\frac{3}{2} & \frac{1}{2} & -350 \\ \frac{3}{2} & -\frac{1}{2} & 1 & 600 \end{array} \\&-R_1 \quad \begin{array}{ccc|c} 1 & \frac{3}{2} & -\frac{1}{2} & 350 \\ \frac{3}{2} & -\frac{1}{2} & 1 & 600 \end{array} \\&R_2 - \frac{3}{2}R_1 \quad \begin{array}{ccc|c} 1 & \frac{3}{2} & -\frac{1}{2} & 350 \\ 0 & -\frac{11}{4} & \frac{7}{4} & 75 \end{array} \\&-\frac{4}{11}R_2 \quad \begin{array}{ccc|c} 1 & \frac{3}{2} & -\frac{1}{2} & 350 \\ 0 & 1 & -\frac{7}{11} & -\frac{300}{11} \end{array} \\&R_1 - \frac{3}{2}R_2 \quad \begin{array}{ccc|c} 1 & 0 & \frac{5}{11} & \frac{4300}{11} \\ 0 & 1 & -\frac{7}{11} & -\frac{300}{11} \end{array}\end{aligned}$$

This is the last row elimination possible and consequently we can solve for only two of the unknowns, x and y , in terms of the third, z .

$$\begin{aligned}x + \frac{5}{11}z &= \frac{4300}{11} \Rightarrow x = \frac{4300 - 5z}{11} \\y - \frac{7}{11}z &= -\frac{300}{11} \Rightarrow y = \frac{7z - 300}{11}\end{aligned}$$

Hence the stockbroker does not have enough information to calculate the number of shares the investor owns in each company.

But if the investor tells the stockbroker that $z = 200$ then,

$$x = \frac{4300 - 5(200)}{11} = 300$$

$$y = \frac{7(200) - 300}{11} = 100$$

- (3) Given the matrices A , B and C , only $A - B$ is undefined as a 3×2 matrix cannot be subtracted from a 3×3 matrix. The rest are calculated below.

$$AB = \begin{bmatrix} 5 & 3 & -1 \\ 1 & 1 & -1 \\ 3 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 10 & -10 \\ 3 & -2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 50 + 9 - 1 & -50 - 6 + 0 \\ 10 + 3 - 1 & -10 - 2 + 0 \\ 30 + 0 + 1 & -30 + 0 + 0 \end{bmatrix} = \begin{bmatrix} 58 & -56 \\ 12 & -12 \\ 31 & -30 \end{bmatrix}$$

$$AC = \begin{bmatrix} 5 & 3 & -1 \\ 1 & 1 & -1 \\ 3 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 & 3 & -1 \\ 0 & -1 & 0 \\ 4 & 2 & -3 \end{bmatrix} = \begin{bmatrix} 10 + 0 - 4 & 15 - 3 - 2 & -5 + 0 + 3 \\ 2 + 0 - 4 & 3 - 1 - 2 & -1 + 0 + 3 \\ 6 + 0 + 4 & 9 + 0 + 2 & -3 + 0 - 3 \end{bmatrix}$$

$$= \begin{bmatrix} 6 & 10 & -2 \\ -2 & 0 & 2 \\ 10 & 11 & -6 \end{bmatrix}$$

$$CB = \begin{bmatrix} 2 & 3 & -1 \\ 0 & -1 & 0 \\ 4 & 2 & -3 \end{bmatrix} \cdot \begin{bmatrix} 10 & -10 \\ 3 & -2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 20 + 9 - 1 & -20 - 6 + 0 \\ 0 - 3 + 0 & 0 + 2 + 0 \\ 40 + 6 - 3 & -40 - 4 + 0 \end{bmatrix} = \begin{bmatrix} 28 & -26 \\ -3 & 2 \\ 43 & -44 \end{bmatrix}$$

$$A + 3C = \begin{bmatrix} 5 & 3 & -1 \\ 1 & 1 & -1 \\ 3 & 0 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 9 & -3 \\ 0 & -3 & 0 \\ 12 & 6 & -9 \end{bmatrix} = \begin{bmatrix} 11 & 12 & -4 \\ 1 & -2 & -1 \\ 15 & 6 & -8 \end{bmatrix}$$

$$AC - CA = \begin{bmatrix} 6 & 10 & -2 \\ -2 & 0 & 2 \\ 10 & 11 & -6 \end{bmatrix} - \begin{bmatrix} 2 & 3 & -1 \\ 0 & -1 & 0 \\ 4 & 2 & -3 \end{bmatrix} \cdot \begin{bmatrix} 5 & 3 & -1 \\ 1 & 1 & -1 \\ 3 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 6 & 10 & -2 \\ -2 & 0 & 2 \\ 10 & 11 & -6 \end{bmatrix} - \begin{bmatrix} 10 + 3 - 3 & 6 + 3 + 0 & -2 - 3 - 1 \\ 0 - 1 + 0 & 0 - 1 + 0 & 0 + 1 + 0 \\ 20 + 2 - 9 & 12 + 2 + 0 & -4 - 2 - 3 \end{bmatrix}$$

$$= \begin{bmatrix} 6 & 10 & -2 \\ -2 & 0 & 2 \\ 10 & 11 & -6 \end{bmatrix} - \begin{bmatrix} 10 & 9 & -6 \\ -1 & -1 & 1 \\ 13 & 14 & -9 \end{bmatrix} = \begin{bmatrix} -4 & 1 & 4 \\ -1 & 1 & 1 \\ -3 & -3 & 3 \end{bmatrix}$$

- (4) Since A is a 2×2 matrix and $\det A \neq 0$,

$$A^{-1} = \frac{1}{2 \times 2 - 1 \times 2} \begin{bmatrix} 2 & -1 \\ -2 & 2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 & -1 \\ -2 & 2 \end{bmatrix} = \begin{bmatrix} 1 & -\frac{1}{2} \\ -1 & 1 \end{bmatrix}$$

To find B^{-1} we consider the following augmented matrix and perform the row reductions below. Note that we are simply solving the system given by $BB^{-1} = I$.

$$\begin{array}{c}
\begin{array}{ccc|ccc}
1 & 2 & 3 & 1 & 0 & 0 \\
1 & 1 & 2 & 0 & 1 & 0 \\
0 & 1 & 2 & 0 & 0 & 1
\end{array} \\
\\
\begin{array}{c} R_1 - 2R_3 \end{array} \begin{array}{ccc|ccc}
1 & 0 & -1 & 1 & 0 & -2 \\
1 & 1 & 2 & 0 & 1 & 0 \\
0 & 1 & 2 & 0 & 0 & 1
\end{array} \\
\\
\begin{array}{c} R_2 - R_1 \end{array} \begin{array}{ccc|ccc}
1 & 0 & -1 & 1 & 0 & -2 \\
0 & 1 & 3 & -1 & 1 & 2 \\
0 & 1 & 2 & 0 & 0 & 1
\end{array} \\
\\
\begin{array}{c} R_3 - R_2 \end{array} \begin{array}{ccc|ccc}
1 & 0 & -1 & 1 & 0 & -2 \\
0 & 1 & 3 & -1 & 1 & 2 \\
0 & 0 & -1 & 1 & -1 & -1
\end{array} \\
\\
\begin{array}{c} R_1 - R_3 \end{array} \begin{array}{ccc|ccc}
1 & 0 & 0 & 0 & 1 & -1 \\
0 & 1 & 3 & -1 & 1 & 2 \\
0 & 0 & -1 & 1 & -1 & -1
\end{array} \\
\\
\begin{array}{c} -R_3 \end{array} \begin{array}{ccc|ccc}
1 & 0 & 0 & 0 & 1 & -1 \\
0 & 1 & 3 & -1 & 1 & 2 \\
0 & 0 & 1 & -1 & 1 & 1
\end{array} \\
\\
\begin{array}{c} R_2 - 3R_3 \end{array} \begin{array}{ccc|ccc}
1 & 0 & 0 & 0 & 1 & -1 \\
0 & 1 & 0 & 2 & -2 & -1 \\
0 & 0 & 1 & -1 & 1 & 1
\end{array}
\end{array}$$

Thus,

$$B^{-1} = \begin{bmatrix} 0 & 1 & -1 \\ 2 & -2 & -1 \\ -1 & 1 & 1 \end{bmatrix}$$

We perform the same procedure on following system to find C^{-1} .

$$\begin{array}{c}
\begin{array}{ccc|ccc}
3 & 1 & 0 & 1 & 0 & 0 \\
1 & -1 & 2 & 0 & 1 & 0 \\
1 & 1 & 1 & 0 & 0 & 1
\end{array} \\
\\
\begin{array}{c} -R_2 \end{array} \begin{array}{ccc|ccc}
3 & 1 & 0 & 1 & 0 & 0 \\
-1 & 1 & -2 & 0 & -1 & 0 \\
1 & 1 & 1 & 0 & 0 & 1
\end{array}
\end{array}$$

$$R_1 - 2R_3 \quad \begin{array}{ccc|ccc} 1 & -1 & -2 & 1 & 0 & -2 \\ -1 & 1 & -2 & 0 & -1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array}$$

$$R_3 - R_1 \quad \begin{array}{ccc|ccc} 1 & -1 & -2 & 1 & 0 & -2 \\ -1 & 1 & -2 & 0 & -1 & 0 \\ 0 & 2 & 3 & -1 & 0 & 3 \end{array}$$

$$R_1 + \frac{1}{2}R_3 \quad \begin{array}{ccc|ccc} 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 & -\frac{1}{2} \\ -1 & 1 & -2 & 0 & -1 & 0 \\ 0 & 2 & 3 & -1 & 0 & 3 \end{array}$$

$$R_2 + R_1 \quad \begin{array}{ccc|ccc} 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 & -\frac{1}{2} \\ 0 & 1 & -\frac{3}{2} & \frac{1}{2} & -1 & -\frac{1}{2} \\ 0 & 2 & 3 & -1 & 0 & 3 \end{array}$$

$$\frac{1}{8}(R_3 - 2R_2) \quad \begin{array}{ccc|ccc} 1 & 0 & -\frac{1}{2} & \frac{1}{2} & 0 & -\frac{1}{2} \\ 0 & 1 & -\frac{3}{2} & \frac{1}{2} & -1 & -\frac{1}{2} \\ 0 & 0 & 1 & -\frac{1}{4} & \frac{1}{4} & \frac{1}{2} \end{array}$$

$$\begin{array}{l} R_1 - \frac{1}{2}R_3 \\ R_2 + \frac{3}{2}R_3 \end{array} \quad \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{3}{8} & \frac{1}{8} & -\frac{1}{4} \\ 0 & 1 & 0 & -\frac{1}{8} & -\frac{3}{8} & \frac{3}{4} \\ 0 & 0 & 1 & -\frac{1}{4} & \frac{1}{4} & \frac{1}{2} \end{array}$$

Thus,

$$C^{-1} = \begin{bmatrix} \frac{3}{8} & \frac{1}{8} & -\frac{1}{4} \\ -\frac{1}{8} & -\frac{3}{8} & \frac{3}{4} \\ -\frac{1}{4} & \frac{1}{4} & \frac{1}{2} \end{bmatrix} = \frac{1}{8} \begin{bmatrix} 3 & 1 & -2 \\ -1 & -3 & 6 \\ -2 & 2 & 4 \end{bmatrix}$$

(5) We calculate the determinant by an expansion on the first row where

$$\det = c_{11} \times 3 + c_{12} \times 0 + c_{13} \times 2$$

$$c_{11} = (-1)^{1+1} \begin{vmatrix} 1 & 1 \\ 1 & 4 \end{vmatrix} = 1(1 \times 4 - 1 \times 1) = 3$$

$$c_{13} = (-1)^{1+3} \begin{vmatrix} -1 & 1 \\ 0 & 1 \end{vmatrix} = 1(-1 \times 1 - 1 \times 0) = -1$$

$$\Rightarrow \det = 3 \times 3 + 0 - 1 \times 2 = 7$$

Expanding by the third row we find

$$\det = c_{31} \times 0 + c_{32} \times 1 + c_{33} \times 4$$

$$\begin{aligned}
c_{32} &= (-1)^{3+2} \begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} = -1(3 \times 1 + 2 \times 1) = -5 \\
c_{33} &= (-1)^{3+3} \begin{vmatrix} 3 & 0 \\ -1 & 1 \end{vmatrix} = 1(3 \times 1 + 0 \times 1) = 3 \\
&\Rightarrow \det = 0 - 5 \times 1 + 3 \times 4 = 7
\end{aligned}$$

Thus we have verified that the two expansions yield the same determinant.

- (6) We construct the following system to find an interpolating polynomial of the form $p(x) = a_0 + a_1x + a_2x^2$ for the given data points.

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \cdot \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 16 \end{bmatrix}$$

We then solve for the unknown coefficients a_0, a_1 and a_2 using Cramer's Rule. Let

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 12 \\ 15 \\ 16 \end{bmatrix}$$

Then,

$$A_0 = \begin{bmatrix} 12 & 1 & 1 \\ 15 & 2 & 4 \\ 16 & 3 & 9 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 1 & 12 & 1 \\ 1 & 15 & 4 \\ 1 & 16 & 9 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 1 & 1 & 12 \\ 1 & 2 & 15 \\ 1 & 3 & 16 \end{bmatrix}$$

The determinants of each 3×3 matrix are given below.

$$|A| = 2 (\neq 0), \quad |A_0| = 14, \quad |A_1| = 12, \quad |A_2| = -2$$

Hence,

$$\begin{aligned}
a_0 &= \frac{|A_0|}{|A|} = \frac{14}{2} = 7 \\
a_1 &= \frac{|A_1|}{|A|} = \frac{12}{2} = 6 \\
a_2 &= \frac{|A_2|}{|A|} = -\frac{2}{2} = -1
\end{aligned}$$

The interpolating polynomial is therefore

$$p(x) = 7 + 6x - x^2$$