

# 67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H ratio

K. Altwegg,<sup>1\*</sup> H. Balsiger,<sup>1</sup> A. Bar-Nun,<sup>2</sup> J. J. Berthelier,<sup>3</sup> A. Bieler,<sup>1,4</sup> P. Bochsler,<sup>1</sup> C. Briois,<sup>5</sup> U. Calmonte,<sup>1</sup> M. Combi,<sup>4</sup> J. De Keyser,<sup>6</sup> P. Eberhardt,<sup>1†</sup> B. Fiethe,<sup>7</sup> S. Fuselier,<sup>8</sup> S. Gasc,<sup>1</sup> T. I. Gombosi,<sup>4</sup> K. C. Hansen,<sup>4</sup> M. Hässig,<sup>1,8</sup> A. Jäckel,<sup>1</sup> E. Kopp,<sup>1</sup> A. Korth,<sup>9</sup> L. LeRoy,<sup>1</sup> U. Mall,<sup>9</sup> B. Marty,<sup>10</sup> O. Mousis,<sup>11</sup> E. Neefs,<sup>6</sup> T. Owen,<sup>12</sup> H. Rème,<sup>13,14</sup> M. Rubin,<sup>1</sup> T. Sémon,<sup>1</sup> C.-Y. Tzou,<sup>1</sup> H. Waite,<sup>8</sup> P. Wurz<sup>1</sup>

The provenance of water and organic compounds on Earth and other terrestrial planets has been discussed for a long time without reaching a consensus. One of the best means to distinguish between different scenarios is by determining the deuterium-to-hydrogen (D/H) ratios in the reservoirs for comets and Earth's oceans. Here, we report the direct in situ measurement of the D/H ratio in the Jupiter family comet 67P/Churyumov-Gerasimenko by the ROSINA mass spectrometer aboard the European Space Agency's Rosetta spacecraft, which is found to be  $(5.3 \pm 0.7) \times 10^{-4}$ —that is, approximately three times the terrestrial value. Previous cometary measurements and our new finding suggest a wide range of D/H ratios in the water within Jupiter family objects and preclude the idea that this reservoir is solely composed of Earth ocean-like water.

The delivery of water and organic compounds to Earth and other terrestrial planets is still under debate (1–4). Existing scenarios range from negligible (1, 2) to substantial (3, 4) cometary contributions to terrestrial water. Hence, the comparison of the deuterium-to-hydrogen ratio (D/H) in water between the different populations of comets and Earth's oceans is crucial if one wants to distinguish among

these scenarios. Previous D/H measurements have been made for a dozen comets from the Oort cloud and the Jupiter family [for example, (5) and references therein]. So far, only one measurement has been made in situ: in the coma of the Oort-cloud comet 1P/Halley, via the mass spectrometers present aboard the European Space Agency (ESA) Giotto spacecraft, and based on an assumption made on the oxygen isotopic composition (6, 7). Here, we report the direct in situ measurement of the D/H ratio in the Jupiter family comet (JFC) 67P/Churyumov-Gerasimenko.


The mass spectrometer ROSINA-DFMS (Rosetta Orbiter Sensor for Ion and Neutral Analysis, Double Focusing Mass Spectrometer) on the Eu-

ropean cometary space mission Rosetta is designed to measure isotopic ratios (8). Its mass resolution and high dynamic range enable it to detect very rare species such as HD<sup>18</sup>O relative to the most abundant isotope H<sub>2</sub><sup>16</sup>O (9). ROSINA has the capability to measure all isotopic ratios in water independently (D/H, <sup>17</sup>O/<sup>16</sup>O, and <sup>18</sup>O/<sup>16</sup>O), and the D/H ratio can be deduced from two different species, namely HD<sup>16</sup>O/H<sub>2</sub><sup>16</sup>O and HD<sup>18</sup>O/H<sub>2</sub><sup>18</sup>O.

Rosetta has a neutral gaseous background due to spacecraft outgassing. The permanent particle density in the close vicinity of the spacecraft far away from the comet is  $\sim 10^6$  cm<sup>-3</sup>, consisting mostly of water but also of organic material, fragments of hydrazine, and vacuum grease (fluorine). Even after 10 years in space and after hibernation, the background from Rosetta can be measured and characterized with ROSINA (10). The D/H ratio in water outgassed from the Rosetta spacecraft is compatible with the terrestrial value of  $1.5 \times 10^{-4}$ , as expected (9), and did not vary with time of degassing, indicating negligible isotope fractionation. These observations demonstrate the capability of DFMS and of the analysis method. On 1 August 2014, Rosetta was within 1000 km from comet 67P/Churyumov-Gerasimenko at 3.6 astronomical units (AU) from the Sun, and the coma was still hidden beneath the spacecraft background for in situ measurements. However, a few days later it approached the comet to within 100 km. The factor of 100 larger densities at this distance surpassed the spacecraft background by more than a factor of 2; the difference between background (26 May) and coma (22 August) at mass 19 dalton is evident (Fig. 1). There are four peaks on mass 19 dalton: fluorine, which is due to background (vacuum grease) from the spacecraft (10); H<sup>18</sup>O, a fragment from H<sub>2</sub><sup>18</sup>O due to electron impact ionization in the instrument and a minor contribution from photodissociation of water in the coma; H<sub>2</sub><sup>17</sup>O; and HD<sup>16</sup>O. All mass peaks have the same shape. H<sub>2</sub><sup>17</sup>O is hidden in the shoulder of


<sup>1</sup>Physikalisches Institut, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland. <sup>2</sup>Department of Geosciences, Tel-Aviv University, Ramat-Aviv, Tel-Aviv, Israel. <sup>3</sup>Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), 4 Avenue de Neptune, F-94100 Saint-Maur, France. <sup>4</sup>Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, 2455 Hayward, Ann Arbor, MI 48109, USA. <sup>5</sup>Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR 6115 CNRS–Université d'Orléans, France. <sup>6</sup>Space Physics Division, Belgisch Instituut voor Ruimte-Aëronomie (BIRA)–Institut d'Aéronomie Spatiale de Belgique (IASB), Ringlaan 3, B-1180 Brussels, Belgium. <sup>7</sup>Institute of Computer and Network Engineering (IDA), Technische Universität Braunschweig, Hans-Sommer-Straße 66, D-38106 Braunschweig, Germany. <sup>8</sup>Department of Space Science, Southwest Research Institute, 6220 Culebra Road, San Antonio, TX 78228, USA. <sup>9</sup>Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany. <sup>10</sup>Centre de Recherches Pétrographiques et Géochimiques, Centre de Recherches Pétrographiques et Géochimiques (CRPG)–CNRS, Université de Lorraine, 15 rue Notre Dame des Pauvres, BP 20, 54501 Vandœuvre lès Nancy, France. <sup>11</sup>Université de Franche-Comté, Institut Univers, Transport, Interfaces, Nanostructures, Atmosphère et Environnement, Molécules (UTINAM), CNRS/Institut National des Sciences de l'Univers (INSU), UMR 6213 Besançon Cedex, France. <sup>12</sup>Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA. <sup>13</sup>Université de Toulouse, Université Paul Sabatier (UPS)–Observatoire Midi-Pyrénées (OMP), Institut de Recherche en Astrophysique et Planétologie (IRAP), Toulouse, France. <sup>14</sup>CNRS, IRAP, 9 Avenue du Colonel Roche, BP 44346, F-31028 Toulouse cedex 4, France. \*Corresponding author. E-mail: altwegg@space.unibe.ch †Deceased.

**Fig. 1. Typical mass/charge 19 daltons/e spectra from DFMS data are shown for 26 May 2014, when the spacecraft was at 800,000 km from the nucleus, and for 22 August, with the spacecraft at 60 km from the nucleus.** The integration time was 20 s. The number of ions detected are plotted as a function of  $m/z$ . The peak fits for 26 August are also shown for fluorine (F), H<sup>18</sup>O, H<sub>2</sub><sup>17</sup>O, and HDO, using the same peak width for all species.



**Fig. 2. Ratios of D/H,  $^{17}\text{O}/^{16}\text{O}$ , and  $^{18}\text{O}/^{16}\text{O}$  for 4 and 5 September.**

The period of 12 hours corresponds almost to a comet rotation (12.4-hour period). The times given are those for the measurement of  $m/z$  18 ( $\text{H}_2^{16}\text{O}$ ),  $m/z$  19 ( $\text{HDO}$  and  $\text{H}_2^{17}\text{O}$ ) and 20 ( $\text{H}_2^{18}\text{O}$ ) are measured 30 s and 1 min later, respectively. The error bars represent statistical errors from the low count rates and errors from the fit. Errors due to background and due to uncertainties in the detector gain, which are of a systematic nature, are only considered for the mean ratios given in the text.



the much larger peak of HDO. Because the positions of the masses are known, the only fit parameters remaining are the amplitudes. In this way,  $\text{H}_2^{17}\text{O}$  can clearly be separated from  $\text{HD}^{16}\text{O}$ .

Before 5 August, all four species had a very similar intensity. However, on 8 August the intensity of  $\text{HD}^{16}\text{O}$  was more than double the height of the  $^{15}\text{OH}$  peak, whereas fluorine stayed constant. This can only be attributed to a much higher D/H ratio in water. By 22 August, the background was almost negligible because the spacecraft was now within 50 km from the nucleus. However, the signal on mass 21 ( $\text{HD}^{18}\text{O}$ ) was still very low. Water peaks on mass 18 and 19 dalton were therefore analyzed by using more than 50 spectra taken between 8 August and 5 September, leading to a derivation of D/H from  $\text{HD}^{16}\text{O}/\text{H}_2^{16}\text{O}$  [analysis is described in (9)]. Uncertainties of the measurements are carefully estimated by error propagation, taking into account statistical uncertainties in the measured signal and uncertainties originating from calibration, background subtraction, and fitting methods. The biggest contribution is probably the uncertainty for the background because Rosetta is now permanently in the cometary coma, and background corrections have to be done with data from before August 2014. Analysis of the spacecraft background over the 10 years of the cruise phase, however, has shown that for a stable spacecraft attitude, the background remains stable over very long times outside of reaction wheel offloadings (10). The value derived from our analysis for D/H is  $(5.3 \pm 0.7) \times 10^{-4}$  (2 $\sigma$  error, where  $\sigma$  is the SD as described above).

By deriving HDO relative to  $\text{H}_2^{16}\text{O}$ , we also found the  $^{17}\text{O}/^{16}\text{O}$  ratio. Additionally, the ratio of  $^{18}\text{O}/^{16}\text{O}$  follows from the analysis of mass-to-charge ratio ( $m/z$ ) = 20 daltons, which contains two well-separated peaks:  $\text{H}_2^{18}\text{O}$  (20.0148 daltons) and HF (20.0062 daltons) (Fig. 2). HF is almost entirely due to spacecraft background. The results for the oxygen isotopic ratios in cometary water are compatible with solar system values,

with  $^{17}\text{O}/^{16}\text{O} = (3.7 \pm 0.9) \times 10^{-4}$  and  $^{18}\text{O}/^{16}\text{O} = (1.8 \pm 0.2) \times 10^{-3}$ . Although from the figure one might see a small modulation for the ratios as a function of rotation of the comet, statistics are too poor to come to a conclusion. Once the comet activity increases, it should be feasible to narrow down the values for the heavy isotopes of water.


The D/H ratio shows dramatic variations among solar system reservoirs of water (Fig. 3). The protosolar nebula (PSN) D/H value is estimated to be  $(2.1 \pm 0.5) \times 10^{-5}$  based on measurements of  $\text{H}_2$  in the atmosphere of Jupiter (11) and ( $^3\text{He}+\text{D}$ )/H in the solar photosphere (12). This value is close to interstellar D/H ratios of  $\text{H}_2$  around  $2.0 \times 10^{-5}$  to  $2.3 \times 10^{-5}$  (13). In contrast, most solar system objects are enriched in deuterium (Fig. 3), with an enrichment factor  $f$  (defined as the ratio  $[\text{D}/\text{H}]_{\text{object}}/[\text{D}/\text{H}]_{\text{PSN}}$ ) averaging 6 for the inner solar system (including Earth, the Moon, and volatile-rich primitive meteorites such as carbonaceous chondrites). Comets analyzed so far, mostly long-period ones, display higher  $f$  values, typically in the 10 to 20 range. The cause of the deuterium enrichment in solar system bodies is usually attributed to water-ice rich in deuterium infalling from the presolar cloud onto the nebula disk (5). Thereafter, because comets may have accreted ice with various chemical histories (14), several mechanisms have been proposed that would induce a deuterium fractionation in the early PSN. Because a part of the ice accreted by comets could have vaporized and recondensed within the PSN, an isotopic exchange could have occurred between the initially deuterium-rich water and molecular hydrogen in the warm regions of the disk (15). At low temperatures, this reaction favors the concentration of deuterium in HDO, but the extremely slow kinetics tend to inhibit the reaction. In these models, isotopic exchange occurs as long as  $\text{H}_2\text{O}$  does not crystallize, implying that the observed D/H ratios should be representative of the local values where and when the building blocks of the host objects condensed (16). Alter-

natively, a part of the ice accreted by comets could have remained pristine (14). Under these circumstances, gas-grain reactions could have induced deuterium fractionation in the cold outer part of the PSN (17, 18). Regardless of the fractionation mechanism, all of these models are consistent with a deuterium enrichment profile following a radial increase throughout the PSN from low values close to the Sun to high values in the outer part of the disk (16–18).

Most D/H ratio measurements in water in comets come from long-period comets, presumably originating from the Oort cloud [Oort Cloud Comets (OCCs)]. Population of this cometary reservoir is attributed to the scattering of icy bodies originally located in the Uranus-Neptune formation region between  $\sim 10$  and 15 AU in the PSN (19), although a nonsolar, external origin for a large fraction of OCCs has recently been proposed (20). In contrast, JFCs are expected to have formed in the Kuiper Belt region beyond Neptune (21). D/H ratios of OCCs, analyzed either in situ by means of mass spectrometry in the case of comet Halley (6, 7) or spectroscopy (5), show a range varying from  $\sim 1.3$  to 2.9 times the terrestrial value ( $f \sim 9.8$  to 21.9) (Fig. 3). In addition, the D/H ratio was found to be similar to the comet Halley in in situ measurements in the water plume of Saturn's satellite Enceladus (22). These values support the predicted D/H radial increase with distance from the Sun and the origin of OCCs from a common, localized region of the disk (16–18).

Recent D/H measurements in water in the two JFCs analyzed so far—namely,  $(1.61 \pm 0.24) \times 10^{-4}$  ( $f \sim 7.8$ ) for 103P/Hartley 2 (23) and an upper limit of  $2.0 \times 10^{-4}$  ( $f < 9.5$ ) for 45P/Honda–Mrkos–Pajdušáková (5)—contradict this view. The hydrogen isotope composition of 103P/Hartley 2 is closer to the terrestrial value than the OCCs average, reviving the possibility of a cometary, rather than asteroidal, origin for the oceans. These data lead to two possible conclusions: Either JFCs originate from the Kuiper Belt and the chemical models developed so far (16–18) are not representative, or these comets formed over a wide range of heliocentric distances in the outer part of the PSN. With regard to the first possibility, a recent chemical model leading to a non-monotonic  $f$  profile throughout the PSN (24) matches these observations if the JFCs were formed in the Kuiper Belt (21). In contrast with previous PSN models evolving as closed systems, this model assumes that the disk continues to be fed by material infalling from the presolar cloud. Alternatively, it has been proposed that JFCs and OCCs could originate from the same extended outer region of the PSN (25), so that 103P/Hartley 2 and 45P/Honda–Mrkos–Pajdušáková (45P/HMP) may simply have formed in the inner part of this common reservoir. In this case, the range of D/H ratios measured in JFCs should be similar to the one found in OCCs, as suggested by observations.

The new D/H value of  $(5.3 \pm 0.7) \times 10^{-4}$  ( $f \sim 25.2$ ) from comet 67P/Churyumov–Gerasimenko is not consistent with previous JFC data and is even higher than values characteristic of OCCs ( $\sim 30$



**Fig. 3. D/H ratios in different objects of the solar system.** Data are from (1, 2, 5–7, 26–28) and references therein. Diamonds represent data obtained by means of in situ mass spectrometry measurements, and circles refer to data obtained with astronomical methods.

to 120% higher than that of comet Halley). In contrast to previous JFC measurements, this estimate matches models that predict a monotonic radial increase of the enrichment profile (16–18). From the ROSINA measurements on comet 67P/Churyumov-Gerasimenko, we conclude that the D/H values of JFCs may be highly heterogeneous, possibly reflecting the diverse origins of JFCs. If this is the case, then the new measurement supports models advocating an asteroidal (carbonaceous chondrite-like) rather than cometary origin for the oceans, and by extension for the terrestrial atmosphere (1, 2).

## REFERENCES AND NOTES

1. B. Marty, The origins and concentrations of water, carbon, nitrogen and noble gases on Earth. *Earth Planet. Sci. Lett.* **313**, 56–66 (2012). doi: [10.1016/j.epsl.2011.10.040](https://doi.org/10.1016/j.epsl.2011.10.040)
2. C. M. O. Alexander *et al.*, The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets. *Science* **337**, 721–723 (2012). doi: [10.1126/science.1223474](https://doi.org/10.1126/science.1223474); pmid: 22798405
3. N. Dauphas, The dual origin of the terrestrial atmosphere. *Icarus* **165**, 326–339 (2003). doi: [10.1016/S0019-1035\(03\)00198-2](https://doi.org/10.1016/S0019-1035(03)00198-2)
4. T. Owen, A. Bar-Nun, I. Kleinfeld, Possible cometary origin of heavy noble gases in the atmospheres of Venus, Earth and Mars. *Nature* **358**, 43–46 (1992). doi: [10.1038/358043a0](https://doi.org/10.1038/358043a0); pmid: 11536499
5. C. Ceccarelli, P. Caselli, D. Bockelée-Morvan, O. Mousis, S. Pizzarello, F. Robert, D. Semenov, Deuterium fractionation: the Ariadne's thread from the pre-collapse phase to meteorites and comets today. *Protostars and Planets VI* (2014); available at <http://arxiv.org/pdf/1403.7143v1.pdf>.

6. H. Balsiger, K. Altwegg, J. Geiss, D/H and O-18/O-16 ratio in the hydronium ion and in neutral water from in situ ion measurements in comet Halley. *J. Geophys. Res.* **100** (A4), 5827–5834 (1995). doi: [10.1029/94JA02936](https://doi.org/10.1029/94JA02936)
7. P. Eberhardt, M. Reber, D. Krankowsky, R. R. Hodges, The D/H and  $^{18}\text{O}/^{16}\text{O}$  ratios in water from comet P/Halley. *Astron. Astrophys.* **302**, 301 (1995).
8. H. Balsiger *et al.*, ROSINA Rosetta Orbiter Spectrometer for Ion and Neutral Analysis. *Space Sci. Rev.* **128**, 745–801 (2002). doi: [10.1007/s11214-006-8335-3](https://doi.org/10.1007/s11214-006-8335-3)
9. M. Hässig *et al.*, ROSINA/DFMS capabilities to measure isotopic ratios in water at comet 67P/Churyumov-Gerasimenko. *Planet. Space Sci.* **84**, 148–152 (2013). doi: [10.1016/j.pss.2013.05.014](https://doi.org/10.1016/j.pss.2013.05.014)
10. B. Schläppli *et al.*, Influence of spacecraft outgassing on the exploration of tenuous atmospheres with in situ mass spectrometry. *J. Geophys. Res.* **115** (A12), A12313 (2012). doi: [10.1029/2010JA015734](https://doi.org/10.1029/2010JA015734)
11. H. F. Levison, M. J. Duncan, R. Brasser, D. E. Kaufmann, Capture of the Sun's Oort cloud from stars in its birth cluster. *Science* **329**, 187–190 (2010). doi: [10.1126/science.1187535](https://doi.org/10.1126/science.1187535); pmid: 20538912
12. P. R. Mahaffy, T. M. Donahue, S. K. Atreya, T. Owen, H. B. Niemann, Galileo Probe measurements of D/H and  $3\text{He}/4\text{He}$  in Jupiter's atmosphere. *Space Sci. Rev.* **84**, 251–263 (1998). doi: [10.1023/A:1005091806594](https://doi.org/10.1023/A:1005091806594)
13. J. Geiss, G. Gloeckler, Abundances of deuterium and helium-3 in the protosolar cloud. *Space Sci. Rev.* **84**, 239–250 (1998). doi: [10.1023/A:1005039822524](https://doi.org/10.1023/A:1005039822524)
14. T. Prodanovic, G. Steigman, B. D. Fields, The deuterium abundance in the local interstellar medium. *Mon. Not. R. Astron. Soc.* **406**, 1108–1115 (2010).
15. R. Visser, S. D. Doty, E. F. van Dishoeck, The chemical history of molecules in circumstellar disks. II. Gas-phase species. *Astron. Astrophys.* **534**, A132 (2011). doi: [10.1051/0004-6361/201117249](https://doi.org/10.1051/0004-6361/201117249)
16. J. Geiss, H. Reeves, Deuterium in the solar system. *Astron. Astrophys.* **93**, 189–199 (1981).

17. J. J. Kavelaars, O. Mousis, J.-M. Petit, J. A. Weaver, On the formation location of Uranus and Neptune as constrained by dynamical and chemical models of comets. *Astrophys. J.* **734**, L30 (2011). doi: [10.1088/2041-8205/734/2/L30](https://doi.org/10.1088/2041-8205/734/2/L30)
18. K. Furuya, Y. Aikawa, H. Nomura, F. Hersant, V. Wakelam, Water in protoplanetary disks: Deuteration and turbulent mixing. *Astrophys. J.* **779**, 11 (2013). doi: [10.1088/0004-637X/779/1/11](https://doi.org/10.1088/0004-637X/779/1/11)
19. T. Albertsson, D. Semenov, T. Henning, Chemodynamical deuterium fractionation in the early solar nebula: The origin of water on Earth and in asteroids and comets. *Astrophys. J.* **784**, 39 (2014). doi: [10.1088/0004-637X/784/1/39](https://doi.org/10.1088/0004-637X/784/1/39)
20. L. Dones, P. R. Weissman, H. F. Levison, M. J. Duncan, Oort cloud formation and dynamics. *Comets II*, 153–174 (2004).
21. M. J. Duncan, H. F. Levison, A disk of scattered icy objects and the origin of Jupiter-family comets. *Science* **276**, 1670–1672 (1997). doi: [10.1126/science.276.5319.1670](https://doi.org/10.1126/science.276.5319.1670); pmid: 9180070
22. J. H. Waite Jr. *et al.*, Liquid water on Enceladus from observations of ammonia and  $40\text{Ar}$  in the plume. *Nature* **460**, 487–490 (2009). doi: [10.1038/nature08153](https://doi.org/10.1038/nature08153)
23. P. Hartogh *et al.*, Ocean-like water in the Jupiter-family comet 103P/Hartley 2. *Nature* **478**, 218–220 (2011). doi: [10.1038/nature10519](https://doi.org/10.1038/nature10519); pmid: 21976024
24. L. Yang, F. J. Ciesla, C. M. O. Alexander, The D/H ratio of water in the solar nebula during its formation and evolution. *Icarus* **226**, 256–267 (2013). doi: [10.1016/j.icarus.2013.05.027](https://doi.org/10.1016/j.icarus.2013.05.027)
25. R. Brasser, A. Morbidelli, Oort cloud and Scattered Disc formation during a late dynamical instability in the Solar System. *Icarus* **225**, 40–49 (2013). doi: [10.1016/j.icarus.2013.03.012](https://doi.org/10.1016/j.icarus.2013.03.012)
26. J. Aléon, J., Multiple origins of nitrogen isotopic anomalies in meteorites and comets. *Astrophys. J.* **722**, 1342–1351 (2010). doi: [10.1088/0004-637X/722/2/1342](https://doi.org/10.1088/0004-637X/722/2/1342)
27. D. Bockelée-Morvan *et al.*, Herschel measurements of the D/H and  $^{16}\text{O}/^{18}\text{O}$  ratios in water in the Oort-cloud comet C/2009 P1 (Garradd). *Astron. Astrophys.* **544**, L15 (2012). doi: [10.1051/0004-6361/201219744](https://doi.org/10.1051/0004-6361/201219744)
28. H. Feuchtgruber *et al.*, The D/H ratio in the atmospheres of Uranus and Neptune from Herschel PACS observations. *Astron. Astrophys.* **551**, A26 (2013).

## ACKNOWLEDGMENTS

A. Morbidelli is acknowledged for helpful discussion. The authors thank the following institutions and agencies, which supported this work: Work at the University of Bern was funded by the State of Bern, the Swiss National Science Foundation, and the ESA PRODEX Program. Work at Max-Planck-Institut für Sonnensystemforschung was funded by the Max Planck Society and BMWI under contract 50QP1302. Work at Southwest Research Institute was supported by subcontract 1496541 from the Jet Propulsion Laboratory. Work at BIRA-IASB was supported by the Belgian Science Policy Office via PRODEX/ROSINA PEA 90020. This work was supported by Centre National d'Études Spatiales grants at IRAP, LATMOS, LPC2E, UTINAM, CRPG, and by the the European Research Council (grant 267255). A.B.-N. thanks the Ministry of Science and the Israel Space agency. Work at the University of Michigan was funded by NASA under contract JPL-1266313. ROSINA would not give such outstanding results without the work of the many engineers and technicians involved in this instrument and in the mission over the years whose contributions are gratefully acknowledged. Rosetta is an ESA mission with contributions from its member states and NASA. The work of the ESA Rosetta team is herewith acknowledged. All ROSINA data are available on request until they are released to the PSA archive of ESA and to the PDS archive of NASA.

1 October 2014; accepted 1 December 2014  
Published online 11 December 2014;  
[10.1126/science.1261952](https://doi.org/10.1126/science.1261952)



**67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H ratio**

K. Altwegg, H. Balsiger, A. Bar-Nun, J. J. Berthelier, A. Bieler, P. Bochslers, C. Briois, U. Calmonte, M. Combi, J. De Keyser, P. Eberhardt, B. Fiethe, S. Fuselier, S. Gasc, T. I. Gombosi, K.C. Hansen, M. Hässig, A. Jäckel, E. Kopp, A. Korth, L. LeRoy, U. Mall, B. Marty, O. Mousis, E. Neefs, T. Owen, H. Rème, M. Rubin, T. Sémon, C.-Y. Tzou, H. Waite and P. Wurz (December 10, 2014) *Science* **347** (6220), . [doi: 10.1126/science.1261952] originally published online December 10, 2014

Editor's Summary

---

This copy is for your personal, non-commercial use only.

---

- |                      |  |
|----------------------|--|
| <b>Article Tools</b> | Visit the online version of this article to access the personalization and article tools:<br><a href="http://science.sciencemag.org/content/347/6220/1261952">http://science.sciencemag.org/content/347/6220/1261952</a> |
| <b>Permissions</b>   | Obtain information about reproducing this article:<br><a href="http://www.sciencemag.org/about/permissions.dtl">http://www.sciencemag.org/about/permissions.dtl</a>  |

*Science* (print ISSN 0036-8075; online ISSN 1095-9203) is published weekly, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. Copyright 2016 by the American Association for the Advancement of Science; all rights reserved. The title *Science* is a registered trademark of AAAS.