



# Methane spikes, background seasonality and non-detections on Mars: A geological perspective

Giuseppe Etiope<sup>a,b,c,\*</sup>, Dorothy Z. Oehler<sup>d</sup>

<sup>a</sup> Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, Roma, Italy

<sup>b</sup> Faculty of Environmental Science and Engineering, Babes-Bolyai University, Cluj-Napoca, Romania

<sup>c</sup> Istituto di Astrofisica e Planetologia Spaziali (INAF-IAPS), Rome, Italy

<sup>d</sup> Planetary Science Institute, Tucson, AZ, USA

## ARTICLE INFO

### Keywords

Mars  
Methane  
Seepage  
Biotic and abiotic origin

## ABSTRACT

The story of methane (CH<sub>4</sub>) on Mars continues to grow. A long series of CH<sub>4</sub> detections (spikes, plumes, seasonal variations) and non-detections in the martian atmosphere suggest that CH<sub>4</sub> is occasionally released from subsurface rocks, but also that an unknown process of rapid CH<sub>4</sub> removal in the atmosphere must exist. The observations can be interpreted integrating concepts of subsurface geological gas source variability (CH<sub>4</sub> seepage flux patterns), atmospheric chemistry (CH<sub>4</sub> removal processes), and atmospheric circulation (CH<sub>4</sub> transport and dilution). While the last two are widely studied, potential gas seepage patterns on Mars have never been discussed. In this work, we use the factors controlling gas seepage flux patterns observed on Earth to address key questions about the significance of the martian detections and non-detections. As on Earth, CH<sub>4</sub> seepage on Mars must occur mainly through advection (non-diffusion) mechanisms, driven by pressure gradients and permeability, typically related to faults and fracture networks, which control release patterns, in terms of intensity, duration and variability. CH<sub>4</sub> release may occur, therefore, whenever gas pressure and/or rock permeability increase (e.g., via destabilization of storage bodies such as clathrates or zeolites; breaching of seals such as permafrost; planetary stresses, impacts, or gas pressure build-up along faults). Rapid atmospheric pressure changes, due to storms and winds, may also play a role. All these forcings can lead to local, episodic, irregular or seasonal gas releases, which may explain the observations acquired so far if relatively rapid (month or day time-scales) mechanisms of CH<sub>4</sub> removal exist in the lower atmosphere or near-surface. The release patterns cannot reveal whether CH<sub>4</sub> is biologic or abiotic. If CH<sub>4</sub> sensors operate discontinuously and only in one site or over limited regions, then, non-detections will be the rule and detections will be relatively rare. The chances of CH<sub>4</sub> detection may increase if the observations are focused and repeated on specific, potential areas of gas release determined by geological analysis.

## 1. Introduction

On Earth, most of the methane (CH<sub>4</sub>) in the atmosphere is a result of biochemical processes, and the possibility that methane in the atmosphere of Mars, so far detected only episodically, might reflect martian microbial life has been a subject of great interest. Since 2003, there have been multiple reports of methane in the martian atmosphere, with detections from orbit around Mars (Formisano et al., 2004; Fonti and Marzo, 2010), from terrestrial telescopes (Krasnopolsky et al., 2004; Mumma et al., 2009) and ground-based measurements in Gale crater (by the Tunable Laser Spectrometer, TLS, on the *Curiosity* rover; Webster et al., 2015, 2018). In particular, *Curiosity* observed methane spikes (up

to ~9 ppbv; Webster et al., 2015; Roos-Serote et al., 2016) and apparently seasonal CH<sub>4</sub> changes (~0.2–0.7 ppbv; Webster et al., 2018). Telescopic measurements detected up to ~45 ppbv, equivalent to plumes of 19,000 metric tons of CH<sub>4</sub>; Mumma et al. (2009). There is a continuing debate on the validity of such detections (Zahnle et al., 2011; Villanueva et al., 2013; Webster et al., 2015), as CH<sub>4</sub> signals have been attributed to erroneous retrieval processes or contamination. However, non-detections have also been reported from ground-based observations by *Curiosity* (Webster et al., 2013), from telescopic follow-ups by Villanueva et al. (2013) to the earlier detections of methane plumes over Mars (Mumma et al., 2009), and from more recent analyses using the Echelon-Cross-Echelle Spectrograph onboard the Stratospheric

\* Corresponding author. Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, Roma, Italy.

E-mail address: [giuseppe.etiope@ingv.it](mailto:giuseppe.etiope@ingv.it) (G. Etiope).

<https://doi.org/10.1016/j.pss.2019.02.001>

Received 28 September 2018; Received in revised form 19 December 2018; Accepted 1 February 2019

Available online 14 February 2019

0032-0633/© 2019 Elsevier Ltd. All rights reserved.

Observatory for Infrared Astronomy (Aoki et al., 2018). Martian atmospheric chemistry predicts that methane is destroyed by excited oxygen atoms,  $O(^1D)$ , and OH, resulting in a methane lifetime of about 300 years (Atreya et al., 2007), but this is not rapid enough to explain the  $CH_4$  variations observed so far. Faster removal processes could be related to  $CH_4$  adsorption onto quartz grains (Knak-Jensen et al., 2014), interactions with zeolites in atmospheric dust (Holmes et al., 2015) or oxidation by hydrogen peroxide in the regolith, with a resulting shorter lifetime of 200 days or even a few hours near the surface (Lefevre and Forget, 2009).  $O_2$  depletions temporally correlated with  $CH_4$  would support this latter model (McConnochie et al., 2014). This is a subject of continuing research. Concerning the source of methane, most of authors agree that  $CH_4$  observed in the martian atmosphere is produced in the subsurface, within rocks (e.g., Max et al., 2013; Blamey et al., 2015; Viscardy et al., 2016; Oehler and Etiope, 2017; Webster et al., 2018), and not in the atmosphere by exogenous processes, as suggested, for example, by hypotheses of meteor showers (Fries et al., 2016), electrical discharge (Robledo-Martinez et al., 2012) and ultraviolet irradiation of organics (Keppler et al., 2012).

The  $CH_4$ -observation history for Mars includes non-detections, variable background concentrations with apparently seasonal oscillations, and episodic spikes. What does all this mean? Methane that is observable in the atmosphere must be a result of a competition between  $CH_4$  source and atmospheric destruction/consumption, modulated by atmospheric transport and dilution. Assuming that martian methane is produced in the subsurface, the variability of the observations can be interpreted only by integrating three types of study: (a)  $CH_4$  release modeling (i.e., geological seepage patterns, in terms of flux intensity, duration, and variability), (b) atmospheric  $CH_4$  sinks (i.e., methane consumption), and (c) atmospheric circulation modeling ( $CH_4$  transport and dilution). Items (b) and (c) have widely been addressed in previous studies (e.g., Atreya et al., 2007; Lefevre and Forget, 2009; Holmes et al., 2015; Viscardy et al., 2016; Pla-Garcia et al., 2018; Neary and Daerden, 2018, and references therein). In this work we focus on item (a), answering the question whether the variability of methane in the martian atmosphere is compatible with geological seepage mechanisms. Specifically, we use factors controlling gas seepage patterns observed on Earth to address two key questions: are the variable and episodic detections on Mars, as well as the non-detections, compatible with geological mechanisms of gas release from subsurface rocks? Can the release variability (seasonality, in particular) provide insight into the origin (microbial, thermogenic or abiotic) of methane?

Subsurface  $CH_4$  generation and seepage phenomena on Earth, and potentially on Mars, were reviewed in Oehler and Etiope (2017). Here, we complete the discussion of possible gas seepage patterns, in the light of the several types of martian  $CH_4$  observations, including the recent report of seasonal variations (Webster et al., 2018). A basic descriptive model of seepage variability is also proposed, which can help in developing new models of the fate of  $CH_4$  in the martian atmosphere. Our analysis will show that all  $CH_4$  observations reported so far, including the non-detections, are compatible with geological gas seepage processes that may occur on Mars, regardless of the  $CH_4$  origin and providing that a mechanism exists for rapid removal of  $CH_4$  from the atmosphere.

## 2. Key concepts of geological $CH_4$ seepage on earth

An extensive discussion of potential methane seepage on Mars is reported in Oehler and Etiope (2017), and we refer the reader to that work also for definitions and terminology on gas origin and seepage (e.g., microbial, thermogenic, biotic, abiotic, macro-seeps, mud volcanoes, microseepage). Here, we review concepts of methane seepage on Earth that are critical for discussion of potential seepage patterns on Mars and interpretation of the atmospheric  $CH_4$  data acquired so far. Specifically, advection (driven by pressure gradients and rock permeability) as the main mechanism of gas motion from subsurface rocks to the surface (Section 2.1), the geological controls on pressure gradients (Section 2.2),

controls on permeability (i.e., faults and fractures; Section 2.3) and the resulting seepage flux patterns in terms of intensity duration and variability (Section 2.4). The application of these processes to Mars is discussed in Section 3.

### 2.1. Mechanism of $CH_4$ seepage: advection or diffusion?

A key point in the discussion on methane source variability on Mars is the recognition that the major gas migration mechanism in subsurface rocks is not diffusion, as often assumed, but advection. This is well known on Earth. Diffusion, the gas motion driven by concentrations gradients, is important only at small spatial scales and over long times in low-permeability, low-porosity rocks (Etiope and Martinelli, 2002). Modeling work has indicated that diffusion cannot be a key mechanism for methane release on Mars and cannot explain the plumes and concentration variations (Stevens et al., 2015, 2017). In contrast, advection is driven by permeability and gas pressure gradients; it is the mechanism leading to relatively fast and long-distance gas migration, including gas seepage to the surface, either as focused emissions (macro-seeps, including gas vents and mud volcanoes) or as weak, diffuse exhalations (microseepage) (Malmqvist and Kristiansson, 1985; Brown, 2000; Etiope and Martinelli, 2002; Etiope, 2015 and references therein).

In geological formulations, advection in a one-dimensional form, along the  $z$  axis is expressed as

$$v = k \Delta P / (\mu Z) \quad (1)$$

Where  $v$  is the gas velocity ( $m \text{ sec}^{-1}$ ),  $k$  is the intrinsic permeability ( $m^2$ ),  $\mu$  is the dynamic gas viscosity ( $kg \text{ m}^{-1} \text{ sec}^{-1}$ ) and  $\Delta P$  is the pressure difference ( $kg \text{ m}^{-1} \text{ sec}^{-2}$ ) between two points spaced at a distance  $Z$  (m). Estimates of gas velocity through faults and fracture systems may be obtained applying the “planar fissure” and “cubic law” relationships, where  $k$  can be expressed as  $b^2/12$  (with  $b$  the fissure width) and  $b^3/6d$  (where  $d$  is the mean distance between intersecting fissures), respectively (Etiope, 2015 and references therein). The relationship between gas seepage and faults, and related geological conditions, is discussed in Section 2.2.

The gas flux ( $F$ ) can then be derived knowing the gas concentration  $C$  ( $kg \text{ m}^{-3}$ ):

$$F = C v \quad (2)$$

Gas advection fluxes can be quite variable over time and can explain the rapid changes observed in surface seeps on Earth, whereas such changes are not compatible with diffusion mechanisms. Importantly, gas advection increases the potential for seepage of gas generated in deep geological sources. It has been considered, for example, that serpentinization on Mars, likely located several kilometers below the surface, could not produce localized gas pulses by diffusion to the surface (Yung and Chen, 2015). But on Earth, methane generated in deep serpentinized rocks clearly reaches the surface through advection along faults (e.g., Etiope et al., 2011, 2016). To be rigorous, then, serpentinization produces  $H_2$ , and  $CH_4$  is produced by subsequent reactions of that  $H_2$  with  $CO_2$  ( $CO_2$  hydrogenation); thus,  $CH_4$  production may occur in rocks that are shallower than those where serpentinization occurs (e.g., Oehler and Etiope, 2017; Etiope et al., 2018).

### 2.2. Controls on pressure gradients

For gas seepage, the pressure gradient in Eq (1) is the key factor that can change over short-time intervals (from instantaneously to a few days or months) and is responsible for seepage variability discussed in Section 2.4. Pressure gradients and their changes can be induced by both endogenous (geological) and exogenous (atmospheric) factors.

#### 2.2.1. Geological controls

Endogenous, geological factors refer to tectonic stresses, variations in

lithostatic loading, rock fracturing, localized gas generation, recharge and discharge of aquifers or fluid reservoirs. Sedimentary basins and their constituent rocks, in particular, are subject to basin loading, compaction, extensional and compressional stresses that change the gas pressure gradients. In the presence of water-saturated rocks, gas can migrate to the surface if its pressure ( $P_G$ ) is above the sum of hydrostatic pressure ( $P_w$ ), given by the height of the piezometric surface, plus capillary pressure ( $P_C$ ) controlled by the interfacial tension of water and pore size. The gas can also fracture the rock if its pressure,  $P_G$ , overcomes the lithostatic pressure,  $P_L$  (Etiope and Martinelli, 2002). Gas seepage is then the result of an interplay between  $P_G$  and  $P_L$ , and  $P_w + P_C$  in presence of aquifers. The build-up of  $P_G$ , the main requisite for gas seepage, implies gas accumulation, and this is the reason why methane seeps are generally linked to subsurface pressurised reservoirs. Earthquakes, through the passage of seismic waves, can then suddenly modify the gas pressure confinement and therefore may trigger advective gas pulses (Mazzini and Etiope, 2017).

### 2.2.2. Atmospheric controls

Methane emission from microseepage (the slow, weak, areal exhalation of gas from the soil, without evident morphological features or fluid manifestations) can be significantly controlled by exogenous (atmospheric) factors, such as barometric pressure and wind intensity. Rapid changes of atmospheric pressure as well as wind velocity and direction are known to affect the release of gases stored in the soil and subsoil, down to depths of several meters. As barometric pressure decreases, gases are drawn upward out of the ground, and when pressure increases, gases are pushed downward, a phenomenon known as “atmospheric pumping” (e.g., Nilson et al., 1991; Massmann and Farrier, 1992). Changes in atmospheric pressure at a site may produce vertical pressure gradients in subsurface gases (Massmann and Farrier, 1992). Barometric changes of 1000–2000 Pa over a period of 1–2 days may produce advective velocities of the order of  $10^{-4}$  cm/sec within a soil with permeability of  $10^{-12}$  m<sup>2</sup> (Clements and Wilkening, 1974). A wind gust can trigger a pulse of gas release via the Bernoulli effect, creating a low-pressure event above the ground (e.g. Reimer and Bowles, 1979; Klusman, 1993). Winds may however also suppress gas release by modifying pressure gradients in the ground in a way that retards vertical gas flow to the surface (Lewicki et al., 2007).

### 2.3. Geological controls on permeability: the role of faults and fracture networks

On Earth, gas seepage is always related to tectonic faults and fracture networks. It occurs especially in fault intersections, regardless of tectonic regime, fault activity and seismicity (Macgregor, 1993; Bonini, 2013; Etiope, 2015; Mazzini and Etiope, 2017). In other words, methane can be released to the surface even in tectonically non-active regions and areas far from seismic zones. Tectonic regimes can be characterized by compressional (reverse and thrust faults), extensional (normal faults), uplift (radial faults) or shear (strike-slip faults) stress. The literature reports numerous examples of methane seepage in different types of faults and tectonic settings on Earth. For example, gas seepage along normal faults occurs in the Katakolo Bay in Greece (Etiope et al., 2013b) and in the Tiber Delta in Italy (Ciotoli et al., 2016); seepages along thrust faults are those typical (but not exclusive) of mud volcanoes worldwide (Bonini, 2013; Mazzini and Etiope, 2017). The Giswil seep (Switzerland) is an example of gas exhalation along a strike-slip fault (Etiope et al., 2010). And gas migration studies demonstrate that non-active, ancient faults in stable terrains (e.g., in granites, within cratons) can be gas-bearing (e.g., Malmqvist and Kristiansson, 1984, 1985). In fact, while compressional and extensional stresses may play a role in controlling water and oil migration (i.e. hydraulic conductivity), they do not significantly affect the gas-bearing property of faults, as a fault that is impermeable to water can be permeable to gas, as happens in argillaceous rocks and granites (e.g., Etiope and Lombardi, 1995; Horseman

et al., 1999; Ciotoli et al., 2005; Etiope and Martinelli, 2002). And as discussed above, gas pressure is the main factor driving seepage even in poorly permeable and non-active (and non-seismic) fracture networks related to faults (Bonini, 2013; Etiope, 2015).

However, seismic events and fault activity may enhance seepage, leading to more intense and episodic emissions. Numerous examples are known of degassing events or variations related to earthquakes, either in mud volcanoes (e.g., Mellors et al., 2007; Kopf et al., 2009; Mazzini and Etiope, 2017 and references therein) or other types of surface seeps (Heinicke and Koch, 2000; Yang et al., 2006). In addition to a modification of gas pressure confinement due to the passage of seismic waves (discussed in Section 2.2), earthquakes can also increase rock permeability through fracturing (Mazzini and Etiope, 2017). This will result in sudden events of degassing.

### 2.4. CH<sub>4</sub> seepage flux patterns: intensity and variability

#### 2.4.1. Methane seepage flux intensity


On Earth, the intensity of methane emission (flux) may span a wide range of values, depending on the type of the seeping structure (gas-oil seep, mud volcano, microseepage from the soil) and the gas pressure and spatial extent (area) of the seep (e.g., Etiope, 2015). Methane release from single emission points (macro-seeps) is typically in the order of  $10^{-1}$  and  $10^4$  tonnes year<sup>-1</sup> (the flux from gas vents with a diameter <1 m is typically between  $10^{-1}$  and  $10^2$  tonnes year<sup>-1</sup>). Individual release events from single macro-seeps may emit orders of grams to several kg of gas in a few minutes (e.g., Etiope et al., 2011; Etiope et al., 2013b). Mud volcanoes can release several tonnes of gas in a few hours (e.g., Mazzini and Etiope, 2017). Diffuse exhalations from the ground (microseepage, over km<sup>2</sup> scale areas) is generally expressed in terms of grams per m<sup>2</sup> per day (typically mg m<sup>-2</sup> d<sup>-1</sup>) and is frequently in the range  $10^0$ – $10^3$  mg m<sup>-2</sup> d<sup>-1</sup> (equivalent to orders of  $10^{-1}$ – $10^2$  tonnes km<sup>-2</sup> year<sup>-1</sup>; e.g., Etiope and Klusman, 2010). These fluxes have been measured for both biotic (thermogenic or microbial) methane in sedimentary basins and abiotic methane in igneous (mafic-ultramafic) rocks. Therefore, significant pressure gradients, and thus gas accumulations (as discussed in section 2.2), can occur in all types of rocks, especially when they are fractured.

#### 2.4.2. Methane seepage flux variability

Methane emission in active gas-oil seeps and mud volcanoes (bubbling pools, gas vents) may significantly change over a wide range of time scales, from minutes to days and years, depending on variations in subsurface gas pressure, which may be cyclic or irregular, and often induced by seismic activity (e.g., Delisle et al., 2010; Yang et al., 2006).

Microbial methanotrophic activity in the soil, modulated by soil temperature, humidity and organic content, determines seasonal variation of methane microseepage, which is higher in dry-winter periods, and lower in wet-summer periods (Etiope and Klusman, 2010). In permafrost regions, where methane of biological or geological origin can accumulate within or below the permafrost layer, seasonal ice-thawing produces summer increases of methane flux (e.g., Christensen et al., 2004; Portnov et al., 2013; Wang et al., 2014). On the basis of numerous observations on macro-seeps and microseepage, the patterns of gas emissions from the ground can be characterized by three main types: continuous, intermittent and episodic, as depicted in Fig. 1.

- (a) continuous emission over long-term (years to geological time scales), with hourly, daily and seasonal weak oscillations; the emission can be one of either
  - high flux, continuous venting (e.g., macro-seeps of hydrocarbons, active mud volcanoes, hydrothermal-volcanic vents, mofettes, fumaroles; e.g., Chiodini et al., 2010; Etiope et al., 2013b) or
  - low flux diffuse microseepage, with weak exhalations over wide areas and with low oscillations (e.g., Klusman et al., 2000; Etiope and Klusman, 2010).



**Fig. 1.** Main methane emission patterns observed in seepage processes on Earth. The patterns are illustrative, summarized from numerous observations reported in the literature (e.g., Klusman et al., 2000; Yang et al., 2006; Chiodini et al., 2010; Delisle et al., 2010; Etiope and Klusman, 2010; Etiope et al., 2013b; Mazzini and Etiope, 2017; Krabbenhöft et al., 2010; Embriaco et al., 2014).

- (b) intermittent emission over a long-term, with stronger and sudden, short-term (seconds, minutes, hours), variations (e.g., repeated gas eruptions, irregular venting in mud volcanoes and gas seeps, geysers; e.g., Delisle et al., 2010; Mazzini and Etiope, 2017). A continuous, background exhalation always accompanies this type of emission.
- (c) episodic emission, with low or strong exhalations, spaced out by periods of days, months, or years with very low or nil emission between the episodes. This pattern is typical of expulsion of gas pockets in shallow subsoil, driven by seismic activity or the Bernoulli effect induced by episodic strong winds and storms, or episodic migration of gas slugs along faults driven by seismic activity, overpressure pulses, and the so-called “fault burps” (e.g., Haney et al., 2005; Krabbenhöft et al., 2010; Embriaco et al., 2014). The events can be singular (one pulse) or multiple (sequence of sub-events), both lasting minutes or hours; a weak continuous, background (quiescent) exhalation may persist.

Patterns (a) and (b) generally imply large and active gas sources (with highly pressurised reservoir rocks). Pattern (c) is more typical of nearly drained seepage systems (generally characterized by old and dying seeps). We consider this pattern as the most likely for Mars, and in the next section, we show that Mars has geological features compatible with an episodic type of seepage.

### 3. Potential seepage intensity and variability on mars

Oehler and Etiope (2017) described in detail the relevant geology on Mars that could affect abiotic methane generation, conduits for gas migration, storage of methane in the subsurface, and triggers for methane release. To summarize, methane generated by any process in the martian subsurface will migrate along permeable conduits, most often provided by fault and fracture systems. That methane will either be released to the atmosphere or stored in clathrates, zeolites, or conventional stratigraphic/structural traps, where it will accumulate. Conventional traps consist of reservoir rocks with effective methane seals, such as shales,

evaporites (salt or anhydrite) or permafrost (Oehler and Etiope, 2017). Anything that destabilizes the clathrates, breaks down the zeolites, or breaches the seals of conventional methane traps, has the potential to contribute to episodic releases. Regions on Mars identified as potential sites for methane seepage in Oehler and Etiope (2017) are summarized in Table 1 of that paper. Of those sites, the mud-volcano-like mounds in deep basins of the lowlands and the faulted and serpentinized areas with mafic-ultramafic rocks (e.g., Nili Fossae) were highlighted as key locations for potential methane release.

Seepage intensity is controlled by the entire generation, accumulation and migration system. In areas where little methane has been generated or accumulated, seepage intensity would be low. In other areas, where methane has accumulated and faults to the surface provide conduits for release, methane seepage may be episodic and of moderate to high intensity when tectonism or atmospheric processes (described in Section 2.2.2) create conditions that enhance seepage. It is also important to note that, since gas accumulations would be expected to be more or less long-lasting in a specific site, a site that may have episodically seeped methane to the martian atmosphere would be expected to repeat that type of release through time. This would apply particularly to sites with visible, macro-seepage phenomena that have modified the ground morphology, such as the mud-volcano-like mounds in the northern lowlands (e.g., Skinner and Mazzini, 2009; Oehler and Allen, 2010; Ivanov et al., 2014). Sites like these could be major release locations suitable for potential follow-up studies.

Below we focus on geological and atmospheric factors that, through rock permeability and gas pressure gradients, would control intensity and variability of gas seepage on Mars. Fig. 2 summarizes these factors. We additionally discuss implications of the lower gravity on Mars (compared to Earth) on fracture density and openness as well as the effects of Mars' comparatively low atmospheric pressure on the abundance of recent impactors that could contribute to episodic pulses of methane.

#### 3.1. Potential methane accumulation and release mechanisms

Once generated by biotic or abiotic processes, methane can migrate





faults that form wrinkle ridges, which may overlie thrust faults (discussed in detail in Golombek and Phillips, 2009).

Impacts add to these fault systems, as does fracturing from processes such as giant polygon-formation in the northern plains (Oehler and Allen, 2012; Allen et al., 2013), serpentinization of olivine-rich rocks (e.g., Ehlmann et al., 2010), which may enhance gas accumulation in igneous rocks whose primary permeability would otherwise be quite low (Etiope et al., 2013a), and sedimentary diapirism, which can produce structures on Mars that resemble terrestrial mud volcanoes (Oehler and Allen, 2010) as well as both sedimentary dykes (Grotzinger et al., 2014) and pipes (Rubin et al., 2016). In addition, it is important to note that the lower gravity of Mars induces lower lithostatic pressure than on Earth, resulting in higher fracture density, at any depth (Heap et al., 2017). The lower gravity may also result in greater depth of fault-penetration and may enhance formation of dilational pit crater chains, some of which may even be forming today (Ferrill et al., 2004; Smart et al., 2011). Therefore, even igneous rocks (mafic and ultramafic) can be more fractured than those on Earth, forming excellent reservoir rocks. Taken together, these processes have created a martian subsurface that is heavily fractured (e.g., Rodríguez et al., 2005) and thus likely to provide numerous conduits for gas migration and release to the atmosphere (e.g., De Toffoli et al., 2018).

Faults from the dichotomy-forming event, the build-up of Tharsis, and major impacts would penetrate deep into the subsurface, providing conduits for methane generated and accumulated at great depth. Although most of these faults would have developed early in Mars' history, they are likely to have been reactivated many times by continuing tectonic stresses, with subsequent, episodic methane releases associated with such reactivations (e.g., Plaza-Faverola et al., 2015). On late Mars, tectonic stresses may be due to planetary cooling, evolution of the dichotomy, basin subsidence, ice sublimation, volcanism, sedimentary diapirism, and continuing impacts. This type of reactivation could keep the older deeper faults open, particularly in cases where they have been cemented (e.g., as in the many faults with cement fills in Gale crater; Siebach et al., 2015; Yen et al., 2017; Kronyak et al., 2018).

It is fault movement and pressure changes associated with the most recent planetary stresses that could contribute to current episodic emissions, accounting for both temporal and spatial variability in methane releases. Data expected from the upcoming Mars InSight Lander should provide new insights to current seismicity and tectonism on Mars.

### 3.3. The role of meteorite impacts in shallow gas release

On Mars, episodic gas releases may be induced by recent/current meteoritic impacts. The loss of atmosphere on Mars particularly enhances the abundance of small and recent impactors hitting the martian surface. These not only could contribute to fault-mediated seepage of methane from the subsurface, but their associated seismic waves could change gas pressures, triggering advective pulses of methane.

To understand the frequency of recent impactors, Daubar et al. (2013) evaluated 248 impact sites formed within the last few decades using images from the Context Camera (CTX) and High Resolution Imaging Science Experiment (HiRISE) on NASA's Mars Reconnaissance Orbiter by comparing the most recent images with previous images taken of the same sites. From these observations, they calculate that meter-to decameter-sized craters are forming on Mars at a rate of  $1.65 \times 10^{-6}$  craters/km<sup>2</sup>/yr, for craters with effective diameters  $\geq 3.9$  m, which amounts to  $> 200$  of these meteorite strikes per year.

However, most recent impact craters are small. Forty-four craters were highgraded by Daubar et al. (2013), as those had both before- and after-impact images of the impact sites. These craters range from  $\sim 2$  to 34 m in diameter and all but one are  $< 15$  m in diameter. Accordingly, the depths of penetration of most of the recent craters would be expected to be from  $< 1$  m to  $\sim 8$  m (Daubar et al., 2014).

So the question is: can these recent impacts penetrate subsurface clathrates or ice (that may seal methane reservoirs), thereby accounting

for episodic methane releases? First, various lines of evidence show that permafrost/ice is present in the near surface of mid to high latitudes of Mars. The Phoenix mission, which landed in the northern plains, at 68.2°N, 234.3°E, showed patches of water ice below the lander (mean depth of 4.6 cm), which had been exposed by the spacecraft thrusters during landing (Mellon et al., 2009; Smith et al., 2009), and permanent water ice has been detected by the High Resolution Stereo Camera (HRSC) on the European Space Agency (ESA) Mars Express spacecraft on the floor of a 35 km crater at  $\sim 70.5^\circ\text{N}$ ,  $103^\circ\text{E}$  (<https://mars.nasa.gov/resources/5266/mars-crater-ice/>). At mid-latitudes, HiRISE image data have shown several examples of small, recent craters that expose water ice, which fades over a few months, presumably from sublimation (Byrne et al., 2009), and potentially extensive ice is suggested by Dundas et al. (2018) in their report of deposits of ice exposed in cliffs at  $\sim 55^\circ\text{S}$  and  $55^\circ\text{N}$ . Those deposits, which can be  $> 100$  m thick, extend downwards from depths of 1–2 m. And, finally, there is the possibility of equatorial ice (Wilson et al., 2018) in the uppermost meter or so of the Medusae Fossae Formation, along the dichotomy. So shallow ice in parts of the mid and higher latitudes (and possibly equatorially) could seal methane trapped in shallow lithologic reservoirs. If a recent impactor hits one of these ice seals, it could breach that seal, accounting for a pulse of methane released to the atmosphere.

In contrast, as noted, methane clathrates may be restricted to depths greater than 15 m in equatorial latitudes (Max et al., 2013) though they are likely to exist at shallower depths at latitudes  $> 60^\circ$  (Chastain and Chevrier, 2007). So clathrates at latitudes  $> 60^\circ$  may well be destabilized by physical disruption caused by shallow recent impacts. Deeper clathrates, at lower latitudes, would be more difficult to destabilize by small impactors and might only be affected by some of the largest of the recent impacts.

### 3.4. The role of atmospheric pressure and temperature

Atmospheric factors (described in Section 2.2) can be quite important for subsurface methane release. Mars is a windy and stormy environment, with wind speeds up to around 30 m per second (e.g., Cantor, 2007). Winds and pressure changes, also related to the strong daily thermal variations, may therefore be important contributors to episodic release of methane that may have reached near-surface horizons, as observed on Earth. The same processes inducing martian dust uplift, such as thermal creep airflow or “dust devils”, characterized by a pressure drop in their center (e.g., de Beule et al., 2014; Küpper and Wurm, 2015) could produce gas exhalation “pulses”. And, in a dry, water-free (and thus presumably microbial-free) subsurface, possible seasonal (cyclic) variations of microseepage can be related to seasonal variations of atmospheric temperature, pressure, and winds.

Atmospheric processes may additionally overprint on any methane releases that occur from subsurface geological processes. For example, in a preliminary study, Pla-Garcia et al. (2018) suggest that variable background methane detections at Gale crater might be a result of changes in local ground temperatures caused by atmospheric circulation. They suggest that during the period on Mars of Ls (solar longitude) 225–315, which correlates with low background methane abundances (Webster et al., 2018), airflow would originate from the cold northern hemisphere, whereas in other seasons, the airflow into Gale crater would be from more tropical areas, closer to Gale and consequently warmer. If this were the case, then warmer temperatures in Gale crater beyond the Ls 225–315 period might affect clathrate releases in the region.

Thermal oscillations, caused by the variation of the obliquity or by seasons, can destabilize clathrates at relatively large depths (several hundred meters) depending on latitude and thermal conductivity of the ground (Gloesener et al., 2018) or cause melting of permafrost seals for conventionally trapped methane. The same temperature changes may control microbial activity, if any, as observed on Earth, so that warmer periods may induce higher methane consumption by methanotrophs and thus lower surface seepage rates (Etiope and Klusman, 2010). This

possibility, though hypothetical, is clearly an area of continuing research and these types of mechanisms deserve additional experimental and modeling work assuming different scenarios of near-surface methane occurrence, as well as seasonal changes in martian circulation processes.

#### 4. Can seepage variability provide insight into biotic vs abiotic methane origin?

The answer is no. Based on what has been discussed above, gas flux intensity and variability at the ground-atmosphere interface are solely determined by physical factors (permeability and pressure gradient), which operate on any type of subsurface methane, regardless its origin. The variability of gas release to the atmosphere is more linked to variability of factors affecting gas accumulations (especially those shallow), rather than to variability of gas production in a deep source rock. Both microbial and abiotic methane can accumulate in gas pockets, in clathrates, in zeolites, or in conventional reservoirs trapped below ice, after migration from a source rock. Different from diffusion, where  $\text{CH}_4$  isotopic composition affects the diffusion coefficient and thus  $\text{CH}_4$  migration, in seepage, which is mostly driven by advection (see Section 2),  $\text{CH}_4$  isotopic composition, and thus its origin, has no influence on gas flow rates or responses to pressure stress.


Even seasonal variations of atmospheric methane on Mars (e.g., Webster et al., 2018) cannot be attributed to biotic or abiotic origins. On Earth, microbes (methanotrophs) in the soil determine microseepage seasonality, by consuming more methane in summer and wetter seasons, so that fluxes are higher in winter and dry seasons (Etiope and Klusman, 2010). However, on Mars, potential methane-consuming and/or generating bacteria could only reside in the subsurface, in the presence of liquid water, in endolithic ecosystems within pore spaces of rocks and in ice and permafrost (Oehler and Etiope, 2017 and references therein). Although seasonal temperature variations in these systems could control

such hypothetical microbial activity (and thus methane fluxes at the surface), as discussed above,  $\text{CH}_4$  seasonal variations on Mars can also be due to atmospheric changes (winds and storms), melting/sublimation of permafrost and ice, and destabilization of shallower clathrates, all of which could operate on non-microbial methane, as well. Therefore,  $\text{CH}_4$  seasonality on Mars is not diagnostic of methane origin.

#### 5. Summary model and concluding remarks

By analogy with gas seepage on Earth, subsurface gas pressure gradients on Mars (induced by gas accumulations and permeability of fracture/fault networks) represent the main determinant of methane seepage occurrence, intensity and variability. Mars has a brittle and potentially highly fractured and faulted lithosphere, which may widely host gas accumulations and gas migration routes for seepage to the surface. Episodic, occasional or recurrent, methane releases will occur when specific geological and atmospheric processes induce sudden or progressive changes of pressure and permeability. It is the sum of generation location, subsurface accumulation points, conduits for release to the atmosphere, and triggers for release that control the sites of methane release on the surface, accounting for spatial variability. Atmospheric processes (e.g., barometric pressure drops related to winds and storms) will overprint on the spatial variability and may contribute to seasonal variations. Fig. 4 summarizes the entire system.

Although it might still be argued that some of the methane detections on Mars may not be real, the methane observations acquired so far, including the spikes, seasonal oscillations and non-detections (see references in the Introduction), appear compatible with geological seepage dynamics. In addition, the very existence of methane seepage appears logical and expected given the abundance of methane-generating mechanisms and the great potential for gas storage and migration to the surface (Oehler and Etiope, 2017). Obviously, the variability of



**Fig. 4.** Summary sketch of endogenous (geological, subsurface) and exogenous (atmospheric) processes (all marked in red) that can produce episodic (irregular or seasonal) release of methane on Mars. Gas seepage spikes may result from rapid changes in the gas pressure gradient (reservoir and atmospheric pumping) or sudden increase of rock permeability (opening of pathways due to seismicity, impact, ice melting/sublimation, or local tectonism). Squiggly arrows: methane release; thin blue dashed arrows = near-ground air circulation (winds, storms, “dust devils”) inducing low pressure above the ground (Bernoulli effect). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)



methane seepage discussed in this work needs to be integrated in a unifying model with simulations of atmospheric circulation and proposed, additional processes of rapid atmospheric methane removal (e.g., Lefèvre and Forget, 2009; Knak-Jensen et al., 2014; Holmes et al., 2015), which can be tested against all methane observations acquired so far.

Given the very low (ppbv) amounts of methane only episodically detected in the martian atmosphere, net of the atmospheric removal process, the average level of seepage on Mars is likely orders of magnitude lower than that on Earth. This fact coupled with the known variability of seepage on Earth, argue that the source locations for the methane detected in the martian atmosphere may be restricted to specific regions with relatively high potential for methane generation and seepage (see detailed discussion of potential seepage sites in Oehler and Etiope, 2017). And weak, localized, sporadic seepage fluxes (as in the seepage pattern “c” discussed in Section 2.4.2) could result into very low global average CH<sub>4</sub> concentrations, eventually below detection limits or not requiring very fast removal processes. However, it is important to take into account that the “lifetime” of a seepage site is generally quite long or perennial, at least lasting centuries. Therefore, a site that may have episodically released methane on Mars is expected to repeat the release sooner or later.

Allowing for the oscillating character of methane seepage and the potential for additional atmospheric removal processes (as referenced above), non-detections can be more frequent than detections. Thus, if methane sensors operate discontinuously and only in one site or over a limited number of regions, then non-detections will be the rule and detections will be relatively rare. The chances of methane detection may increase if the observations are focused and repeated on specific, high-graded areas of gas release determined by geological analysis (Oehler and Etiope, 2017). Seepage variability, including seasonality, being controlled mainly by physical migration factors, is not related to methane origin. Thus, although methane of biologic origin is conceivable, to date there is no definitive evidence for past or present life on Mars, and the methane data are not sufficient, in themselves, to conjecture about any relevance to life on Mars.

## Acknowledgments

This work stemmed from collaborations and valuable discussions with Marco Giuranna, Frank Daerden, Sebastien Viscardy and Lori Neary. We thank Steven A. Bowman and Gulf PetroLink for providing permission to adapt a figure from Bowman (2011), originally published in GeoArabia, for Fig. 3 of this paper. Seismic data in Fig. 3 were provided courtesy of CGG.

## References

- Allen, C.C., Oehler, D., Etiope, G., Van Rensbergen, P., Baci, C., Feyzullayev, A., Martinelli, G., Tanaka, K., Van Rooij, D., 2013. Fluid expulsion in terrestrial sedimentary basins: a process providing potential analogs for giant polygons and mounds in the martian lowlands. *Icarus* 224, 424–432.
- Anderson, R.C., Dohm, J.M., Golombek, M.P., Haldemann, A.F.C., Franklin, B.J., Tanaka, K.L., Lias, J., Peer, B., 2001. Primary centers and secondary concentrations of tectonic activity through time in the western hemisphere of Mars. *J. Geophys. Res.* 106, 20,563–20,585.
- Aoki, S., Richter, M.J., DeWitt, C., Boegert, A., Encrenaz, T., Sagawa, H., Nakagawa, H., Vandaele, A.C., Giuranna, M., Greathouse, T.K., Fouchet, T., Gemina, A., Sindoni, G., McKelvey, M., Case, M., Kasaba, Y., 2018. Stringent upper limit of CH<sub>4</sub> on Mars based on SOFIA/EXES observations. *Astron. Astrophys.* 610, A78. <https://doi.org/10.1051/0004-6361/201730903>.
- Atreya, S.K., Mahaffy, P.R., Wong, A.-S., 2007. Methane and related trace species on Mars: origin, loss, implications for life, and habitability. *Planet. Space Sci.* 55, 358–369.
- Blamey, N., Parnell, J., McMahon, S., Mark, D.F., Tomkinson, T., Lee, M., Shivak, J., Izawa, M.R.M., Banerjee, N.R., Flemming, R.L., 2015. Evidence for methane in Martian meteorites. *Nat. Commun.* 6, 7399. <https://doi.org/10.1038/ncomms8399>.
- Bonini, C., 2013. Fluid seepage variability across the external Northern Apennines (Italy): structural controls with seismotectonic and geodynamic implications. *Tectonophysics* 590, 151–174.
- Bowman, S.A., 2011. Regional seismic interpretation of the hydrocarbon prospectivity of offshore Syria. *GeoArabia* 16, 95–124.
- Brown, A., 2000. Evaluation of possible gas microseepage mechanisms. *Am. Assoc. Petrol. Geol. Bull.* 84, 1775–1789.
- Byrne, S., Dundad, C.M., Kennedy, M.R., Mellon, M.T., McEwen, A.S., Cull, S.C., Daubar, I.J., Shean, D.E., Seelos, K.D., Murchier, S.L., et al., 2009. Distribution of mid-latitude ground ice on Mars from new impact craters. *Science* 325, 1674–1676.
- Cantor, B.A., 2007. MOC observations of the 2001 Mars planet-encircling dust storm. *Icarus* 186, 60–96.
- Chastain, B.K., Chevrier, V., 2007. Methane clathrate hydrates as a potential source for Martian atmospheric methane. *Planet. Space Sci.* 55, 1246–1256.
- Chiodini, G., Granieri, D., Avino, R., Caliro, S., Costa, A., Minopoli, C., Vilardo, G., 2010. Non-volcanic CO<sub>2</sub> Earth degassing: case of Mefite d'Ansanto (southern Apennines), Italy. *Geophys. Res. Lett.* 37, L11303. <https://doi.org/10.1029/2010GL042858>.
- Ciotoli, G., Etiope, G., Guerra, M., Lombardi, S., Duddridge, G.A., Grainger, P., 2005. Migration and behaviour of gas injected into a fault in low-permeability ground. *Quart. J. Engineer. Geol. Hydrog.* 38, 305–320.
- Ciotoli, G., Etiope, G., Marra, F., Florindo, F., Giraudi, C., Ruggiero, L., 2016. Tiber delta CO<sub>2</sub>-CH<sub>4</sub> degassing: a possible hybrid, tectonically active Sediment-Hosted Geothermal System near Rome. *J. Geophys. Res.* - Solid Earth 121, 48–69.
- Clements, W.E., Wilkening, M.H., 1974. Atmospheric pressure effects on radon-222 transport across the Earth–air interface. *J. Geophys. Res.* 79, 5025–5029.
- Christensen, T.R., Johansson, T., Akerman, H.J., Mastepanov, M., Malmer, N., Friberg, T., Crill, P., Svensson, B.H., 2004. Thawing sub-arctic permafrost: effects on vegetation and methane emissions. *Geophys. Res. Lett.* 31, L04501. <https://doi.org/10.1029/2003GL018680>.
- Daubar, I.J., McEwen, A.S., Byrne, S., Kennedy, M.R., Ivanov, B., 2013. The current martian cratering rate. *Icarus* 225, 506–516.
- Daubar, I.J., Atwood-Stone, C., Byrne, S., McEwen, A.S., Russell, P.S., 2014. The morphology of small fresh craters on Mars and the Moon. *J. Geoph. Res. - Planets* 119, 2620–2639.
- de Beule, C., Wurm, G., Kelling, T., Küpper, M., Jankowski, T., Teiser, J., 2014. The Martian soil as a planetary gas pump. *Nat. Phys.* 10, 17–20.
- Delisle, G., Teschner, M., Faber, E., Panahi, B., Guliev, I., Aliev, C., 2010. First approach in quantifying fluctuating gas emissions of methane and radon from mud volcanoes in Azerbaijan. In: Wood, L. (Ed.), *Shale Tectonics*, vol. 93. Amer. Assoc. Petrol. Geol. Memoir, pp. 209–222.
- De Toffoli, B., Pozzobon, R., Mazzarini, F., Orgel, C., Massironi, M., Giacomini, L., Mangold, N., Cremonese, G., 2018. Estimate of depths of source fluids related to mound fields on Mars. *Planet. Space Sci.* 164, 164–173.
- Dundas, C.M., Bramson, A.M., Ojha, L., Wray, J.J., Mellon, M.T., Byrne, S., McEwen, A.S., Putzig, N.E., Viola, D., Sutton, S., Clark, E., Hold, J.W., 2018. Exposed subsurface ice sheets in the Martian mid-latitudes. *Science* 359, 199–201.
- Ehlmann, B.L., Mustard, J.F., Murchie, S.L., 2010. Geologic setting of serpentine deposits on Mars. *Geophys. Res. Lett.* 37, L06201. <https://doi.org/10.1029/2010GL042596>.
- Embriaco, D., Marinaro, G., Frugoni, F., Monna, S., Etiope, G., Gasperini, L., Polonia, A., Del Bianco, F., Çagatay, M.N., Ulgen, U.B., Favali, P., 2014. Monitoring of gas and seismic energy release by multi-parametric benthic observatory along the North Anatolian Fault in the Sea of Marmara (NW Turkey). *Geophys. J. Int.* 196, 850–866.
- Etiope, G., 2015. Natural Gas Seepage. The Earth's Hydrocarbon Degassing. Springer, Switzerland, p. 199.
- Etiope, G., Christodoulou, D., Kordella, S., Marinaro, G., Papatheodorou, G., 2013b. Offshore and onshore seepage of thermogenic gas at Katakolo Bay (Western Greece). *Chem. Geol.* 339, 115–126.
- Etiope, G., Ehlmann, B., Schoell, M., 2013a. Low temperature production and exhalation of methane from serpentinized rocks on Earth: a potential analog for methane production on Mars. *Icarus* 224, 276–285.
- Etiope, G., Ifandi, E., Nazzari, M., Procesi, M., Tsikouras, B., Ventura, G., Steele, A., Tardini, R., Szatmari, P., 2018. Widespread abiogenic methane in chromitites. *Sci. Rep.* 8, 8728. <https://doi.org/10.1038/s41598-018-27082-0>.
- Etiope, G., Klusman, R.W., 2010. Microseepage in drylands: flux and implications in the global atmospheric source/sink budget of methane. *Glob. Planet. Chang.* 72, 265–274.
- Etiope, G., Lombardi, S., 1995. Evidence for radon transport by carrier gas through faulted clays in Italy. *J. Radioanal. Nucl. Chem.* 193, 291–300.
- Etiope, G., Martinelli, G., 2002. Migration of carrier and trace gases in the geosphere: an overview. *Phys. Earth Planet. In.* 129 (3–4), 185–204.
- Etiope, G., Schoell, M., Hosgormez, H., 2011. Abiotic methane flux from the Chimaera seep and Tekirova ophiolites (Turkey): understanding gas exhalation from low temperature serpentinization and implications for Mars. *Earth Planet. Sci. Lett.* 310, 96–104.
- Etiope, G., Vadillo, I., Whitar, M.J., Marques, J.M., Carreira, P.M., Tiago, I., Benavente, J., Jimenez, P., Urresti, B., 2016. Abiotic methane seepage in the Ronda peridotite massif, southern Spain. *Appl. Geochem.* 66, 101–113.
- Etiope, G., Zwahlen, C., Anselmetti, F.S., Kipfer, R., Schubert, C.J., 2010. Origin and flux of a gas seep in the northern Alps (Giswil, Switzerland). *Geofluids* 10, 476–485.
- Ferrill, D.A., Wyrick, D.Y., Morris, A.P., Sims, D.W., Franklin, N.M., 2004. Dilational fault slip and pit chain formation on Mars. *GSA Today* 14, 4–12.
- Fonti, S., Marzo, G.A., 2010. Mapping the methane on Mars. *Astron. Astrophys.* 512, A51. <https://doi.org/10.1051/0004-6361/200913178>.
- Formisano, V., Atreya, S., Encrenaz, T., Ignatiev, N., Giuranna, M., 2004. Detection of methane in the atmosphere of Mars. *Science* 306, 1758–1761.
- Fries, M., Christou, A., Archer, D., Conrad, P., Cooke, W., Eigenbrode, J., ten Kate, I.L., Matney, M., Niles, P., Sykes, M., Steele, A., Treiman, A., 2016. A cometary origin for martian atmospheric methane. *Geochim. Perspect. Lett.* 2, 10–23. <https://doi.org/10.7185/geochemlet.1602>.
- Golombek, M.P., Phillips, R.J., 2009. Mars tectonics. In: Watters, T.R., Schultz, R.A. (Eds.), *Planetary Tectonics*. Cambridge University Press, Cambridge, UK, pp. 183–232.



- Gloesener, E., Karatekin, O., Dehant, V., 2018. Influence of subsurface properties on ice and methane clathrate stability on Mars. In: Scientific Workshop "From Mar Express to ExoMars", 27–28 February 2018. ESAC, Madrid.
- Grotzinger, J.P., Sumner, D.Y., Kah, L.C., Stack, K., Gupta, S., Edgar, L., Rubin, D., Lewis, K., Schieber, J., Mangold, N., Milliken, R., Conrad, P.G., Des Marais, D., Farmer, J., Siebach, K., Calef, F., Hurowitz, J., McLennan, S.M., Ming, D., Vaniman, D., Crisp, J., Vasavada, A., Edgett, K.S., Malin, M., Blake, D., Geliert, R., Mahaffy, P., Wiens, R.C., Maurice, S., Grant, J.A., Wilson, S., Anderson, R.C., Beegle, L., Arvidson, R., Hallet, B., Sletten, R.S., Rice, M., Bell, J., Griffes, J., Ehlmann, B., Anderson, R.B., Bristow, T.F., Dietrich, W.E., Dromart, G., Eigenbrode, J., Fraeman, A., Hardgrove, C., Herkenhoff, K., Jandura, L., Kocurek, G., Lee, S., Leshin, L.A., Leveille, R., Limonadi, D., Maki, J., McCloskey, S., Meyer, M., Minitti, M., Newsom, H., Oehler, D., Okon, A., Palucis, M., Parker, T., Rowland, S., Schmidt, M., Squyres, S., Steele, A., Stolper, E., Summons, R., Treiman, A., Williams, R., Yingst, A., the MSL Science Team, 2014. A habitable fluvio-lacustrine environment at yellowknife Bay, Gale crater, Mars. *Science* 343. <https://doi.org/10.1126/science.1242777>.
- Haney, M.M., Snieder, R., Sheiman, J., Losh, S., 2005. A fault caught in the act of burping. Consortium project on seismic inverse methods for complex structures. *CWP* 513, 219–224.
- Hauber, E., Grott, M., Kronberg, P., 2010. Martian rifts: structural geology and geophysics. *Earth Planet. Sci. Lett.* 294, 393–410. <https://doi.org/10.1016/j.epsl.2009.11.005>.
- Heap, M.J., Byrne, P.K., Mikhail, S., 2017. Low surface gravitational acceleration of Mars results in a thick and weak lithosphere: implications for topography, volcanism, and hydrology. *Icarus* 281, 103–114.
- Heinicke, J., Koch, U., 2000. Slug flow - a possible explanation for hydrogeochemical earthquake precursors at Bad Brambach, Germany. *Pure Appl. Geophys.* 157, 1621–1641.
- Holmes, J.A., Lewis, S.R., Patel, M.R., 2015. Analysing the consistency of martian methane observations by investigation of global methane transport. *Icarus* 257, 23–32.
- Horseman, S.T., Harrington, J.F., Sellin, P., 1999. Gas migration in clay barriers. *Eng. Geol.* 54, 139–143.
- Ivanov, M., Hiesinger, H., Erkeling, G., Reiss, D., 2014. Mud volcanism and morphology of impact craters in Utopia Planitia on Mars: evidence for the ancient ocean. *Icarus* 228, 121–140.
- Keppler, F., Vigano, I., McLeod, A., Ott, U., Fruchtl, M., Rockmann, T., 2012. Ultraviolet-radiation-induced methane emissions from meteorites and the martian atmosphere. *Nature* 486, 93–96.
- Klusman, R.W., 1993. Soil-gas and Related Methods for Natural Resource Exploration. Wiley, Chichester, p. 483.
- Klusman, R.W., Leopold, M.E., LeRoy, M.P., 2000. Seasonal variation in methane fluxes from sedimentary basins to the atmosphere: results from chamber measurements and modeling of transport from deep sources. *J. Geophys. Res.* 105D, 24,661–24,670.
- Knæk Jensen, S.J., Skibsted, J., Jakobsen, H.J., ten Kate, I.L., Gunnlaugsson, H.P., Merrison, J.P., Finster, K., Bak, E., Iversen, J.J., Kondrup, J.C., Nørnberg, P., 2014. A sink for methane on Mars? The answer is blowing in the wind. *Icarus* 236, 24–27.
- Knapmeyer, M., Oberst, J., Hauber, E., Wählisch, M., Deuchler, C., Wagner, R., 2006. Working models for spatial distribution and level of Mars' seismicity. *J. Geophys. Res.* 111. <https://doi.org/10.1029/2006JE002708>.
- Kopf, A., Delisle, G., Faber, E., Panahi, B., Aliyev, C.S., Guliyev, I., 2009. Long-term in situ monitoring at Dashgil mud volcano, Azerbaijan: a link between seismicity, pore-pressure transients and methane emission. *Int. J. Earth Sci.* 99, 227–240.
- Krabbenhöft, A., Netzeband, G.L., Bialas, J., Papenberg, C., 2010. Episodic methane concentrations at seep sites on the upper slope Opouawe Bank, southern Hikurangi Margin, New Zealand. *Mar. Geol.* 272, 71–78.
- Krasnopolsky, V.A., Maillard, J.P., Owen, T.C., 2004. Detection of methane in the Martian atmosphere: evidence for life? *Icarus* 172, 537–547.
- Kronyak, R.E., Kah, L.C., Miklusciak, N.B., Edgett, K.S., Nachon, M., Fedo, C.M., 2018. Formation of Fracture Networks in the Siccar Point Group: Implications for Timing of Post-depositional Fluid Flow in Gale Crater, Mars. 49th LPSC. Abs. # 1371.
- Küpper, M., Wurm, G., 2015. Thermal creep-assisted dust lifting on Mars: wind tunnel experiments for the entrainment threshold velocity. *J. Geophys. Res. - Planets* 120, 1346–1356. <https://doi.org/10.1002/2015JE004848>.
- Lefèvre, F., Forget, F., 2009. Observed variations of methane on Mars unexplained by known atmospheric chemistry and physics. *Nature* 460, 720–723.
- Lewicki, J.L., Hilley, G.E., Tosha, T., Aoyagi, R., Yamamoto, K., Benson, S.M., 2007. Dynamic coupling of volcanic CO<sub>2</sub> flow and wind at the Horseshoe Lake tree kill, Mammoth Mountain, California. *Geophys. Res. Lett.* 34, L03401. <https://doi.org/10.1029/2006GL028848>.
- Macgregor, D.S., 1993. Relationships between seepage, tectonics and subsurface petroleum reserves. *Mar. Petrol. Geol.* 10, 606–619.
- Malmqvist, L., Kristiansson, K., 1984. Experimental evidence for an ascending microflow of geogas in the ground. *Earth Planet. Sci. Lett.* 70, 407–416.
- Malmqvist, L., Kristiansson, K., 1985. A physical mechanism for the release of free gases in the lithosphere. *Geoscientific Exploration* 23, 447–453.
- Massmann, J., Farrier, D.F., 1992. Effects of atmospheric pressures on gas transport in the vadose zone. *Water Resour. Res.* 28, 777–791.
- Max, M.D., Clifford, S.M., Johnson, A.H., 2013. Hydrocarbon system analysis for methane hydrate exploration on Mars. *Am. Assoc. Pet. Geol. Mem.* 101, 99–114. <https://doi.org/10.1306/13361573M1013546>.
- Mazzini, A., Etiope, G., 2017. Mud volcanism: an updated review. *Earth Sci. Rev.* 168, 81–112.
- McConnochie, T.H., Smith, M.D., Bender, S., Wolff, M.J., Johnson, J.R., Lemmon, M.T., Wiens, R.C., Maurice, S., Gasnault, O., Blaney, D., DeFlores, L.P., Harri, A.-M., Kemppinen, O., Genzer, M., Moores, J.E., Wong, M.H., Trainer, M.G., Martin-Torres, F.J., Zorzano, M.-P., de la Torre Juárez, M., 2014. ChemCam passive spectroscopy of atmospheric O<sub>2</sub> and H<sub>2</sub>O. In: Eighth International Conference on Mars, Pasadena, CA, Jul. 2014.
- Mège, D., Cook, A.C., Garel, E., Lagabriele, Y., Cormier, M.-H., 2003. Volcanic rifting at Martian grabens. *J. Geophys. Res. - Planets* 108. <https://doi.org/10.1029/2002JE001852>.
- Mellon, M.T., Arvidson, R.E., Sizemore, H.G., Searis, M.L., Blaney, D.L., Cull, S., Hecht, M.H., Heet, T.L., Uwe Keller, H., Lemmon, M.T., Markiewicz, W.J., Ming, D.W., Morris, R.V., Thomas Pike, W., Zent, A.P., 2009. Ground ice at the Phoenix landing site: stability state and origin. *J. Geophys. Res. - Planets* 114, E00E07. <https://doi.org/10.1029/2009JE003417>.
- Mellors, R., Kilb, D., Aliyev, A., Gasanov, A., Yetimishli, G., 2007. Correlations between earthquakes and large mud volcano eruptions. *J. Geophys. Res.* 112, B04304. <https://doi.org/10.1029/2006JB004489>.
- Mousis, O., Simon, J.-M., Bellat, J.-P., Schmidt, F., Bouley, S., Chassefière, E., Sautter, V., Quesnel, Y., Picaut, S., Lécuyer, S., 2016. Martian zeolites as a source of atmospheric methane. *Icarus* 278, 1–7.
- Mumma, M.J., Villanueva, G.L., Novak, R.E., Hewagame, T., Bonev, B.P., DiSanti, M.A., Mandell, A.M., Smith, M.D., 2009. Strong release of methane on Mars in northern summer 2003. *Science* 323, 1041–1045.
- Nearly, L., Daerden, F., 2018. The GEM-Mars general circulation model for Mars: description and evaluation. *Icarus* 300, 458–476.
- Nilson, R.H., Peterson, E.W., Lie, K.H., Burkhard, N.R., Hearst, J.R., 1991. Atmospheric pumping: a mechanism causing vertical transport of contaminated gases through fractured permeable media. *J. Geophys. Res.* 96, 21,933–21,948.
- Oehler, D.Z., Allen, C.C., 2010. Evidence for pervasive mud volcanism in acidalia planitia, Mars. *Icarus* 208, 636–657.
- Oehler, D.Z., Allen, C.C., 2012. Giant polygons and mounds in the lowlands of Mars: signatures of an ancient ocean? *Astrobiology* 12, 601–615.
- Oehler, D.Z., Etiope, G., 2017. Methane seepage on Mars: where to look and why. *Astrobiology* 17, 1233–1264. <https://doi.org/10.1089/ast.2017.1657>.
- Pla-Garcia, J., Rafkin, S.C.R., Webster, C., Mahaffy, P.R., Karatekin, O., Gloesener, E., Moores, J.E., 2018. Martian winds drive seasonal methane variations observed by MSL-SAM. *Geophys. Res. Abstr.* 20. EGU2018-18277.
- Plaza-Faverola, A., Bünz, S., Johnson, J.W., Chand, S., Knies, J., Mienert, J., Franek, P., 2015. Role of tectonic stress in seepage evolution along the gas hydrate-charged Vestnesa Ridge, Fram Strait. *Geophys. Res. Lett.* 42, 733–742. <https://doi.org/10.1002/2014GL062474>.
- Portnov, A., Smith, A.J., Mienert, J., Cherkashov, G., Reikant, P., Semenov, P., Serov, P., Vanshtein, B., 2013. Offshore permafrost decay and massive seabed methane escape in water depths > 20 m at the South Kara Sea shelf. *Geophys. Res. Lett.* 40, 3962–3967.
- Reimer, G.M., Bowles, C.G., 1979. Soil-gas helium concentrations in the vicinity of a uranium deposit, Red esert, Wyoming. *U.S. Geol. Surv. Open-File Rept.* 79–975.
- Robledo-Martínez, A., Sobral, H., Ruiz-Meza, A., 2012. Electrical discharges as a possible source of methane on Mars: lab simulation. *Geophys. Res. Lett.* 39. <https://doi.org/10.1029/2012GL053255>.
- Rodríguez, J.A.P., Sasaki, S., Dohm, J.M., Tanaka, K.L., Strom, B., Kargel, J.S., Kuzmin, R., Miyamoto, H., Spary, J.G., Fairén, A.G., Komatsu, G., Kurita, K., Baker, V.R., 2005. Control of impact crater fracture systems on subsurface hydrology, ground subsidence, and collapse, Mars. *J. Geophys. Res.* 110. <https://doi.org/10.1029/2004JE002365>.
- Roos-Serote, M., Atreya, S.K., Webster, C.R., Mahaffy, P.R., 2016. Cometary origin of atmospheric methane variations on Mars unlikely. *J. Geophys. Res. - Planets* 121. <https://doi.org/10.1002/2016JE005076>.
- Rubin, D.M., Fairén, A.G., Martínez-Frías, J., Frydenvang, J., Gasnault, O., Gelfenbaum, G., Goetz, W., Grotzinger, J.P., Le Mouélis, S., Mangold, N., Newsom, H., Oehler, D.Z., Rapin, W., Schieber, J., Wiens, R.C., 2016. Fluidized-sediment pipes in Gale crater, Mars, and possible Earth analogs. *Geology* 45. <https://doi.org/10.1130/G38339.1>.
- Ruppel, C.D., Kessler, J.D., 2017. The interaction of climate change and methane hydrates. *Rev. Geophys.* 44, 126–168.
- Siebach, K.L., Grotzinger, J.P., McLennan, S.M., Hurowitz, J.A., Ming, D.W., Vaniman, D.T., Rampe, E.B., Blaney, D.L., Kah, L.C., MSL Science Team, 2015. Constraining the Texture and Composition of Pore-Filling Cements at Gale Crater, Mars. 46th LPSC. Abs. # 2234.
- Skinner, J., Mazzini, A., 2009. Martian mud volcanism: terrestrial analogs and implications for formational scenarios. *Mar. Petrol. Geol.* 26, 1866–1878.
- Smart, K.J., Wyrick, D.Y., Ferrill, D.A., 2011. Discrete element modeling of Martian pit crater formation in response to extensional fracturing and dilational normal faulting. *J. Geophys. Res. - Planets* 116. <https://doi.org/10.1029/2010JE003742>.
- Smith, P., Tamppari, L.K., Arvidson, R.E., Bass, D., Blaney, D., Boynton, W.V., Carswell, A., Catling, D.C., Clark, B.C., Duck, T., DeJong, E., Fisher, D., Goetz, W., Gunnlaugsson, H.P., Hecht, M.H., Hipkin, V., Hoffman, J., Hviid, S.F., Keller, H.U., Kounaves, S.P., Lange, C.F., Lemmon, M.T., Madsen, M.B., Markiewicz, W.J., Marshall, J., McKay, C.P., Mellon, M.T., Ming, D.W., Morris, R.V., Pike, W.T., Renno, N., Staufer, U., Stoker, C., Taylor, P., Whiteway, J.A., Zent, A.P., 2009. H<sub>2</sub>O at the Phoenix landing site. *Science* 325, 58–61.
- Stevens, A., Patel, M.R., Lewis, S.R., 2015. Numerical modelling of the transport of trace gases including methane in the subsurface of Mars. *Icarus* 250, 587–594.
- Stevens, A.H., Patel, M.R., Lewis, S.R., 2017. Modelled isotopic fractionation and transient diffusive release of methane from potential subsurface sources on Mars. *Icarus* 281, 240–247.

- Thornton, B.F., Geibel, M.C., Crill, P.M., Humborg, C., Mörtz, C.-M., 2016. Methane fluxes from the sea to the atmosphere across the Siberian shelf seas. *Geophys. Res. Lett.* 43, 5869–5877.
- Villanueva, G.L., Mumma, M.J., Novak, R.E., Radeva, Y.L., Käufel, H.U., Smette, A., Tokunaga, A., Khayat, A., Encrenaz, T., Hartogh, P., 2013. A sensitive search for organics (CH<sub>4</sub>, CH<sub>3</sub>OH, H<sub>2</sub>CO, C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>2</sub>, C<sub>2</sub>H<sub>4</sub>), hydroperoxyl (HO<sub>2</sub>), nitrogen compounds (N<sub>2</sub>O, NH<sub>3</sub>, HCN) and chlorine species (HCl, CH<sub>3</sub>Cl) on Mars using ground-based high-resolution infrared spectroscopy. *Icarus* 222, 11–27.
- Viscardy, S., Daerden, F., Neary, L., 2016. Formation of layers of methane in the atmosphere of Mars after surface release. *Geophys. Res. Lett.* 43, 1868–1875.
- Wang, P., Zhang, X., Zhu, Y., Li, B., Huang, X., Pang, S., Zhang, S., Lu, C., Xiao, R., 2014. Effect of permafrost properties on gas hydrate petroleum system in the Qilian Mountains, Qinghai, Northwest China. *Envir. Sci. Proc. Impacts* 16, 2711–2720.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Farley, K.A., MSL Science Team, 2013. Low upper limit to methane abundance on Mars. *Science* 342, 355–357.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Moores, J.E., Flesch, G.J., Malespin, C., McKay, C.P., Martinez, G., Smith, C.L., Martin-Torres, J., Gomez-Elvira, J., Zorzano, M.-P., Wong, M.H., Trainer, M.G., Steele, A., Archer Jr., D., Sutter, B., Coll, P.J., Freissinet, C., Meslin, P.-Y., Gough, R.V., House, C.H., Pavlov, A., Eigenbrode, J.L., Glavin, D.P., Pearson, J.C., Keymeulen, D., Christensen, L.E., Schwenzer, S.P., Navarro-Gonzalez, R., Pla-García, J., Rafkin, S.C.R., Vicente-Retortillo, A., Kahanpää, H., Viudez-Moreiras, D., Smith, M.D., Harri, A.-M., Genzer, M., Hassler, D.M., Lemmon, M., Crisp, J., Sander, S.P., Zurek, R.W., Vasavada, A.R., 2018. Background levels of methane in Mars' atmosphere show strong seasonal variations. *Science* 360, 1093–1096.
- Webster, C.R., Mahaffy, P.R., Atreya, S.K., Flesch, G.J., Farley, K.A., Meslin, P.Y., Farley, K.A., Conrad, P.G., Christensen, L.E., Pavlov, A.A., Martin-Torres, J., Zorzano, M.P., McConnochie, T.H., Owen, T., Eigenbrode, J.L., Glavin, D.P., Steele, A., Malespin, C.A., Archer Jr., P.D., Sutter, B., Coll, P., Freissinet, C., McKay, C.P., Moores, J.E., Schwenzer, S.P., Bridges, J.C., Navarro-Gonzalez, R., Gellert, R., Lemmon, M.T., MSL Science Team, 2015. Mars methane detection and variability at Gale Crater. *Science* 247, 415–417.
- Wilson, J.T., Eke, V.R., Massey, R.J., Elphic, R.C., Feldman, W.C., Maurice, S., Teodoro, L.F.A., 2018. Equatorial locations of water on Mars: improved resolution maps based on Mars Odyssey neutron spectrometer data. *Icarus* 299, 148–160.
- Yang, T.F., Fu, C.C., Walia, V., Chen, C.H., Chyi, L.L., Liu, T.K., Song, S.R., Lee, M., Lin, C.W., Lin, C.C., 2006. Seismo-geochemical variations in SW Taiwan: multi-parameter automatic gas monitoring results. *Pure Appl. Geophys.* 163, 693–709.
- Yen, A.S., Ming, D.W., Vaniman, D.T., Gellert, R., Blake, D.F., Morris, R.V., Morrison, S.M., Bristow, T.F., Chipera, S.J., Edgett, K.S., Treiman, A.H., Clark, B.C., Downs, R.T., Farmer, J.D., Grotzinger, J.P., Rampe, E.B., Schmidt, M.E., Sutter, B., Thompson, L.M., MSL Science Team, 2017. Multiple stages of aqueous alteration along fractures in mudstone and sandstone strata in Gale Crater, Mars. *Earth Planet. Sci. Lett.* 471, 186–198.
- Yung, Y.L., Chen, P., 2015. Methane on Mars. *Astrobiol. Outreach* 3, 1. <https://doi.org/10.4172/2332-2519.1000125>.
- Zahnle, K., Freedman, R.S., Catling, D.C., 2011. Is there methane on Mars? *Icarus* 212, 493–503.