

Why does repeatedly multiplying a vector by a square matrix cause the vector to converge on or along the matrix's eigenvector?

<http://setosa.io/ev/eigenvectors-and-eigenvalues/>

2 Answers



Peter Elliott, UCLA Applied Math '13; CMU Statistics PhD Student

Updated Sep 19 2016 · Author has 57 answers and 82k answer views

First let's assume the matrix A has a full set of eigenvectors v_1, \dots, v_m with associated eigenvalues $\lambda_1, \dots, \lambda_m$. Further I'm going to assume that $\lambda_1 = 1$ and for all other k , $|\lambda_k| < 1$.

(Now this might seem like a strong assumption. However, if the largest magnitude eigenvalue were anything else we could simply factor it out to get a matrix $\tilde{A} = \frac{1}{\lambda_1} A$. This matrix has the same eigenvectors as A , and multiplying it by a vector gives only a scaled version of what you get multiplying A by the same vector. As a result, repeated multiplication by \tilde{A} will converge along the same vector as repeated multiplication by A .)

For any vector x we can write it in terms of the eigenvectors, $x = \sum_{k=1}^m a_k v_k$. Now let's try repeatedly multiplying x by A .

$$A^n x = \sum_{k=1}^m a_k A^n v_k = \sum_{k=1}^m a_k \lambda_k^n v_k$$

So the component of x corresponding to each eigenvector is repeatedly multiplied by the associated eigenvalue. For the first component, since the eigenvalue is 1 it just stays the same. For all other components, $\lambda_k^n \rightarrow_{n \rightarrow \infty} 0$, so they eventually disappear. That means we're left with

$$A^n x \rightarrow_{n \rightarrow \infty} a_1 v_1.$$

The intuition here is that when you perform repeated multiplication by the same matrix, you're really repeatedly scaling the coefficients corresponding to the eigenvector basis by the size of the eigenvalue. When you do this a large number of times, the largest eigenvalue/eigenvector will come to dominate.

1k Views · View Upvoters

 Your feedback is private.



Is this answer useful?

Yes

No

Promoted by Swagbucks.com

What's the fastest way to find coupon codes online?

The price they show you is never the best price. Luckily, this tool finds the best coupon for your cart.

 Learn more at swagbucks.com

Related Questions

More Answers Below

How can I recover the vector from the matrix (generated from the vector multiplied by its transpose but contaminated by a noise matrix)?

How does multiplying a column vector with a matrix differ from the same with a row vector? How do you know which form a vector is in?

Is it necessary to find the inverse of a given matrix? Should that matrix be a square matrix?

Like a matrix multiplied by a vector can be written as a linear combination of columns of the matrix, what does a matrix multiplied by a matrix...

If vector A and vector B are unit vectors, then what is the angle between them so that $\sqrt{2} \text{A vector} - \text{B vector}$ is a unit vector?

[+ Ask New Question](#)



Nadeem Rojenko, Ms. student of Theoretical Physics.

Answered Sep 18 2016 · Author has **399** answers and **242.9k** answer views

Pretty simple, eigen vector by definition is a vector that will stay it self (same direction, but may have different length) after performing matrix transformation.

So applying a matrix many times, will keep transforming, rotating, shrinking and expanding the vector till at some stage it will become one of the eigenvectors, then it will not change any more (change in direction).

343 Views

Related Questions

[How is the gradient of a column vector a matrix?](#)

Why do we use a column vector in Dirac matrix?

What is a square matrix?

How come a matrix is a vector?

Is every vector position vector?

If the augmented matrix from a set of vectors could be reduced to the identity matrix, can I always imply the vectors are linearly independent...

Why can we find the eigenvector by cross multiplying two linearly independent row vectors in a matrix $A - \lambda I$?

Can a matrix have a vector in each element?

Can we multiply a $n \times n$ square matrix with an $N \times 1$ matrix in Java?

What is a row-vector matrix product?

What is the Hessian matrix? What is it used for and for what reason?

What is PAM matrix?

Why is a 2×2 matrix a four component vector?

Why does (1) cannot be square matrix?

Why is A vector cross B vector not equal to B vector cross A vector?

+ Ask New Question