

# Problems in Mathematics

You solved 0 problems!!

[Solved Problems](#) / [Solve later Problems](#)

LINEAR ALGEBRA



## Eigenvalues of a Stochastic Matrix is Always Less than or Equal to 1

BY YU · PUBLISHED 11/17/2016 · UPDATED 07/06/2017



# LINEAR ALGEBRA

## Problem 185

Let  $A = (a_{ij})$  be an  $n \times n$  matrix.

We say that  $A = (a_{ij})$  is a **right stochastic matrix** if each entry  $a_{ij}$  is nonnegative and the sum of the entries of each row is 1. That is, we have

$$a_{ij} \geq 0 \quad \text{and} \quad a_{i1} + a_{i2} + \cdots + a_{in} = 1$$

for  $1 \leq i, j \leq n$ .

Let  $A = (a_{ij})$  be an  $n \times n$  right stochastic matrix. Then show the following statements.

**(a)** The stochastic matrix  $A$  has an eigenvalue 1.

**(b)** The absolute value of any eigenvalue of the stochastic matrix  $A$  is less than or equal to 1.

♡ Add to solve later

### Contents [\[hide\]](#)

Problem 185

Proof.

(a) The stochastic matrix  $A$  has an eigenvalue 1.

(b) The absolute value of any eigenvalue of the stochastic matrix  $A$  is less than or equal to 1.

Remark.

Proof.

(a) The stochastic matrix  $A$  has an eigenvalue 1.

We compute that

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} = \begin{bmatrix} a_{11} + a_{12} + \dots + a_{1n} \\ a_{21} + a_{22} + \dots + a_{2n} \\ \vdots \\ a_{n1} + a_{n2} + \dots + a_{nn} \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}.$$

Here the second equality follows from the definition of a right stochastic matrix.

(Each row sums up to 1.)

This computation shows that  $\mathbf{1}$  is an eigenvector of  $A$  and  $\begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$  is an eigenvector corresponding to the eigenvalue 1.

(b) The absolute value of any eigenvalue of the stochastic matrix  $A$  is less than or equal to 1.

Let  $\lambda$  be an eigenvalue of the stochastic matrix  $A$  and let  $\mathbf{v}$  be a corresponding eigenvector.

That is, we have

$$A\mathbf{v} = \lambda\mathbf{v}.$$

Comparing the  $i$ -th row of the both sides, we obtain

$$a_{i1}v_1 + a_{i2}v_2 + \cdots + a_{in}v_n = \lambda v_i \quad (*)$$

for  $i = 1, \dots, n$ .

Let

$$|v_k| = \max\{|v_1|, |v_2|, \dots, |v_n|\},$$

namely  $v_k$  is the entry of  $\mathbf{v}$  that has the maximal absolute value.

Note that  $|v_k| > 0$  since otherwise we have  $\mathbf{v} = \mathbf{0}$  and this contradicts that an eigenvector is a nonzero vector.

Then from (\*) with  $i = k$ , we have

$$\begin{aligned} |\lambda| \cdot |v_k| &= |a_{k1}v_1 + a_{k2}v_2 + \cdots + a_{kn}v_n| \\ &\leq a_{k1}|v_1| + a_{k2}|v_2| + \cdots + a_{kn}|v_n| && \text{(by the triangle inequality and } a_{ij} \geq 0\text{)} \\ &\leq a_{k1}|v_k| + a_{k2}|v_k| + \cdots + a_{kn}|v_k| && \text{(since } |v_k| \text{ is maximal)} \\ &= (a_{k1} + a_{k2} + \cdots + a_{kn})|v_k| = |v_k|. \end{aligned}$$

Since  $|v_k| > 0$ , it follows that

$$\lambda \leq 1$$

as required.

**Remark.**

A stochastic matrix is also called probability matrix, transition matrix, substitution matrix, or Markov matrix.

Click here if solved ☆ 4

B! 0

Like 1

Share 1

Tweet

♥ Add to solve later

## More from my site



Stochastic Matrix (Markov Matrix) and its Eigenvalues and Eigenvectors (a) Let

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

be a matrix such that  $a_{11} + a_{12} = 1$  and  $a_{21} + a_{22} = 1$ . Namely, the sum of the entries in each row is 1. (Such a matrix is called (right) stochastic matrix (also termed [...]



Find the Limit of a Matrix Let

$$A =$$

be  $3 \times 3$  matrix. Find

.

(Nagoya University Linear [...]




### Basis with Respect to Which the Matrix for Linear Transformation is Diagonal Let

be the vector space of all real polynomials of degree 1 or less. Consider the linear transformation  $T : \quad \rightarrow \quad$  defined by


$$T(ax + b) = (3a + b)x + a + 3,$$

for any  $ax + b \in \quad$ . (a) With respect to the basis  $B = \{1, x\}$ , find the matrix of the linear transformation [...]




### If Every Vector is Eigenvector, then Matrix is a Multiple of Identity Matrix Let $A$ be an

$n \times n$  matrix. Assume that every vector  $\mathbf{x}$  in  $\quad$  is an eigenvector for some eigenvalue of  $A$ . Prove that there exists  $\lambda \in \mathbb{R}$  such that  $A = \lambda I$ , where  $I$  is the  $n \times n$  identity matrix. Proof. Let us write [...]



### Given All Eigenvalues and Eigenspaces, Compute a Matrix Product Let $C$ be a $4 \times 4$

matrix with all eigenvalues  $\lambda = 2, -1$  and eigenspaces  $E_2 = \text{Span}\left\{\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}\right\}$  and  $E_{-1} = \text{Span}\left\{\begin{bmatrix} 1 \\ 2 \\ 1 \\ 1 \end{bmatrix}\right\}$  [...]




### Eigenvalues of a Hermitian Matrix are Real Numbers Show that eigenvalues of a

Hermitian matrix  $A$  are real numbers. (The Ohio State University Linear Algebra Exam Problem) We give two proofs. These two proofs are essentially the same. The second proof is a bit simpler and concise compared to the first one. [...]



## Determine Eigenvalues, Eigenvectors, Diagonalizable From a Partial Information of a Matrix

Suppose the following information is known about a  $3 \times 3$  matrix  $A$ .  
$$\begin{bmatrix} 1 & 2 & 1 \\ 6 & 1 & 2 & 1 \\ 1 & -1 & 1 & \dots \end{bmatrix}, \quad A \begin{bmatrix} 1 & 2 & 1 \\ 1 & -1 & 1 & \dots \end{bmatrix}$$



## Given Eigenvectors and Eigenvalues, Compute a Matrix Product (Stanford University Exam)

Suppose that  $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$  is an eigenvector of a matrix  $A$  corresponding to the eigenvalue 3 and that  $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$  is an eigenvector of  $A$  corresponding to the eigenvalue  $-2$ . Compute  $A^2 \begin{bmatrix} 4 \\ \dots \end{bmatrix}$

Tags:

eigenvalue

eigenvector

linear algebra

Markov matrix

matrix

probability matrix

stochastic matrix

substitution matrix

transition matrix

triangle inequality

vector

PREVIOUS STORY



Eigenvalues of Squared Matrix and Upper Triangular Matrix

NEXT STORY

Is the Determinant of a Matrix Additive?



YOU MAY ALSO LIKE...





## Symmetric Matrix and Its Eigenvalues, Eigenspaces, and Eigenspaces

08/02/2016



## Find the Matrix Representation of $T(f)(x) = f(x^2)$ if it is a Linear Transformation

01/21/2018



## Rank and Nullity of Linear Transformation From $\mathbb{R}^3$ to $\mathbb{R}^2$

04/06/2017

### LEAVE A REPLY

Comment

Name \*

Email \*

Website

### Post Comment

- ☐ **Notify me of follow-up comments by email.**
- ☐ **Notify me of new posts by email.**

### TAGS

abelian group augmented matrix **basis** basis for a  
vector space **characteristic polynomial**  
commutative ring **determinant** determinant of a  
matrix diagonalization diagonal matrix  
**eigenvalue** **eigenvector** elementary row  
operations **exam** field theory finite group **group**  
group homomorphism **group theory**  
homomorphism ideal inverse matrix  
invertible matrix kernel **linear**  
**algebra** linear combination linearly  
independent **linear transformation**  
**matrix** matrix representation **nonsingular matrix**  
normal subgroup null space **Ohio State Ohio**  
**State.LA** rank ring ring theory subgroup

### SUBSCRIBE TO BLOG VIA EMAIL

Enter your email address to subscribe to this  
blog and receive notifications of new posts  
by email.

Join 250 other subscribers

Subscribe

### SEARCH MORE PROBLEMS

### CREATE AN ACCOUNT

Username

Email

Password

Repeat

Password

First Name

Last Name

Membership Free

Level

Register

If you are a member, Login here.

subspace symmetric matrix system of linear  
equations transpose vector vector space



Problems in Mathematics © 2018. All Rights Reserved.

