Chapter 8: Analysis of Variance (ANOVA)

Statistics for Environmental Science

Mallory Barnes

Rendered October 23, 2026

This PDF was rendered on October 23, 2026. The book is updated during the semester; the latest version of this chapter is at https://malloryb.github.io/statistics_E538/08-ANOVA.html.

Statistics for Environmental Science · CC BY-SA 4.0 · Credits and license: https://malloryb.github.io/statistics_E538/

Table of contents

ANOVA is analysis of variance **2**
- Why not run multiple t-tests? . . . 2
- ANOVA is an omnibus test . . . 3
- Kinds of ANOVA . . . 4

The F ratio **4**

F when H_0 is true **5**
- Making decisions . . . 6
- What group means look like when F exceeds the critical value . . . 7

Computing F by hand **8**
- Nine litter counts . . . 8
- SS Total . . . 9
- SS Between . . . 10
- SS Within . . . 10
- Degrees of freedom . . . 11
- Mean squares . . . 11
- Calculate F . . . 12
- The ANOVA table . . . 12

The full ANOVA workflow **13**
- Salamander abundance across forest types . . . 13
- Checking ANOVA's assumptions . . . 14
- Effect size: eta-squared . . . 16
- Post-hoc comparisons with Tukey HSD . . . 17
- Nonparametric alternative: the Kruskal-Wallis test . . . 19

Repeated-measures ANOVA **20**
- Fine particulate matter at four monitoring stations . . . 21

Splitting off the stations 21

Chapter Summary **23**

ANOVA is analysis of variance

ANOVA stands for analysis of variance. It tests whether the means of three or more groups differ by more than chance alone would produce. Chapters 6 and 7 compared two groups, and with two groups a t-test answers this question. Environmental studies often compare more than two, such as several sites, treatments, or time periods, and ANOVA extends the t-test's logic to them.

This chapter starts with why running a t-test for every pair of groups goes wrong, and with the single question ANOVA asks instead. It then builds ANOVA's test statistic, F, as a ratio, shows what values F takes when there are no real differences, and computes F by hand for a small example. The full workflow on one example follows, through post-hoc comparisons and a rank-based (nonparametric) alternative, and the chapter ends with repeated-measures ANOVA.

> **Note: Where ANOVA came from**
>
> Sir Ronald Fisher invented the ANOVA (Salsburg 2001) and wanted to publish it in the journal Biometrika. The editor at the time was Karl Pearson, of Pearson's r, and the two men disliked each other, so Pearson refused to publish the new test. Fisher published it in the Journal of Agricultural Science instead. The feud carried into the next generation. Years later, Karl Pearson's son Egon Pearson, working with Jerzy Neyman, revamped Fisher's ideas and re-cast them into what we now call null vs. alternative hypothesis testing. Fisher objected to that too.

Why not run multiple *t*-tests?

With three groups, A, B and C, there are three possible pairwise differences:

- Group A vs. Group B
- Group A vs. Group C
- Group B vs. Group C

We could test each with a separate t-test, asking whether that pair's means differ significantly.

The problem is that every test carries its own 5% chance of a false positive (a Type I error, Chapter 5), and those chances accumulate across tests. This follows from the independence multiplication rule (Section 3.5.2). Each test has a $1 - \alpha = 0.95$ chance of avoiding a false positive. If the m tests are independent, the chance that all of them avoid one is 0.95 multiplied by itself m times, or $(1 - \alpha)^m$. The chance of at least one false positive is 1 minus that, so for m independent tests at $\alpha = 0.05$,

$$P(\text{at least one false positive}) = 1 - (1 - \alpha)^m.$$

With three groups, that's three tests. The **familywise error rate** is the chance of at least one false positive across a set of tests. For three tests it is 14%, compared with 5% for a single test:

$$1 - (1 - 0.05)^3 = 0.14.$$

With k groups there are $k(k-1)/2$ pairs to compare, so the number of tests grows faster than the number of groups. Four groups give $4 \times 3/2 = 6$ tests, and five groups give $5 \times 4/2 = 10$ (Table 1).

Table 1: Familywise error rate for independent tests at $\alpha = 0.05$.

Number of groups (k)	Pairwise tests (m)	Familywise error rate
3	3	$1 - 0.95^3 = 0.14$
4	6	$1 - 0.95^6 = 0.26$
5	10	$1 - 0.95^{10} = 0.40$

Even with only five groups, there's a 40% chance of turning up at least one "significant" difference somewhere purely by chance, even if every population mean is identical. Figure 1 shows how the chance of at least one false positive grows as the number of independent t-tests grows.




Figure 1: Familywise Type I error rate as the number of independent t-tests increases, at $\alpha = 0.05$ per test.

Pairwise t-tests that share a group are not fully independent, so for pairwise comparisons these rates run a little high. Simulating three groups of 10 with no real differences gives about 12% for three pairwise tests, 20% for four groups, and 29% for five. The rate still climbs quickly as groups are added.

ANOVA solves this by replacing the separate t-tests with one test, so the chance of a false positive stays at α.

ANOVA is an omnibus test

An **omnibus test** is a single test that covers several comparisons at once, and ANOVA is one. Instead of the three pairwise t-tests listed above, ANOVA asks one more general question about all three means:

- **Null (H_0):** no differences anywhere, $\mu_A = \mu_B = \mu_C$.
- **Alternative (H_a):** a difference somewhere, in one of four patterns:
  - $\mu_A \neq \mu_B = \mu_C$ (A differs)
  - $\mu_B \neq \mu_A = \mu_C$ (B differs)
  - $\mu_C \neq \mu_A = \mu_B$ (C differs)

- $\mu_A \neq \mu_B$, $\mu_B \neq \mu_C$, and $\mu_A \neq \mu_C$ (all three differ)

A large p-value means we don't reject the hypothesis of no differences. Because the differences between the sample means could easily be due to chance, we have no evidence that the three population means differ. A small p-value (below α) means we reject that hypothesis and conclude at least one group mean differs from the others. The omnibus test tells us *whether* the means differ, and a separate step tells us *which* ones.

Kinds of ANOVA

As with the t-test, different research designs call for different kinds of ANOVA.

The one-way ANOVA is the simplest version of the test. In ANOVA an independent variable is called a **factor**, and the groups it defines are its **levels**. A one-way ANOVA has one factor with two or more levels.

Table 2 sets the one-way ANOVA beside the two other kinds of ANOVA this book covers.

Table 2: Three kinds of ANOVA.

ANOVA type	Factors and units	Design
One-way ANOVA	One factor with two or more levels, with different units in each group	Between-subjects
Repeated-measures ANOVA	One factor with two or more levels, with the *same* units measured at every level	Within-subjects
Two-way (factorial) ANOVA	Two factors, each with two or more levels, fully crossed (Chapter 10)	Factorial

The *F* ratio

ANOVA's test statistic is F, named for Fisher, who invented ANOVA. Like z and t, it divides a measure of effect by the variation that sampling error alone would produce (Chapter 4).

$$F = \frac{\text{measure of effect}}{\text{measure of error}}$$

The t-test puts a difference between two means over its standard error. F uses variances for both parts. It divides the variation **between** groups (how far the group means are from one another) by the variation **within** groups (how far individual scores are from their own group's mean).

$$F = \frac{\text{Variation between groups}}{\text{Variation within groups}}$$

When the two are about the same size, $F \approx 1$, and the group means differ no more than ordinary sampling noise would make them. When the between-group variation is much larger than the within-group variation, F climbs well above 1, which is evidence that the groups differ. When the between-group variation is smaller, F drops below 1. An F of 5 means the group means vary five times as much as within-group noise alone would make them vary, and an F of 0.6 means they vary less than that noise alone would typically produce. Both parts of F are variances, so "five times as much" compares squared distances, not raw distances.

With two groups, F is exactly the square of the pooled t (Section 6.5), $t^2 = F$, which is why the two tests give the same p-value.

Figure 2 shows both situations with the same noise. Both panels are simulated, so we know the population means behind them. In the left panel all three population means are 10, so any differences among the group means are chance. In the right panel the population means are 10, 12.2 and 15. The dashed line in each panel is the **grand mean**, the mean of all the scores across every group.




Figure 2: Simulated scores with the same within-group noise. Left: all three population means are 10, so there are no real differences. Right: the population means are 10, 12.2 and 15. Three groups of 12; small points are individual scores, large dots are group means, bars are 95% confidence intervals. The blue bracket marks within-group variation, the spread of Group A's scores. The orange arrows mark between-group variation, each group mean's distance from the grand mean (dashed line).

The within-group variation is the same in both panels. Group A's scores span about 9 units, the length of the blue bracket. In the "No real differences" panel (left), the three group means are all within about half a unit of the grand mean, so the between-group variation is small compared with the within-group variation, and $F = 0.64$. In the "Real differences" panel (right), Group C's mean is about 3 units above the grand mean, and $F = 21.4$. The within-group variation did not change, so the larger F comes entirely from the larger between-group variation.

F when H_0 is true

Before computing F by hand, we need to know what values it takes when the null hypothesis is true, because that is what an observed F is compared against. The p-value for F means the same thing as for every other test we've covered. It is the probability of an F this large or larger if the null hypothesis (no real group differences) were true.

Since F is a sample statistic, we can find out how it behaves the same way we did for t, by simulating it many times under the null. Consider a monitoring study with three site types (A, B, C) and 10 sampling plots per type, 30 in all, measuring species abundance at each plot. All 30 plots are drawn from one null population distribution (population mean $\mu = 100$, $\sigma = 10$), so site type has no real effect, and any differences between the group means come from sampling error alone. We repeat this study 10,000 times and compute F each time. The histogram of those 10,000 F-values, in Figure 3, is the null distribution of F for this design. The

null population distribution describes individual abundance scores. The null distribution of F describes the F-values, one per experiment.

Before you look. Each bar in Figure 3 counts F-values from these null studies. Decide whether most of them will fall near 0, near 1, or well above 1.




Figure 3: A simulation of 10,000 studies from a null population distribution with no group differences. The histogram shows the 10,000 F-values, one per study, all produced by sampling error alone. The blue dot-dash curve is the F distribution with 2 and 27 degrees of freedom. The red line is the critical value at $\alpha = 0.05$, and the shaded bars beyond it hold 5% of the F-values.

Most simulated F-values here fall near or below 1, because under the null the numerator and denominator of F estimate the same thing. All 30 plots come from one null population distribution with $\sigma = 10$. The denominator of F, the within-group variation, estimates σ^2 from the spread of scores inside each group. The numerator, the between-group variation, estimates the same σ^2 from how far the group means stray from the grand mean, since with no real effect, sampling variation is the only thing moving them. Two estimates of one variance give a ratio near 1. When the population means really differ, the numerator picks up those differences on top of σ^2, and F grows. (Section shows how each part is computed.)

The F of 5 from Section is rare under the null. Values that large occur in about 1% of the simulations.

The null distribution of F never goes below 0, because F is a ratio of two variances and variances can't be negative. It is right-skewed.

From simulation to the F distribution. The blue dot-dash curve over the histogram is the F **distribution**, the theoretical distribution of F when the null hypothesis is true. Its shape is set by two degrees of freedom, the numerator df ($k-1 = 2$ for the three site types) and the denominator df ($N-k = 30-3 = 27$, where N is the total number of plots). It is worked out from theory with no simulation, and it follows the histogram closely, so from here on we use the curve. Software does the same, and the p-value it reports for F comes from the F distribution for the design's two dfs.

Making decisions

To decide whether a real study's F is too large to be explained by chance:

1. Set $\alpha = 0.05$.
2. Find the critical value of F for the design's two dfs (in R, `qf(0.95, 2, 27)`), the red line in Figure 3.
3. Reject H_0 if the observed F is at or beyond it.

For 2 and 27 degrees of freedom the critical value is 3.35, and only 5% of the F distribution lies at or beyond it. The simulation agrees, with 4.8% of the 10,000 simulated F-values at or beyond 3.35. The F-test has one critical value, in the upper tail. Differences among the means in any direction make the numerator of F larger, so they all push F up, and a small F is consistent with H_0.

If we ran the same design for real (30 plots, 10 per site type), an observed F at or beyond 3.35 would lead us to reject H_0 and conclude that mean abundance differs among site types. The critical value and the p-value give the same decision. F is at or beyond the critical value exactly when its p-value is at or below α.

What group means look like when *F* exceeds the critical value

The 10,000 simulated studies kept only each F-value, not the group means behind it. To see those means, we reran the same null simulation and kept the first 10 studies with $F > 3.35$, shown in Figure 4. Every one of them was still drawn from the same null population distribution, with no real group differences. Rejecting the null whenever F clears 3.35 is the right long-run decision rule, but every one of these particular simulations would be a Type I error if we did.




Figure 4: Group means from 10 null simulations in which F exceeded the critical value. Error bars are 95% confidence intervals. Every one of them is a Type I error.

In four of the ten panels in Figure 4, at least one pair of 95% confidence intervals doesn't overlap, which looks like a convincing real difference. In the others, the means are still spread out enough for F to clear

the critical value. When chance produces a pattern like these, the F-value is large and the p-value is small, so in the output a Type I error looks exactly like a real effect.

Studies like these ten are the 5% the critical value allows. One F-test keeps that chance at 5% for the whole set of groups, where three separate pairwise t-tests would have raised it to about 12%. The same risk returns after a significant F, when post-hoc comparisons test each pair of groups, and that is why those comparisons are adjusted (Section).

Computing *F* by hand

ANOVA splits the total variation in the data into two sources: variation between groups, which group membership can explain, and variation within groups, which it can't. If we can measure those two parts separately, we can turn them into a ratio and get an F-value.

$$\text{Total variation} = \text{Variation between groups} + \text{Variation within groups}$$

To make that precise we need an actual measure of variation. As in Chapter 2, the measure is the sum of squares.

$$SS_{\text{Total}} = SS_{\text{Between}} + SS_{\text{Within}}$$

The next sections compute each piece for a small example with nine scores. We find SS_{Total} and SS_{Between}, get SS_{Within} by subtraction, and check it directly. Then we divide each sum of squares by its degrees of freedom, take the ratio to get F, and compare it with its critical value.

Nine litter counts

Suppose a city counts the pieces of litter along nine stretches of sidewalk, three in each of three neighborhoods, A, B, and C. The counts are invented. Each count is one score. Table 3 lists them by neighborhood, with each neighborhood's mean and the grand mean, the mean of all nine scores.

Table 3: Litter counts on nine stretches of sidewalk, three per neighborhood.

Neighborhood	Scores	Group mean
A	20, 11, 2	11
B	6, 2, 7	5
C	2, 11, 2	5
All nine		7 (grand mean)

Figure 5 plots the same nine scores. Neighborhood A's mean is above the grand mean, B's and C's are below it, and within each neighborhood the scores spread widely around their own mean. The tables in the next sections list them in long format, which means each row is a single score, the way R stores them.




Figure 5: The nine litter counts. Points are individual scores, short solid bars are the neighborhood means, and the dashed line is the grand mean, 7.

SS Total

SS_{Total} measures all of the variation in the data, how far every score is from the grand mean. Each score's distance from the grand mean is its deviation. Table 4 lists every deviation and its square.

Table 4: Deviations from the grand mean.

Neighborhood	Score	Deviation from grand mean	Squared deviation
A	20	13	169
A	11	4	16
A	2	-5	25
B	6	-1	1
B	2	-5	25
B	7	0	0
C	2	-5	25
C	11	4	16
C	2	-5	25
Sums	63	0	302

The deviations sum to zero, because the mean is the balancing point of the data, so their sum can't measure spread. Squaring turns every deviation positive and, as a side effect, weights larger deviations more heavily.

Adding up the squared deviations gives the sum of squares (SS) from Table 2.3. Writing Y_{ij} for score i in group j and $\bar{Y}_{\text{grand}}$ for the grand mean,

$$SS_{\text{Total}} = \sum (Y_{ij} - \bar{Y}_{\text{grand}})^2 = 302.$$

That is the first piece of the partition:

$$SS_{\text{Total}} = SS_{\text{Between}} + SS_{\text{Within}}, \qquad 302 = SS_{\text{Between}} + SS_{\text{Within}}.$$

We only need to compute one of the two remaining pieces directly. The other follows by subtraction.

SS Between

SS_{Total} captured all of the variation in the data. SS_{Between} is the part of the total variation that group membership can explain. It measures how far each group's mean is from the grand mean.

To find it, replace every score with its own group's mean (Table 5), then square and sum each group mean's distance from the grand mean. Writing $\bar{Y}_j$ for the mean of group j and n_j for its number of scores,

$$SS_{\text{Between}} = \sum_j n_j(\bar{Y}_j - \bar{Y}_{\text{grand}})^2.$$

The n_j is there because every score in group j is replaced by the same group mean, so each group's squared distance counts once per score. Here every group has 3 scores. With unequal groups, larger groups count more.

Table 5: Group means replace the scores.

Neighborhood	Score	Group mean	Group mean minus grand mean	Squared
A	20	11	4	16
A	11	11	4	16
A	2	11	4	16
B	6	5	-2	4
B	2	5	-2	4
B	7	5	-2	4
C	2	5	-2	4
C	11	5	-2	4
C	2	5	-2	4
Sums	63	63	0	72

In the group mean column, Group A's mean is 11, so there are three 11s, one per score in group A. Groups B and C both average to 5, so the rest of the column is 5s.

Within a group every value is now identical, so this column throws away the individual variation inside each group and keeps only the differences between groups, which is what SS_{Between} measures.

Plugging the group means into the formula gives

$$SS_{\text{Between}} = 3(11-7)^2 + 3(5-7)^2 + 3(5-7)^2 = 48 + 12 + 12 = 72.$$

SS_{Within} is a sum of squares, so it cannot be negative. That means SS_{Between} can never be larger than SS_{Total}, and here 72 is smaller than 302.

SS Within

We already have $SS_{\text{Total}} = 302$ and $SS_{\text{Between}} = 72$, so we can solve for SS_{Within} by subtraction:

$$SS_{\text{Within}} = SS_{\text{Total}} - SS_{\text{Between}} = 302 - 72 = 230.$$

We can also calculate SS_{Within} directly from the data (Table 6), both to see where it comes from and to check the subtraction.

$$SS_{\text{Within}} = \sum (Y_{ij} - \bar{Y}_j)^2$$

Table 6: Deviations from each group's own mean.

Neighborhood	Score	Group mean	Score minus group mean	Squared
A	20	11	9	81
A	11	11	0	0
A	2	11	-9	81
B	6	5	1	1
B	2	5	-3	9
B	7	5	2	4
C	2	5	-3	9
C	11	5	6	36
C	2	5	-3	9
Sums	63	63	0	230

The fourth column is the difference between each score and its own group mean, and the last column squares it. Summing the last column gives $SS_{\text{Within}} = 230$, the same value we got by subtraction.

SS_{Within} has several other names. It is also called the **error sum of squares**, since it is the variation left over after group membership has explained what it can. R labels its row `Residuals`. In the next sections we divide it by its degrees of freedom to get MS_{Within}, which some articles report as MSE (mean squared error).

Degrees of freedom

Degrees of freedom work the same way here as for the t-test. Every time we estimate something from the data, we use up a degree of freedom.

$df_{\text{Between}} = k - 1$, where k is the number of groups. With $k = 3$, $df_{\text{Between}} = 3 - 1 = 2$. Once we fix the grand mean, two of the three group means are free to be anything, but the third is determined by the other two.

$df_{\text{Within}} = N - k$, where N is the total number of scores. With $N = 9$ and $k = 3$, $df_{\text{Within}} = 9 - 3 = 6$. Estimating the 3 group means uses up 3 degrees of freedom, one per group. Within each group, the deviations from the group mean must sum to zero, so only 2 of a group's 3 deviations are free to vary. Three groups with 2 free deviations each give $df_{\text{Within}} = 6$.

$df_{\text{Total}} = N - 1$, and it splits the same way the sums of squares do: $df_{\text{Total}} = df_{\text{Between}} + df_{\text{Within}}$, here $8 = 2 + 6$.

Mean squares

We're building toward $F = \dfrac{\text{measure of effect}}{\text{measure of error}}$, and the first idea is to divide SS_{Between} by SS_{Within} directly. The problem is that each sum of squares grows with the number of free pieces that go into it, its degrees of freedom. SS_{Between} comes from 3 group means, which have 2 degrees of freedom, and SS_{Within} comes from 9 deviations with 6 degrees of freedom. Even with no real differences, SS_{Within} would tend to be about three times as large as SS_{Between} for that reason alone. Dividing each one by its degrees of freedom turns it into an average, and two averages can be compared directly.

A **mean square** is a sum of squares divided by its degrees of freedom. We divide by degrees of freedom rather than by n, because each group mean we estimate uses up a degree of freedom, as the previous section showed.

$$MS_{\text{Between}} = \frac{SS_{\text{Between}}}{df_{\text{Between}}} = \frac{72}{2} = 36$$

$$MS_{\text{Within}} = \frac{SS_{\text{Within}}}{df_{\text{Within}}} = \frac{230}{6} = 38.33$$

Both are variances. MS_{Within} measures how much individual scores vary around their own group's mean. MS_{Between} measures how much the group means vary, multiplied by the number of scores per group. Here the group means 11, 5 and 5 have variance

$$\frac{(11-7)^2 + (5-7)^2 + (5-7)^2}{3-1} = \frac{24}{2} = 12,$$

and $3 \times 12 = 36$, which is MS_{Between}. The multiplication is needed because a mean of n scores varies less than a single score does. The variance of a mean is σ^2/n, the squared standard error from Section 3.10. Multiplying by n puts MS_{Between} on the same scale as MS_{Within}.

Calculate *F*

With MS_{Between} and MS_{Within} in hand, F is their ratio:

$$F = \frac{MS_{\text{Between}}}{MS_{\text{Within}}} = \frac{36}{38.33} = 0.94$$

The ANOVA table

An **ANOVA table** lists the sums of squares, degrees of freedom, mean squares, and F for each source of variation. Table 7 gives the formulas, and Table 8 fills them in for the toy data.

Table 7: The one-way ANOVA table in formulas.

Source	df	SS	MS	F
Between groups	$k-1$	$\sum_j n_j(\bar{Y}_j - \bar{Y}_{\text{grand}})^2$	$SS_{\text{Between}}/df_{\text{Between}}$	$MS_{\text{Between}}/MS_{\text{Within}}$
Within groups	$N-k$	$\sum(Y_{ij} - \bar{Y}_j)^2$	$SS_{\text{Within}}/df_{\text{Within}}$	
Total	$N-1$	$\sum(Y_{ij} - \bar{Y}_{\text{grand}})^2$		

Table 8: R's ANOVA table for the toy data.

Source	df	SS	MS	F	p
groups	2	72	36.00	0.94	0.442
Residuals	6	230	38.33		

R labels the two rows `groups` (the between-groups row: what the means can explain) and `Residuals` (the within-groups row: what they can't). Other software uses slightly different labels for the same two rows. R leaves out the Total row. Apart from the p-value in the last column, the table contains only numbers we already computed, and MS_{Between} (36) divided by MS_{Within} (38.33) gives the F-value.

With $F = 0.94$, the group means vary no more than the noise within groups would make them vary. The critical value of F at $\alpha = 0.05$ for 2 and 6 degrees of freedom is 5.14 (`qf(0.95, 2, 6)`), and 0.94 is far below it, so we fail to reject H_0, as in Section . R's p-value, 0.44, gives the same decision. These nine scores give no evidence that the neighborhoods differ.

The full ANOVA workflow

In practice, software computes the ANOVA table. We computed one by hand so that each number in the table has a meaning when we read it.

The following example applies the full ANOVA workflow, from descriptive summaries through the omnibus F test to post-hoc comparisons, to a salamander study.

Salamander abundance across forest types

Plethodontid salamanders are sensitive indicators of forest condition, because they depend on moist microhabitats, abundant coarse woody debris, and stable soil temperatures, all features that decline with forest disturbance. A field ecologist surveys salamander abundance across three forest types, laid out in Figure 6. The counts in this example are invented. We built them so that old-growth plots average about 14 salamanders, secondary forest about 7, and managed plantation about 5, with 8 plots of each type.




Figure 6: The salamander study as a design schematic. Structurally this is the two-group design from Chapters 4 and 6 with one more box added.

More groups mean more pairwise comparisons, which raises the familywise error rate from the start of the chapter.

The hypotheses are:

- **Null (H_0):** mean salamander abundance is the same in all three forest types, $\mu_{\text{Old-growth}} = \mu_{\text{Secondary}} = \mu_{\text{Managed}}$.
- **Alternative (H_a):** at least one forest type's mean differs.




Figure 7: Mean salamander abundance (individuals per plot) across three forest types. Points show individual plot counts, and error bars show 95% confidence intervals.

The bar graph shows a gradient. Old-growth plots have the highest mean abundance, managed plantations the lowest, and secondary forest is in between. The individual plot counts (dots) spread widely within each forest type. Old-growth plots range from 10 to 19 salamanders and secondary plots from 3 to 13, so even where a real difference exists, individual plots from different forest types overlap.

Checking ANOVA's assumptions

Before running the ANOVA, check whether its assumptions hold. Section 6.3.5 introduced independence as the assumption every test in this book shares, so we do not list it again here. ANOVA adds two assumptions of its own.

The first is **homogeneity of variance**, which means the groups have roughly equal spread around their own means. The second is approximately normal residuals. In ANOVA the model predicts each score with its group mean, so a residual is the score minus its group mean, the same idea as the residuals in Figure 2.7. Here each residual is a plot's count minus its forest type's mean, the same deviations that make up SS_{Within}, and the residuals should be roughly bell-shaped.

Both matter because the F-test pools the within-group spread of every group into one error variance, MS_{Within}, and then compares F to a null distribution that assumes normal residuals. Pooling only makes sense when the groups have similar spread.

Levene's test (Levene 1960; Brown and Forsythe 1974) checks homogeneity of variance. Its null hypothesis is that every group has the same variance, so a small p-value is evidence that the spreads differ. It reports its own F statistic, separate from the ANOVA's F. The Shapiro-Wilk test from Section 6.3.5 checks normality, here applied to the residuals.

```
car::leveneTest(abundance ~ forest_type, data = salamander_df)
```

Table 9: Levene's test for homogeneity of variance.

Statistic (F)	df	df residual	p
0.52	2	21	0.599

The residuals come from the fitted model, so we fit the ANOVA first and set its table aside until the checks are done.

```
aov_out <- aov(abundance ~ forest_type, data = salamander_df)
shapiro.test(residuals(aov_out))
```

Table 10: Shapiro-Wilk test on ANOVA residuals.

Statistic (W)	p
0.93	0.078

Levene's test gives $F(2, 21) = 0.52$, $p = 0.599$, so there is no evidence of unequal variance across forest types. The Shapiro-Wilk test (Shapiro and Wilk 1965) on the model's residuals gives $W = 0.93$, $p = 0.078$, so there is no evidence against normality either. Both checks agree with the individual plots in Figure 7, which show similar spread across groups and no outliers far from the rest.

Formal assumption tests are evidence, not a gate. Section 6.3.5 made this point about Shapiro-Wilk, and it holds for Levene's test too. It's tempting to treat Levene's test as a strict gate, where significant means "stop" and not significant means "proceed." With a small sample (exactly when a violated assumption matters most), these tests have low power and will often say "no problem" even when there is one. With a very large sample (where ANOVA is fairly robust to mild violations anyway), they have high power and will flag trivial differences as "significant." Look at the data too, as we did with Figure 7 above.

In practice, one-way ANOVA's p-values stay close to correct under mild departures from these assumptions, especially when the design is **balanced**, with the same number of observations in every group, as here (8 plots per forest type). Be most cautious when small, unbalanced samples come with visibly skewed data or very different spreads. For very different spreads, use Welch's ANOVA, and for skewed residuals in a small or unbalanced sample, use the Kruskal-Wallis test. Both are described below.

Table 11 lists each assumption, how to check it, and what to use if it fails.

Table 11: ANOVA's assumptions, how to check them, and what to do if one fails.

Assumption	How to check	If it fails
Approximately normal residuals	Shapiro-Wilk test on the residuals, and a look at the scores in each group (Figure 7)	Kruskal-Wallis test
Homogeneity of variance	Compare the spread of each group in a plot, and Levene's test	Welch's ANOVA

With the assumptions reasonable, we run the one-way ANOVA to test whether these differences are larger than expected by chance alone.

```
summary(aov_out)
```

Table 12: One-way ANOVA table for salamander abundance by forest type.

Source	df	SS	MS	F	p
forest_type	2	312.58	156.29	14.60	< 0.001
Residuals	21	224.75	10.70		

Table 12 gives the result. With 3 forest types and 24 plots, the degrees of freedom are $3 - 1 = 2$ and $24 - 3 = 21$, and the critical value at $\alpha = 0.05$ is 3.47, a little higher than the 3.35 for 30 plots because the error term has fewer degrees of freedom. Our F of 14.60 is more than four times that cutoff. Values this large arise by chance less than 0.1% of the time, so we reject the null hypothesis that all three group means are equal.

Effect size: eta-squared

The F-test answers whether the group means differ by more than chance would produce. It says nothing about *how much* of the variation in salamander abundance forest type explains. Chapter 5 made the same distinction, where p told us whether a difference was detectable and the effect size told us how large it was, in the study's units or, with Cohen's d, in standard deviations. ANOVA adds a third kind, the share of the total variation that the groups explain.

ANOVA's effect size is **eta-squared** (η^2), and we already computed both of its ingredients when we partitioned the sums of squares:

$$\eta^2 = \frac{SS_{\text{Between}}}{SS_{\text{Total}}} = \frac{SS_{\text{Between}}}{SS_{\text{Between}} + SS_{\text{Within}}}$$

SS_{Between} is the variation the group means explain and SS_{Total} is all the variation there is. So η^2 is the proportion of the total variation in the response that the factor explains. It runs from 0 to 1.

Table 13: Eta-squared for the effect of forest type on salamander abundance.

SS Between	SS Within	SS Total	Eta-squared
312.58	224.75	537.33	0.582

Here $\eta^2 = 0.58$ (Table 13), so forest type explains about 58% of the variation in salamander abundance across these 24 plots. The remaining 42% is within-group variation, the plot-to-plot differences that forest type alone cannot explain.

Regression (Chapter 9, the next chapter) uses the same quantity under the name R^2. It also divides the total variation into a part the model explains and a part it leaves over, and reports the explained part as a proportion of the total.

Interpreting the number. Cohen's conventional benchmarks are $\eta^2 = 0.01$ (small), 0.06 (medium), and 0.14 (large), so 0.58 is a large effect by these benchmarks. Treat those cutoffs as rough orientation. What counts as a large effect depends on the system you are studying. An η^2 of 0.10 would be unremarkable in a controlled greenhouse experiment and notable in a field survey where dozens of uncontrolled factors also drive the response.

p depends on sample size, and η^2 does not. Survey enough plots and a difference of half a salamander per plot will clear $p < 0.05$ while explaining almost none of the variation. Reporting both keeps a statistically detectable result from being mistaken for an ecologically meaningful one.

Post-hoc comparisons with Tukey HSD

The omnibus F test tells us that at least one group mean differs, but not which pairs, and the ANOVA table does not report the group means at all. **Post-hoc comparisons** are tests run after a significant omnibus test to find which pairs of groups differ. This is a different use of "post-hoc" from post-hoc power (Section 5.4.4). To identify which forest types differ, we use **Tukey's Honestly Significant Difference (HSD)** test (Tukey 1949). Tukey HSD compares every pair of groups while keeping the familywise error rate across all of the comparisons at α, so it is the standard choice after a significant ANOVA.

```
TukeyHSD(aov_out)
```

Table 14: Tukey HSD post-hoc comparisons for salamander abundance by forest type.

Comparison	diff	lwr	upr	p adj
Secondary-Old-growth	-6.88	-11.00	-2.75	0.001
Managed-Old-growth	-8.25	-12.37	-4.13	< 0.001
Managed-Secondary	-1.37	-5.50	2.75	0.683

Reading Table 14, column by column. Each row is one pairwise comparison, and there are three of them because three groups make three pairs.

- `diff` is the difference in group means, computed as *the first group in the row label minus the second*: `Secondary-Old-growth` is secondary minus old-growth.
- `lwr` and `upr` are the lower and upper bounds of the 95% confidence interval for that difference.
- `p adj` is the p-value *after* adjusting for the fact that you are making three comparisons at once. Because of this adjustment, the chance of at least one false positive across all three comparisons stays at 5%, compared with about 12% for three unadjusted pairwise tests.

Watch the sign. The row `Secondary-Old-growth` has a diff of -6.88, and that negative sign does *not* mean secondary forest has more salamanders. It means secondary *minus* old-growth is negative, so old-growth is the higher of the two, by 6.88 individuals per plot. Always translate the sign back into a sentence naming which group is larger before you write anything down.

The confidence interval and the p-value say the same thing. A pair differs significantly at $\alpha = 0.05$ exactly when its 95% interval excludes zero. Check both columns against each other as a habit: `Secondary-Old-growth` runs from -11.00 to -2.75, entirely below zero, and its p is 0.001. `Managed-Secondary` runs from -5.50 to 2.75, straddling zero, and its p is 0.683. If they seem not to agree, reread that row.

So the three comparisons give:

- **Old-growth vs. Secondary.** Old-growth is higher by 6.88 individuals/plot, $p = 0.001$, a significant difference.
- **Old-growth vs. Managed.** Old-growth is higher by 8.25 individuals/plot, $p < 0.001$, a significant difference.
- **Secondary vs. Managed.** They differ by 1.37 individuals/plot, $p = 0.683$, not a significant difference.

A non-significant difference is one the test could not detect. Secondary and managed plantation differ by 1.37 salamanders per plot in this sample. The test says that with eight plots per group, we cannot distinguish it from sampling noise. A larger study might well separate them. Report it as "we did not detect a difference," never as "there is no difference."

Compact letter display. A compact letter display summarizes post-hoc results with a letter above each group. With more than three groups, a list of pairwise sentences gets long, so this is the usual summary. Groups that share a letter were not significantly different, and groups that share no letter were. Figure 8 puts the salamander results in that form.




Figure 8: Mean salamander abundance with Tukey letters. Error bars show 95% confidence intervals.

Old-growth gets its own letter because it differs from both of the others. Secondary and managed share a letter because the test could not separate them. Sometimes a group carries *two* letters, written "ab". Suppose group 1 is labeled "a" and group 3 is labeled "b" because those two differ significantly. Group 2 has a mean between them. If group 2 does not differ significantly from group 1 or from group 3, it shares a letter with each, so it is labeled "ab".

We built a difference of about 2 salamanders per plot between secondary forest and managed plantation into these invented counts, and the test could not detect it with 8 plots of each type.

! Results reporting: one-way ANOVA

Report the direction in plain language, the test statistic with both degrees of freedom, p, and the effect size, then which groups the post-hoc comparisons separated.

> Salamander abundance differed significantly among forest types, $F(2, 21) = 14.60$, $p < 0.001$, $\eta^2 = 0.58$. Tukey HSD comparisons showed more salamanders in old-growth plots than in secondary forest, by 6.88 per plot, 95% CI [2.75, 11.00], $p = 0.001$, and than in managed plantation, by 8.25 per plot, 95% CI [4.13, 12.37], $p < 0.001$. Secondary forest and managed plantation did not differ significantly, $p = 0.683$.

Table 15 collects the one-way ANOVA on one card.

Table 15: Test card: one-way ANOVA.

Purpose	Do the means of two or more separate groups differ? One factor, different units in each group
Hypotheses	$H_0 : \mu_1 = \mu_2 = \cdots = \mu_k$ H_a: at least one mean differs
Assumptions	Approximately normal residuals Similar spread in every group (homogeneity of variance)
Test statistic	$F = MS_{\text{Between}}/MS_{\text{Within}}$ $df_{\text{Between}} = k - 1$ $df_{\text{Within}} = N - k$
Report	$F(df_{\text{Between}}, df_{\text{Within}})$ and p $\eta^2 = SS_{\text{Between}}/SS_{\text{Total}}$ After a significant F, Tukey HSD comparisons naming which groups differ

Nonparametric alternative: the Kruskal-Wallis test

Our assumption checks above came back clean, so the ANOVA result stands as-is. Suppose instead the groups had visibly unequal spread. Then Welch's ANOVA (`oneway.test()` in R) compares the means without assuming equal variances, the same fix Welch's t-test makes (Table 6.4). If the residuals were clearly skewed in a small or unbalanced sample, the **Kruskal-Wallis test** (Kruskal and Wallis 1952) is the rank-based, nonparametric alternative to one-way ANOVA. It has the same relationship to ANOVA that the Mann-Whitney-Wilcoxon test has to the independent-samples t-test.

How it works, conceptually. Rank every observation across all groups together, from smallest to largest, ignoring group membership. Then check whether the average rank differs more across groups than chance alone would produce. If abundance differs among forest types, one group's observations should occupy systematically higher or lower ranks than the others. In the salamander data, the old-growth plots have a mean rank of 19.7 out of 24, against 10.1 for secondary and 7.8 for managed. If forest type had no effect, all three mean ranks would be close to 12.5, the middle of 1 to 24.

Its assumption. Kruskal-Wallis assumes the groups' distributions have roughly the same shape, and it does not require normality. When the shapes match, a significant result means the groups differ in location, meaning where the center of each distribution is, measured by the median, as with the Mann-Whitney-Wilcoxon test. If the shapes differ a lot, a significant result tells you the groups differ, but not whether that difference is in location, spread, or shape.

```
kruskal.test(abundance ~ forest_type, data = salamander_df)
```

Table 16: Kruskal-Wallis test on salamander abundance by forest type.

Statistic (χ^2)	df	p
13.09	2	0.001

R reports the Kruskal-Wallis statistic as χ^2 because, when the null hypothesis is true, it approximately follows a chi-square distribution with $k - 1 = 2$ degrees of freedom. Here $\chi^2(2) = 13.09$, $p = 0.001$, a significant result and the same conclusion as the ANOVA ($F(2, 21) = 14.60$, $p < 0.001$).

Post-hoc comparisons. Just as a significant ANOVA needs Tukey HSD to identify *which* groups differ, a significant Kruskal-Wallis result needs its own follow-up. We run a Mann-Whitney-Wilcoxon test on each

pair of groups and adjust the p-values with the Holm correction (Holm 1979), which keeps the familywise error rate across the comparisons at α.

```
pairwise.wilcox.test(salamander_df$abundance, salamander_df$forest_type,
                     p.adjust.method = "holm")
```

Table 17: Pairwise Mann-Whitney-Wilcoxon comparisons (Holm-adjusted).

Group 1	Group 2	p
Secondary	Old-growth	0.011
Managed	Old-growth	0.004
Managed	Secondary	0.486

The pattern matches Tukey HSD exactly. Old-growth forest differs significantly from both secondary forest ($p = 0.011$) and managed plantation ($p = 0.004$), while secondary forest and managed plantation do not differ from each other ($p = 0.486$). The two approaches use different assumptions and different calculations, and they agree here because the parametric assumptions held.

Table 18 collects the Kruskal-Wallis test on one card.

Table 18: Test card: Kruskal-Wallis test.

Purpose	Do two or more separate groups differ? Skewed residuals in a small or unbalanced sample
Hypotheses	H_0: group has no effect, so every group's mean rank is close to the middle rank H_a: at least one group ranks systematically higher or lower
Assumptions	Normality not required Reads as a difference in location (median) only when the groups' distributions have similar shape
Test statistic	Rank all observations together Compare the mean ranks across groups R reports the statistic as χ^2 $df = k - 1$
Report	$\chi^2(df)$ and p After a significant result, pairwise Mann-Whitney-Wilcoxon tests with a Holm correction

Repeated-measures ANOVA

Every unit in the salamander study was measured once: 24 plots, 8 per forest type. Many environmental studies instead measure the same monitoring stations in every season or the same streams before, during, and after a drought. A **repeated-measures** (within-subjects) design measures the same units under every condition, and it calls for a repeated-measures ANOVA.

This is ANOVA's version of the paired t-test (Section 6.4). The paired t-test subtracted each plot's before score from its own after score, so differences between plots dropped out and only the change was left. With three or more conditions there is no single difference to take, so repeated-measures ANOVA does the equivalent with sums of squares, measuring how much the units differ from one another and removing that variation from the error term.

Fine particulate matter at four monitoring stations

An air quality agency has four PM2.5 monitoring stations, ordered from the most rural (A) to the most urban (D). The concentrations in this example are invented. Table 19 gives each station's mean PM2.5 concentration in winter, spring, and summer. The question is whether concentrations differ by season.

- **Null (H_0):** mean PM2.5 is the same in every season, $\mu_{\text{Winter}} = \mu_{\text{Spring}} = \mu_{\text{Summer}}$.
- **Alternative (H_a):** at least one seasonal mean differs.

Table 19: Seasonal mean PM2.5 (µg/m³) at four monitoring stations, with each station's mean, each season's mean, and the grand mean (bottom right).

Station	Winter	Spring	Summer	Station mean
A	9	2	4	5
B	9	8	7	8
C	16	11	12	13
D	18	11	13	14
Season mean	13	8	9	10




Figure 9: Seasonal mean PM2.5 at four stations, one line per station.

Figure 9 draws the same numbers, one line per station. Winter is the highest season at every station. But the stations themselves differ far more than the seasons do. Station A averages 5 µg/m³ and station D averages 14.

Splitting off the stations

A one-way ANOVA on these 12 numbers would split the variation into two pieces, as before. The grand mean is 10 and the season means are 13, 8, and 9. Each season mean counts once per station, so

$$SS_{\text{Between}} = 4\left[(13-10)^2 + (8-10)^2 + (9-10)^2\right] = 4(14) = 56.$$

Table 20 gives each value's distance from the grand mean of 10.

Table 20: Distance of each PM2.5 value from the grand mean of 10.

Station	Winter	Spring	Summer
A	−1	−8	−6
B	−1	−2	−3
C	6	1	2
D	8	1	3

Squaring and summing the 12 distances, one season at a time, gives

$$SS_{\text{Total}} = (1 + 1 + 36 + 64) + (64 + 4 + 1 + 1) + (36 + 9 + 4 + 9) = 230,$$

which leaves $SS_{\text{Within}} = 230 - 56 = 174$.

A one-way ANOVA would also break the independence assumption, because each station contributes three of the 12 numbers, and readings from the same station tend to be alike.

Repeated-measures ANOVA splits SS_{Within} one more time. Part of it is station-to-station variation. The units measured repeatedly are called subjects, even when they are stations, so this piece is SS_{Subjects}, computed exactly like SS_{Between} but from the station means instead of the season means. Each station mean counts once per season, so

$$SS_{\text{Subjects}} = 3\left[(5-10)^2 + (8-10)^2 + (13-10)^2 + (14-10)^2\right] = 3(54) = 162.$$

Whatever is left is the error term:

$$SS_{\text{Error}} = SS_{\text{Within}} - SS_{\text{Subjects}} = 174 - 162 = 12,$$

so the full split is

$$SS_{\text{Total}} = SS_{\text{Between}} + SS_{\text{Subjects}} + SS_{\text{Error}}, \qquad 230 = 56 + 162 + 12.$$

SS_{Error} is the part of the within-season variation that the stations' own levels can't explain. It measures how much each station's seasonal pattern departs from the average seasonal pattern. Its degrees of freedom are the one-way df_{Within} minus the degrees of freedom the four station means use up, where n is the number of stations:

$$\begin{aligned} df_{\text{Within}} &= N - k = 12 - 3 = 9 \\ df_{\text{Subjects}} &= n - 1 = 4 - 1 = 3 \\ df_{\text{Error}} &= 9 - 3 = 6 = (k-1)(n-1) \end{aligned}$$

F uses the same numerator as before, $MS_{\text{Between}} = 56/2 = 28$. Only the denominator changes:

$$\text{One-way: } F = \frac{28}{174/9} = \frac{28}{19.33} = 1.45$$

$$\text{Repeated measures: } F = \frac{28}{12/6} = \frac{28}{2} = 14.00$$




Figure 10: The same 12 PM2.5 values split two ways. Both bars total 230. The one-way analysis leaves the station differences in the within-groups error. The repeated-measures analysis removes them (dashed gray piece), so the error term shrinks from 174 to 12 and F grows from 1.45 to 14.00.

Figure 10 draws the two splits of the same 230, and Table 21 sets the two analyses side by side.

Table 21: The same 12 PM2.5 values analyzed as a one-way ANOVA and as a repeated-measures ANOVA. Error is the within-groups row in the one-way analysis, and F is the season mean square divided by the error mean square.

Source	One-way df	One-way SS	One-way MS	Repeated-measures df	Repeated-measures SS	Repeated-measures MS
Season	2	56	28	2	56	28
Stations				3	162	54
Error	9	174	19.33	6	12	2
Total	11	230		11	230	
F			1.45 ($p = 0.285$)			14.00 ($p = 0.005$)

The repeated-measures test finds a significant seasonal effect, $F(2, 6) = 14.00$, $p = 0.005$. The one-way ANOVA would give $F(2, 9) = 1.45$, $p = 0.285$, and miss it. Removing the stations shrank the error mean square from 19.33 to 2, and F grew by the same factor. A repeated-measures design has more power than a between-subjects design with the same number of measurements when the units differ from one another, as monitoring stations usually do.

Chapter Summary

Why it matters. Comparing more than two groups with a separate t-test for each pair raises the chance of at least one false positive with every group added. ANOVA solves that by asking one omnibus question first, and its variance-partitioning logic is the same idea that reappears in regression, factorial designs, and ANCOVA.

Core ideas

- **Many t-tests inflate the false-positive rate.** Each comparison carries its own chance of a false positive, and they accumulate. With m comparisons, the familywise error rate for independent tests is $1 - (1 - \alpha)^m$, which is already 14% for three tests.
- **Partition the variation.** Total variation splits into a piece explained by group membership (SS_{Between}) and unexplained residual variation (SS_{Within}).
- **F is a ratio of the two.** After dividing each sum of squares by its degrees of freedom to get a mean square, $F = MS_{\text{Between}}/MS_{\text{Within}}$. A large F means between-group differences are large relative to within-group noise.
- **Under H_0, F is near 1.** Both mean squares estimate the same σ^2, so F follows an F distribution with df_{Between} and df_{Within}. Differences in any direction make F larger, so the test has one critical value, in the upper tail.
- **Run the omnibus test first, then post-hoc comparisons.** The omnibus test asks whether *any* group means differ. If it is significant, run post-hoc comparisons (Tukey HSD) to find out *which* pairs differ.
- **η^2 is the effect size.** $SS_{\text{Between}}/SS_{\text{Total}}$ is the proportion of variation explained by the factor. It is the same quantity regression calls R^2 (Chapter 9), and it does not depend on sample size the way p does, so report it alongside F.
- **Check the assumptions.** ANOVA needs approximately equal variances across groups (Levene's test) and roughly normal residuals. Treat formal tests as evidence rather than a gate, since they have least power exactly when small samples make violations matter most.
- **When assumptions fail.** Welch's ANOVA handles unequal spread. The Kruskal-Wallis test is the rank-based alternative for skewed residuals in small samples, with `pairwise.wilcox.test()` as its post-hoc counterpart.
- **Repeated measures.** When the same units are measured under every condition, repeated-measures ANOVA splits SS_{Within} into SS_{Subjects} and a smaller SS_{Error}. The smaller error term is why it has more power, the same reason the paired t-test does.

Brown, M. B., and A. B. Forsythe. 1974. "Robust Tests for Equality of Variances." *Journal of the American Statistical Association* 69: 364–67.

Holm, S. 1979. "A Simple Sequentially Rejective Multiple Test Procedure." *Scandinavian Journal of Statistics* 6: 65–70.

Kruskal, W. H., and W. A. Wallis. 1952. "Use of Ranks in One-Criterion Variance Analysis." *Journal of the American Statistical Association* 47: 583–621. https://doi.org/10.1080/01621459.1952.10483441.

Levene, H. 1960. "Robust Tests for Equality of Variances." In *Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling*, edited by I. Olkin et al. Stanford University Press.

Salsburg, David. 2001. *The Lady Tasting Tea: How Statistics Revolutionized Science in the Twentieth Century*. Macmillan.

Shapiro, S. S., and M. B. Wilk. 1965. "An Analysis of Variance Test for Normality (Complete Samples)." *Biometrika* 52: 591–611.

Tukey, John W. 1949. "Comparing Individual Means in the Analysis of Variance." *Biometrics* 5 (2): 99–114.