

A Prickly Situation: Contrasting frequency and morphological measures in three cactus species

Malory Owen¹ | Dr. Chris Lortie¹ | York University¹ | www.ecoblender.org¹

Question

What measurable characteristics determine ecologically functional differences in three species of cacti?

Double Mutualism & Facilitation

- Positive interactions drive ecosystem infrastructure¹
- Birds are nectarivores and frugivores of cacti^{2, 3}
 - Double mutualism: two positive interactions between interspecifics⁴
 - Harsh environments promote double mutualism⁵
- Cacti are desert foundational species⁶




Figure 1: Do birds pollinate and disperse seeds of foundational plants?




Figure 2: Nectarivores visit higher and showier floral displays.⁷

Hypotheses and Predictions

Different species of cacti occupy different ecological and facilitating niches.

- Different cactuses will have different sizes and health which impacts interactor visitation.

Methods

- Transects or haphazard sampling
- Major axis, minor axis, vertical axis
- Health index 1-5
 - Scarification, rot, branch death

Results




Figure 4: *C. acanthocarpa* (1.04 meters) > *C. echinocarpa* (0.55 meters) > *O. basilaris* (0.17 meters)

Each cactus species had significantly different mean heights (*Kruskal-Wallis*, Chi-square = 3.71, $p > 0.0001$, $df = 52$).




Figure 5: *C. acanthocarpa* and *C. echinocarpa* had more individuals with health scores of 4 or 5, whereas *O. basilaris* had a even distribution of health scores.

C. acanthocarpa and *C. echinocarpa* are healthier than *O. basilaris* (Pearson's Chi-squared Test, X-squared = 27.325, $df = 8$, $p > 0.001$).




Figure 3: *C. acanthocarpa* was the most abundant and *O. basilaris* was the least abundant

Table 1: *C. acanthocarpa* had the largest size class bins.

Species	Small	Medium	Large
Cyllindropuntia acanthocarpa	<85cm	86cm - 152cm	>153cm
Cyllindropuntia echinocarpa	<45cm	46cm - 72cm	>73cm
Opuntia basilaris	<15cm	16cm - 22cm	>23cm

Conclusions/Future Research

Because health and size may be strong predictors of reproductive output, all three cactus species will likely fulfill different ecological interaction niches. Additional studies surveying reproductive outputs against these traits will provide further insight into the way phenotypical differences impact positive interactions, mutualism, and eventual facilitation. These easily reproducible studies can be expanded upon by monitoring nectarivorous and frugivorous interactors at different phenological stages.

Works Cited

1. Callaway, Ragan M. 1997. "Positive interactions in plant communities and the individualistic-continuum concept." *Oecologia* 112: 143–49.
2. Montiel, Salvador, and Carlos Montaña. 2000. "Vertebrate Frugivory and Seed Dispersal of a Chihuahuan Desert Cactus" 146 (2): 221–29. doi:10.1111/jpb.12297.
3. Gorostigue, P., and P. Ortega-Baes. 2016. "How specialised is bird pollination in the Cactaceae?" *Plant Biology* 18 (1): 63–72.
4. Kelly, Dave, Jenny J Ladley, Alastair W Robertson, and Jenny J Ladley. 2004. "Is dispersal easier than pollination? Two tests in New Zealand Loranthaceae." *New Zealand Journal of Botany* 42: 89–103. doi:10.1080/0028825X.2004.9512892.
5. Garcia, Maria B., Xavier Espadaler, and Jens M. Olesen. 2012. "Extreme Reproduction and Survival of a True Cliffhanger: The Endangered Plant *Boronia choudardii*." *PLOS One* 7 (9): 1–7. doi:10.1371/journal.pone.0044657.
6. Filazzola, A., Lortie, C. J. 2014. "A systematic review and conceptual framework for the mechanistic pathways of nurse plants." *Global Ecology and Biogeography* 23 (12): 1335–1345.
7. Wolf, L. L., and Hainsworth, A.R. 1990. "Non-Random Foraging by Hummingbirds: Patterns of Movement Between Ipomopsis." *Functional Ecology* 4 (2): 149–57.

