

# Mathematical Methods - III

## Week 2

Jayampathy Ratnayake

Faculty of Science  
Department of Mathematics  
Room 209(B)  
jratnaya@indiana.edu

September 23, 2016

In this lesson, we will ...

In this lesson, we will ...

- 1 derive and use **Mid Point**, **Trapezoidal** and **Simpson's** rules to estimate the integral of a function (on a finite interval),

In this lesson, we will ...

- 1 derive and use **Mid Point**, **Trapezoidal** and **Simpson's** rules to estimate the integral of a function (on a finite interval),

Last week, we considered two types of estimates to the area under the *continuous function*  $f(x)$ , between  $x = a$  and  $x = b$

Last week, we considered two types of estimated to the area under the *continuous function*  $f(x)$ , between  $x = a$  and  $x = b$

**Right sums:** Uses the right end point of each subdivision

**Left sums :** Uses the left end point of each subdivision

These are called **Riemann sums**

Last week, we considered two types of estimated to the area under the *continuous function*  $f(x)$ , between  $x = a$  and  $x = b$

**Right sums:** Uses the right end point of each subdivision

**Left sums :** Uses the left end point of each subdivision

These are called **Riemann sums**

### Remark

*Reimann, with other mathematician, studied the Theory of Integration and derived many techniques to find it, some of which we now learn in the high school.*

*His definition of Integral is based on the two types of approximations mentioned above.*

Consider the following steps



Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts

Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts
- 2 find the suitable height of the each rectangle

Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts
- 2 find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do

Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts
- 2 find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the

Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts
- 2 find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the right end point of each division

Consider the following steps

- 1 divide the interval  $[a, b]$  in to  $n$  equal parts
- 2 find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the right end point of each division  
for Left Sums, this will be the left end point of each division.

Consider the following steps

- ① divide the interval  $[a, b]$  in to  $n$  equal parts
- ② find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the right end point of each division  
for Left Sums, this will be the left end point of each division.
- ③ All divisions have the same width( $\Delta x$ ), equal to

Consider the following steps


- ① divide the interval  $[a, b]$  in to  $n$  equal parts
- ② find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the right end point of each division  
for Left Sums, this will be the left end point of each division.
- ③ All divisions have the same width( $\Delta x$ ), equal to  $\frac{b - a}{n}$ .




Consider the following steps

- ① divide the interval  $[a, b]$  in to  $n$  equal parts
- ② find the suitable height of the each rectangle  
this will depend on the type of sums you intend to do  
for Right Sums, this will be the right end point of each division  
for Left Sums, this will be the left end point of each division.
- ③ All divisions have the same width( $\Delta x$ ), equal to  $\frac{b - a}{n}$ .
- ④ Find the Riemann sum

Consider the division of the interval  $[a, b]$  in to  $n$  sub-intervals, as shown in the following picture




Consider the division of the interval  $[a, b]$  in to  $n$  sub-intervals, as shown in the following picture



$$\Delta x = \frac{b - a}{n}$$


Consider the division of the interval  $[a, b]$  in to  $n$  sub-intervals, as shown in the following picture



$$\Delta x = \frac{b - a}{n}$$

$$x_i = a + i \cdot \Delta x$$

Consider the division of the interval  $[a, b]$  in to  $n$  sub-intervals, as shown in the following picture




$$\Delta x = \frac{b - a}{n}$$

$$x_i = a + i \cdot \Delta x$$

Right Hand Sum with  $n$  divisions (  $RHS(n)$  ) is then given by

Consider the division of the interval  $[a, b]$  in to  $n$  sub-intervals, as shown in the following picture



$$\Delta x = \frac{b - a}{n}$$

$$x_i = a + i \cdot \Delta x$$

Right Hand Sum with  $n$  divisions (  $RHS(n)$ ) is then given by

$$\begin{aligned} RHS(n) &= \sum_{i=1}^n \Delta x f(x_i) = \left( \frac{b - a}{n} \right) \sum_{i=1}^n f(x_i) \\ &= \left( \frac{b - a}{n} \right) (f(x_1) + f(x_2) + \dots + f(x_n)) \end{aligned}$$

Similarly, the Left Hand Sum with  $n$  divisions (  $LHS(n)$  ) is given by

Similarly, the Left Hand Sum with  $n$  divisions (  $LHS(n)$  ) is given by

$$\begin{aligned} LHS(n) &= \sum_{i=0}^{n-1} \Delta x f(x_i) = \left( \frac{b-a}{n} \right) \sum_{i=0}^{n-1} f(x_i) \\ &= \left( \frac{b-a}{n} \right) (f(x_0) + f(x_1) + \dots + f(x_{n-1})) \end{aligned}$$



Similarly, the Left Hand Sum with  $n$  divisions (  $LHS(n)$  ) is given by

$$\begin{aligned} LHS(n) &= \sum_{i=0}^{n-1} \Delta x f(x_i) = \left( \frac{b-a}{n} \right) \sum_{i=0}^{n-1} f(x_i) \\ &= \left( \frac{b-a}{n} \right) (f(x_0) + f(x_1) + \dots + f(x_{n-1})) \end{aligned}$$

Where

Similarly, the Left Hand Sum with  $n$  divisions (  $LHS(n)$  ) is given by

$$\begin{aligned} LHS(n) &= \sum_{i=0}^{n-1} \Delta x f(x_i) = \left( \frac{b-a}{n} \right) \sum_{i=0}^{n-1} f(x_i) \\ &= \left( \frac{b-a}{n} \right) (f(x_0) + f(x_1) + \dots + f(x_{n-1})) \end{aligned}$$

Where

**Note that the difference in the two sums  $RHS(n)$  and  $LHS(n)$  is in the summation index.**

Similarly, the Left Hand Sum with  $n$  divisions ( $LHS(n)$ ) is given by

$$\begin{aligned} LHS(n) &= \sum_{i=0}^{n-1} \Delta x f(x_i) = \left( \frac{b-a}{n} \right) \sum_{i=0}^{n-1} f(x_i) \\ &= \left( \frac{b-a}{n} \right) (f(x_0) + f(x_1) + \dots + f(x_{n-1})) \end{aligned}$$

Where

**Note that the difference in the two sums  $RHS(n)$  and  $LHS(n)$  is in the summation index.**

For  $LHS$  we add the first  $n$  values of the function (out of  $n+1$  values).

Similarly, the Left Hand Sum with  $n$  divisions ( $LHS(n)$ ) is given by

$$\begin{aligned} LHS(n) &= \sum_{i=0}^{n-1} \Delta x f(x_i) = \left( \frac{b-a}{n} \right) \sum_{i=0}^{n-1} f(x_i) \\ &= \left( \frac{b-a}{n} \right) (f(x_0) + f(x_1) + \dots + f(x_{n-1})) \end{aligned}$$

Where

**Note that the difference in the two sums  $RHS(n)$  and  $LHS(n)$  is in the summation index.**

For  $LHS$  we add the first  $n$  values of the function (out of  $n+1$  values).

For  $RHS$  we add the last  $n$  values of the function (out of  $n+1$  values).

## Theorem (Riemann Integration)

*For nice functions ( like Continuous functions )*

$$\lim_{n \rightarrow \infty} RHS(n) = \lim_{n \rightarrow \infty} LHS(n)$$

## Theorem (Riemann Integration)

*For nice functions ( like Continuous functions )*

$$\lim_{n \rightarrow \infty} RHS(n) = \lim_{n \rightarrow \infty} LHS(n)$$

*This common limit is defined to be the integral of the function  $f$  in the interval  $[a, b]$ ,*

## Theorem (Riemann Integration)

*For nice functions ( like Continuous functions )*

$$\lim_{n \rightarrow \infty} RHS(n) = \lim_{n \rightarrow \infty} LHS(n)$$

*This common limit is defined to be the integral of the function  $f$  in the interval  $[a, b]$ , and denote it by*

## Theorem (Riemann Integration)

*For nice functions ( like Continuous functions )*

$$\lim_{n \rightarrow \infty} RHS(n) = \lim_{n \rightarrow \infty} LHS(n)$$

*This common limit is defined to be the integral of the function  $f$  in the interval  $[a, b]$ , and denote it by*

$$\int_a^b f(x) dx$$



## Theorem (Riemann Integration)

*For nice functions ( like Continuous functions )*

$$\lim_{n \rightarrow \infty} RHS(n) = \lim_{n \rightarrow \infty} LHS(n)$$

*This common limit is defined to be the integral of the function  $f$  in the interval  $[a, b]$ , and denote it by*

$$\int_a^b f(x) dx$$

We understand this limit as the "Area Under the Curve".

## Remark

## Remark

- 1 *Theorem is true even if we choose the height of the  $i^{\text{th}}$  rectangle to be  $f(\zeta_i)$  for any  $\zeta_i \in [x_{i-1}, x_i]$ , instead of the end points.*

## Remark

- ① *Theorem is true even if we choose the height of the  $i^{\text{th}}$  rectangle to be  $f(\zeta_i)$  for any  $\zeta_i \in [x_{i-1}, x_i]$ , instead of the end points.*

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \Delta x f(\zeta_i) = \int_a^b f(x) dx$$

Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

### Definition

A function  $F$  is called an anti-derivative for  $f$  if

Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

### Definition

A function  $F$  is called an anti-derivative for  $f$  if

$$\frac{dF}{dx} = F' = f$$

Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

### Definition

A function  $F$  is called an anti-derivative for  $f$  if

$$\frac{dF}{dx} = F' = f$$

### Theorem (Fundamental Theorem of Calculus)

*If  $f$  is continuous in the interval  $[a, b]$  and  $F$  is an anti-derivative of  $f$ ,*



Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

### Definition

A function  $F$  is called an anti-derivative for  $f$  if

$$\frac{dF}{dx} = F' = f$$

### Theorem (Fundamental Theorem of Calculus)

*If  $f$  is continuous in the interval  $[a, b]$  and  $F$  is an anti-derivative of  $f$ , then*

$$\int_a^b f(x)dx = F(b) - F(a)$$

Before we go into the numerical schemes, here is the theorem which relates the Riemann Integral with the Anti-derivative.

### Definition

A function  $F$  is called an anti-derivative for  $f$  if

$$\frac{dF}{dx} = F' = f$$

### Theorem (Fundamental Theorem of Calculus)

*If  $f$  is continuous in the interval  $[a, b]$  and  $F$  is an anti-derivative of  $f$ , then*

$$\int_a^b f(x)dx = F(b) - F(a)$$

- 1 In the previous slides we saw the Integration is the Limit of a process of taking "finite" approximations.

- 1 In the previous slides we saw the Integration is the Limit of a process of taking "finite" approximations.
- 2 Integrals can be approximated by using one of these finite approximations.

- ① In the previous slides we saw the Integration is the Limit of a process of taking "finite" approximations.
- ② Integrals can be approximated by using one of these finite approximations.
- ③ We can increase the accuracy by increasing the  $n$  (in this case the "number of divisions").

- ① In the previous slides we saw the Integration is the Limit of a process of taking "finite" approximations.
- ② Integrals can be approximated by using one of these finite approximations.
- ③ We can increase the accuracy by increasing the  $n$  (in this case the "number of divisions").
- ④ Different ways of taking finite approximations gives rise to various numerical methods for integration.

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval



## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval
- 3 mid point of the sub interval  $[x_{i-1}, x_i]$  is

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval
- 3 mid point of the sub interval  $[x_{i-1}, x_i]$  is

$$\xi = x_{i-1} + \Delta/2 = \frac{x_{i-1} + x_i}{2}$$

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval
- 3 mid point of the sub interval  $[x_{i-1}, x_i]$  is

$$\zeta_i = x_{i-1} + \Delta/2 = \frac{x_{i-1} + x_i}{2}$$

- 4 height of the rectangle is  $f(\zeta_i)$ .

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval
- 3 mid point of the sub interval  $[x_{i-1}, x_i]$  is

$$\zeta_i = x_{i-1} + \Delta/2 = \frac{x_{i-1} + x_i}{2}$$

- 4 height of the rectangle is  $f(\zeta_i)$ .

### Remark

*We can think of this as*

## Mid-Point method.

- 1 divide the interval  $[a, b]$  to  $n$  sub-intervals
- 2 in each sub-interval the area is approximated by the area of the rectangle whose height is the function value at the mid-point of the sub-interval
- 3 mid point of the sub interval  $[x_{i-1}, x_i]$  is

$$\zeta_i = x_{i-1} + \Delta/2 = \frac{x_{i-1} + x_i}{2}$$

- 4 height of the rectangle is  $f(\zeta_i)$ .

### Remark

*We can think of this as*

- *in each sub-interval  $i$  the function is approximated by the horizontal line passing through the point  $(\zeta_i, f(\zeta_i))$ .*

Some standard numerical methods, that we will discuss to day, to estimate an integral.

## 1 Mid-Point Rule

Some standard numerical methods, that we will discuss today, to estimate an integral.

### 1 Mid-Point Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by a horizontal line passing through  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .

Some standard numerical methods, that we will discuss today, to estimate an integral.

1 Mid-Point Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by a horizontal line passing through  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .

2 Trapezoidal Rule



Some standard numerical methods, that we will discuss today, to estimate an integral.

1 Mid-Point Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by a horizontal line passing through  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .

2 Trapezoidal Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by the straight line joining  $(x_i, f(x_i))$  and  $(x_{i+1}, f(x_{i+1}))$ .

Some standard numerical methods, that we will discuss today, to estimate an integral.

1 Mid-Point Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by a horizontal line passing through  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .

2 Trapezoidal Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by the straight line joining  $(x_i, f(x_i))$  and  $(x_{i+1}, f(x_{i+1}))$ .

3 Simpson's Rule

Some standard numerical methods, that we will discuss today, to estimate an integral.

1 Mid-Point Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by a horizontal line passing through  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .


2 Trapezoidal Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by the straight line joining  $(x_i, f(x_i))$  and  $(x_{i+1}, f(x_{i+1}))$ .

3 Simpson's Rule

Approximates the function in each sub-interval  $[x_i, x_{i+1}]$  by the quadratic polynomial passing through  $(x_i, f(x_i))$ ,  $(x_{i+1}, f(x_{i+1}))$  and  $\left(\frac{x_i + x_{i+1}}{2}, f\left(\frac{x_i + x_{i+1}}{2}\right)\right)$ .


# Mid-Point Rule



Mid-point rule for  $n$  division is

$$\int_a^b f(x)dx \approx \Delta x (f(a_0) + f(a_1) + \dots + f(a_{n-1}))$$

# Mid-Point Rule



Mid-point rule for  $n$  division is

$$\int_a^b f(x) dx \approx \Delta x (f(a_0) + f(a_1) + \dots + f(a_{n-1}))$$

Where

$$\Delta x = \frac{b-a}{n} \text{ and } a_i = \frac{x_i + x_{i+1}}{2}$$

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.



# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.
- 4 what is the connection between this value and the left and the right approximations corresponding to this division.

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.
- 4 what is the connection between this value and the left and the right approximations corresponding to this division.
- 5 divide the interval into 3 equal parts to approximate the integral as before.

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.
- 4 what is the connection between this value and the left and the right approximations corresponding to this division.
- 5 divide the interval into 3 equal parts to approximate the integral as before.
- 6 what is the connection between this value and the left and the right approximations corresponding to this division.

# Trapezoidal Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .


- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.
- 4 what is the connection between this value and the left and the right approximations corresponding to this division.
- 5 divide the interval into 3 equal parts to approximate the integral as before.
- 6 what is the connection between this value and the left and the right approximations corresponding to this division.
- 7 what is the rule for  $n$  divisions.

# Trapezoidal Rule


## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .


- 1 approximate the function by a straight line connecting  $(0, f(0))$  and  $(2, f(2))$ . What is the approximation for the area with this approximation for the function.
- 2 what is the connection between this value and the left and the right approximations corresponding to this division.
- 3 divide the interval  $[0, 2]$  into 2 equal parts. In each sub-interval, do as in the previous part and find an approximation for the integral.
- 4 what is the connection between this value and the left and the right approximations corresponding to this division.
- 5 divide the interval into 3 equal parts to approximate the integral as before.
- 6 what is the connection between this value and the left and the right approximations corresponding to this division.
- 7 what is the rule for  $n$  divisions.



- 1 divide the interval  $[a, b]$  to  $n$  equal parts.
- 2 what is the  $i$ -th interval




- 1 divide the interval  $[a, b]$  to  $n$  equal parts.
- 2 what is the  $i$ -th interval  $[x_{i-1}, x_i]$ .




- 1 divide the interval  $[a, b]$  to  $n$  equal parts.
- 2 what is the  $i$ -th interval  $[x_{i-1}, x_i]$ .
- 3 what is the area of the trapezoid approximating the area under the function in this sub-interval.






- 1 divide the interval  $[a, b]$  to  $n$  equal parts.
- 2 what is the  $i$ -th interval  $[x_{i-1}, x_i]$ .
- 3 what is the area of the trapezoid approximating the area under the function in this sub-interval.
- 4 hence write down the trapezoidal rule.



Trapezoidal rule for  $n$  division is

$$\int_a^b f(x) dx \approx \frac{\Delta x}{2} (f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n))$$



Trapezoidal rule for  $n$  division is

$$\int_a^b f(x) dx \approx \frac{\Delta x}{2} (f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n))$$

Where  $\Delta x = \frac{b - a}{n}$

## Example

What is the connection between the trapezoidal rule and the left and the right approximations corresponding to a subdivision.

## Example

- 1 what is the quadratic curve passing through the points  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$ .

## Example

- ① what is the quadratic curve passing through the points  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$ .
- ② What is the quadratic curve passing through the points  $(x_1, f(x_1))$ ,  $(x_2, f(x_2))$  and  $\left(\frac{x_1 + x_2}{2}, f\left(\frac{x_1 + x_2}{2}\right)\right)$ .

## Example

- ① what is the quadratic curve passing through the points  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$ .
- ② What is the quadratic curve passing through the points  $(x_1, f(x_1))$ ,  $(x_2, f(x_2))$  and  $\left(\frac{x_1 + x_2}{2}, f\left(\frac{x_1 + x_2}{2}\right)\right)$ .
- ③ What is the area bounded by this parabola in the interval  $[x_1, x_2]$ .

# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.



# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.

# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.
- 3 Use this curve to approximate the integral.

# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.
- 3 Use this curve to approximate the integral.
- 4 divide the interval  $[0, 2]$  in to 2 equal parts. In each sub-interval, do as in the previous part by taking the two end points and the mid point of the each subinterval to find an approximation for the integral.

# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .

- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.
- 3 Use this curve to approximate the integral.
- 4 divide the interval  $[0, 2]$  in to 2 equal parts. In each sub-interval, do as in the previous part by taking the two end points and the mid point of the each subinterval to find an approximation for the integral.
- 5 divide the interval in to 3 equal parts to approximate the integral as before.

# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .


- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.
- 3 Use this curve to approximate the integral.
- 4 divide the interval  $[0, 2]$  in to 2 equal parts. In each sub-interval, do as in the previous part by taking the two end points and the mid point of the each subinterval to find an approximation for the integral.
- 5 divide the interval in to 3 equal parts to approximate the integral as before.
- 6 what is the rule for  $n$  divisions.


# Simpson's Rule

## Example

Consider the integral  $\int_0^2 x \sin(x) dx$ .


- 1 Recall that in the trapezoidal rule, we approximate the function in each subinterval by a straight line connecting the end points of the curve in that interval.
- 2 Consider the two points  $(0, f(0))$ ,  $(2, f(2))$  and  $(1, f(1))$ . What is the quadratic curve passing through these three points.
- 3 Use this curve to approximate the integral.
- 4 divide the interval  $[0, 2]$  in to 2 equal parts. In each sub-interval, do as in the previous part by taking the two end points and the mid point of the each subinterval to find an approximation for the integral.
- 5 divide the interval in to 3 equal parts to approximate the integral as before.
- 6 what is the rule for  $n$  divisions.





Approximate the function in each interval  $[x_i, x_{i+1}]$  by the quadratic  $q_i$ , passing through  $x_i$ ,  $x_{i+1}$  and the midpoint  $m_i = \frac{x_i + x_{i+1}}{2}$ , for  $i = 0, 1, \dots, n - 1$ .





Approximate the function in each interval  $[x_i, x_{i+1}]$  by the quadratic  $q_i$ , passing through  $x_i$ ,  $x_{i+1}$  and the midpoint  $m_i = \frac{x_i + x_{i+1}}{2}$ , for  $i = 0, 1, \dots, n - 1$ .

$$q_i = f(x_i) \frac{(x - m_i)(x - x_{i+1})}{(x_i - m_i)(x_i - x_{i+1})} + f(x_{i+1}) \frac{(x - x_i)(x - m_i)}{(x_{i+1} - x_i)(x_{i+1} - m_i)} + f(m_i) \frac{(x - x_i)(x - x_{i+1})}{(m_i - x_i)(m_i - x_{i+1})}$$

In each interval  $[x_i, x_{i+1}]$ ,

In each interval  $[x_i, x_{i+1}]$ ,

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx \int_{x_i}^{x_{i+1}} q_i(x) dx = \frac{\Delta x}{6} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

In each interval  $[x_i, x_{i+1}]$ ,

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx \int_{x_i}^{x_{i+1}} q_i(x) dx = \frac{\Delta x}{6} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

Simpson's Rule with  $n$  divisions.

In each interval  $[x_i, x_{i+1}]$ ,

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx \int_{x_i}^{x_{i+1}} q_i(x) dx = \frac{\Delta x}{6} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

Simpson's Rule with  $n$  divisions.

$$\int_a^b f(x) dx \approx \frac{\Delta x}{6} \sum_{i=0}^{n-1} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

In each interval  $[x_i, x_{i+1}]$ ,

$$\int_{x_i}^{x_{i+1}} f(x) dx \approx \int_{x_i}^{x_{i+1}} q_i(x) dx = \frac{\Delta x}{6} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

Simpson's Rule with  $n$  divisions.

$$\int_a^b f(x) dx \approx \frac{\Delta x}{6} \sum_{i=0}^{n-1} [f(x_i) + 4f(m_i) + f(x_{i+1})]$$

Where

$$\Delta x = \frac{b - a}{n}$$
$$m_i = \frac{x_i + x_{i+1}}{2}$$

## Remark

## Remark

- 1 *It is not necessary to divide the interval in to equal pieces. What is important is that the "sizes of the intervals" should go to zero.*



## Remark

- 1 *It is not necessary to divide the interval in to equal pieces. What is important is that the "sizes of the intervals" should go to zero. This fact can be used to estimate the integral more efficiently. For example*

## Remark


- ① *It is not necessary to divide the interval in to equal pieces. What is important is that the "sizes of the intervals" should go to zero. This fact can be used to estimate the integral more efficiently. For example*
- in parts where the function is varying rapidly, one can use smaller divisions*

## Remark

- ① *It is not necessary to divide the interval in to equal pieces. What is important is that the "sizes of the intervals" should go to zero. This fact can be used to estimate the integral more efficiently. For example*
- *in parts where the function is varying rapidly, one can use smaller divisions*
  - *and for intervals the function does not change much, one can use larger divisions,*

## Remark

- ① *It is not necessary to divide the interval in to equal pieces. What is important is that the "sizes of the intervals" should go to zero. This fact can be used to estimate the integral more efficiently. For example*
- *in parts where the function is varying rapidly, one can use smaller divisions*
  - *and for intervals the function does not change much, one can use larger divisions,*
  - *increasing the speed and decreasing the resources used without compromising the accuracy.*



$xy$

When using a numerical method to estimate a value, it is the common practice to generate sequence of solutions with decreasing step sizes to while the solution has reached a desired accuracy

When using a numerical method to estimate a value, it is the common practice to generate sequence of solutions with decreasing step sizes to while the solution has reached a desired accuracy (while the numerical scheme converge).

## Algorithm

```

Integrate( $f$ ,  $a$ ,  $b$ ,  $\epsilon$ )
 $f$  : function to integrate
 $[a, b]$  : interval
 $\epsilon$  : precision ( required accuracy )
integer  $n = 10$  : number of divisions
 $sum\_new$  = estimate with  $n$  divisions
do
     $sum\_old = sum\_new$ 
     $n = 3n$  : increase the number of divisions
     $sum\_new$  = estimate with  $n$  divisions : now  $n$  has increased!
while |  $sum\_new - sum\_old$  |  $< \epsilon$ 
return  $sum\_new$ 
    
```

Absolute error =  $|\text{true value} - \text{estimated value}|$



Absolute error =  $| \text{true value} - \text{estimated value} |$

Relative error =  $\frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$

Absolute error =  $| \text{true value} - \text{estimated value} |$

$$\text{Relative error} = \frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$$

Sources of Errors in Numerical Calculations

$$\text{Absolute error} = | \text{true value} - \text{estimated value} |$$

$$\text{Relative error} = \frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$$

## Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.

Absolute error =  $| \text{true value} - \text{estimated value} |$

$$\text{Relative error} = \frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$$

### Sources of Errors in Numerical Calculations


- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due finite approximations.

$$\text{Absolute error} = | \text{true value} - \text{estimated value} |$$

$$\text{Relative error} = \frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$$

### Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due finite approximations.



- It is important to know the error estimates of our numerical schemes.

- It is important to know the error estimates of our numerical schemes.
- Errors will give us an exact idea of what we compute, compared to the exact solution.

- It is important to know the error estimates of our numerical schemes.
- Errors will give us an exact idea of what we compute, compared to the exact solution.
- We will do the error estimates of the three methods we have discussed so far in the next class.



Using the definition of the definite integral, you can easily verify that in all three methods we discussed, the answer converge to the exact solution as the number of divisions,  $n$ , gets larger and larger (i.e. as the step size goes to zero).

Using the definition of the definite integral, you can easily verify that in all three methods we discussed, the answer converge to the exact solution as the number of divisions,  $n$ , gets larger and larger (i.e. as the step size goes to zero).

$$\lim_{n \rightarrow \infty} |\text{Error with } n \text{ division}| = 0$$

Using the definition of the definite integral, you can easily verify that in all three methods we discussed, the answer converge to the exact solution as the number of divisions,  $n$ , gets larger and larger (i.e. as the step size goes to zero).

$$\lim_{n \rightarrow \infty} |\text{Error with } n \text{ division}| = 0$$

### Definition (Convergence)

Numerical solution approaches the exact solution as the step size goes to zero. i.e. errors go to zero

- Consistency
- Stability