

Category Theory

Malthe Sparring

May 22, 2022

1 Basic notions

Definition 1.1. A **category** C consists of a collection ¹ $ob(C)$ of objects, and for every two objects $a, b \in ob(C)$ a collection $C(a, b)$ of maps satisfying the following:

- If $f \in C(a, b)$ and $g \in C(b, c)$ then there exists $g \circ f \in C(a, c)$.
- There is a unique identity map $id_C \in C(c, c)$ such that $f \circ id_c = id_c \circ f = f$ for every f .

If $ob(C)$ is a finite set then we say C is **small**. If $C(a, b)$ is a set for every $a, b \in ob(C)$ then we say C is **locally small**

Definition 1.2. A (covariant) **functor** $F : C \rightarrow D$ between two categories is a mapping taking every $c \in ob(C)$ to an object $Fc \in D$ and every $f \in C(a, b)$ to a map $F(f) \in D(Fa, Fb)$ such that $F(g \circ f) = F(g) \circ F(f)$.

A **contravariant functor** F is a functor $F : C^{op} \rightarrow D$ from the opposite category. In other words, F reverses the direction of arrows in C .

Definition 1.3. A **natural transformation** $\eta : F \rightarrow G$ between two functors $F, G : C \Rightarrow D$ is, for every $x \in C$ a mapping $\eta_x : Fx \rightarrow Gx$ (called the component at x) such that the following diagram commutes.

$$\begin{array}{ccc} F(x) & \xrightarrow{\eta_x} & D(x) \\ \downarrow F(f) & & \downarrow G(f) \\ F(y) & \xrightarrow{\eta_y} & D(y) \end{array}$$

If each component η_x is an isomorphism, then we say η is a **natural isomorphism**.

¹These can be bigger than sets. I don't know the technical definition

Lets give many examples.

Example 1.4 (Categories). The following are categories.

- The category **Set** of sets and functions.
- The category **Cat** of small categories and functors.
- The category **CAT** of locally small categories and functors.
- The category **Ab** of abelian groups and group homomorphisms.
- The category **Grp** of groups and group homomorphisms.
- The category **Ring** of rings and ring homomorphisms.
- The category **R – Mod** of R –modules and module homomorphisms.
- The category **Top** of topological spaces and continuous maps.
- The category **Top*** of pointed topological spaces and pointed maps.
- The category **hTop** of pointed topological spaces and classes of homotopy equivalences.
- For a category C , the opposite category C^{op} where arrow directions are reversed.
- For a category C and an object x , the **slice category** or **over category** C/x whose elements are maps into x and maps commutative diagrams

$$\begin{array}{ccc} a & \longrightarrow & x \\ \downarrow f & \nearrow & \\ b & & \end{array}$$

- The similarly defined **under category** x/C of morphisms out of x and maps that form commutative triangles.
- The trivial category with a single object and a single identity map.
- For every group G the category C_G with one object and one map for every $g \in G$ with composition defined by the group law.
- A **poset** P is a category with a unique map $f : a \rightarrow b$ between any two objects, interpreted as a relation $a \leq b$.

- A **discrete category** is a category whose only maps are identity maps.
- For any two categories C, D , there is a category $Fun(C, D)$ of functors between them and natural transformations.

Example 1.5 (Functors). The following are functors.

- There is a large class of "forgetful" functors that forget extra structure. For example, the functors $U : \mathbf{Grp} \rightarrow \mathbf{Set}$, $U : \mathbf{Top} \rightarrow \mathbf{Set}$, $U : \mathbf{Ab} \rightarrow \mathbf{Grp}$, $U : \mathbf{Top}^* \rightarrow \mathbf{Top}$ which in each case forget the extra structure.
- There are also a large class of "free" functors, for example $F : \mathbf{Set} \rightarrow \mathbf{Ab}$ mapping a set to the free abelian group on it, $\delta : \mathbf{Set} \rightarrow \mathbf{Top}$ giving a set the discrete topology, $\tau : \mathbf{Set} \rightarrow \mathbf{Top}$ giving a set the chaotic topology, and the abelianisation functor $F : \mathbf{Grp} \rightarrow \mathbf{Ab}$ sending a group to the free abelian group on it.

Definition 1.6. Two functors $F : C \rightarrow D$ and $G : D \rightarrow C$ are said to be an **equivalence of categories** if there exist natural isomorphisms $FG \cong id_C$ and $GF \cong id_D$.

2 Abelian categories

Loosely, an **additive category** is one where morphisms can be added, and where we generally act like the category of abelian groups.

Definition 2.1. An **additive category** is a category C such that

- $Hom(a, b)$ is an abelian group under $+$, where \circ distributes over $+$:

$$f \circ (g + h) = (f + g) \circ (f + h)$$

$$(f + g) \circ h = (f \circ h) + (g \circ h)$$

- C has a **zero object**² 0 .
- C has finite (co)products and these two coincide.³

The abelian group structure will be different for each pair $a, b \in C$. In the case of Ab , the group structure on $Hom(a, b)$ is given by the group structure on b , but b needn't have a group structure in general. As an example, if J and A are categories with J small and A abelian, then $Fun(J, A)$ is an abelian (in particular additive) category.

²A **zero object** is an initial and final object.

³In fact, is enough to demand it has finite products - the rest will follow.

We want to extend this definition to encompass all categories of "abelian things". In particular, we want to be able to take kernels and cokernels of maps - this will allow us to do homological algebra. Notice that the (co)kernel has a universal property as respectively a pullback and a pushout.

$$\begin{array}{ccc}
 \ker(f) & \longrightarrow & A \\
 \downarrow & \lrcorner & \downarrow f \\
 0 & \longrightarrow & B
 \end{array}
 \qquad
 \begin{array}{ccc}
 A & \xrightarrow{f} & B \\
 \downarrow & & \downarrow \\
 0 & \longrightarrow & \operatorname{coker}(f)
 \end{array}$$

Definition 2.2. An **abelian** category is an additive category with kernels and cokernels. In addition, we require that every monomorphism is the kernel of its cokernel and every epimorphism is the cokernel of its kernel.

The additional requirement correspond to how we expect injective and surjective homomorphisms to act. If f is injective, then $\pi : B \rightarrow B/\operatorname{im} f$ has kernel $\operatorname{im}(f) \cong A$. If f is surjective, then $A/\ker(f) \cong \operatorname{im}(f) = B$. These are not an immediate property of the (co)kernels as (co)limits⁴. Kernels and cokernels allow us to define images as $\operatorname{im}(f) = \ker(\operatorname{coker}(f))$. These have pleasing properties: $A \rightarrow B$ factors uniquely through $\operatorname{im} f \rightarrow B$ and $A \rightarrow \operatorname{im}(f)$ is an epimorphism.

Cool fact alert!

Theorem 2.3 (Freyd-Mitchell Embedding Theorem). *Every abelian category s.t. $\operatorname{Hom}(X, Y)$ is a set embeds fully faithfully into Mod_R for some (possibly non-commutative) ring R .*

Proof. [1] □

Which abelian categories correspond to non-commutative rings? Also, is R unique?

Definition 2.4. A covariant functor $F : A \rightarrow B$ is right-exact if it preserves right-exact sequences. Similar for left-exact.

$$A \longrightarrow A' \longrightarrow A'' \longrightarrow 0$$

$$FA \longrightarrow FA' \longrightarrow FA'' \longrightarrow 0$$

A contravariant functor is left-exact if it takes right-exact sequences to left-exact sequences. Similar for right-exact.

⁴Can I give an example?

Lemma 2.5. *Limits commute with limits and right adjoints. In particular, in an abelian category, because kernels are limits, both right adjoints and limits are left-exact.*

Dually, colimits commute with colimits and left adjoints. In particular, because cokernels are colimits, both left adjoints and colimits are right-exact.

References

- [1] Charles A Weibel, *An introduction to homological algebra*, Univ. Pr, 1995.