The information cost of boundary-free weak estimation under correlated noise

Danil Romanovich Malyshev

Independent researcher, Almaty, Kazakhstan

Corresponding author: malyshevdanil04@gmail.com

ORCID: https://orcid.org/0009-0007-0947-3866

MSC 2020: Primary 62F12; secondary 62M10, 34A55.

Abstract

Boundary-free weak equations discard information about known initial states under correlated measurement error. For deterministic ODEs observed with stationary Ornstein–Uhlenbeck noise, we identify their closed adjoint influence class and sharp nuisance-adjusted information bound. The restriction raises the local RMSE reference by 29% for Van der Pol and almost doubles it for Lorenz. For polynomial libraries satisfying a nonzero trajectory-jet condition, we construct a feasible full-library estimator with smoothed-data instruments, vanishing regularization and bounded output. Pilot rank, covariance-weight consistency and uniform integrability are derived; the estimator attains finite-span covariance and reaches the weak-class bound in sequential small-noise and dense-span limits. Exact symbolic calculation verifies the condition for the eighteen-coefficient Van der Pol library. Known-initial traces restore the full closed influence class. This established adaptation, due to Brunel and coauthors, recovers about 97% and 95% of the empirical small-noise gaps in Van der Pol and Lorenz. Matched covariance benchmarks support constructive attainability while showing excess risk and incomplete boundary recovery at moderate noise.

Keywords: weak-form identification; correlated measurement error; fixed-domain asymptotics; Fisher information; polynomial covariance; errors in variables.

1. Introduction

Boundary-vanishing weak equations can lose information even when an initial state is known, as recognized by orthogonality-condition estimation [1] and recent weak identifiability work [2]. We quantify this effect: the closed class of boundary-free weak influences has exactly the local information available when the initial state is treated as a nuisance parameter in the full path likelihood. For an unknown initial state, this restriction causes no additional first-order information loss; for a known initial state, the lost information is the initial-sensitivity projection, and the retained information is its Schur complement. The equivalence concerns local information.

Known initial states also permit a remedy. Initial-trace conditions are established in orthogonality estimation [1]. We characterize their complete local influence class, quantify finite-family recovery under physical temporal correlation, and distinguish their contribution from an initial-noise control.

The control reveals a correlated error whose covariance with interior moments can reduce variance. The ablation determines its practical contribution rather than presuming it drives recovery.

Sparse identification of nonlinear dynamics represents a vector field through a library and selects active coefficients [3]. Integral formulations reduce sensitivity to numerical differentiation [4], and weak formulations transfer differentiation to test functions [5, 6]. Their regression matrices still contain noisy states. Statistical consistency depends jointly on feature correction, residual covariance and the acquisition limit.

The distinction between fixed-domain and increasing-domain observations is classical in Gaussian process inference [7]. Ying established fixed-domain results for exponential covariance parameters [8], and Zhang used equivalence of Gaussian measures to explain inconsistent covariance estimation in the Matérn family [9]. Those results concern covariance parameters. Here the covariance is common across parameters, and the parameter enters the deterministic mean through an ODE. The same measure-equivalence mechanism supplies an estimator-independent limit, with an explicit norm determined by dynamical sensitivities.

Optimal estimating equations and their sandwich information are classical [10]. Efficient scores remove nuisance-score directions by orthogonal projection [11, 12]; the initial-state Schur complement is the finite-dimensional instance here. Continuous-time BLUE and endpoint-weighted estimation likewise precede weak ODE analysis [13, 14]. Our contribution identifies the adjoint range of the restricted weak class and quantifies its cost and recovery in this dynamical experiment.

The continuum-data theory of WSINDy studies uncorrelated measurement errors and library-dependent consistency [15]. Fixed physical correlation time defines a different experiment: adjacent correlations approach one as the mesh is refined. Correlation induced by filtering independently sampled errors also requires distinguishing a physical filter scale from a fixed number of sample indices. Our analysis concerns a fixed covariance process observed at increasingly dense times.

The paper has three contributions: exact boundary-free and known-initial adjoint-range characterizations; a constructive efficient full-library estimator under a checkable jet condition, verified exactly for Van der Pol (Theorem 3); and quantitative correlated-noise benchmarks separating information cost, established boundary recovery and prespecified testing. Polynomial Hermite covariance supplies the feasible weights. Independent-record GMM, fission and cross-fitting are preserved in a separate research archive.

2. Related estimation methods

ODE estimation has long used trajectory smoothing and integral matching: spline least squares [16], nonparametric gradient matching [17], measurement-error regression [18], smooth-and-match estimation [19], and direct integral estimation [20]. Generalized profiling can be efficient under its approximation and independent-sampling assumptions [21, 22]; one-step likelihood refinement of an ODE pilot is also established [23]. These provide full-observation references. Our question is the information attainable by restricted weak equations under physical temporal correlation.

WENDy treats weak parameter estimation as an errors-in-variables problem, retaining coupled feature and response perturbations and iteratively updating their linearized covariance [24]. WENDy-MLE uses an approximate Gaussian weak-residual likelihood with parameter-dependent covariance and analytic derivatives [25]. Its log determinant and covariance-derivative terms distinguish its finite-noise objective from frozen-weight GLS. Polynomial Hermite correction and exact residual covariance here extend that coupled calculation to all Gaussian orders for supplied OU noise. They

do not alone correct the nonlinear coefficient estimator. Gaussian correction of noisy polynomial features has classical precedents [26, 27], and WENDy finite-noise bias and uncertainty remain active topics [28].

Test design also has substantial precedents. Tran and Bortz choose local radii through quadrature-error estimates [29], while Fourier Weak SINDy selects sinusoidal frequencies spectrally [30]. Wm-SINDy combines weak regression with dynamics and noise reconstruction [31]. Ensemble-SINDy quantifies feature inclusion by bootstrap aggregation [32]. Our author-method checks preserve their distinct structural inputs and are documented in the research archive. The new class characterization and constructive full-library result complement these methods; the supplied-candidate Wald comparison establishes no general superiority for structure selection.

2.1. Complementarity with orthogonality-condition estimation

Brunel et al. [1] combine a nonparametric proxy, orthogonality conditions and GMM in an independent-observation, increasing-point experiment, including delay equations. They recognize boundary-information loss and adapt moments to known boundary values. Our deterministic polynomial setting is narrower. It instead treats fixed-horizon physical OU error, identifies the complete adjoint influence classes, gives their sharp initial-state Schur-complement relation and an explicit OU optimal test, and constructs an attainable full-library estimator under a checkable jet condition. The small-noise and sequential dense-span limits differ from increasing-point asymptotics. The results complement each other.

3. Observation model and weak equations

Let $x_\theta : [0,T] \to \mathbb{R}^d$ be a deterministic ODE trajectory. In the polynomial setting,

$$x_i' = f_i(x) = \sum_{j=1}^{p} \xi_{ji}^* \vartheta_j(x), \qquad i = 1, \dots, d.$$

The library contains all true terms. Observations are $Y(t) = x_\theta(t) + \eta_C(t)$, with independent stationary OU coordinates [33] and common-across-parameter covariance

$$C_j(s,t) = \sigma_j^2 \exp(-|s-t|/\tau_j), \qquad \sigma_j > 0, \quad \tau_j > 0.$$

The continuous record is a statistical reference experiment; actual data are evaluations $y_n = Y(t_n)$ at mesh h. The initial state $u = x(0)$ is known in the baseline acquisition experiment; Sections 5 and 7 compare known and unknown u explicitly. The symbol Z denotes its fundamental sensitivity matrix, h the observation mesh, ϵ a scalar noise amplitude, and C the continuous covariance operator. Finite weak covariance matrices are denoted by Σ. In Proposition 3, C is the unit-amplitude covariance shape and the physical covariance is $\epsilon^2 C$.

Fix K test functions $\phi_k \in C_c^2(0,T)$. Their positions and full support widths L remain fixed as h decreases. Let $\mathcal{Q}_h$ denote composite trapezoidal quadrature with weights w_n in the formulas; matched endpoint confirmations use composite Simpson quadrature as specified in their frozen protocols. Clean weak quantities are

$$G^0_{kj} = \int_0^T \phi_k \vartheta_j(x)\, dt, \qquad b^0_{ki} = -\int_0^T \phi'_k x_i\, dt, \qquad b^0 = G^0 \xi^*.$$

Noisy matrices use corrected polynomial features $\tilde{\vartheta}_j$, defined in Section 6:

$$(G_h)_{kj} = \mathcal{Q}_h(\phi_k \tilde{\vartheta}_j(y)), \qquad (b_h)_{ki} = -\mathcal{Q}_h(\phi'_k y_i).$$

Physical OU infill sets $\rho(h) = \exp(-h/\tau)$ with fixed T, τ. An iid discrete control has independent error of fixed per-sample variance. Fixed-index AR(1) instead keeps $\rho < 1$ constant, making the physical correlation time $-h/\log\rho \to 0$. These models have different continuum limits.

4. Fixed-horizon information limits

4.1. OU translation geometry

The stationary OU process satisfies $d\eta_{C,j} = -\eta_{C,j} dt/\tau_j + \sqrt{2\sigma_j^2/\tau_j}\, dW_j$, with an independent initial value $\eta_{C,j}(0) \sim N(0, \sigma_j^2)$. Its Cameron–Martin space is $H^1([0,T]; \mathbb{R}^d)$ as a set, with norm

$$\|g\|^2_{\mathcal{H}} = \sum_j \left\{ \frac{g_j(0)^2}{\sigma_j^2} + \frac{\tau_j}{2\sigma_j^2} \int_0^T (g'_j + g_j/\tau_j)^2\, dt \right\}.$$

Transforming a path into its initial value and innovation path $z(t) - z(0) + \tau_j^{-1} \int_0^t z(s) ds$ separates the normal initial observation from scaled Brownian motion. Translation by g gives an initial shift and Brownian drift $g' + g/\tau_j$. The classical translation theorem [34] gives the norm. Equivalently,

$$\|g\|^2_{\mathcal{H}} = \sum_j \left\{ \frac{\tau_j}{2\sigma_j^2} \int_0^T (g'_j)^2 dt + \frac{1}{2\sigma_j^2 \tau_j} \int_0^T g_j^2 dt + \frac{g_j(0)^2 + g_j(T)^2}{2\sigma_j^2} \right\}.$$

The endpoint terms encode stationary initial noise. Zero initial error would define a different experiment.

Lemma 1 (fixed-horizon two-point bound; proof in S1.1). Suppose the mean trajectories belong to $\mathcal{H}$ and share the stated positive OU covariance. For $\delta = \|\theta - \eta\| > 0$ and $d = \|x_\theta - x_\eta\|_{\mathcal{H}}$, every finite-grid estimator satisfies

$$\max_{v \in \{\theta, \eta\}} P_v\{\|\hat{\theta}_h - v\| \geq \delta/2\} \geq \Phi(-d/2) > 0.$$

Its maximal two-point squared risk is at least $(\delta^2/4)\Phi(-d/2)$. Thus dense-grid consistency throughout a nontrivial fixed-horizon family is excluded by Gaussian path equivalence [9, 35].

4.2. Continuous-record information

Proposition 1 (continuous-record information). If $\theta \mapsto x_\theta$ is differentiable in $\mathcal{H}$ and $s_r = \partial_{\theta_r} x_\theta$, continuous-record Fisher information is

$$\mathcal{I}_{rs}(\theta) = \langle s_r, s_s \rangle_{\mathcal{H}}.$$

The Gaussian mean-score identity proves the assertion: the directional score has variance equal to the squared Cameron-Martin norm of its sensitivity. Its inverse is the Cramér–Rao covariance bound for regular unbiased estimators, when nonsingular. Biased estimators require the corresponding bias derivative; finite-horizon maximum likelihood need not attain the unbiased bound.

Exact discrete OU innovations converge to this finite path information under grid refinement. S6.1 gives the scalar calculation and shows why unguarded weak ratios can have infinite error moments despite a finite Gaussian influence variance. Information bounds concern local regular influences; finite-noise estimator risk requires separate control.

5. Boundary-free weak class and efficiency

5.1. Exact boundary-free influence class

Along $x_{\theta,u}$, put $J(t) = D_x f(x_{\theta,u}(t), \theta)$ and $Z(t) = \partial_u x_{\theta,u}(t)$. Assume $J \in C^1([0,T]; \mathbb{R}^{d\times d})$, and let the invertible fundamental matrix satisfy $Z' = JZ$, $Z(0) = I_d$. Put $\mathcal{D}v = v' - Jv$ and $\mathcal{D}^*\phi = -\phi' - J^T\phi$. Sensor coordinates have independent stationary OU covariance with fixed positive scales. Its Cameron–Martin space $\mathcal{H}$ is H^1 with the OU inner product. Write $\mathcal{E} = \mathcal{H}'$ for its continuous dual, equipped with the covariance norm

$$\|\lambda\|_C^2 = \langle \lambda, C\lambda \rangle, \qquad \langle C^{-1}g, v \rangle = \langle g, v \rangle_{\mathcal{H}}.$$

This definition includes ordinary integrated influences and endpoint evaluation measures. General dual influences are defined by $L^2(P)$ limits of integrated Gaussian observations. Their variance norm is $\|\lambda\|_C$; the noise path itself need not belong to $\mathcal{H}$.

Proposition 2 (exact range and closure). In L^2,

$$\mathcal{D}^* H_0^1 = \left\{ \ell \in L^2 : \int_0^T Z(t)^T \ell(t) dt = 0 \right\}.$$

The range of compactly supported smooth tests is dense in this space in L^2. Its closure in $\mathcal{E}$ is exactly

$$\mathcal{M} = \overline{\mathcal{D}^* C_c^\infty}^{\|\cdot\|_C} = \{\lambda \in \mathcal{E} : \langle \lambda, Z_j \rangle = 0, j = 1, \dots, d\}.$$

Consequently $\mathcal{M}$ has codimension d.

Proof. Since $Z^T \mathcal{D}^* \phi = -(Z^T \phi)'$, zero endpoint values imply the moment constraint. Conversely, for an L^2 influence with zero moment,

$$\phi(t) = -Z(t)^{-T} \int_0^t Z(s)^T \ell(s) ds$$

belongs to H_0^1 and satisfies $\mathcal{D}^*\phi = \ell$. Approximation in H_0^1 by C_c^∞ and continuity of $\mathcal{D}^* : H_0^1 \to L^2$ give L^2 density. For the stronger closure statement, a function $v \in \mathcal{H}$ annihilates every $\mathcal{D}^*\phi$ if and only if $\mathcal{D}v = 0$ distributionally, hence $v = Zc$. The Hahn–Banach annihilator characterization of a closed subspace then gives the stated closure in $\mathcal{H}'$. The columns of Z are linearly independent elements of $\mathcal{H}$, so their dual constraints have rank d. □

The exact range uses zero-endpoint Sobolev tests; literal compact tests require closure. The covariance norm is equivalent to the dual Sobolev norm for this positive OU model.

Lemma 2 (a concrete dense test family). Let V_m be vector-valued continuous piecewise-linear splines on the nested uniform dyadic partitions of $[0, T]$, with zero values at both endpoints. Then

$$\overline{\bigcup_m \mathcal{D}^* V_m}^{\|\cdot\|_C} = \mathcal{M}.$$

Proof. Given $v \in H_0^1$, choose $\psi \in C_c^\infty$ close to v in H^1. Its nodal interpolant $I_m\psi \in V_m$ satisfies $\|I_m\psi - \psi\|_{H^1} \le c\Delta_m \|\psi''\|_{L^2}$, where Δ_m is the mesh width. The union of V_m is therefore dense in H_0^1. Continuity of $\mathcal{D}^* : H_0^1 \to L^2$ and of the embedding $L^2 \hookrightarrow \mathcal{H}'$ transfers this approximation to the covariance norm. Proposition 2 identifies the closure of $\mathcal{D}^* H_0^1$ with $\mathcal{M}$. □

These splines are admissible zero-endpoint Sobolev tests; they need not vanish in an endpoint neighborhood. If literal smooth compact tests are required, approximate each spline in H_0^1 by such tests. Thus Lemma 2 provides an explicit family for the density condition in Proposition 3. The experiments use finite multiscale approximations.

5.2. Sharp information and optimal influence

Let $S = [S_1, \dots, S_p] \in \mathcal{H}^p$ be coefficient sensitivities, and form the joint information blocks for S and Z. Assume $\mathcal{I}_{uu}$ and the Schur complement are positive definite. Set

$$R = S - Z\mathcal{I}_{uu}^{-1}\mathcal{I}_{u\theta}, \qquad \mathcal{I}_e = \mathcal{I}_{\theta\theta} - \mathcal{I}_{\theta u}\mathcal{I}_{uu}^{-1}\mathcal{I}_{u\theta}.$$

The nuisance-score projection and efficient influence are classical [11, 12], as is continuous-record BLUE [13, 14]. Here the adjoint-range characterization identifies exactly which dynamical weak influences obey that projection.

Theorem 1 (sharp boundary-free influence covariance). Let $\Lambda = [\lambda_1, \dots, \lambda_p] \in \mathcal{M}^p$ satisfy $\langle \lambda_r, S_s \rangle = \delta_{rs}$. Its Gaussian influence covariance obeys

$$V(\Lambda) \succeq \mathcal{I}_e^{-1}.$$

The bound is attained in $\mathcal{M}$ by the matrix-valued dual influence

$$\Lambda_* = C^{-1} R\, \mathcal{I}_e^{-1}.$$

It is the infimum for normalized compact-test influences. It is a minimum among literal compact tests only when Λ_* belongs to their adjoint range.

Proof. The $\mathcal{H}$ projection makes $\langle C^{-1}R, Z\rangle = 0$ and $\langle C^{-1}R, S\rangle = \mathcal{I}_e$. Thus Λ_* meets both constraints and has covariance $\mathcal{I}_e^{-1}$. The difference $\Delta = \Lambda - \Lambda_*$ pairs to zero with both S and Z, hence with R. Covariance cross terms vanish and

$$V(\Lambda) = \mathcal{I}_e^{-1} + V(\Delta).$$

For attainability as an infimum, approximate each column of Λ_* by compact-test influences in covariance norm. Their normalization matrices converge to I_p by continuous pairing with S; right-multiplication by their inverse transpose enforces exact normalization while preserving the range and covariance convergence. □

For an unknown initial state, $\mathcal{I}_e$ is the full likelihood nuisance-adjusted information, so this weak restriction has no further first-order cost. When the initial state is known, the information loss is exactly $\mathcal{I}_{\theta u}\mathcal{I}_{uu}^{-1}\mathcal{I}_{u\theta}$; covariance inflation is $\mathcal{I}_e^{-1} - \mathcal{I}_{\theta\theta}^{-1}$. There is no direction-independent scalar loss percentage in general. When u is known, $Y(0) - u$ also reveals the initial stationary measurement error, which remains correlated with later observations. The unrestricted score can exploit that observation while the weak influence must retain its initial-sensitivity annihilation constraint.

Known-boundary adaptations have a direct precedent: Brunel et al. [1] allow a nonzero initial test trace when the initial state is known and retain a zero terminal trace. Their construction motivates endpoint-enabled moments. The information comparison above concerns the endpoint-vanishing class; restoring endpoint information changes that class. Recent test construction work also treats numerical integration error as a central design criterion [29]. An implementation of endpoint-enabled tests therefore needs a separate clean quadrature check before its empirical accuracy can be compared with the known-initial likelihood reference.

5.3. Explicit optimal OU test

For a scalar coordinate with $g \in H^2$, integration by parts in the stationary OU norm gives the dual measure

$$C^{-1}g = \frac{g - \tau^2 g''}{2\sigma^2\tau}\,dt + \frac{g(0) - \tau g'(0)}{2\sigma^2}\delta_0 + \frac{g(T) + \tau g'(T)}{2\sigma^2}\delta_T.$$

For independent coordinates the formula applies coordinatewise, using their respective scales. If only $g \in H^1$, the Riesz representative remains well-defined in $\mathcal{H}'$ but need not be a density plus finite endpoint measures.

For a smooth optimal representative write $\Lambda_* = m(t)dt + m_0\delta_0 + m_T\delta_T$. On the open interval an adjoint test matrix is

$$\Phi_*(t) = -Z(t)^{-T}\left[Z(0)^T m_0 + \int_0^t Z(s)^T m(s)ds\right].$$

Extend it by zero outside the observation interval. Its endpoint jumps supply $m_0 = -\Phi_*(0+)$ and $m_T = \Phi_*(T-)$, and distributionally $\mathcal{D}^*\Phi_* = \Lambda_*$. These formulas include the initial atom; omitting

it in the cumulative integral is incorrect. The annihilation condition supplies the terminal jump. Cutoff and smoothing approximate this generalized test in covariance norm. These measures describe the boundary-free statistical reference. Section 7.2 separately evaluates an endpoint-enabled extension for a known initial state.

The scalar exponential specialization and fitted-pilot formula are in S6.5. Figure 1 shows how the generalized optimum and its compact cutoff differ near the endpoints.

5.4. Correct-covariance GLS and dense spans

For tests ϕ_k, put $\lambda_k = \mathcal{D}^*\phi_k$, $D_{kr} = \langle \lambda_k, S_r \rangle$ and $\Sigma_{k\ell} = \langle \lambda_k, C\lambda_\ell \rangle$. Remove exact linear dependencies. If D has full column rank, covariance-weighted weak regression has local influence and covariance

$$P = (D^T\Sigma^{-1}D)^{-1}D^T\Sigma^{-1}, \qquad V_K = (D^T\Sigma^{-1}D)^{-1}.$$

WENDy's covariance iteration motivates this construction [24], and physical-OU covariance changes its temporal weights. Under polynomial features Proposition 4 supplies the complete coupled residual weights. Its order-one term determines the small-noise efficient influence; higher orders correct residual uncertainty at finite amplitude without by themselves removing coefficient-estimator bias.

This is the Godambe-information construction for optimally weighted estimating equations [10]: the local information is $D^\top\Sigma^{-1}D$. WENDy-MLE has the same leading small-noise map under regularity and correctly specified covariance; finite-noise covariance-derivative terms distinguish its objective (S6.3).

Proposition 3 (guarded fixed-family weak GLS). Fix a finite deterministic boundary-free test family, a correctly specified polynomial vector field linear in its coefficients, known initial state and supplied independent-coordinate OU covariance shape with a known common correlation time. Suppose the true coefficient is interior to a compact final box; clean and nearby pilot trajectories exist in a common bounded region; the clean pilot and final designs have full column rank; and the first-order residual covariance Σ_1 is positive definite. Quadrature weights have bounded total mass, converge to their integrals, and clean quadrature and pilot-model numerical defects are $o(\epsilon)$ with the local derivative control in (G4) of S7.2.

Use a declared pilot family containing the true vector field and a pilot box containing its true coefficients. Apply corrected least squares, clip the pilot, evaluate the full Hermite covariance on its integrated trajectory, and apply the vanishing-ridge, guarded and box-projected GLS routine described below. Then its pilot satisfies $\widetilde{\theta} - \theta = O_P(\epsilon)$, its normalized weights converge, and

$$\epsilon^{-1}(\hat{\theta}_{K,\epsilon} - \theta) \Rightarrow N(0, V_K), \qquad \epsilon^{-2}\,\mathrm{Cov}(\hat{\theta}_{K,\epsilon}) \to V_K.$$

The squared scaled errors are uniformly integrable. These conclusions are derived from the stated structural and numerical conditions; an asymptotic expansion, weight consistency and integrability are not assumed. S7.2 proves the proposition. For nested dense families satisfying these conditions separately, the subsequent family limit gives the bound of Theorem 1. Theorem 3 supplies a full-library smoothed-data construction whose clean ranks follow from a jet condition instead of being assumed.

The archived solvers in Tables 1–6 use fixed relative ridge and an unbounded final estimate. Their local matrix geometry and empirical recovery are separate from the risk guarantee above. For fixed design D and residual shape Σ_1, their regularized linear map has

$$W_\rho = [\Sigma_1 + \rho \operatorname{tr}(\Sigma_1) I/q]^{-1}, \quad P_\rho = (D^\top W_\rho D)^{-1} D^\top W_\rho, \quad V_\rho = P_\rho \Sigma_1 P_\rho^\top.$$

Here q is the number of moments, and V_0 is ideal GLS covariance. The exact identity

$$V_\rho - V_0 = (P_\rho - P_0)\Sigma_1 (P_\rho - P_0)^\top \succeq 0$$

quantifies ridge cost without any estimator-limit assumption. On eight actual 480-interior-window archived spans and their endpoint/control extensions, including Lorenz, the largest computed relative RMSE increment is 6.7×10^{-15}; S7.3 gives the matrices and checks. Thus the stated ridge has negligible local matrix cost in these examples. This check cannot establish moments of an unbounded nonlinear final estimate. Table S8 separates those archived benchmarks, the constructive estimator and supplied-candidate testing.

5.5. Extended class and recovery with a known initial state

Throughout this subsection, u is known exactly, covariance is supplied and independent of θ, J is continuous, and S and Z belong to the OU Cameron–Martin space $\mathcal{H}$. Assume $\mathcal{I}_{\theta\theta}$ is positive definite. The coordinate OU scales and correlation times are positive. Consequently $\mathcal{E}$ is the covariance-dual space of vector-valued $H^1([0,T])$, with continuous endpoint evaluation. The initial sensitivity is normalized by $Z(0) = I_d$ and the parameter sensitivity satisfies $S(0) = 0$.

For $\phi \in H^1([0,T]; \mathbb{R}^d)$, use the known-initial residual

$$m_\phi(\theta) = -\int_0^T [\phi'^T Y + \phi^T f_\theta^H(Y)]dt + \phi(T)^T Y(T) - \phi(0)^T u.$$

Its first-order influence is $L\phi = (-\phi' - J^T\phi)dt + \phi(T)\delta_T$. The initial boundary term contains the exact state rather than its noisy observation. This adaptation of the weak identity is established [1]. The result below specifies its covariance-dual geometry and its relation to the correlated-noise control.

Theorem 2 (known-initial extended influence class). Under these assumptions,

$$\overline{L(H^1)}^{\mathcal{E}} = \overline{L\{\phi \in H^1 : \phi(T) = 0\}}^{\mathcal{E}} = \mathcal{E}.$$

In contrast, allowing only a terminal trace preserves the original class:

$$\overline{L\{\phi \in H^1 : \phi(0) = 0\}}^{\mathcal{E}} = \mathcal{M}.$$

Alternatively, adding the initial noise controls to the closed boundary-free class gives the exact equality

$$\mathcal{M} + \{a\delta_0 : a \in \mathbb{R}^d\} = \mathcal{E}.$$

The sharp normalized influence covariance in either full extended closure is $\mathcal{I}_{\theta\theta}^{-1}$, with representative $\Lambda_{\text{full}} = C^{-1}S\mathcal{I}_{\theta\theta}^{-1}$. For ordinary tests it is an infimum; an exact smooth-test minimum additionally requires the representative to belong to the specified range. Explicit initial observation controls can realize the initial atom.

Proof. For $g \in \mathcal{H}$, the continuous dual pairing obeys

$$\langle g, L\phi\rangle = \int_0^T \phi^T(g' - Jg)dt + \phi(0)^T g(0).$$

If this vanishes for all permitted tests, compact interior tests first imply $g' = Jg$ in the weak sense. Such solutions are $g = Zc$. For all H^1 tests, or for tests with $\phi(T) = 0$, the initial trace is arbitrary, so $g(0) = c = 0$. Their annihilator is trivial and their closure is $\mathcal{E}$. For tests with $\phi(0) = 0$, every Zc annihilates the range and there are no further annihilators. Its closure is therefore $\mathcal{M}$, as characterized in Theorem 1.

For any $\lambda \in \mathcal{E}$, define $a_j = \langle Z_{\cdot j}, \lambda\rangle$. Continuity of this pairing follows from $Z \in \mathcal{H}$. Since $Z(0) = I_d$, $\lambda - a\delta_0$ annihilates every initial-sensitivity column and belongs to $\mathcal{M}$. This proves the exact sum identity. For covariance optimality, $\langle \Lambda_{\text{full}}, S\rangle = I_p$ and $\text{Cov}(\Lambda_{\text{full}}) = \mathcal{I}_{\theta\theta}^{-1}$. Any other normalized influence has a covariance-orthogonal remainder, giving a positive-semidefinite covariance increment. Density, continuous pairing with S and normalization by the convergent sensitivity matrix give attainability as an infimum, by the same approximation argument as Theorem 1. □

Correct-covariance GLS on dense universal extended spans approaches this bound under the sequential small-noise conditions of Proposition 3, with the full rather than nuisance-adjusted information. The theorem concerns the closed local influence class; performance of finite families and finite-noise covariance pilots requires numerical assessment. In particular, finite-span gains of different endpoint additions can overlap.

For compact tests, let Ω be the exact Hermite residual covariance, $\Sigma_0 = \text{diag}(\sigma_j^2)$ and $B = \text{Cov}(m_\phi, Y(0) - u)$. The optimal linear control replaces the residual covariance by

$$\Omega - B\Sigma_0^{-1}B^T.$$

Only the first Hermite order contributes to B. In coordinate j, its cross-covariance is the integral of the linear residual weight against $\sigma_j^2 e^{-t/\tau_j}$. In a sampled iid model the compact-test weights and derivatives vanish at the initial node; hence this cross-covariance is exactly zero. This distinction explains the correlated-noise control mechanism. It does not establish an additional finite-family gain when initial-trace tests already approximate the full efficient directions.

5.6. Constructive full-library attainability

This is a constructive attainability result: the algorithm witnesses the sharp small-noise bound. Its moderate-noise precision is assessed separately in Section 7.6. The following procedure estimates every supplied library coefficient. Let B be the compact coefficient box and use uniform grids with an even number of intervals. Smooth the linear interpolant of Y with

$$K_b(t) = b^{-1}(315/256)[1-(t/b)^2]_+^4, \quad b = \epsilon^{1/10}/4,$$
$$w(t) = [1-((t-T/2)/(T/2-1/2))^2]_+^4.$$

Reflect the interpolant at the endpoints. The proxy z supplies the m feature tests $wg_j(z)$ and their analytic derivatives. Their Hermite-corrected weak equations give a full-library least-squares pilot, clipped to B. A fixed deterministic-window pilot on a declared correct structural family gives the alternative construction in Proposition 3.

Append the cubic splines $B_3(t/a-j)$, $a = T/2^k$, $j = 2, \dots, 2^k-2$, where B_3 is the normalized cardinal cubic spline supported on $[-2,2]$. Whiten the combined H^1 Gram matrix, retaining eigenvalues above $10^{-10}\epsilon^{1/10}$ times its largest eigenvalue. The exact Hermite covariance (Proposition 4), evaluated at the proxy and pilot, gives $\widehat{\Sigma}_1 = \widehat{\Omega}/\epsilon^2$. Let r_k be the number of retained scalar tests and $q_k = dr_k = \dim(\widehat{\Sigma}_1)$ the number of stacked residual moments. Add ridge $\epsilon^2 \operatorname{tr}(\widehat{\Sigma}_1) I_{q_k}/q_k$, then solve corrected GLS and project the result to B. Relative minimum-eigenvalue guards use $10^{-8}\epsilon^2$; singular, nonfinite or failed calculations return the box midpoint. S8 specifies the column scaling and covariance guard. No generating support or pilot ODE solution is supplied to this full-library procedure.

The technical restriction $T > 1$ makes the fixed half-unit interior margins of w nonempty. Proportional margins and a rescaled kernel give an equivalent construction on any positive horizon. Write $\mathcal{J}_{e,C}$ for the initial-state-adjusted information of Theorem 1 computed with the unit-amplitude covariance C of η.

Theorem 3 (constructive attainability under a jet condition). Consider an analytic trajectory x_0 of a polynomial ODE on $[0,T]$, $T > 1$, with vector field linear in a polynomial library $g_1, \dots, g_m$ in each of d equations. Let the dm true coefficients lie strictly inside a supplied compact box. Observe $Y = x_0 + \epsilon\eta$ with known positive independent-coordinate stationary OU covariance and common correlation time $\tau > 0$. Assume that, at some t_* in the observation interval,

$$\det[\partial_t^r g_j(x_0(t_*))]_{r=0,\dots,m-1;\ j=1,\dots,m} \neq 0.$$

For the full-library procedure just defined, let $h^2 = o(\epsilon)$. For each fixed spline level k, the pilot satisfies $\tilde{\theta} - \theta_0 = O_P(\epsilon)$, the normalized covariance weights are consistent, and

$$\epsilon^{-1}(\hat{\theta}_{k,\epsilon} - \theta_0) \Rightarrow N(0, V_k), \qquad \epsilon^{-2} E[(\hat{\theta}_{k,\epsilon} - \theta_0)(\hat{\theta}_{k,\epsilon} - \theta_0)^\top] \to V_k.$$

The squared scaled errors are uniformly integrable, and $V_k \to \mathcal{J}_{e,C}^{-1}$ as $k \to \infty$ after the fixed-level small-noise limit. The coefficient pilot uses all library terms; no generating support is supplied. S8.1 proves the general result, including rank, weight consistency and moment control; S8.2 verifies it for Van der Pol with the full 18-coefficient cubic library and $u = (2,0)$, where the jet determinant is -65265167726360395776. This theorem concerns the specified guarded estimator and sequential limits, not finite-noise Wald calibration or the archived fixed-ridge fits.

6. Exact polynomial weak-residual covariance

6.1. Finite Hermite expansion

Assume a common τ for independent coordinates, and set $v_j = \sigma_j^2$. Define $H_0(z;v) = 1$, $H_1(z;v) = z$, $H_{q+1}(z;v) = zH_q(z;v) - qvH_{q-1}(z;v)$ and corrected monomials $\tilde{\vartheta}_\beta(y) = \prod_j H_{\beta_j}(y_j; v_j)$. Their expectation given x is x^β. This Gaussian debiasing construction [27] centers the polynomial residual at the true coefficients.

For continuum residual $r_{ki} = b_{ki} - (G\xi^*)_{ki}$, polynomial expansion gives

$$r_{ki} = \sum_{1 \le |\alpha| \le p_*} \int_0^T c_{ki,\alpha}(t) H_\alpha(\eta_C(t); v) dt,$$

where $p_* \ge 1$ is the relevant maximum degree, $H_\alpha = \prod_j H_{\alpha_j}$ and

$$c_{ki,\alpha} = -\mathbf{1}_{\{\alpha = e_i\}} \phi_k' - \phi_k \frac{\partial^\alpha f_i(x)}{\alpha!}.$$

The response and feature perturbations share this coefficient and retain their cross-covariance.

Proposition 4 (exact covariance). Under these assumptions,

$$\mathrm{Cov}(r_{ki}, r_{\ell m}) = \sum_{1 \le |\alpha| \le p_*} \alpha! \prod_j v_j^{\alpha_j} \int_0^T \int_0^T c_{ki,\alpha}(s) c_{\ell m,\alpha}(t) e^{-|\alpha||s-t|/\tau} \, ds \, dt.$$

Proof. The generating function $\exp(az - va^2/2)$ yields $E[H_q(U;v)H_r(V;v)] = \mathbf{1}_{q=r} q! v^q \rho^q$ for jointly Gaussian variables of variance v and correlation ρ. This is the Hermite form of classical Gaussian moment identities [36]. Coordinate independence gives orthogonality between different multi-indices and factor $\alpha! \prod_j v_j^{\alpha_j}$. Substitute $\rho = e^{-|s-t|/\tau}$ and integrate the finite expansion. □

The order-one covariance supplies the GLS weights required by Proposition 3. Higher orders refine residual weights at finite amplitude; coefficient bias also depends on random-design terms. Order q has correlation time τ/q. Higher orders need not be negligible at moderate amplitude. Different windows and equations remain correlated despite independent sensor coordinates. The covariance is exact for a supplied clean trajectory and true polynomial; plugging in estimates introduces another approximation.

6.2. Implementation and covariance misspecification

The finite Hermite sum can be evaluated on a grid without storing a dense temporal matrix. S6.4 gives the recursion and the extension to incorrectly corrected raw features. Under temporal weights W, leading covariance is $P_W \Sigma_{\mathrm{true},1} P_W^\top$; changing correction variances can also change the moment mean. The present comparisons supply the sensor covariance. Sensitivity to supplied and fitted correlation times is reported in S5.4.

7. Numerical benchmarks: cost, recovery and attainability

The benchmarks use fully observed polynomial systems, supplied structure and sensor covariance, and archived paired observations. Feasible covariance pilots receive no generating coefficients. S2–S5 preserve protocols, seed ranges, fingerprints and additional checks. Local references and finite-noise performance are reported separately.

7.1. Information cost and finite-span efficiency

For Van der Pol, we fit four independent coefficients in $x' = av$, $v' = bx + cv + dx^2v$, with truth $(1, -1, 1, -1)$, initial state $(2, 0)$, $T = 20$, $h = 0.005$ and OU range 0.3. Relative errors are divided by the true coefficient norm, two. The known-initial local RMSE reference at 5% noise is 0.013093, versus the boundary-free reference 0.016904: a 29% cost. Both scale linearly with noise amplitude.

Table 1. Oracle Gaussian influence RMSE under actual OU noise at 5%. Both methods use the same test span and clean design; they differ only in the covariance used for GLS weights. Actual variance uses the OU covariance for both.

Prescribed windows	Retained tests	OU-weight RMSE	iid-weight RMSE	OU RMSE / class reference
30	30	0.019605	0.020296	1.1597
120	120	0.017043	0.017839	1.0082
480	479	0.016938	0.017734	1.0020

Rich spans produce most of the improvement: 480-window OU GLS is within 0.2% of the boundary-free local reference, while iid weights retain a roughly 5% gap. Exact residual-covariance calculations verify the stated local geometry. The original 128-path finite-span check is in S5.2; the larger paired comparison follows.

Table 2. Van der Pol on the same 2000 paths at 0.5% noise. Error norms are divided by the true four-coefficient norm, two. The reference concerns the leading local Gaussian influence. All paths are retained.

Estimator	RMSE (95% interval)	Bias	Dispersion	Local reference
One step, known initial	0.001301 [0.001269, 0.001332]	0.000085	0.001298	0.001309
One step, unknown initial	0.001838 [0.001790, 0.001887]	0.000469	0.001777	0.001690
Three steps, unknown initial	0.001669 [0.001630, 0.001708]	0.000068	0.001668	0.001690
Weak GLS, OU	0.001685 [0.001645, 0.001723]	0.000083	0.001683	0.001690
Weak GLS, iid	0.001759 [0.001718, 0.001801]	0.000103	0.001756	0.001690

All 2000 prespecified paths at 0.5% noise are retained, including the original 128-path prefix once. Feasible weak GLS uses a bounded 30-window covariance pilot and 480 prescribed windows. Likelihood refinements use exact OU innovations and analytical ODE sensitivities. Paired iid-minus-OU RMSE is 0.00007426, with bootstrap interval [0.00005576, 0.00009327], a 4.22% benefit from temporal weighting. The initial unknown-state scoring step has an appreciable nonlinear initialization bias; three steps approach the nuisance-adjusted reference. The known-initial estimate has lower error because it observes the information discarded by compact tests.




Figure 1: Scalar optimal generalized test for $a = -1$, $x_0 = 2$, $T = 4$ and $\tau = 0.4$, alongside its compact cutoff with width 0.1 and an ordinary width-three bell. All tests use the same clean-moment normalization. Endpoint panels show nonzero one-sided traces and jumps of the zero extension; their adjoint derivatives supply endpoint atoms. The curves describe influences annihilating the initial sensitivity, under the unknown-initial information reference.




Figure 2: Van der Pol on 2000 shared paths at 0.5% noise: one scoring step with known initial state, one or three joint steps with unknown initial state, and weak GLS with OU or iid weights. Whiskers give 95% bootstrap intervals for RMSE over independent paths, not parameter confidence intervals or error standard deviations. The known-initial information reference is lower than the unknown-initial reference. The finite-step unknown-initial estimates are identified separately; reference agreement concerns local small-noise behavior.

7.2. Recovery with universal endpoint tests

For known u, allow nonzero test traces and use the corrected residual

$$m_\phi(\theta) = -\int_0^T [\phi'(t)^T Y(t) + \phi(t)^T f_\theta^H(Y(t))]dt + \phi(T)^T Y(T) - \phi(0)^T u.$$

The universal family adds twelve time-only endpoint half-windows and $Y(0) - u$ controls to 480 interior windows. Covariance weights still require an integrated 30-window pilot. Theorem 2 identifies the full local influence class; the known-initial trace adaptation itself is established by Brunel et al. [1].

Table 3. Known-initial endpoint confirmation on 1968 paired Van der Pol paths at 0.5% noise. The likelihood reference is one Fisher-scoring step. All errors use the four-coefficient norm, two.

Method	RMSE	Bias norm	Dispersion
Boundary-free, 480	0.00168754	0.00008679	0.00168531
Universal endpoints	0.00130936	0.00007049	0.00130746
Model-adjoint endpoints	0.00129741	0.00006771	0.00129564
Known-initial scoring	0.00129800	0.00008753	0.00129505

These 1968 records were unused for endpoint fitting, while their compact and scoring references were archived. After inspecting the 32-path pilot, we replaced the absolute RMSE gate of 0.00140 with paired relative criteria because known-initial likelihood itself failed that gate; the amendment was frozen before any new endpoint outcomes on the 1968 confirmation records were inspected. The frozen confirmation criterion requires at least 50% gap recovery. Universal endpoints recover 97.1%, with paired interval 95.5–98.6%, and lie 0.88% above scoring. Model-adjoint tests realize score directions and provide an implementation check. S2 and S5.5 record the criterion amendment and quadrature checks.

7.3. Attribution to initial traces

The 500-path ablation holds Simpson quadrature and the covariance pilot fixed. Component gains are measured against the matched compact branch and known-initial scoring.

Table 4. Endpoint-component ablation on 500 paired paths at 0.5% noise. Recovery is measured against the matched Simpson boundary-free branch and known-initial scoring. Intervals resample common record indices.

Added component	RMSE	Gap closed	Paired 95% interval
Initial half-windows	0.00132445	98.9%	[95.1%; 102.7%]
Terminal half-windows	0.00164493	-0.3%	[-1.8%; 1.2%]
Initial controls only	0.00159795	14.3%	[7.0%; 21.4%]
All components	0.00132453	98.9%	[95.1%; 102.6%]
Matched boundary-free	0.00164395	0%	—
Known-initial scoring	0.00132100	100%	—

Initial traces alone recover essentially the entire gap, quantifying the Brunel mechanism under OU noise. Initial controls give a smaller standalone gain and terminal traces give none resolved. Combining all components changes RMSE relative to initial traces by 7.46×10^{-8}, with paired interval $[-7.18 \times 10^{-7}, 8.55 \times 10^{-7}]$. Their contributions overlap and cannot be added. The control's standalone improvement is specific to correlation between initial and interior error, rather than evidence for an additional gain after trace adaptation.




Figure 3: Known-initial information recovery on 1968 paired Van der Pol records at 0.5% OU noise. Whiskers are Monte Carlo 95% RMSE intervals from shared record bootstrap resamples. Universal time-only endpoint windows recover 97.1% of the original weak-to-scoring gap; model-adjoint tests serve as a score-realization check. The reference is a single known-initial Fisher-scoring step.




Figure 4: Paired endpoint-component attribution on 500 Van der Pol records. Whiskers show 95% bootstrap intervals for the fraction of the matched weak-to-scoring RMSE gap recovered. The dashed line marks the prespecified 50% material-contribution rule. Initial traces recover the gap; initial-noise controls give a smaller standalone gain, and terminal traces give no resolved improvement. Contributions overlap.

7.4. Moderate-noise limits and Lorenz confirmation

The 5% Van der Pol check reuses all 128 matched observations, with known initial state and the same feasible pilot. The likelihood reference optimizes exact OU innovations.

Table 5. Moderate-noise endpoint confirmation on 128 paired Van der Pol paths at 5%. The known-initial reference uses full nonlinear likelihood fitting.

Method	RMSE	Bias norm	Dispersion
Matched boundary-free	0.0255965	0.0081646	0.0242594
Universal endpoints	0.0218790	0.0056954	0.0211247
Known-initial MLE	0.0142088	0.0007255	0.0141902
Unknown-initial MLE	0.0171952	0.0006504	0.0171829

Universal endpoints recover 32.6% of the gap (18.3–45.8%), failing the frozen 50% criterion. Clean-design first-order and full-Hermite references are 0.013094 and 0.013636, below empirical dispersion 0.021125. Restored local geometry therefore leaves finite-noise design, pilot and fitting effects. The matched author-WENDy comparison is in S5.3.

For Lorenz, we fit $(s, r, b) = (10, 28, 8/3)$ in its supplied physical structure, with $u = (1, 1, 1)$, $T = 4$, $h = 0.001$, OU range 0.1 and 0.5% coordinate noise. The unknown-initial local relative RMSE reference is 0.000411, versus 0.000209 when the initial state is known: almost a factor two.

Table 6. Lorenz endpoint confirmation on 256 paired paths. Errors are relative to the norm of $(10, 28, 8/3)$. The common horizon is four and the sampling interval is 0.001.

Method	RMSE	Bias norm	Dispersion
Matched boundary-free	0.00040994	0.00002170	0.00040936
Universal endpoints	0.00021686	0.00002153	0.00021579
Known-initial MLE	0.00020784	0.00002355	0.00020650

All 256 paths and all converged fits are retained. Universal endpoint windows recover 95.5% of the compact-to-likelihood gap (92.8–98.0%), passing the same 50% rule. The full-Hermite clean-design reference is 0.00020926, close to the observed endpoint dispersion. These two supplied structures support small-noise recovery; moderate-noise universality remains unestablished.

7.5. Direction-specific information cost

For the prespecified negative cubic perturbation δx^3 in the second Van der Pol equation, profiling all seventeen other library coefficients gives local full-information and boundary-free two-point reference magnitudes 0.01928 and 0.01941 at 0.5% noise. The initial-state information cost for this direction is only 0.68%, despite its larger aggregate coefficient-estimation cost. Supplied-candidate Wald testing and library-wide sparse selection have different inputs and targets; their detailed comparison and calibration studies are supplementary (S3, S4 and S9).

7.6. Matching the constructive estimator to its finite-span reference

The interior proxy smoothing bias is $O(b^2) = O(\epsilon^{1/5})$ (Lemma S8.A); instrument convergence can therefore be slow even though the clean weak identity gives $\tilde{\theta} - \theta_0 = O_P(\epsilon)$. This instrument rate is not a claimed rate for the empirical RMSE ratio. Conditioning amplifies small perturbations: the clean raw Gram has minimum eigenvalue 6.07×10^{-10} and condition number at least 7.1×10^9, from a cubic-test diagonal Rayleigh bound. The column-scaled GLS normal matrix has condition

number 1.40×10^6. Positivity at fixed level suffices for Theorem 3, but its constants depend on these gaps; no uniform growing-span rate is established.

The full-library estimator of Theorem 3 uses nine smoothed-data feature instruments and 125 time-only cubic splines. The deterministic limiting covariance for this 134-test span has $\sqrt{\operatorname{tr} V_k} = 9.8144$. The calculation retains all rows and uses neither ridge nor rank truncation. Refining the finest grid from $h = 0.00015625$ to 0.000078125 changes this reference by 4.7×10^{-8} relatively. It concerns all eighteen coefficients, unlike the earlier supplied four-coefficient Van der Pol benchmarks.

Table 7. Constructive full-library estimator on 100 independent paths at each amplitude. Intervals resample archived coefficient errors 10,000 times and are pointwise; no new observations are generated for this comparison.

Noise	RMSE/ϵ	RMSE/$(\epsilon\sqrt{\operatorname{tr} V_k})$	95% interval for ratio
5%	11.6265	1.1846	1.1140–1.2563
0.5%	10.3505	1.0546	0.9883–1.1202
0.1%	9.9850	1.0174	0.9452–1.0885
0.05%	9.5899	0.9771	0.8994–1.0528
0.01%	9.3516	0.9528	0.8891–1.0147

All four lower-noise ratio intervals include one; the 5% interval lies above it. There are no guard fallbacks or final clipping on these 500 paths. The reference uses the full limiting span of 134 tests. At the four larger amplitudes every fit retains 134 tests; at 0.01% noise, 65 fits retain 133 and 35 retain 132. Finite-amplitude cutoff and adaptive-instrument effects therefore remain, and the ratios establish compatibility with the limiting reference rather than numerical proof of its attainment. At 5% noise the total eighteen-coefficient RMSE is 0.5813, or root-mean-square coordinate error 0.1370. At this amplitude the coordinate error limits practical precision: Theorem 3 establishes attainability in the small-noise limit, and this construction is an asymptotic benchmark rather than a recommended moderate-noise fitting method. Each level has only 100 paths; the intervals leave the apparent lower-noise trend unresolved. Supplement S9.2 records matrix checks and covariance diagnostics; Table S8 distinguishes the estimators.

8. Discussion and limitations

Boundary-free equations retain exactly the local information available when the initial state is a nuisance parameter. Their restriction has a quantified cost when the state is known: 29% in the Van der Pol aggregate RMSE reference and almost a factor two for Lorenz. Rich spans and correct temporal weights approximate the sharp class reference. Theorem 2 supplies the complementary full-information class for known-initial boundary adaptations. The exact adjoint-range equalities, rather than classical projection formulas or the Brunel remedy, are the theoretical contribution.

The recovery experiment and ablation distinguish class theory from method novelty. Universal endpoint windows recover 97.1% of the weak-to-scoring gap on 1968 Van der Pol records and 95.5% of the weak-to-likelihood gap on 256 Lorenz records. On the 500-record Van der Pol ablation, initial traces alone explain essentially all that gain, reproducing the established Brunel adaptation. The correlated-noise initial control is useful separately, recovering 14.3% of the gap, but contributes no resolved improvement after initial traces. Thus the remedy is an established boundary mechanism with a quantitative OU benchmark. At 5% Van der Pol noise only 32.6% of the likelihood gap is recovered. Covariance weights still require an integrated pilot model; computational superiority has not been demonstrated.

Information loss is direction specific: initial-state projection raises the local cubic-candidate detec-

tion reference by only 0.68%, compared with the larger aggregate coefficient-estimation cost. The supplementary testing study distinguishes prespecified candidates from unrestricted structure discovery. The main empirical attainability check instead matches the full-library estimator to its own finite-span covariance, with uncertainty reported for the limited 100-path samples. At moderate noise this large library remains imprecise, and the local theorem does not establish finite-noise inferential calibration.

The scope is fully observed deterministic polynomial ODEs with known positive sensor covariance. Small-noise influence efficiency, finite-noise feasible accuracy and support selection are separate claims. Exact residual covariance does not by itself give exact finite-noise estimator variance when weights and designs depend on the record. On the matched moderate-noise benchmark, likelihood is more accurate than the original boundary-free weak estimator. Independent-record GMM, fission and cross-fitting are preserved in the separate research archive and lie outside the main claims. Joint span/noise rates, unknown-covariance calibration with heterogeneous states, partial observation and process noise require further analysis.

Data availability

The current code and synthetic-data archive, v1.3.1, is publicly available in the versioned reproducibility repository. Configurations, seed ranges, source fingerprints, coefficient estimates, covariance summaries and SHA256 manifests permit exact input reconstruction and paired audit. Full synthetic time series are regenerated from the frozen seed protocol. Pinned download specifications retrieve original author sources. The separate research archive preserves earlier extensions. Moderate-noise bootstrap validation remains incomplete after a recorded compute-budget decision. The preceding v1.2.0 Zenodo snapshot remains unchanged; a DOI for the current version has not yet been publicly verified. No external experimental dataset is used.

Declaration of generative AI and AI-assisted technologies

Codex assisted with research code, literature searches, Russian-to-English translation, and mathematical derivations and proof development. The author verified the results and takes full responsibility for the manuscript.

References

[1] Brunel NJ-B, Clairon Q, d'Alche-Buc F. Parametric estimation of ordinary differential equations with orthogonality conditions. *Journal of the American Statistical Association* 109 (2014) 173–185. https://doi.org/10.1080/01621459.2013.841583.

[2] Heitzman-Breen N, Dukic V, Bortz DM. A practical identifiability criterion leveraging weak-form parameter estimation. *Bulletin of Mathematical Biology* 88 (2026) 70. https://doi.org/10.1007/s11538-026-01639-x.

[3] Brunton SL, Proctor JL, Kutz JN. Discovering governing equations from data by sparse identification of nonlinear dynamical systems. *Proceedings of the National Academy of Sciences* 113 (2016) 3932–3937. https://doi.org/10.1073/pnas.1517384113.

[4] Schaeffer H, McCalla SG. Sparse model selection via integral terms. *Physical Review E* 96 (2017) 023302. https://doi.org/10.1103/PhysRevE.96.023302.

[5] Messenger DA, Bortz DM. Weak SINDy: Galerkin-based data-driven model selection. *Multiscale Modeling and Simulation* 19 (2021) 1474–1497. https://doi.org/10.1137/20M1343166.

[6] Reinbold PAK, Gurevich DR, Grigoriev RO. Using noisy or incomplete data to discover models of spatiotemporal dynamics. *Physical Review E* 101 (2020) 010203. https://doi.org/10.1103/PhysRevE.101.010203.

[7] Stein ML. Interpolation of spatial data: Some theory for kriging. Springer (1999) https://doi.org/10.1007/978-1-4612-1494-6.

[8] Ying Z. Asymptotic properties of a maximum likelihood estimator with data from a gaussian process. *Journal of Multivariate Analysis* 36 (1991) 280–296. https://doi.org/10.1016/0047-259X(91)90062-7.

[9] Zhang H. Inconsistent estimation and asymptotically equal interpolations in model-based geostatistics. *Journal of the American Statistical Association* 99 (2004) 250–261. https://doi.org/10.1198/016214504000000241.

[10] Godambe VP. An optimum property of regular maximum likelihood estimation. *The Annals of Mathematical Statistics* 31 (1960) 1208–1211. https://doi.org/10.1214/aoms/1177705693.

[11] Bickel PJ, Klaassen CAJ, Ritov Y, Wellner JA. Efficient and adaptive estimation for semiparametric models. Johns Hopkins University Press (1993).

[12] van der Vaart AW. Asymptotic statistics. Cambridge University Press (1998) https://doi.org/10.1017/CBO9780511802256.

[13] Dette H, Pepelyshev A, Zhigljavsky A. Optimal designs in regression with correlated errors. *The Annals of Statistics* 44 (2016) 113–152. https://doi.org/10.1214/15-AOS1361.

[14] Dette H, Konstantinou M, Zhigljavsky A. A new approach to optimal designs for correlated observations. *The Annals of Statistics* 45 (2017) 1579–1608. https://doi.org/10.1214/16-AOS1500.

[15] Messenger DA, Bortz DM. Asymptotic consistency of the WSINDy algorithm in the limit of continuum data. *IMA Journal of Numerical Analysis* 45 (2025) 3264–3312. https://doi.org/10.1093/imanum/drae086.

[16] Varah JM. A spline least squares method for numerical parameter estimation in differential equations. *SIAM Journal on Scientific and Statistical Computing* 3 (1982) 28–46. https://doi.org/10.1137/0903003.

[17] Brunel NJ-B. Parameter estimation of ODE's via nonparametric estimators. *Electronic Journal of Statistics* 2 (2008) 1242–1267. https://doi.org/10.1214/07-EJS132.

[18] Liang H, Wu H. Parameter estimation for differential equation models using a framework of measurement error in regression models. *Journal of the American Statistical Association* 103 (2008) 1570–1583. https://doi.org/10.1198/016214508000000797.

[19] Gugushvili S, Klaassen CAJ. $\sqrt{n}$-consistent parameter estimation for systems of ordinary differential equations: Bypassing numerical integration via smoothing. *Bernoulli* 18 (2012) 1061–1098. https://doi.org/10.3150/11-BEJ362.

[20] Dattner I, Klaassen CAJ. Optimal rate of direct estimators in systems of ordinary differential equations linear in functions of the parameters. *Electronic Journal of Statistics* 9 (2015) 1939–1973. https://doi.org/10.1214/15-EJS1053.

[21] Ramsay JO, Hooker G, Campbell D, Cao J. Parameter estimation for differential equations: A generalized smoothing approach. *Journal of the Royal Statistical Society: Series B* 69 (2007) 741–796. https://doi.org/10.1111/j.1467-9868.2007.00610.x.

[22] Qi X, Zhao H. Asymptotic efficiency and finite-sample properties of the generalized profiling estimation of parameters in ordinary differential equations. *The Annals of Statistics* 38 (2010) 435–481. https://doi.org/10.1214/09-AOS724.

[23] Dattner I, Gugushvili S. Application of one-step method to parameter estimation in ODE models. *Statistica Neerlandica* 72 (2018) 126–156. https://doi.org/10.1111/stan.12124.

[24] Bortz DM, Messenger DA, Dukic V. Direct estimation of parameters in ODE models using WENDy: Weak-form estimation of nonlinear dynamics. *Bulletin of Mathematical Biology* 85 (2023) 110. https://doi.org/10.1007/s11538-023-01208-6.

[25] Rummel N, Messenger DA, Becker S, Dukic V, Bortz DM. Weak form estimation of nonlinear dynamics (WENDy) for nonlinear-in-parameters ODEs. arXiv preprint arXiv:2502.08881v3 (2025).

[26] Fuller WA. Measurement error models. Wiley (1987) https://doi.org/10.1002/9780470316665.

[27] Stefanski LA. Unbiased estimation of a nonlinear function of a normal mean with application to measurement error models. *Communications in Statistics - Theory and Methods* 18 (1989) 4335–4358. https://doi.org/10.1080/03610928908830159.

[28] Chawla A, Bortz DM, Dukic V. Bias and coverage properties of the WENDy-IRLS algorithm. *Handbook of visual, experimental and computational mathematics* Springer Nature Switzerland (2026) 1639–1746. https://doi.org/10.1007/978-3-032-16368-4_86.

[29] Tran A, Bortz DM. Weak form scientific machine learning: Test function construction for system identification. *SIAM Journal on Scientific Computing* 48 (2026) C890–C915. https://doi.org/10.1137/25M1776020.

[30] Chen Z, Fasel U, Bizyaeva A. Fourier weak SINDy: Spectral test function selection for robust model identification. arXiv preprint arXiv:2604.20141v1 (2026).

[31] López C, Naranjo Á, Salazar D, Moore KJ. Weak-form modified sparse identification of nonlinear dynamics. *Journal of Computational Physics* 544 (2026) 114410. https://doi.org/10.1016/j.jcp.2025.114410.

[32] Fasel U, Kutz JN, Brunton BW, Brunton SL. Ensemble-SINDy: Robust sparse model discovery in the low-data, high-noise limit, with active learning and control. *Proceedings of the Royal Society A* 478 (2022) 20210904. https://doi.org/10.1098/rspa.2021.0904.

[33] Uhlenbeck GE, Ornstein LS. On the theory of the brownian motion. *Physical Review* 36 (1930) 823–841. https://doi.org/10.1103/PhysRev.36.823.

[34] Cameron RH, Martin WT. Transformations of weiner integrals under translations. *Annals of Mathematics* 45 (1944) 386–396. https://doi.org/10.2307/1969276.

[35] Feldman J. Equivalence and perpendicularity of gaussian processes. *Pacific Journal of Mathematics* 8 (1958) 699–708. https://doi.org/10.2140/pjm.1958.8.699.

[36] Isserlis L. On a formula for the product-moment coefficient of any order of a normal frequency distribution in any number of variables. *Biometrika* 12 (1918) 134–139. https://doi.org/10.1093/biomet/12.1-2.134.