Methods behind the CVineMarketGen course

Derivations for notebooks 01 to 05

Mamadou Yamar Thioub

September 2026

Abstract

This note carries the derivations the notebooks refer to, so that a notebook can state a rule and show it working without arguing for it. Part I is the generators: four moments matched by a cubic of a normal, a dependence put in the normals, and the quantities used to read a tail. Part II is the specification layer. Part III is dynamics in time: a model as an invertible filter, the two filters used, the test that decides which is adequate, and a moment condition the test cannot see. Part IV collects what none of it does.

Part II is the longest, and it answers this. An analyst who wants scenarios for a universe of assets rarely has a complete description of it. They have a history for some assets, a mean or a volatility for others, exposures for an asset class, a correlation they insist on, a view on a driver. This note states the rule that turns any such partial description into one simulator, and derives the boundaries of what can be asked: the feasible residual variance, the feasible residual shape, the range of a pair correlation, the largest correlation an asset can have with a factor, the completion of the correlations of a factor with no history. It closes with two selection questions: which copula family a pair of near-orthogonal factors can identify, and how the dynamics of a factor are chosen. Numbers appear only where one makes a statement concrete; the notebooks carry the numbers.

Part I
Generators: four moments and a dependence

1 Matching four moments

A target is a number the synthetic data must reproduce. From n observations $r_{i,1}, \dots, r_{i,n}$ of asset i the targets are the sample moments

$$\mu_i = \frac{1}{n}\sum_t r_{i,t}, \quad \sigma_i^2 = \frac{1}{n-1}\sum_t (r_{i,t}-\mu_i)^2, \quad \gamma_{1,i} = \frac{\frac{1}{n}\sum_t (r_{i,t}-\mu_i)^3}{\hat{s}_i^3}, \quad \gamma_{2,i} = \frac{\frac{1}{n}\sum_t (r_{i,t}-\mu_i)^4}{\hat{s}_i^4}, \tag{1}$$

with $\hat{s}_i$ the biased standard deviation, together with the Pearson correlation matrix $\mathbf{\Sigma}^*$. The kurtosis is raw, so the normal has $\gamma_2 = 3$. For N assets that is $4N + N(N-1)/2$ numbers.

Fleishman's (1978) generator writes each asset as a cubic polynomial of a standard normal,

$$X_i = a_i + b_i Z_i + c_i Z_i^2 + d_i Z_i^3, \qquad Z_i \sim \mathcal{N}(0,1). \tag{2}$$

The moments of a polynomial of a normal are polynomials of its coefficients, so requiring $\mathbb{E}X_i = 0$, $\operatorname{Var} X_i = 1$ and the target skewness and kurtosis is three equations in (b_i, c_i, d_i) with $a_i = -c_i$, solved numerically; the mean and volatility are applied afterwards by $r_i = \mu_i + \sigma_i X_i$. A solution exists when

$$\gamma_{2,i} - 3 \;\geq\; \gamma_{1,i}^2 - 2, \tag{3}$$

which every return series met in the work behind this note.

2 Dependence through the normals

Equation (2) fixes each marginal and says nothing about the joint law. Vale and Maurelli (1983) put the dependence in the normals: if (Z_i, Z_k) have correlation ρ_Z, then

$$\mathrm{Corr}(X_i, X_k) = \rho_Z\,(b_i b_k + 3b_i d_k + 3d_i b_k + 9d_i d_k) + \rho_Z^2\,(2c_i c_k) + \rho_Z^3\,(6d_i d_k). \tag{4}$$

For each pair the *intermediate* correlation ρ_Z is the root of that cubic which makes $\mathrm{Corr}(X_i, X_k)$ equal the target $\mathbf{\Sigma}^*_{ik}$. The matrix of intermediate correlations is projected onto the nearest positive-definite one if an edit breaks it, and the normals are drawn from it by a Cholesky factor.

Remark 1. *The construction is exact on the four moments and on every pairwise correlation, and it is silent about everything else. Its copula is the Gaussian one, inherited from* $\mathbf{Z}$ *and never chosen, which is the starting point of §3.*

3 Reading a tail

Four quantities recur in the notebooks and are collected here once.

Value at risk and expected shortfall. For a portfolio with fixed weights $\mathbf{w}$ and return $r_p = \mathbf{w}'\mathbf{r}$,

$$\mathrm{VaR}_\alpha = -q_\alpha(r_p), \qquad \mathrm{ES}_\alpha = -\mathbb{E}\big[r_p \mid r_p \le q_\alpha(r_p)\big], \tag{5}$$

minus the α quantile, and minus the mean of the returns below it.

The quantile implied by four moments. The Cornish-Fisher expansion writes a quantile as a function of the first four moments: with $z = \Phi^{-1}(\alpha)$ and excess kurtosis $\kappa = \gamma_2 - 3$,

$$q_\alpha \approx \mu + \sigma\left[z + \frac{z^2-1}{6}\gamma_1 + \frac{z^3-3z}{24}\kappa - \frac{2z^3-5z}{36}\gamma_1^2\right]. \tag{6}$$

It is the tool that separates two explanations of a missing tail. If a portfolio's own four moments are reproduced, (6) says its quantile is too, so a gap that survives is sampling noise in the estimated quantile rather than a defect of the generator.

Dependence that is not an average. A correlation is one number per pair, an average over the whole distribution. The exceedance correlation conditions on where the first variable is: for a pair (x, y) with x standardised and a threshold z,

$$\hat\rho(z) = \begin{cases} \mathrm{Corr}(x, y \mid x < z) & z < 0, \\ \mathrm{Corr}(x, y \mid x \ge z) & z \ge 0. \end{cases} \tag{7}$$

A profile higher on the left than on the right is dependence concentrated in joint falls. Under a Gaussian copula the profile is flat at the correlation, whatever the marginals, and that flatness is what §16 is about.

Dependence without a direction. Two series can have no asymmetry and still fail to be independent, because the *sizes* of their moves coincide. With s and b the standardised returns of two assets, the co-kurtosis is

$$\mathbb{E}[s^2 b^2] \;=\; 1 + 2\rho^2 \qquad \text{when } s^2 \text{ and } b^2 \text{ are independent}, \tag{8}$$

so the gap between the sample value and $1 + 2\rho^2$ measures co-movement of sizes that no correlation describes. A portfolio inherits it as kurtosis: its variance is then not the same every month, and its tail is fatter than its correlations say.

Where a view lands. A change to one target reaches a portfolio through

$$\sigma_p^2 = \sum_i w_i^2 \sigma_i^2 + \sum_{i \neq k} w_i w_k \, \rho_{ik} \, \sigma_i \sigma_k, \tag{9}$$

which is how the consequence of a stated view can be predicted before the generator is refitted, and therefore how the refit can be checked.

Part II
The specification layer

4 The object

The simulator draws K common factors and maps each simulated factor month to N assets,

$$r_j = \alpha_j + \boldsymbol{\beta}_j' \mathbf{f} + e_j, \qquad j = 1, \ldots, N, \tag{10}$$

with $\mathbf{f}$ the factor vector, e_j independent of $\mathbf{f}$ and, except where a pair is targeted, of the other residuals. Writing $\mathbf{B}$ for the $N \times K$ matrix of betas and $\mathbf{D}$ for the diagonal matrix of residual variances, the implied covariance is

$$\boldsymbol{\Sigma}_r = \mathbf{B} \, \boldsymbol{\Sigma}_f \, \mathbf{B}' + \mathbf{D}, \tag{11}$$

which is what makes the exercise possible at all: $K(K+1)/2$ factor correlations and NK betas in place of $N(N-1)/2$ asset correlations, and an asset with no history reduced to a row of $\mathbf{B}$ and one variance.

The factor vector is drawn by a generator that reproduces the factors' means, volatilities, skewnesses, kurtoses, correlations, and, in the C-vine case, their pairwise tail dependence. The specification layer sits above it: it decides what the generator is asked for, and what each asset carries.

5 The four dimensions of knowledge, and one order

What an analyst can supply falls in four groups: a *history* (of an asset, of a factor, of neither); *moments* (a mean, a volatility, a skewness, a kurtosis, for an asset or for a factor); *exposures* (a beta, or a correlation with a factor, stated for an asset or for a whole class); and *pair correlations* between two assets or two factors. Any subset may be given, in any combination.

The rule is a fixed order, each step taking the previous ones as given:

1. **betas** — stated values; correlations with a factor solved for the beta; regression on the factors marked as fitted; zero elsewhere;
2. **volatility** — the residual variance is the target variance minus the systematic variance, or the history's residual variance when no target is given;
3. **shape** — the residual cumulants are the target cumulants minus the systematic cumulants, or the history's residual shape;
4. **pair correlations** — the residual correlation that delivers the target, inside its feasible range;
5. **mean** — the intercept absorbs it.

The order is forced: the residual's scale cannot be computed before the betas, its shape not before its scale, a pair's room not before both, and the intercept is free of all of them. An empty cell is not a missing value but an instruction: estimate it. What is estimated is the model's own quantity — the beta, the residual, the intercept — and not the asset's moment, which is why a view on a factor reaches an asset whose cells are empty.

6 Betas

6.1 A correlation with a factor as a linear constraint

Let S be the set of factors for which a correlation ρ_k with asset j is stated rather than a beta. Since $\mathrm{Cov}(r_j, f_k) = (\boldsymbol{\Sigma}_f \boldsymbol{\beta}_j)_k$, the constraint $\mathrm{Corr}(r_j, f_k) = \rho_k$ for $k \in S$ reads

$$\boldsymbol{\Sigma}_{SS}\,\boldsymbol{\beta}_S \;=\; \sigma_j \left(\rho_k \sigma_{f_k}\right)_{k\in S} \;-\; \boldsymbol{\Sigma}_{S,-S}\,\boldsymbol{\beta}_{-S}, \tag{12}$$

a linear system in $\boldsymbol{\beta}_S$ once σ_j and the other betas are known. When the asset has a volatility of its own, σ_j is given and (12) is solved once. When it does not, σ_j is itself a function of the betas, and (12) is iterated with

$$\sigma_j^2 \;=\; \boldsymbol{\beta}_j' \boldsymbol{\Sigma}_f \boldsymbol{\beta}_j + \mathrm{Var}(e_j), \qquad e_j \;=\; r_j - \boldsymbol{\beta}_j' \mathbf{f}, \tag{13}$$

the residual being re-estimated at each pass. The iteration converges quickly in practice, but not every target is reachable.

6.2 The saturation bound

Proposition 1. *Let asset j have a history, no volatility target, and one factor f with $\mathrm{Corr}(r_j, f) = \rho_0$ in the sample. If the beta on f is set to b and the residual is the empirical residual $e = r_j - bf$, then the correlation the model delivers is maximised at $b^\star = \sigma_{r_j}/(\rho_0 \sigma_f)$ and equals*

$$\rho_{\max} \;=\; \frac{1}{\sqrt{2 - R_j^2}}, \qquad R_j^2 = \rho_0^2. \tag{14}$$

Proof. Write $x = b\sigma_f/\sigma_{r_j}$. The residual variance is $\mathrm{Var}(e) = \sigma_{r_j}^2 - 2b\,\mathrm{Cov}(r_j, f) + b^2\sigma_f^2$, so by (13) the asset's variance under the imposed beta is $\sigma_j^2 = 2b^2\sigma_f^2 - 2b\,\mathrm{Cov}(r_j, f) + \sigma_{r_j}^2$, and

$$\rho(x) \;=\; \frac{b\sigma_f^2}{\sigma_f \sigma_j} \;=\; \frac{x}{\sqrt{1 - 2\rho_0 x + 2x^2}}.$$

Setting the derivative of ρ^2 to zero gives $2x(1-\rho_0 x) = 0$, hence $x^\star = 1/\rho_0$ and $\rho(x^\star) = (1/\rho_0)/\sqrt{2/\rho_0^2 - 1} = 1/\sqrt{2 - \rho_0^2}$. □

Remark 2. *The bound is the price of leaving the volatility cell empty: raising the beta raises the covariance, but it raises the asset's own variance twice as fast, because the residual no longer fits. With several factors, R_j^2 is the share of variance the factors explain and (14) is an upper bound rather than an identity. A volatility in the assets sheet pins $\mathrm{Var}(e_j)$ through $\sigma_e^2 = v^2 - \boldsymbol{\beta}'\boldsymbol{\Sigma}_f\boldsymbol{\beta}$ and removes the bound: any correlation up to the variance-feasibility limit is then attainable.*

7 Volatility

With a target volatility v and the betas fixed,

$$\sigma_e^2 \;=\; v^2 - \boldsymbol{\beta}'\boldsymbol{\Sigma}_f\boldsymbol{\beta} \;>\; 0, \tag{15}$$

so a target is feasible if and only if it exceeds the volatility the betas alone produce. This is the one refusal an analyst meets most often, and it is informative: it says the exposures and the volatility are inconsistent, and the refusal names the two largest contributors to $\boldsymbol{\beta}'\boldsymbol{\Sigma}_f\boldsymbol{\beta}$ so that one knows which beta to lower. With no target, σ_e is the sample residual standard deviation and the asset's volatility follows from (11): a view on a factor then moves the asset's volatility, as it should.

8 Shape

8.1 The cumulant rule

Cumulants of independent random variables add. With s and k the target skewness and kurtosis of r_j, v its volatility and κ_3, κ_4 the third and fourth cumulants,

$$\kappa_{3,e} \;=\; s\,v^3 - \kappa_3(\boldsymbol{\beta}'\mathbf{f}), \qquad \kappa_{4,e} \;=\; (k-3)\,v^4 - \kappa_4(\boldsymbol{\beta}'\mathbf{f}), \tag{16}$$

and the residual's standardised shape is $s_e = \kappa_{3,e}/\sigma_e^3$, $k_e = 3 + \kappa_{4,e}/\sigma_e^4$. The systematic cumulants are read off a large pilot draw of the fitted factor generator, which costs nothing and avoids assuming anything about the factors' joint law.

Two things follow. First, the residual must carry the whole difference between what is asked and what the factors already give, so the smaller the residual share σ_e^2/v^2, the larger the shape the residual must have: dividing by σ_e^3 and σ_e^4 magnifies the demand. Second, the demand may leave the family of distributions the residual is drawn from.

8.2 The Johnson S_U feasible region

The residual is drawn from a Johnson S_U law, $X = \xi + \lambda \sinh\big((Z-\gamma)/\delta\big)$ with Z standard normal, whose (s,k) region is bounded below by the lognormal line: with $w = e^{1/\delta^2}$,

$$|s| \;=\; (w+2)\sqrt{w-1}, \qquad k \;=\; w^4 + 2w^3 + 3w^2 - 3. \tag{17}$$

Given $|s|$, the first equation has one root $w > 1$ (the left side is increasing) and the second gives the smallest kurtosis the family reaches at that skewness. The floor is 3 at $s = 0$ and grows quickly: about 3.8 at $|s| = 0.6$ and 7.6 at $|s| = 1.5$. A pair (s_e, k_e) below the line is refused, and the refusal reports s_e, k_e, the residual share and the shape the factors already give, which together say whether to soften the target or to lower the exposures.

Remark 3. *The same floor acts silently in the other direction. When no shape is given and the sample residual of an asset is flatter than the family allows — a residual kurtosis of* 2.9 *at a skewness of* -0.3*, say — the kurtosis is raised to the floor and the asset is flagged. The change is small, and hiding it would be worse than making it.*

8.3 How precisely a shape is delivered

A mean and a volatility are matched by construction, up to Monte Carlo error. A shape is not: it is matched through a fitted marginal whose parameters solve (16) at the third and fourth moment, and

the sample skewness and kurtosis of a finite draw of a heavy-tailed variable are themselves noisy. In the notebooks, a target skewness of -0.5 and kurtosis of 5 on an asset whose residual is two thirds of its variance lands within about one tenth in skewness and half a point in kurtosis across seeds. A shape is a target, not an identity, and should be read at that resolution.

9 The correlation between two assets

9.1 The eligible range

From (10), for two assets $i \neq j$,

$$\rho_{ij} \;=\; \underbrace{\frac{\boldsymbol{\beta}_i' \boldsymbol{\Sigma}_f \boldsymbol{\beta}_j}{\sigma_i \sigma_j}}_{\rho_{ij}^{\text{sys}}} + \rho_{e,ij} \sqrt{(1 - R_i^2)(1 - R_j^2)}, \tag{18}$$

where $R_i^2 = \boldsymbol{\beta}_i' \boldsymbol{\Sigma}_f \boldsymbol{\beta}_i / \sigma_i^2$. The betas are already fixed by step 1, so the only free quantity is $\rho_{e,ij} \in [-1, 1]$, and

$$\rho_{ij} \in \Big[\rho_{ij}^{\text{sys}} - w_{ij},\; \rho_{ij}^{\text{sys}} + w_{ij}\Big], \qquad w_{ij} = \sqrt{(1 - R_i^2)(1 - R_j^2)}, \tag{19}$$

with the residual correlation that delivers a feasible target given by

$$\rho_{e,ij} \;=\; \frac{\rho_{ij}\, \sigma_i \sigma_j - \boldsymbol{\beta}_i' \boldsymbol{\Sigma}_f \boldsymbol{\beta}_j}{\sigma_{e,i}\, \sigma_{e,j}}. \tag{20}$$

Two well-explained assets have almost no room; two assets the factors barely touch have almost the whole interval. This is the honest answer to "can I set the correlation between these two funds": partly, and how much is a number you can compute before asking.

9.2 Several pairs at once

Each target writes one off-diagonal entry of the residual correlation matrix $\mathbf{R}_e$ through (20). Individually feasible targets may be jointly infeasible, and the condition is the one the matrix imposes: $\mathbf{R}_e \succeq 0$. The smallest eigenvalue is checked once every pair has been written, and a negative one is reported with the assets involved rather than projected away, because a projection would silently move targets the analyst stated.

10 A factor with no history

A factor known only by a view needs a mean and a volatility — there is nothing to estimate them from — and its skewness and kurtosis fall back to the normal. Its correlations are the real question: an analyst will state one or two, and K are needed.

Proposition 2. *Let $\mathbf{R}$ be the correlation matrix of the K existing factors, S the set of factors whose correlation $\mathbf{r}_S$ with the new factor is stated, and U the rest. Among all $\mathbf{r}_U$ making the extended matrix positive semidefinite, the determinant of that matrix is maximised by*

$$\mathbf{r}_U \;=\; \mathbf{R}_{US}\, \mathbf{R}_{SS}^{-1}\, \mathbf{r}_S, \tag{21}$$

which is also the unique completion whose partial correlations between the new factor and U given S are zero. A completion exists if and only if $\mathbf{r}_S' \mathbf{R}_{SS}^{-1} \mathbf{r}_S \leq 1$.

Proof. The extended matrix has determinant $\det(\mathbf{R})\,(1-\mathbf{r}'\mathbf{R}^{-1}\mathbf{r})$ with $\mathbf{r}=(\mathbf{r}_S,\mathbf{r}_U)$, so maximising it is minimising the quadratic form $\mathbf{r}'\mathbf{R}^{-1}\mathbf{r}$ over $\mathbf{r}_U$. Writing $\mathbf{Q}=\mathbf{R}^{-1}$ in blocks, the first-order condition is $\mathbf{Q}_{US}\mathbf{r}_S+\mathbf{Q}_{UU}\mathbf{r}_U=0$, hence $\mathbf{r}_U=-\mathbf{Q}_{UU}^{-1}\mathbf{Q}_{US}\mathbf{r}_S$; the block-inverse identities $\mathbf{Q}_{UU}=(\mathbf{R}_{UU}-\mathbf{R}_{US}\mathbf{R}_{SS}^{-1}\mathbf{R}_{SU})^{-1}$ and $\mathbf{Q}_{US}=-\mathbf{Q}_{UU}\mathbf{R}_{US}\mathbf{R}_{SS}^{-1}$ turn this into (21). The conditional covariance of the new factor with U given S is $\mathbf{r}_U-\mathbf{R}_{US}\mathbf{R}_{SS}^{-1}\mathbf{r}_S$, zero exactly at (21). At the optimum $\mathbf{r}'\mathbf{R}^{-1}\mathbf{r}=\mathbf{r}_S'\mathbf{R}_{SS}^{-1}\mathbf{r}_S$, so the extended matrix is positive semidefinite for some $\mathbf{r}_U$ if and only if that quantity does not exceed one. □

The completion adds nothing to what was said: the new factor is related to the factors nobody mentioned only through the ones that were. The feasibility condition is also the whole story behind the refusal an analyst meets here. If two existing factors correlate 0.61 with each other, a new factor cannot correlate $+0.9$ with one and -0.9 with the other: the quadratic form is then 4.6, far above one.

Once the correlations are settled, the new factor is attached to the fitted generator by a Gaussian conditional draw. Let $\mathbf{z}$ be the normal scores of the simulated existing factors and $\mathbf{r}$ the vector of target correlations. The new factor's normal score is drawn as

$$z_{new}\mid\mathbf{z}\;\sim\;\mathcal{N}\big(\mathbf{r}'\mathbf{R}^{-1}\mathbf{z},\;1-\mathbf{r}'\mathbf{R}^{-1}\mathbf{r}\big), \tag{22}$$

and mapped through the quantile function of its own marginal. The existing factors keep the joint law the vine gave them — the attachment does not touch them — and the new factor is joined to them by a Gaussian dependence, which is the least one can assume when only correlations were stated.

Part III
Dynamics in time

11 A model as a filter, and its inverse

Everything up to here produces exchangeable observations: a draw has the right moments, the right correlations and the right tails, and no order in time. A history has order, because large moves arrive in runs.

A model of the dynamics is used here not to forecast but as a change of variable in both directions,

$$r_t\;\longmapsto\;\varepsilon_t\quad\text{(the filter)},\qquad\varepsilon_t\;\longmapsto\;r_t\quad\text{(its inverse)}. \tag{23}$$

Forwards, the model removes what its own past implies and leaves a *generalized error* that is i.i.d. over time. Backwards, given i.i.d. errors and the state at the end of the sample, the same equations rebuild returns one period at a time.

That is the whole construction, and it is what makes the two halves of the course compose. The errors are exchangeable, so they can be drawn by the generators of Part I, including jointly across several series; the returns they produce are not exchangeable, because each filter puts the clustering and the starting state back.

What has to be removed. Two questions decide what a filter must take out, and both are diagnostics rather than selection rules. Is the level predictable from its past? The autocorrelations $\hat\rho_j$ of r_t, summarised over m lags by the Ljung-Box statistic

$$Q_m=n(n+2)\sum_{j=1}^{m}\frac{\hat\rho_j^2}{n-j},\qquad Q_m\sim\chi^2_m\text{ under independence}. \tag{24}$$

Is the size predictable? The same statistic on r_t^2, and Engle's ARCH-LM test, the nR^2 of the regression of r_t^2 on its m lags, also χ^2_m. On monthly equity the answer is consistently that the level is unpredictable and the size is not, so a filter has one job.

12 The GARCH family

The return is a mean plus an innovation whose variance depends on the past:

$$r_t = a + \sigma_t \varepsilon_t, \qquad \sigma_t^2 = \omega + \alpha\, e_{t-1}^2 + \beta\, \sigma_{t-1}^2, \qquad e_t = r_t - a, \tag{25}$$

the GARCH(1,1) of Bollerslev. A large move raises tomorrow's variance through α, the variance decays at β, the unconditional level is $\omega/(1-\alpha-\beta)$ and the persistence $\alpha+\beta$. Three variants are used: an AR(1) mean $a + b\, r_{t-1}$; the GJR term $\gamma\, e_{t-1}^2 \mathbf{1}\{e_{t-1} < 0\}$, which lets a fall raise the variance more than a rise; and the EGARCH, the same idea on the log variance. Estimation is by maximum likelihood with a Gaussian innovation, on returns times 100, with the log-likelihood put back on the data's scale so that information criteria are comparable across candidates.

The generalized error is the standardised residual,

$$\varepsilon_t = \frac{r_t - a}{\sigma_t}, \tag{26}$$

and the filter is invertible because ε_t with the state $(e_{t-1}, \sigma_{t-1}^2)$ returns r_t exactly. From the last observed state the inverse runs forward as

$$\sigma_{T+h}^2 = \omega + \alpha\, e_{T+h-1}^2 + \beta\, \sigma_{T+h-1}^2, \qquad r_{T+h} = a + \sigma_{T+h}\, \varepsilon_{T+h}, \qquad h = 1, 2, \ldots \tag{27}$$

13 The hidden Markov model

The second filter says the return comes from one of K regimes, $r_t \mid s_t = k \sim \mathcal{N}(\mu_k, \sigma_k^2)$, with s_t a Markov chain of transition matrix $\mathbf{Q}$. Estimation is by the EM algorithm.

The state is never observed. What the data give is its filtered probability $\eta_{t|t-1,k} = \Pr(s_t = k \mid r_1, \ldots, r_{t-1})$, updated by Bayes' rule at each observation, so the predictive distribution of the next return is the mixture

$$F_t(r) = \sum_{k=1}^{K} \eta_{t|t-1,k}\, \Phi\!\left(\frac{r - \mu_k}{\sigma_k}\right). \tag{28}$$

Its generalized error is a different object from the GARCH's: the Rosenblatt transform of the observation,

$$v_t = F_t(r_t), \qquad \varepsilon_t = \Phi^{-1}(v_t), \tag{29}$$

i.i.d. uniform and i.i.d. normal under the model. It inverts the same way, $r_{T+h} = F_{T+h}^{-1}(\Phi(\varepsilon_{T+h}))$ by bisection on the mixture, with the probabilities updated after each step.

So two filters, two different errors, one a standardised residual and one a probability transform, with the same two properties: i.i.d. by construction, and a recursion that gives the return back.

14 Choosing the dynamics of a factor

The filters of §12 and §13 are candidates, not conclusions. One is chosen from a grid — a constant or an AR(1) mean, a constant variance, a GARCH, a GJR-GARCH or an EGARCH at orders up to two, a Gaussian hidden Markov model with two or three regimes — by one rule in two steps.

Step 1: goodness of fit. Every candidate is filtered, and its generalized error is mapped to a uniform,

$$v_t \;=\; F\Big(\varepsilon_t \mid \mathcal{F}_{t-1};\hat{\boldsymbol{\theta}}\Big)\,, \tag{30}$$

the Rosenblatt transform of the observation under the fitted model. If the model is correctly specified, (v_t) is i.i.d. uniform on $(0,1)$. The discrepancy is measured by the Cramér-von Mises statistic

$$S_n \;=\; \frac{1}{12n} + \sum_{i=1}^{n}\left(v_{(i)} - \frac{i-1/2}{n}\right)^2, \tag{31}$$

whose null distribution depends on the estimated parameters, so the p-value is obtained by parametric bootstrap: B series are simulated from the fitted model, each is refitted, its own statistic computed, and the p-value is the share of bootstrap statistics above the observed one. A candidate is admissible when that p-value is at least the level. For a GARCH filter whose innovation is a Johnson S_U — what the generator actually simulates — the uniforms are taken through that marginal, and the bootstrap refits the marginal as well; testing such a model against a Gaussian F would reject it for a defect it does not have.

Step 2: BIC among the admissible. Among the candidates that pass, the one with the lowest BIC is kept. The order matters and is not interchangeable: the test asks whether a model is adequate, an information criterion only ranks models against each other and will happily return the best of a set of bad ones. Ljung-Box statistics on the generalized error and on its square, and an ARCH-LM statistic, are reported alongside, as diagnostics of *what* a rejected model got wrong. They do not enter the rule.

If no candidate passes, the lowest BIC overall is returned with a warning saying so, which is the honest output: the grid does not contain a model this series cannot reject.

What the Johnson S_U error costs. The GARCH literature fits a Gaussian or a Student t innovation; the Johnson S_U is available in software but rarely used, and two of the reasons matter here. First, its tails cannot be power-law: it is a sinh of a normal, so every moment is finite and no parameter value produces the regularly varying tail that a Student t with ν degrees of freedom gives by construction. Second, its shape is matched by moments rather than estimated by likelihood, which is the Johnson system's own tradition and the same mechanism as (16): the variance parameters are still fitted under normality, and the two shape parameters are solved from the generalized error's skewness and kurtosis. They are weakly identified in a short sample, so a shape read off 230 observations is a rough quantity.

Both costs are accepted for the same reason: one marginal family carries the vine, the asset residuals and the error layer, so a scenario crosses no change of family. The finite fourth moment is also what makes the condition $\mathbb{E}[(\alpha\varepsilon^2+\gamma\varepsilon^2\mathbf{1}\{\varepsilon<0\}+\beta)^2]<1$ computable at all; with a Student t of four or fewer degrees of freedom the fourth moment of the process is infinite whatever the parameters, and the check says nothing.

15 Does the recursion have a finite kurtosis?

A filter can pass the test of §14 and still be unusable, because the test looks at n standardised errors and says nothing about a moment of the process. For the GARCH family with a GJR term the fourth moment of the recursion is finite, and so is the unconditional kurtosis, only if

$$\mathbb{E}\big[(\alpha\,\varepsilon^2+\gamma\,\varepsilon^2\mathbf{1}\{\varepsilon<0\}+\beta)^2\big] \;<\; 1, \tag{32}$$

an expectation taken under the innovation's own law, so it depends on the error's kurtosis as well as on the variance parameters. Nothing in the likelihood or in the selection rule enforces it: the likelihood is finite either way.

Above one, any pooled estimate of the kurtosis of simulated paths grows with the number of paths, and individual paths can leave the range a return can occupy at all. In the course this is not hypothetical. Of seven monthly factors, three selected models violate (32), one of them by a factor of six, and that one produces a simulated month below -100 percent.

The remedy used is a second filter on the selection: among the candidates that pass the goodness-of-fit test, keep the lowest BIC whose (32) is below one. It is not free. On those seven factors the stable choice costs 15 BIC units on one series and 72 and 101 on two others, where the only admissible alternative is a constant variance, so the price of a finite kurtosis is the clustering itself.

Remark 4. *Persistence is the wrong diagnostic here. The series whose condition is 5.9 has a persistence of 0.85, well inside the stationary region; what breaks it is the size of γ interacting with a heavy-tailed error. The two conditions are different inequalities and only the second bounds a kurtosis.*

Part IV
What the rules do not do

16 Why the copula layer is quiet on near-orthogonal factors

A factor model is built so that its factors are nearly uncorrelated, and a copula is then hard to identify. The reason is a matter of orders. Expand a bivariate copula density in the Hermite basis of the normal scores; for the Gaussian copula, Mehler's formula gives

$$c_\rho(u,v) \;=\; \sum_{m\ge 0} \frac{\rho^m}{m!}\, H_m(x)\, H_m(y), \qquad x=\Phi^{-1}(u),\; y=\Phi^{-1}(v). \tag{33}$$

Every one-parameter family reduces to the independence copula at zero dependence, and matching two families on the Pearson correlation ρ forces their first Hermite coefficient to agree. Families therefore differ only from the second coefficient on: their densities differ by $O(\rho^2)$, and the expected log-likelihood gap per observation, a Kullback-Leibler divergence between two densities at distance $O(\rho^2)$, is $O(\rho^4)$.

Put numbers on it. At $\rho = 0.1$ and $n = 230$ monthly observations, the expected gap is of order $n\rho^4 \approx 0.02$ in log-likelihood, while the BIC penalty for one extra parameter is $\frac{1}{2}\log n \approx 2.7$ and the sampling standard deviation of the gap is larger still. No selection rule can separate the families there, and none should pretend to. The consequence for practice is not that the copula is useless but that it earns its place elsewhere: between asset classes, where correlations are 0.5 and above and a Student t or a rotated Gumbel is chosen against a Gaussian with a clear margin, and in the tails of a regime model. At the premia level, a Gaussian dependence with the right marginals is an honest answer.

Whether a pair's dependence is asymmetric is a separate question, and the package answers it with a statistic rather than a family fit: the mean exceedance correlation below -1 standard deviation minus the mean above $+1$, computed on normal scores, with a bootstrap interval. A pair whose interval contains zero is given a symmetric family and nothing else is read into it.

17 Three limits of the asset rule

Three limits are worth stating plainly. The betas are estimated by ordinary least squares on the factors marked as fitted, with no shrinkage and no variable selection: an asset with a short history and many factors will overfit, and the remedy in the specification layer is to state fewer factors per class rather than to regress more. The residuals are independent except where a pair is targeted, so a cluster of assets that move together for a reason no factor captures needs one pair per link, and the joint positive-semidefinite condition of §6.2 will eventually bind. And the scenarios are exchangeable months: the layer above, a filter per factor with its errors drawn jointly, is what gives them an order in time.