Age structure of well-mixed compartmental systems

Department Biogeochemical Processes

Overview

We find general formulas for the transit time and age distributions in well-mixed compartmental systems. As an example we consider a simple three-pool global carbon model with nonlinearities in two different scenarios:

steady state pre-industrial system before 1765

perturbed adding fossil fuels to the atmosphere, considering land use change, 1765-2500

Main quantities of interest

transit time time that a particle needs to travel through the system
exit time − entry time

system age for particles in the system
current time − entry time

compartment age system age of particles in a compartment

Mathematical description

Well-mixed compartmental systems can be descibed by a system of generally **nonlinear** ordinary differential equations of the form

$$\frac{d}{dt}\mathrm{x}(t) = \mathrm{A}(\mathrm{x}(t),t)\,\mathrm{x}(t) + \mathrm{u}(t), \quad t > t_0, \qquad (1)$$

with a given initial value x_0.

Reduction to linear nonautonomous systems

We assume to know (at least numerically) the unique solution of (1) and denote it by x. We then plug it into $\mathrm{A}(\mathrm{x},t))$ and obtain the **linear** system of ordinarzy differential equations.

$$\frac{d}{dt}\mathrm{x}(t) = \mathrm{A}(t)\mathrm{x}(t) + \mathrm{u}(t), \quad t > t_0$$

This equation has the general solution formula

$$\mathrm{x}(t) = \underbrace{\Phi(t,t_0)\mathrm{x}_0}_{\text{age}(t)=t-t_0+\text{initial age}} + \int_{t_0}^{t} \underbrace{\Phi(t,\tau)\mathrm{u}(\tau)}_{\text{age}(t)=t-\tau}\,d\tau,$$

where Φ is the so-called state transition matrix. This leads immediately to the vector of age densities

$$\rho(a,t) = \begin{cases} \Phi(t,t_0)\rho_0(a-(t-t_0)), & a \geq t-t_0, \\ \Phi(t,t-a)\mathrm{u}(t-a), & a < t-t_0, \end{cases}$$

where ρ_0 is the initial age distribution.

Pre-industrial (<1765), system in steady state

Transit time density
$p_T(t) = \mathrm{z}^T\, e^{t\,\mathrm{A}}\,\mathrm{u}$

System age density
$p_A(a) = \sum_j \left(e^{a\,\mathrm{A}}\,\mathrm{u}\right)_j$

Compartmental age density
$p_j(a) = \left(e^{a\,\mathrm{A}}\,\mathrm{u}\right)_j$



Model

Pre-industrial (<1765) + perturbed (1765-2500)



External input fluxes

$u_S = 45, \quad u_A(t) =$ fossil-fuels(t)

$$F_{AT} = 60\,(x_A/700)^{0.2}$$
$$F_{TA}(t) = 60\,x_T/3000 + \text{land-use-change}(t)$$
$$F_{AS} = 100\,x_A/700$$
$$F_{SA} = 100\,(x_S/1000)^{10}$$
$$F_{SD} = 45\,x_S/1000$$

State transition matrix

one-dimensional	$\Phi(t,t_0) = \exp\left[(t-t_0)\,A(t)\right]$	exponential function
multi-dimensional, autonomous	$\Phi(t,t_0) = \exp\left[(t-t_0)\,A(t)\right]$	matrix exponential
multi-dimensional, nonautonomous	$\frac{d}{dt}\Phi(t,t_0) = \mathrm{A}\Phi(t,t_0)$	only numerical solution

Perturbed (1765-2500)

Anthropogenic perturbations



Mean age and transit time



Atmosphere



Transit time

If we define the (backward) transit time of a particle as the age of the particle at the time when it exits the system, we can easily derive explicit formulas for its distribution from the age distribution.

The simple approach stock/flux for the transit time is valid only for a system in steady state. Out of steady state, stock(t)/flux(t) cannot be interpreted as a transit time.

Evolution

Year	Author	System	Quantity	Geometric structure	Example
1983	Anderson	steady state	mean	point	
2009	Manzoni et al.	steady state (simple structure)	density	curve over age	
2016	Rasmussen et al.	linear time dependent	mean	curve over time	
2016	Metzler & Sierra	steady state (all structures)	density	curve over age	
2017	Metzler et al.	linear/nonlinear time dependent	density	surface over age and time	

increasing information

Classification

$\mathrm{A}(\mathrm{x}(t),t)$	nonlinear,
$\mathrm{A}(\mathrm{x}(t)),\mathrm{u}(t)$	nonautonomous
$\mathrm{A}(\mathrm{x}(t)),\mathrm{u}$	nonlinear, autonomous
$\mathrm{A}(t),\mathrm{u}$	linear,
$\mathrm{A},\mathrm{u}(t)$	nonautonomous
$\mathrm{A}(t),\mathrm{u}(t)$	
A,u	linear, autonomous

$\mathrm{x}(t)$ vector of compartment content (e.g. C) at time t

A compartmental matrix, describes fluxes between compartments

$\mathrm{z}(t)$ external outflux vector at time t

$\mathrm{u}(t)$ external input vector at time t

Bibliography

Anderson DH (1983) Compartmental modeling and tracer kinetics, vol 50. Springer Science & Business Media

Manzoni S, Katul GG, Porporato A (2009) Analysis of soil carbon transit times and age distributions using network theories. Journal of Geophysical Research 114(G4):1âĂŞ14, DOI10.1029/2009JG001070

Metzler H, MÃijller M, Sierra CA Age and transit time distributions of well-mixed compartmental systems, PNAS

Metzler H, Sierra CA (2016, in review) Linear autonomous compartmental models as continuous-time Markov chains: transit time and system age distributions

Rasmussen M, Hastings A, Smith MJ, Agusto FB, Chen-Charpentier BM, Hoffman FM, Jiang J, Todd-Brown KEO, Wang Y, Wang YP, Luo Y (2016) Transit times and mean ages for nonautonomous and autonomous compartmental systems. J Math Biol


Holger Metzler


Markus MÃijller


Carlos Sierra