

# Application of Input to State Stability to ecological reservoir models

## Department Biogeochemical Processes

### Challenge

Many models in ecology and biogeochemistry, in particular models of the global carbon cycle, can be generalized as systems of non-autonomous ordinary differential equations (ODEs). For many applications, it is important to determine the stability properties for this type of systems, but most methods available for autonomous systems are not necessarily applicable for the non-autonomous case. We discuss here stability notions for non-autonomous nonlinear models represented by systems of ODEs explicitly dependent on time and a time-varying input signal. Is there a stability concept that is:

- 1 broad enough to encompass these models
- 2 rigorous enough to be proved analytically
- 3 interpretable in ecologically meaningful terms ?

### Example: general soil model

$$\begin{aligned} C &= I(t) + T(C, t) \cdot N(t, C) \cdot C(t) \\ N_{i,i}(C, t) &\geq 0 \quad \forall i \\ T_{i,i}(C, t) &= -1 \quad \forall i \\ T_{i,j}(C, t) &\geq 0 \quad \forall i \neq j \\ \sum_i T_{i,j}(C, t) &= 1 \quad \forall j \end{aligned}$$

This model structure generalizes most SOM decomposition models with any arbitrary number of pools, including those describing nonlinear interactions among state variables. It enforces mass balance and substrate dependence of decomposition, and it is flexible enough to describe:

- 1 Heterogeneity of decomposition rates
  - 2 Transformations of organic matter
  - 3 Environmental variability effects
  - 4 Organic matter interactions
- Examples for nonlinear models are:

- 1 Exoenzyme models [Schimel and Weintraub, 2003, Sinsabaugh and Follstad Shah, 2012]
- 2 AWB [Allison et al., 2010]
- 3 Bacwave [Zelenev et al., 2000]
- 4 MEND [Wang et al., 2013]
- 5 Manzoni [Manzoni and Porporato, 2007]

Also linear models fit into the general framework

- 1 Henin's model [Henin and Dupuis, 1945, Henin et al., 1959]
- 2 ICBM [Andren and Katterer, 1997]
- 3 RothC [Jenkinson and Rayner, 1977, Coleman and Jenkinson, 1999]
- 4 Century [Parton et al., 1987]
- 5 Fontaine 1-4 [Fontaine and Barot, 2005]

One general concept to encompass especially these nonlinear models is clearly desirable.

### Results II, ISS like behavior and proof for example system

The graphs show the reactions of a prototypical class of non-linear two pool soil models to a disturbing time varying signal. This model is a technically simple place holder for ecologically motivated nonlinear systems like the soil models mentioned above to be analyzed in the future. It is given by:

$$\dot{C}_x = I_x(t) - (C_x^2 + C_x) k_x(t) \quad (1)$$

$$\dot{C}_y = I_y(t) - (C_y^2 + C_y) k_y(t) \quad (2)$$

where  $C_x, C_y$  are the carbon contents of two unconnected pools and the bounded periodic functions  $k_x(t)$  and  $k_y(t)$  with:

$$k_{xmin} \leq k_x(t) \leq k_{xmax} \quad (3)$$

$$k_{ymin} \leq k_y(t) \leq k_{ymax} \quad (4)$$

describe the seasonal changes in decomposition speed. e.g.:

$$k_x = \frac{k_{xmax}}{2} + \frac{k_{xmin}}{2} + \frac{1}{2} (k_{xmax} - k_{xmin}) \sin(4t) \quad (5)$$


$$k_y = \frac{k_{ymax}}{2} + \frac{k_{ymin}}{2} + \frac{1}{2} (k_{ymax} - k_{ymin}) \sin(4t) \quad (6)$$

The system can have a fixed point:

$$C_f = \begin{pmatrix} C_{fx} \\ C_{fy} \end{pmatrix}$$

if the input streams have the same period and phase as the decomposition rates. For constant input streams it stays in a predictable region (an invariant set in the phase plane) The plots show the typical behavior of an ISS system: The changes in the state variables will asymptotically converge to a region of stability around an invariant set, whose size is a monotone function of the size of the disturbance (denoted by  $|u|_\infty$ ). For this particular system we proved the ISS property rigorously. The proof relies on the construction of an ISS Lyapunov function whose choice is *not determined but inspired* by a property of the system interpretable in ecologically terms. Expressed casually: "The system can counteract supply changes fast enough". This situation seems to be typical: The problem of establishing ISS for e.g. all the I, T, N models based on the ecologic principles they follow, is too hard. But bio-chemical

of biophysical restrictions could provide clues to ISS proofs for a particular system.




- 1 The four plots on the top show the disturbances.
- 2 The next four plots in the middle show the effect of this disturbances on the solutions for a system with fixed point.
- 3 The next four plots in the middle show the effect of this disturbances on the solutions for the system which no longer has a fixed point, but at least an invariant set, the dark blue square in the middle.

### Conclusion

We propose Input to State Stability (ISS) as candidate for the necessary generalization of the established analysis with respect to equilibria or invariant sets for autonomous systems, and showed for example systems its usefulness by applying it to reservoir models typical for element cycling in ecosystem, e.g. in soil organic matter decomposition. In a forthcoming paper we also showed how ISS generalizes existent concepts formerly only available for Linear Time Invariant (LTI) and Linear Time Variant (LTV) systems to the nonlinear case.

### Results I, ISS as generalization of available stability concepts



The graph shows different stability concepts one could try to establish for the general soil model mentioned above depending on properties of its components I, T and N. The hardest to prove is Input to State Stability for time varying systems (ISStv) in the lower left corner. It turns out that ISStv also generalizes all the other concepts mentioned:

- In the case of Linear Time Invariant (LTI) systems ISS follows from the properties of the matrix.(eigenvalues)
- In the case of Linear Time Variant (LTV) systems it can be established if sufficient information about the state transition operator allows to prove uniform asymptotic stability UAS.
- For input free system (on the blue right-hand side) ISS reduces to Global Asymptotic Stability (GAS)
- ...

### Bibliography

I. S. D. Allison, M. D. Wallenstein, and M. A. Bradford. Soil-carbon response to warming dependent on microbial physiology. *Nature Geosci.* 3(5):336–340, 2010. 10.1038/ngeo846.

O. Andren and T. Katterer. Icbm: The introductory carbon balance model for exploration of soil carbon balances. *Ecological Applications*, 7(4):1226–1236, 1997.

K. Coleman and D. Jenkinson. Rothc-26.3 a model for the turnover of carbon in soil: model description and windows user guide (modified 2008). Technical report, IACR Rothamsted, 1999.

S. Fontaine and S. Barot. Size and functional diversity of microbe populations control plant persistence and long-term soil carbon accumulation. *Ecology Letters*, 8(10):1075–1087, 2005. ISSN 1461-0248. doi: 10.1111/j.1461-0248.2005.00813.x. URL <http://dx.doi.org/10.1111/j.1461-0248.2005.00813.x>.

S. Henin and M. Dupuis. Essai de bilan de la matière organique du sol. *Annales Agronomique*, 15:17–29, 1945.

S. Henin, G. Monnier, and L. Turc. Un aspect de la dynamique des matières organiques du sol. *Comptes rendus hebdomadaires des séances de l'Académie des sciences*, 248(1):138–141, 1959.

D. S. Jenkinson and J. H. Rayner. The turnover of soil organic matter in some of the rothamsted classical experiments. *Soil Science*, 123(6):239–305, 1977.

S. Manzoni and A. Porporato. A theoretical analysis of nonlinearities and feedbacks in soil carbon and nitrogen cycles. *Soil Biology and Biochemistry*, 39(7):1542–1556, 2007.


W. J. Parton, D. S. Schimel, C. V. Cole, and D. S. Ojima. Analysis of factors controlling soil organic matter levels in great plains grasslands I. *Soil Sci. Soc. Am. J.*, 51(5):1173–1179, 1987. URL <https://www.ssis.org/publications/ssaaj/abstracts/51/5/1173>.

J. P. Schimel and M. N. Weintraub. The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biology and Biochemistry*, 35(4):549–563, 2003.

R. L. Sinsabaugh and J. J. Follstad Shah. Ecological stoichiometry and ecological theory. *Annual Review of Ecology, Evolution, and Systematics*, 2012. doi: 10.1146/annurev-ecolsys-071112-124414.

G. Wang, W. M. Post, and M. A. Mayes. Development of microbial-enzyme-mediated decomposition model parameters through steady-state and dynamic analyses. *Ecological Applications*, 23(1):255–272, 2013/09/20 2013. doi: 10.1890/12-0681.1. URL <http://dx.doi.org/10.1890/12-0681.1>.

V. Zelenev, A. van Bruggen, and A. Semenov. "BACWAVE", a spatial-temporal model for traveling waves of bacterial populations in response to a moving carbon source in soil. *Microbial Ecology*, 40(2):260–272, 2000. ISSN 0095-3628. doi: 10.1007/s002480000029. URL <http://dx.doi.org/10.1007/s002480000029>.



Get me out, I do not belong here....

