
a
LARRY lineage agreement (S14)
Spearman, predicted vs clonal displacement (◆) and 5 nulls (●)
refer- ence
selected
held-out day 6 (hatched = selected; 95 % CI)
energy distance to held-out day 6
flow
mean displ.
no move
b
S11 velocity agreement
density
per-cell velocity cosine
LARRY cell-type λ₁ (S11; n = OT cells)
Undifferentiated (5962)
Monocyte (748)
Neutrophil (692)
Baso (319)
Lymphoid (74)
Meg (69)
Mast (67)
Erythroid (23)
Eos (21)
Ccr7_DC (18)
pDC (7)
median λ₁ (flow J)
LARRY OT vs lineage (median 0.69)
scNT labelling vs dynamo (median -0.05)
scNT labelling vs scVelo (median 0.03)
scNT scVelo vs dynamo (median 0.15)
OT flow J
lineage flow J
c
S20 runtime
OT
drift fit + J
wall time (s)
cells (3 days)
S20 memory
dense
minibatch
lowrank_r50
lowrank_r200
peak RSS, OT stage (GB)
cells (3 days)
S20 agreement vs dense
agreement with dense OT
minibatch
lowrank_r50
lowrank_r200
cosine (median)
λ₁ Spearman
cells (3 days)
d
full LARRY vs subsample (Spearman 0.82)
λ₁, all 130,887 cells (S20 minibatch)
median λ₁, 4,000 cells/day (S11)
Lymphoid
pDC
Undifferentiated
Ccr7_DC
Erythroid
Meg
Eos
Monocyte
Mast
Baso
Neutrophil