# Validating the Dietary Inflammatory Index: A Comprehensive Review for Biomedical Research and Clinical Application

Generated at: 2026-03-31
Source URL: https://www.nutribiosci.com/posts/validating-the-dietary-inflammatory-index-a-comprehensive-review-for-biomedical-research-and-clinical-application

# Validating the Dietary Inflammatory Index: A Comprehensive Review for Biomedical Research and Clinical Application

## Abstract

This article provides a systematic review of the validation methodologies and applications of the Dietary Inflammatory Index (DII) and its derivatives. It explores the scientific foundation of dietary inflammatory potential scoring, examines the correlation between DII scores and established inflammatory biomarkers across diverse populations, and compares the performance of various dietary indices. Aimed at researchers, scientists, and drug development professionals, this review synthesizes current evidence, addresses methodological challenges, and highlights the potential of these tools for refining clinical nutrition strategies, personalizing dietary interventions, and informing clinical trial designs in chronic disease prevention and management.

## The Science of Quantifying Dietary Inflammation: From Concept to Clinical Tool

The **Dietary Inflammatory Index (DII)** is a literature-derived, quantitative tool designed to assess the inflammatory potential of an individual's overall diet [1]. It was developed to summarize the effect of dietary parameters on specific inflammatory biomarkers, providing researchers and clinicians with a means to evaluate how diet may influence chronic inflammation—a key driver in the pathogenesis of numerous chronic diseases [2] [3]. Unlike dietary patterns derived from the same study in which they are applied, the DII was designed for standardized, reproducible application across diverse populations, enhancing between-study comparability [4] [5] [3]. The DII represents a significant advancement in nutritional epidemiology by creating a hypothesis-driven assessment tool based on a comprehensive review of scientific literature linking dietary components to inflammation [3].

*Table 1: Key Characteristics of Dietary Inflammatory Indices*

| Index Name | Development Basis | Number of Dietary Components | Key Inflammatory Biomarkers | Primary Output |
|---|---|---|---|---|
| Original DII | Literature review through 2010 | 45 | IL-1β, IL-4, IL-6, IL-10, TNF-α, CRP | A continuous score; positive values indicate pro-inflammatory diet, negative values indicate anti-inflammatory diet [1] [3] |
| Empirical DII (EDII) | Reduced rank regression followed by stepwise linear regression | 18 food groups | IL-6, CRP, TNFαR2 | Weighted sum of food groups; predicts inflammatory biomarker concentrations [4] [5] |
| Empirical Anti-inflammatory Diet Index (eADI) | 10-fold feature selection with filtering (Lasso regression) | 17 food groups | hsCRP, IL-6, TNF-R1, TNF-R2 | Summed scores of consumption tertiles (0, 0.5, 1 point); lower concentrations of inflammatory biomarkers [6] |
| China-DII (CHINA-DII) | Adapted from original DII using Chinese dietary intake data | 27 | hs-CRP | A score validated for Chinese population; positively correlated with hs-CRP [7] |
| Food Inflammation Index (FII) | Adapted from DII with weighted algorithm | Various whole foods | Validated by NHANES | Inflammatory effect score for whole foods, revealing heterogeneity within food groups [8] |

### Principles and Development of the DII

#### Foundational Concepts and Theoretical Framework

The development of the DII was motivated by the growing understanding of the role of inflammation in health and the recognition that diet plays a crucial role in modulating inflammatory processes [3]. The DII is grounded in the principle that dietary components can have either pro-inflammatory or anti-inflammatory effects, which can be quantified based on their influence on specific inflammatory biomarkers [1]. The initial DII

development involved a comprehensive literature review spanning from 1950 to 2010, identifying research articles that assessed the role of whole foods and dietary components on six inflammatory biomarkers: interleukins (IL-1β, IL-4, IL-6, and IL-10), tumor necrosis factor-α (TNF-α), and C-reactive protein (CRP) [1]. This evidence-based approach distinguishes the DII from other dietary indices such as the Healthy Eating Index (HEI) or Mediterranean Dietary Index (MDI), which are based on dietary recommendations or specific cuisines rather than direct links to inflammatory pathways [1] [3].

### Methodological Evolution and Algorithm Development

The DII's development has undergone significant methodological refinements since its first version debuted in 2009 [3]. The original DII algorithm relied on raw consumption amounts, which posed challenges due to right-skewing of dietary data and the need for arbitrary adjustments to regulate the influence of certain nutrients [3]. The enhanced DII algorithm addressed these limitations by linking reported dietary intake to global norms of intake derived from 11 datasets from around the world, including countries such as Australia, Denmark, Japan, Mexico, the United States, and others [3]. This global database provides means and standard deviations for the intakes of each food parameter, serving as a comparative reference for calculating individual DII scores [3]. The current DII calculation involves several steps: first, dietary intake data is linked to global norms to compute a z-score for each parameter; these z-scores are then converted to cumulative proportions; finally, centered percentiles are calculated and multiplied by the respective inflammatory effect score for each food parameter [3]. The global inflammatory effect scores for the 45 food parameters range from -1 (maximally anti-inflammatory) to +1 (maximally pro-inflammatory), with the overall DII score representing the sum of these weighted values [3].




*Diagram 1: DII Development Workflow. This diagram illustrates the evidence-based methodology for creating the Dietary Inflammatory Index, from literature review to final score calculation.*

## Comparative Analysis of DII Variants and Validation Studies

### Empirical Adaptations of the DII Framework

Several empirical adaptations of the original DII have been developed to address specific research needs or population characteristics. The **Empirical Dietary Inflammatory Index (EDII)** was developed using reduced rank regression models followed by stepwise linear regression analyses in the Nurses' Health Study to identify a dietary pattern most predictive of three plasma inflammatory markers: IL-6, CRP, and TNFαR2 [4]

[5]. The resulting EDII comprises 18 food groups (9 anti-inflammatory and 9 pro-inflammatory) and has demonstrated strong construct validity in independent samples of women and men [4] [5]. More recently, the **empirical Anti-inflammatory Diet Index (eADI)** was developed using a 10-fold feature selection with filtering (based on Lasso regression) to select food groups most correlated with multiple inflammatory biomarkers (hsCRP, IL-6, TNF-R1, and TNF-R2) [6]. The eADI-17 includes 17 food groups (11 with anti-inflammatory and 6 with pro-inflammatory potential) and was designed to be a user-friendly index with clear scoring criteria for clinical application [6]. Cultural adaptations have also emerged, such as the **China Dietary Inflammatory Index (CHINA-DII)**, which was developed using dietary intake data specific to Chinese adults to better reflect regional dietary patterns [7].

*Table 2: Validation Studies of Dietary Inflammatory Indices Across Populations*

| Study Reference | Population | Sample Size | Key Findings | Statistical Significance |
|---|---|---|---|---|
| Garcia-Arellano et al. [1] | Spanish PREDIMED study (high CV risk) | 7,216 | Adherence to Mediterranean diet inversely associated with DII; higher DII associated with higher BMI | Significant association after median 5-year follow-up |
| Tabung et al. [4] [5] | NHS, NHS-II, HPFS validation | 5,230 (NHS), 1,002 (NHS-II), 2,632 (HPFS) | EDII significantly predicted concentrations of IL-6, CRP, TNFαR2, adiponectin in independent samples | P-trend < 0.0001 to 0.003 for extreme quintile comparisons |
| Frontiers in Nutrition 2025 [9] | US non-diabetic adults (NHANES) | 13,408 | Highest DII quartile associated with elevated all-cause and CV mortality | HR = 1.554 for all-cause mortality; HR = 2.100 for CV mortality |
| CHINA-DII Study [7] | Chinese gastric cancer patients | 256 | CHINA-DII scores positively correlated with hs-CRP levels | r = 0.20, p ≤ 0.001; OR = 1.90 for hs-CRP ≥ 3 mg/L |
| eADI Study [6] | Cohort of Swedish Men | 4,432 | Each 4.5-point eADI-17 increment associated with lower inflammatory biomarkers | 12% lower hsCRP, 6% lower IL-6, 8% lower TNF-R1, 9% lower TNF-R2 |

### Health Outcome Validation and Predictive Performance

The predictive validity of DII scores has been extensively evaluated in relation to various health outcomes across diverse populations. Research has consistently demonstrated that higher (more pro-inflammatory) DII scores are associated with increased risk of numerous inflammation-related conditions. In the Spanish PREDIMED study of high cardiovascular risk participants, those in the highest DII category showed significantly higher average BMI compared to those in the lowest quartile [1]. A recent study of US non-diabetic adults from NHANES (2009-2018) found that participants in the highest DII quartile exhibited elevated risks of both all-cause mortality (HR = 1.554) and cardiovascular mortality (HR = 2.100) compared to those with lower DII scores [9]. The DII has also been shown to predict mental health outcomes, with a study of North American adults revealing that participants consuming the most pro-inflammatory diet had a 24% higher risk of developing depressive symptoms compared to those with the most anti-inflammatory diet [1]. A comparative study in postmenopausal women found that both the DII and a food-based dietary inflammatory index (FDII) were significantly associated with the severity of sexual symptoms and quality of life during menopause, with both indices demonstrating significant predictive power [10].

## Research Reagent Solutions for DII Validation Studies

*Table 3: Essential Research Reagents and Materials for DII Validation Studies*

| Reagent/Material | Manufacturer/Source | Function in DII Research | Example Application |
|---|---|---|---|
| High-sensitivity CRP (hsCRP) Assay | Denka Seiken Company (Architect Ci8200 analyzer) | Quantifies low-grade inflammation using high-sensitivity immunoturbidimetric assay | Primary inflammatory biomarker in DII validation studies [5] |
| IL-6 ELISA Kit | R&D Systems | Measures interleukin-6 concentrations via enzyme-linked immunosorbent assay | Component of inflammatory biomarker panels for DII validation [5] |
| TNF-α Receptors Assay | Olink Proteomics (CVD II and III panels) | Determines TNF-R1 and TNF-R2 concentrations using normalized protein expression (NPX) | Used in development of empirical anti-inflammatory diet indices [6] |

| Reagent/Material | Manufacturer/Source | Function in DII Research | Example Application |
|---|---|---|---|
| Adiponectin Radioimmunoassay | Linco Research | Quantifies adiponectin levels via competitive radioimmunoassay | Anti-inflammatory adipokine included in EDII validation [5] |
| Food Frequency Questionnaire (FFQ) | Study-specific adaptations | Assesses habitual dietary intake for DII calculation | Primary dietary assessment method in cohort studies [4] [6] [7] |
| 24-hour Dietary Recall | Automated Multiple-Pass Method (USDA) | Collects detailed dietary intake data for DII calculation | Used in NHANES and other population studies [9] |

## Detailed Experimental Protocols for DII Validation

### Blood Collection and Biomarker Assessment Protocol

Standardized protocols for blood collection and processing are critical for reliable validation of dietary inflammatory indices. In major cohort studies such as the Nurses' Health Study and Health Professionals Follow-up Study, blood samples are collected in the morning following an overnight fast [5]. For lithium-heparin plasma preparation, blood samples are typically light-protected, and after a delay of 15-20 minutes at room temperature, the samples are centrifuged at approximately 1600 g for 10-15 minutes at 4°C [6] [5]. The plasma is then frozen in multiple aliquots and stored at -80°C until analysis to preserve biomarker integrity [6]. Quality control procedures include interspersing quality-control samples randomly among case-control samples, with laboratory personnel blinded to quality-control and case-control status for all assays [5]. The coefficients of variation for inflammatory biomarkers should be monitored, with typical intra-assay CVs ranging from 1.0% to 12.8% for various inflammatory markers across batches [5]. For studies measuring multiple biomarkers, an overall inflammatory marker score can be derived by computing a z-score for each inflammatory marker and summing these z-scores to create a standardized composite score for each participant [5].

### Dietary Assessment and DII Calculation Methodology

The calculation of DII scores requires comprehensive dietary assessment, typically obtained through Food Frequency Questionnaires (FFQs) or 24-hour dietary recalls [9]. The validity and reliability of the dietary assessment method should be established in the target population prior to DII calculation [5] [7]. For the original DII, intake data for the 45 food parameters are first standardized to a global reference database using z-scores calculated as (individual intake - global mean intake) / global standard deviation [3]. These z-scores are then converted to percentiles and centered by multiplying by 2 and subtracting 1 to achieve a symmetric distribution [3]. Each centered percentile is multiplied by the corresponding inflammatory effect score derived from the literature, and the sum of all values represents the overall DII score [3]. For the Empirical DII (EDII), the weighted sum of 18 food groups is calculated using regression weights derived from reduced rank regression analysis [4] [5]. In the empirical Anti-inflammatory Diet Index (eADI), food groups are scored based on consumption tertiles (0, 0.5, and 1 point), with the total score representing the sum across all food groups [6].




*Diagram 2: DII Validation Protocol. This workflow outlines the key methodological steps for validating dietary inflammatory indices in research studies, from data collection to statistical analysis.*

The Dietary Inflammatory Index represents a significant methodological advancement in nutritional epidemiology, providing a standardized, evidence-based tool for assessing the inflammatory potential of diet across diverse populations. The development of the DII and its empirical variants (EDII, eADI) has enabled researchers to quantitatively evaluate the relationship between diet-associated inflammation and various health outcomes, from all-cause mortality to specific disease risks [9] [5]. The consistent associations observed between higher DII scores and adverse health outcomes across multiple studies and populations underscore the utility of this tool for both research and clinical applications [1] [9] [7]. Future research directions include further validation of DII variants in diverse cultural contexts, refinement of the scoring algorithms as new evidence emerges, and exploration of the DII's potential for guiding personalized nutritional interventions to reduce chronic inflammation and improve public health outcomes [2] [6] [7]. The ongoing development of culture-specific indices like the CHINA-DII highlights the importance of adapting dietary assessment tools to regional eating patterns while maintaining the core principles of the DII framework [7].

In the field of nutritional science, the validation of dietary intervention effects requires precise and reliable measurement of inflammatory status. The **Dietary Inflammatory Index (DII)** was developed as a novel, hypothesis-driven tool to quantify the inflammatory potential of whole diets based on their capacity to influence systemic inflammation [4]. Accurate validation of DII scores in experimental and clinical settings depends on measuring specific inflammatory biomarkers that reflect the diet-induced inflammatory state. Among the numerous biomarkers available, **C-reactive protein (CRP), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNF-α)** have emerged as core biomarkers for this validation process due to their established roles in the inflammatory cascade and responsiveness to dietary modulation.

The selection of appropriate inflammatory biomarkers is complicated by their diverse biological functions, dynamics, and methodological considerations. Inflammation represents a complex physiological response to infection, injury, or harmful stimuli, coordinated by a network of signaling molecules and cellular effectors [11]. While acute inflammation is a protective mechanism, chronic low-grade inflammation has been identified as a secret contributor to numerous age-related diseases and metabolic disorders [11]. This article provides a comprehensive comparison of established and emerging inflammatory biomarkers, with a specific focus on their application in validating dietary inflammatory potential within research contexts.

## Comparative Analysis of Core Inflammatory Biomarkers

### Biological Characteristics and Measurement

**Table 1: Fundamental Characteristics of Core Inflammatory Biomarkers**

| Biomarker | Biological Role | Source Cells/Tissues | Baseline Level | Peak Response | Half-Life |
|---|---|---|---|---|---|
| **CRP** | Acute-phase protein, pathogen recognition, complement activation | Hepatocytes (primarily) | <1 μg/mL [12] | 48 hours [13] [12] | 19 hours [13] |
| **IL-6** | Pro-inflammatory cytokine, stimulates CRP production, metabolic regulation | Macrophages, neutrophils, adipocytes, endothelial cells | Varies by population | 90-120 minutes [14] | Short (minutes) |
| **TNF-α** | Pro-inflammatory cytokine, regulates immune cell function | Macrophages, T-cells, natural killer cells | Varies by population | 90-120 minutes [14] | Short (minutes) |

CRP, IL-6, and TNF-α occupy different positions in the inflammatory cascade. IL-6 and TNF-α are among the **first-line cytokines** released in response to inflammatory stimuli, with TNF-α helping to determine the strength, effectiveness and duration of inflammatory reactions [15]. IL-6 subsequently acts as the **primary stimulator** of hepatic production of acute-phase proteins, including CRP [12]. This temporal relationship creates a coordinated inflammatory response, with cytokine levels rising within hours and acute-phase proteins peaking days after the initial trigger [14].

Beyond their classical inflammatory roles, recent research indicates that both IL-6 and CRP participate in broader physiological processes. IL-6 plays important roles in **metabolic regulation**, tissue maintenance, and energy allocation, with some effects actually being anti-inflammatory rather than pro-inflammatory [12]. Similarly, CRP exists in different isoforms with distinct functions, participating not only in inflammation but also in various **tissue maintenance** processes [12]. These multifaceted roles complicate the interpretation of these biomarkers as simple indicators of inflammation in nutritional studies.

### Performance Characteristics in Disease Prediction

**Table 2: Biomarker Performance in Predicting Clinical Outcomes Across Populations**

| Biomarker | Population | Clinical Endpoint | Effect Size | Reference |
|---|---|---|---|---|
| **IL-6** | Medical inpatients at nutritional risk | 30-day mortality | Adjusted HR 3.5 (95% CI 1.95-6.28) [16] [14] | EFFORT Trial 2025 |
| **IL-6** | Frail community-living elderly | 4-year all-cause mortality | HR 2.18 (95% CI 1.29-3.69) [15] | ilSIRENTE Study |
| **CRP** | Frail community-living elderly | 4-year all-cause mortality | HR 2.58 (95% CI 1.52-4.40) [15] | ilSIRENTE Study |
| **TNF-α** | Frail community-living elderly | 4-year all-cause mortality | HR 1.26 (95% CI 0.74-2.15) [15] | ilSIRENTE Study |
| **IL-6** | Elderly with depression | Cross-sectional comparison | SMD 0.38 (95% CI 0.16-0.60) [17] | Meta-analysis 2018 |
| **Composite Score** | Frail community-living elderly | 4-year all-cause mortality | Highest risk for 3 elevated markers [15] | ilSIRENTE Study |

The predictive performance of inflammatory biomarkers varies significantly across different populations and clinical contexts. In frail, community-living elderly individuals, both IL-6 and CRP demonstrated **strong predictive value** for all-cause mortality over a 4-year follow-up period, while TNF-α did not show a significant association after adjustment for confounders [15]. The ilSIRENTE study further revealed that a **composite summary score** combining all three inflammatory markers identified individuals with the highest mortality risk, suggesting that combined biomarker approaches may enhance predictive power [15].

In hospital settings, IL-6 has demonstrated particular utility for risk stratification. A recent secondary analysis of the EFFORT trial found that medical inpatients at risk of malnutrition with high IL-6 levels (≥11.2 pg/mL) had a **more than 3-fold increase** in 30-day mortality compared to those with lower levels [16] [14]. Interestingly, the same study found that CRP and TNF-α were not independently associated with mortality in this patient population, highlighting the potential superior prognostic value of IL-6 in acute care settings [16] [14].

The predictive capacity of inflammatory biomarkers extends to specific disease contexts. A meta-analysis of elderly populations found significantly elevated IL-6 levels in those with depression compared to controls, while differences in TNF-α and CRP did not reach statistical significance after rigorous adjustment [17]. For Alzheimer's disease, only IL-1β showed significant elevation among the inflammatory markers analyzed [17]. These findings suggest biomarker performance is condition-specific and influenced by underlying pathophysiology.

## Biomarkers in Dietary Intervention and Nutritional Research

### Validation of Dietary Inflammatory Index

The Dietary Inflammatory Index (DII) was developed as a standardized tool to quantify the inflammatory potential of whole diets based on their effects on specific inflammatory biomarkers [4]. The initial development and validation of the DII incorporated **six inflammatory biomarkers** (IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP), with subsequent validations focusing primarily on CRP, IL-6, and TNF-α as the most responsive markers to dietary manipulation [4] [18].

In a cross-sectional study of postmenopausal women, the DII score demonstrated a significant positive association with IL-6 levels, with individuals in the highest DII category (most pro-inflammatory diet) showing significantly elevated IL-6 compared to those in the lowest category [18]. The same study found pro-inflammatory diets were associated with unfavorable lipid profiles, including higher triglycerides and TG/HDL-C ratios [18]. These findings support the **construct validity** of the DII and highlight IL-6 as a responsive biomarker for dietary inflammatory potential.

The DII represents an example of a **hypothesis-driven, empirically derived dietary pattern** that assesses diet quality based on inflammatory potential [4]. Its calculation involves linking reported dietary intakes to a global database, with food parameter-specific z-scores transformed and multiplied by inflammatory effect scores based on the published literature [18]. This method allows standardized calculation across different populations, addressing a major limitation of study-specific dietary patterns [4].

### Predicting Response to Nutritional Therapy

Inflammatory biomarkers not only reflect dietary influences but may also predict response to nutritional interventions. A secondary analysis of the EFFORT trial demonstrated that the mortality benefit of individualized nutritional support was **modulated by baseline inflammation** [16] [14]. Patients with high IL-6 levels showed a less pronounced mortality benefit from nutritional therapy compared to those with lower inflammation (hazard ratio 0.82 vs. 0.32) [16] [14].

Similarly, patients with elevated CRP levels (>100 mg/dL) showed diminished response to nutritional intervention, with hazard ratios of 1.25 compared to 0.47 in those with lower CRP [16] [14]. These findings support the concept that **high inflammatory states** are associated with reduced benefits from nutritional therapy, potentially informing personalized nutritional strategies based on inflammatory profiles [16] [14].

The EFFORT trial analysis further suggested that while both CRP and IL-6 effectively predicted treatment response to nutritional therapy, IL-6 provided additional value as a **prognostic marker for increased mortality** [16] [14]. This dual functionality positions IL-6 as a particularly valuable biomarker in nutritional research and clinical practice.

## Methodological Considerations for Biomarker Measurement

### Analytical Approaches and Protocols

**Table 3: Research Reagent Solutions for Inflammatory Biomarker Analysis**

| Reagent/Assay | Biomarker | Function | Methodology |
|---|---|---|---|
| **U-PLEX Human Assay (MSD)** | IL-6, TNF-α | Multiplex cytokine quantification | Electrochemiluminescence detection [14] |
| **High Sensitivity ELISA** | IL-6, TNF-α | Sensitive cytokine measurement | Enzyme-linked immunosorbent assay [15] |
| **Latex-enhanced immunoturbidimetry** | CRP | High-sensitivity CRP quantification | Automated clinical chemistry analyzers [15] |
| **Nuclear Magnetic Resonance** | GlycA | Composite inflammatory marker | Spectral deconvolution algorithm [19] |

Accurate measurement of inflammatory biomarkers requires appropriate analytical methods with sufficient sensitivity and specificity. For cytokine measurement, **enzyme-linked immunosorbent assays (ELISA)** remain widely used, with high-sensitivity versions enabling detection of low levels characteristic of chronic inflammation [15]. More recently, **multiplex assay systems** such as the MSD Multi-Spot Assay System enable simultaneous measurement of multiple cytokines from small sample volumes, improving efficiency in large-scale studies [14].

CRP measurement typically employs **immunoturbidimetric methods** on automated clinical chemistry platforms, with high-sensitivity (hs-CRP) assays extending detection limits to the lower ranges relevant for cardiovascular risk assessment and chronic inflammation monitoring [15] [13]. For novel biomarkers like GlycA, which reflects integrated concentrations and glycosylation states of several acute-phase proteins, **nuclear magnetic resonance (NMR) spectroscopy** with spectral deconvolution algorithms provides the necessary analytical approach [19].

Sample handling and processing significantly impact biomarker stability and measurement accuracy. Most cytokines have short half-lives and require careful sample processing with immediate centrifugation and storage at **-80°C** to preserve integrity [14]. Blood samples should be collected in the morning after an overnight fast to minimize diurnal variation, and measurements should be performed in duplicate to enhance reliability [15] [14].

### Emerging and Composite Biomarkers

Beyond the core biomarkers, several emerging and composite markers show promise for nutritional research. **GlycA**, a composite biomarker derived from NMR spectroscopy, reflects the integrated concentrations and glycosylation states of several acute-phase proteins [19]. In the Multi-Ethnic

Study of Atherosclerosis, GlycA demonstrated predictive value for total death, cardiovascular events, inflammatory-related events, and total cancer events, independently of established inflammatory markers including hsCRP, IL-6, and d-dimer [19].

Composite scores combining multiple inflammatory markers may provide enhanced predictive power compared to individual biomarkers. The ilSIRENTE study created a **composite inflammation score** based on IL-6, CRP, and TNF-α levels, finding the highest mortality risk in individuals with elevation of all three markers [15]. Similarly, infection probability scores incorporating CRP along with clinical signs and SOFA scores have shown improved diagnostic accuracy for infection in ICU settings [13].

Questionnaire-based measures like the DII offer non-invasive alternatives for assessing inflammatory potential, though they require validation against biological markers. The DII has demonstrated consistent correlations with inflammatory biomarkers including CRP and IL-6 across different populations [4] [11] [18], supporting its utility as a **standardized dietary assessment tool** in research settings.

## Signaling Pathways and Experimental Workflows

### Inflammatory Signaling Cascade




*Figure 1: Inflammatory Signaling Cascade and Biomarker Dynamics. This diagram illustrates the sequential activation of inflammatory mediators in response to stimuli, showing the position of TNF-α, IL-6, and CRP in the inflammatory cascade.*

The inflammatory response follows a coordinated sequence of mediator release and cellular activation. In response to inflammatory stimuli such as infection, tissue injury, or dietary factors, resident immune cells including **macrophages and T-cells** are activated and release early response cytokines including TNF-α [12]. TNF-α then stimulates further production of IL-6 from various cell types [12].

IL-6 circulates to the liver where it triggers hepatocytes to produce and release **acute-phase proteins** including CRP [12]. This cascade results in the characteristic temporal pattern of biomarker appearance: TNF-α and IL-6 levels peak within 90-120 minutes after stimulation, while CRP levels rise more slowly, peaking at approximately 48 hours [14] [13]. The half-lives of these molecules also differ significantly, with cytokines having short half-lives (minutes to hours) while CRP has a longer half-life of approximately 19 hours [13].

Understanding this sequential relationship is crucial for interpreting biomarker measurements in research settings, as different biomarkers reflect different stages and aspects of the inflammatory response. The position of IL-6 as a central mediator connecting early cytokine responses with downstream acute-phase protein production makes it particularly valuable for capturing comprehensive inflammatory information.

### Dietary Inflammatory Index Validation Workflow




*Figure 2: Dietary Inflammatory Index Calculation and Validation Workflow. This diagram outlines the systematic process for calculating DII scores from dietary data and validating them against inflammatory biomarkers.*

The Dietary Inflammatory Index calculation follows a standardized multi-step process. Dietary intake data is first collected using validated instruments such as **food frequency questionnaires (FFQ)** or 24-hour recalls [4] [18]. The reported intake for each dietary parameter is then converted to a z-score by comparing to a global reference database that provides representative means and standard deviations [18].

These z-scores are transformed to proportions (0-1) to minimize right skewing, then converted to a symmetrical distribution centered on zero with a range from -1 to +1 [18]. The resulting values are multiplied by **food parameter-specific inflammatory effect scores** derived from systematic literature review [4] [18]. Finally, all food parameter-specific DII scores are summed to generate the overall DII score for each individual [18].

Validation of the DII involves testing its correlation with established inflammatory biomarkers in population studies. The index has demonstrated consistent associations with CRP, IL-6, and other inflammatory markers across diverse populations [4] [18], supporting its use as a standardized tool for assessing the inflammatory potential of diets in research settings.

CRP, IL-6, and TNF-α represent core inflammatory biomarkers with distinct but complementary roles in validating dietary inflammatory potential in research contexts. IL-6 demonstrates particular utility as both a **sensitive predictor** of clinical outcomes and a **responsive marker** to dietary influences, while providing prognostic information beyond CRP in certain clinical settings. The temporal relationship between these biomarkers, with cytokines preceding acute-phase proteins in the inflammatory cascade, necessitates careful consideration of timing in measurement protocols.

The selection of inflammatory biomarkers for research purposes should be guided by study objectives, population characteristics, and methodological considerations. Composite approaches combining multiple biomarkers may enhance predictive power, while emerging markers like GlycA offer promising alternatives reflecting integrated inflammatory information. Standardized assessment tools like the Dietary Inflammatory Index provide valuable methods for quantifying dietary inflammatory potential, particularly when validated against established inflammatory biomarkers.

As research advances, understanding the multifaceted roles of inflammatory biomarkers beyond their classical pro-inflammatory functions will enable more nuanced interpretation of findings in nutritional science. The integration of inflammatory biomarker assessment with dietary pattern analysis continues to provide important insights into the relationships between diet, inflammation, and health outcomes across diverse populations.

## The Critical Link Between Diet, Chronic Inflammation, and Disease Pathogenesis

Chronic inflammation represents a fundamental pathological process underpinning a wide spectrum of non-communicable diseases (NCDs). The Dietary Inflammatory Index (DII) has emerged as a validated tool to quantify the inflammatory potential of individual diets, providing researchers with a standardized metric for investigating nutrition-inflammation-disease pathways. This review synthesizes current evidence on the association between pro-inflammatory diets and chronic disease pathogenesis, examines the methodological framework of DII development and validation, and explores innovative technologies for dietary assessment. Evidence consistently demonstrates that higher DII scores (indicating pro-inflammatory diets) correlate with increased prevalence and severity of conditions including cardiovascular-kidney-metabolic (CKM) syndrome, coronary artery disease, and various cardiometabolic disorders. These findings validate the DII as a crucial tool for elucidating the mechanistic role of diet-driven inflammation in disease pathogenesis and for developing targeted nutritional interventions.

Chronic inflammation has become increasingly recognized as a significant contributor to the onset and progression of various non-communicable diseases (NCDs) [20]. This sustained inflammatory condition, characterized by elevated levels of pro-inflammatory markers including high-sensitivity C-reactive protein (hs-CRP), IL-6, and TNF-α, results from a dysregulated immune response and leads to tissue impairment and malfunction [20]. The mechanistic bridge connecting chronic inflammation to NCDs is well-established, with inflammation contributing to atherosclerotic plaque formation in cardiovascular disease (CVD), inducing insulin resistance in type 2 diabetes mellitus (T2DM), facilitating tumor progression in cancer, and promoting neuronal damage in cognitive disorders [20].

Dietary components are important determinants of systemic inflammation, serving as either promoters or suppressors of inflammatory pathways [21]. 'Unhealthy' dietary patterns characterized by high intake of fats, refined carbohydrates, and processed foods are typically associated with higher levels of inflammation, whereas 'healthier' diets rich in fruits, vegetables, fish, and whole grains demonstrate anti-inflammatory properties [21]. The Dietary Inflammatory Index (DII) was developed to provide a quantitative means for assessing the role of diet in relation to health outcomes based on empirical evidence from peer-reviewed literature [3]. This review explores the validation of DII scores within the context of diet-induced inflammation and its critical role in disease pathogenesis.

### The Dietary Inflammatory Index: Development and Mechanistic Basis

#### DII Development Methodology

The DII was developed through a systematic review of 1,943 peer-reviewed articles published from 1950 to 2010 that examined associations between dietary components and inflammation [22] [3]. This comprehensive approach identified 45 food parameters with robust evidence regarding their effects on inflammatory biomarkers. The index evaluates dietary components based on their impact on key inflammatory markers including IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP [22].

The current energy-adjusted DII (E-DII) calculation involves linking reported dietary intake of the parameters to global norms of intake derived from 11 datasets worldwide [23] [3]. For each dietary parameter, individual intake is standardized against these global norms to create a z-score, which is then converted to a percentile and centered. These centered percentiles are multiplied by the respective inflammatory effect scores and summed across all parameters to generate the overall DII score [23] [21]. The E-DII is normalized per 1,000 kilocalories to account for energy intake differences, enhancing comparability across populations [23]. Higher DII scores indicate pro-inflammatory diets, while lower (more negative) scores indicate anti-inflammatory potential [3].

#### Inflammatory Potential of Dietary Components

*Table 1: Anti-Inflammatory and Pro-Inflammatory Food Components and Their Mechanisms*

| Category | Key Food Sources | Anti-Inflammatory Effects | Primary Mechanisms |
|---|---|---|---|
| **ω-3 Fatty Acids** | Fatty fish (salmon, mackerel), flaxseed, chia seeds, walnuts | Inhibit leukocyte chemotaxis, reduce adhesion molecule expression, decrease pro-inflammatory eicosanoids, suppress inflammatory cytokines | Compete with ω-6 fatty acids for enzymatic pathways, reduce generation of arachidonic acid-derived eicosanoids [20] |
| **Polyphenols** | Extra virgin olive oil, tea, dark chocolate, red wine, berries, coffee | Antioxidant properties, mitigate blood pressure, lipid profiles, abdominal obesity, blood glucose | Modulate NF-κB pathway, inhibit cyclooxygenases (COX) and cytokine production, scavenge free radicals [20] |
| **Dietary Fiber** | Whole grains, fruits, vegetables, legumes, nuts, seeds | Increases anti-inflammatory short-chain fatty acids (SCFAs), reduces proinflammatory cytokines, maintains healthy gut microbiome | Fermentation by gut microbiota produces SCFAs (e.g., butyrate), which inhibit NF-κB activation and support gut barrier integrity [20] |
| **Antioxidant Vitamins** | Citrus fruits, berries, vegetables, nuts, seeds | Quench free radicals, reduce oxidative stress, protect against DNA damage | Neutralize reactive oxygen species (ROS), decrease oxidative stress underlying inflammation [20] |
| **Saturated Fats** | Red meat, processed foods, full-fat dairy | Promote pro-inflammatory state, increase inflammatory biomarkers | Activate TLR4 signaling, promote inflammation through increased LPS translocation [24] |
| **Refined Carbohydrates** | White bread, sugary drinks, processed snacks | Regulate postprandial blood glucose, increase inflammatory markers | Induce oxidative stress and postprandial inflammation, promote advanced glycation end-products [20] |

The DII incorporates the inflammatory effect scores of various dietary components, with specific values assigned based on the strength of evidence. For example, anti-inflammatory components include fiber (-0.663), β-carotene (-0.584), magnesium (-0.484), vitamin D (-0.446), and omega-3 fatty acids (-0.436), while pro-inflammatory components include saturated fatty acids (0.373), total fat (0.298), and cholesterol (0.110) [23].

#### Pathophysiological Pathways Linking Diet to Inflammation

Dietary components influence inflammatory pathways through multiple interconnected mechanisms. The diagram below illustrates key pathways through which pro-inflammatory and anti-inflammatory dietary patterns influence chronic disease pathogenesis:




The NF-κB pathway represents a central signaling cascade through which pro-inflammatory dietary components exert their effects. This transcription factor regulates the expression of numerous genes involved in inflammation, including cytokines, chemokines, and adhesion molecules [20]. Anti-inflammatory dietary components such as polyphenols from fruits, vegetables, and olive oil can inhibit NF-κB activation, thereby reducing

downstream inflammatory mediators [20]. Additionally, the gut microbiome serves as a critical interface between diet and inflammation, with dietary fiber promoting the growth of beneficial bacteria that produce anti-inflammatory short-chain fatty acids (SCFAs), while Western-style diets promote dysbiosis and increased gut permeability, facilitating translocation of inflammatory bacterial products [24].

## Validation of DII Scores: Association with Clinical Disease Outcomes

### Cardiovascular-Kidney-Metabolic (CKM) Syndrome

A recent cross-sectional study of 7,110 participants from the National Health and Nutrition Examination Survey (NHANES) demonstrated a significant association between higher E-DII scores and increased CKM syndrome prevalence (OR: 1.22, 95% CI: 1.09-1.37) [23]. The relationship exhibited linearity (p for nonlinearity = 0.464), with consistent associations across demographic and socioeconomic subgroups. Component analysis identified alcohol as the dietary factor with the strongest association with CKM syndrome [23]. CKM syndrome represents the complex interplay of metabolic risk factors, chronic kidney disease, and cardiovascular disease, affecting approximately 25% of US adults and imposing a substantial burden on healthcare systems [23].

### Coronary Artery Disease Severity

A cross-sectional study of 1,015 individuals undergoing elective angiography examined the association between DII scores and coronary artery disease (CAD) severity classified by Gensini score [22]. After adjusting for confounding factors, results indicated significantly increased severe CAD risk for higher DII quartiles, with odds ratios of 1.52 (95% CI: 1.05-2.22) and 1.48 (95% CI: 1.01-2.16) for the 3rd and 4th quartiles (most pro-inflammatory), respectively (P for trend: 0.034) compared to the 1st quartile (most anti-inflammatory) [22]. The study also investigated the neutrophil-to-lymphocyte ratio (NLR) as an inflammatory biomarker and found that NLR mediated 24.7% (95% CI: 15.2%-98.3%) of the total effect of DII on severe CAD, providing mechanistic insight into how pro-inflammatory diets influence CAD pathogenesis [22].

### Correlation with Established Dietary Quality Indices

Validation studies demonstrate that the DII correlates with established dietary quality indices while providing specific focus on inflammatory potential. Research from the Energy Balance Study showed that as DII increased (became more pro-inflammatory), scores on the Alternative Healthy Eating Index (AHEI), Healthy Eating Index-2010 (HEI-2010), and Dietary Approaches to Stop Hypertension (DASH) decreased (became more unhealthy, all $p<0.01$) [21]. This inverse relationship confirms that the DII captures meaningful aspects of dietary quality while specifically focusing on inflammatory pathways, offering advantages for research investigating inflammation-mediated diseases.

*Table 2: DII Validation Across Clinical Disease Outcomes*

| Disease Outcome | Study Design | Population | Key Findings | Effect Size (Odds Ratio/Hazard Ratio) |
|---|---|---|---|---|
| **CKM Syndrome** | Cross-sectional | 7,110 NHANES participants | Higher E-DII scores associated with increased CKM prevalence | OR: 1.22 (95% CI: 1.09-1.37) [23] |
| **Severe Coronary Artery Disease** | Cross-sectional | 1,015 angiography patients | Higher DII quartiles associated with severe CAD (Gensini score ≥60) | Q3 OR: 1.52 (95% CI: 1.05-2.22)\nQ4 OR: 1.48 (95% CI: 1.01-2.16) [22] |
| **Chronic Kidney Disease** | Cross-sectional | Various populations | Higher DII scores associated with CKD risk | 29% increased risk [23] |
| **Mortality in CKD Patients** | Prospective | CKD patients | Higher DII scores associated with mortality | All-cause: 33% increased risk\nCVD mortality: 54% increased risk [23] |
| **Metabolic Syndrome** | Prospective (5-year) | 10,138 participants | Higher DII scores associated with MetS development | 29% increased risk [23] |

## Methodological Considerations in DII Research

### Dietary Assessment Methodologies

Accurate dietary assessment presents fundamental challenges in nutritional epidemiology. Conventional methods including 24-hour dietary recalls, food frequency questionnaires (FFQs), and food diaries are limited by recall bias, measurement error, and participant burden [25]. Technological advances offer promising alternatives, with artificial intelligence (AI)-assisted dietary assessment tools emerging as innovative approaches to overcome these limitations [25].

AI-assisted tools can broadly be categorized as "image-based" and "motion sensor-based" systems [25]. Image-based tools utilize food recognition, classification, and volume/weight estimation through computer vision algorithms, while motion sensor-based tools capture eating occasions through

wrist movement, eating sounds, jaw motion, and swallowing [25]. These technologies offer advantages including reduced recall bias, real-time data capture, and objective assessment, potentially enhancing the accuracy of DII calculation in research settings [25].

The diagram below illustrates the workflow for AI-assisted dietary assessment and DII calculation:




#### DII Calculation and Energy Adjustment

The DII calculation involves several methodological considerations that impact its interpretation in research. The energy-adjusted DII (E-DII) addresses the limitation of absolute nutrient intake by standardizing dietary components to 1,000 kilocalories of food consumed [23]. This adjustment is particularly important when comparing populations with varying total energy intakes or when investigating energy-restricted diets. Additionally, researchers must determine which of the 45 possible DII parameters to include based on available dietary data, with studies demonstrating that the predictive capability remains robust when using at least 28 parameters [23].

### Experimental Protocols and Research Applications

#### Standardized DII Research Protocol

A typical protocol for investigating associations between DII and disease outcomes involves the following steps:

- **Dietary Assessment**: Collect dietary intake data using validated methods (e.g., 24-hour recalls, FFQs, or AI-assisted tools). Multiple 24-hour recalls administered by trained personnel following standardized protocols provide the most accurate data [23].
- **DII Parameter Extraction**: Identify and quantify the intake of DII parameters from dietary data. The most commonly used parameters include alcohol, carbohydrates, protein, total fat, saturated fatty acids, monounsaturated fatty acids, polyunsaturated fatty acids, omega-3 and omega-6 fatty acids, cholesterol, dietary fiber, various vitamins (A, B1, B2, B3, B6, B12, C, D, E), folic acid, iron, magnesium, selenium, zinc, and caffeine [23].
- **DII Score Calculation**: Apply the standardized DII algorithm, which involves:
  - Standardizing individual dietary intake against global reference database means and standard deviations
  - Converting z-scores to percentiles and centering
  - Multiplying by respective inflammatory effect scores
  - Summing across all parameters
  - Adjusting for energy intake (for E-DII) [23] [3]
- **Statistical Analysis**: Employ appropriate regression models (logistic, linear, or Cox proportional hazards) to examine associations between DII scores and outcomes, adjusting for relevant covariates including age, sex, BMI, physical activity, smoking status, and medical history [23] [22].
- **Biomarker Validation**: Where possible, correlate DII scores with inflammatory biomarkers (e.g., CRP, IL-6, TNF-α, NLR) to provide mechanistic validation of the dietary-inflammatory relationship [22].

#### Research Reagent Solutions for DII Studies

*Table 3: Essential Research Materials and Tools for DII and Inflammation Studies*

| Research Tool Category | Specific Examples | Application in DII Research | Key Considerations |
|---|---|---|---|
| **Dietary Assessment Platforms** | Automated Self-Administered 24-hour Recall (ASA24), Food Frequency Questionnaires (FFQs), FoodImage Analysis | Standardized collection of dietary intake data for DII parameter extraction | Validation against recovery biomarkers, cultural adaptation of food lists, portion size estimation accuracy [25] |
| **Inflammatory Biomarker Assays** | High-sensitivity CRP (hs-CRP), IL-6, TNF-α, IL-1β ELISA kits, Neutrophil-to-Lymphocyte Ratio (NLR) | Validation of DII scores against objective inflammatory measures | Sensitivity, specificity, reliability, cost-effectiveness for large-scale studies [22] |
| **Nutritional Databases** | USDA FoodData Central, Phenol-Explorer, Food Composition Tables | Conversion of food intake to nutrient values for DII calculation | Completeness of data, regular updates, compatibility with local food varieties [25] |
| **Statistical Analysis Software** | SAS, R, Stata with specialized nutritional epidemiology packages | DII score calculation and association analyses | Handling of complex survey designs, appropriate adjustment for confounding, management of missing data [21] |
| **AI-Assisted Dietary Tools** | Food image recognition apps, Wearable eating detection sensors | Enhanced accuracy of dietary data collection and reduced respondent burden | Integration with nutrient databases, validation across diverse populations and food cultures [25] |

The validation of DII scores through consistent associations with clinically relevant disease outcomes solidifies the role of diet as a critical modulator of chronic inflammation and disease pathogenesis. Future research directions should focus on several key areas: (1) refinement of DII parameters to include emerging bioactive food components; (2) investigation of gene-diet interactions influencing inflammatory responses; (3) development of culturally-specific DII adaptations for global applications; and (4) integration of AI-assisted dietary assessment tools to enhance measurement precision [25] [3].

The consistent demonstration that pro-inflammatory diets (higher DII scores) associate with increased disease risk across multiple populations and conditions provides robust validation of the DII as a research tool and underscores the fundamental role of nutrition in chronic disease prevention and management. As research continues to elucidate the complex pathways linking diet, inflammation, and disease, the DII provides a validated metric for quantifying dietary inflammatory potential and developing targeted anti-inflammatory dietary interventions for specific populations and clinical conditions.

Chronic inflammation is a well-established contributor to the pathogenesis of numerous chronic diseases, including cardiovascular diseases, diabetes, cancer, and metabolic syndrome [26] [27]. As evidence accumulated on the central role of diet in modulating inflammatory processes, the scientific community recognized the need for a standardized tool to quantify the overall inflammatory potential of an individual's diet. Prior to the development of the Dietary Inflammatory Index (DII), most dietary assessment tools fell into one of three categories: those based on dietary recommendations, those related to adherence to a specific cuisine pattern, or those derived from study-specific statistical techniques [3]. These approaches suffered from limitations, including narrow exposure variability and population-specific biases that limited their generalizability across diverse populations.

The original DII, developed in 2009, represented the first attempt to create a literature-derived, population-based dietary index specifically focused on inflammation [26] [3]. However, this initial version had several methodological limitations that hindered its widespread adoption. The subsequent revised DII, published in 2014, incorporated significant improvements and has since become an established research tool with over 200 studies and multiple meta-analyses to its credit [3]. This evolution has continued with the development of population-specific adaptations including an Energy-Adjusted DII (E-DII), a children's DII (C-DII), and other variants designed to address unique research and clinical needs [28].

## The Original DII: Foundation and Limitations

The development of the original DII began in 2004, with the first version debuting in 2009 [28]. This pioneering tool was based on a systematic review of 927 peer-reviewed articles published through 2007 that investigated the relationship between dietary parameters and six specific inflammatory biomarkers: IL-1β, IL-4, IL-6, IL-10, TNF-α, and C-reactive protein (CRP) [3]. The scoring algorithm assigned values to food parameters based on their reported effects on these biomarkers, with "+1" for pro-inflammatory effects, "-1" for anti-inflammatory effects, and "0" for no significant effect [26].

Despite its innovative approach, the original DII faced several critical limitations. The scoring system relied on raw consumption amounts, which required arbitrary arithmetic manipulations to place different nutrients on a comparable scale [3]. Additionally, the index omitted important bioactive compounds with established anti-inflammatory properties, particularly flavonoids [3]. The original scoring direction, with anti-inflammatory diets receiving positive scores and pro-inflammatory diets receiving negative scores, was also counterintuitive to many researchers

[28]. Perhaps most significantly, the original DII failed to gain traction in the scientific community, with no subsequent research studies published using this initial version by its original developers [3].

*Table 1: Key Limitations of the Original DII (2009)*

| Limitation Category | Specific Issue | Impact on Research Application |
|---|---|---|
| **Statistical Methodology** | Use of raw consumption amounts requiring arbitrary adjustments | Introduced potential bias and distortion in scoring |
| **Nutrient Coverage** | Omission of flavonoids and other important bioactive compounds | Incomplete assessment of dietary inflammatory potential |
| **Scaling Direction** | Anti-inflammatory diets scored as positive values | Counterintuitive interpretation of results |
| **Literature Base** | Limited to articles published through 2007 | Less robust evidence foundation |
| **Standardization** | No reference to global intake norms | Limited comparability across populations |

## The Revised DII (2014): Methodological Advancements

### Enhanced Literature Review and Scoring Algorithm

The revised DII, published in 2014, addressed the limitations of the original version through substantial methodological improvements [26]. The literature review was expanded to include research published through 2010, more than doubling the evidence base to 1,943 qualifying articles [26] [3]. The search strategy employed variations of inflammatory biomarker terms combined with food parameter terms using Boolean logic, with strict inclusion criteria requiring primary research on specific food parameter–inflammatory marker relationships [26].

The scoring algorithm was refined to incorporate study quality weights, with human experimental studies receiving the highest weight (10), followed by prospective cohort studies (8), case-control studies (7), cross-sectional studies (6), animal experimental studies (5), and cell culture studies (3) [26]. The food parameter-specific inflammatory effect scores were calculated by subtracting the anti-inflammatory fraction from the pro-inflammatory fraction of the weighted literature, with adjustments made for parameters with less robust literature bases [26].




### Standardization to Global Intake Norms

A critical advancement in the revised DII was the standardization of individual intakes to global referent values [26]. Researchers identified eleven population-based nutrition surveys from around the world to create a composite database representing diverse dietary patterns: Australia, Bahrain, Denmark, India, Japan, Mexico, New Zealand, South Korea, Taiwan, the United Kingdom, and the United States [26] [3]. This approach addressed the arbitrariness of using raw consumption amounts and mitigated the right-skewing commonly observed in dietary intake data [3].

The DII calculation process involves comparing an individual's intake of each food parameter to the global composite database, calculating Z-scores, and converting these to centered percentiles [26]. These values are then multiplied by the respective food parameter-specific overall inflammatory effect scores and summed to generate the overall DII score [26]. The theoretical bounds of the DII are -8.87 (maximally anti-inflammatory) to +7.98 (maximally pro-inflammatory), with a median of +0.23 in the global composite database [26].

*Table 2: Key Methodological Improvements in the Revised DII (2014)*

| Feature | Original DII (2009) | Revised DII (2014) |
|---|---|---|
| **Literature Base** | 927 articles (through 2007) | 1,943 articles (through 2010) |
| **Food Parameters** | Approximately 30 | 45 parameters, including flavonoids |
| **Scaling Reference** | Raw consumption amounts | Global composite database (11 populations) |
| **Scoring Direction** | Anti-inflammatory = positive | Anti-inflammatory = negative (intuitive) |
| **Statistical Issues** | Right-skewing concerns addressed arbitrarily | Percentile-based approach minimizes skewing effects |
| **Flavonoid Inclusion** | Not included | 16 flavonoids across 6 categories added |

## Population-Specific Adaptations and Validation

### Specialized Variants for Research and Clinical Applications

The evolution of the DII has continued with the development of specialized variants designed to address specific research questions and population needs. The Energy-Adjusted DII (E-DII) was created to account for the strong correlation between total energy intake and nutrient consumption, which can confound the interpretation of standard DII scores [3]. The children's DII (C-DII) was developed at the request of the USDA to address the unique dietary patterns and nutritional needs of pediatric populations [28]. Additionally, researchers have created simplified versions such as the empirical DII (eDII), which is based on the frequency of consumption of only 16 foods (8 pro-inflammatory and 8 anti-inflammatory) and does not require detailed nutrient intake estimation [29].

More comprehensive tools have also emerged, such as the Dietary and Lifestyle Inflammation Score (DLIS), which integrates both dietary inflammation scores and lifestyle factors including physical activity, alcohol intake, BMI, and smoking status [30]. This holistic approach recognizes that inflammation is modulated by multiple behavioral factors beyond diet alone.

### Validation Across Diverse Populations and Health Outcomes

The DII has undergone extensive validation across diverse populations and health conditions. Construct validation studies have demonstrated significant correlations between DII scores and various inflammatory markers, including interleukin 1 beta, interleukin 4, interleukin 6, interleukin 10, tumor necrosis factor TNFα-R2, C-reactive protein, and homocysteine, both individually and as combined inflammatory biomarker scores [28].

Recent research has further strengthened the validity of the DII in specific disease contexts. A 2025 cohort study of 13,751 adults with metabolic syndrome from NHANES data found that participants in the highest DII tertile had significantly increased all-cause mortality compared to those in the lowest tertile, with multivariate-adjusted hazard ratios of 1.16 even after controlling for sex, education, smoking, income, BMI, CVD, and alcohol consumption [27]. Similarly, a 2025 case-control study on polycystic ovary syndrome demonstrated that higher DII scores were significantly associated with increased odds of PCOS, with an odds ratio of 2.82 in the highest versus lowest tertile after multivariable adjustment [30].

*Table 3: Selection of DII Validation Studies Across Health Conditions*

| Health Condition | Study Design | Population | Key Finding | Citation |
|---|---|---|---|---|
| **Metabolic Syndrome Mortality** | Cohort Study | 13,751 US adults | Highest DII tertile: 16% increased all-cause mortality | [27] |
| **Polycystic Ovary Syndrome** | Case-Control | 200 Iranian women | Highest DII tertile: 2.82x odds of PCOS | [30] |
| **Cardiovascular Disease** | Various | Multiple populations | Pro-inflammatory diets increase CVD risk | [27] [3] |
| **Cancer** | Multiple studies | Various sites | DII associated with cancers of breast, colon, etc. | [3] |
| **Mental Health** | Observational studies | Diverse populations | Links between DII and depression outcomes | [3] |

## Experimental Protocols for DII Validation

### Cohort Study Methodology

The association between DII and health outcomes is typically evaluated using large-scale cohort studies with extended follow-up periods. The protocol for the NHANES-based study on metabolic syndrome mortality exemplifies this approach [27]. Researchers analyzed data from 13,751 adults with diagnosed metabolic syndrome, with DII scores computed based on 24-hour dietary recall data at baseline. The cohort was followed for a mean duration of 114 months (approximately 9.5 years), with mortality status ascertained through probabilistic matching with the National Death Index [27].

Statistical analysis employed multivariate Cox proportional hazards models, with DII treated as both continuous and categorical (tertile) variables. Models were adjusted for demographic characteristics (age, sex, race, education), socioeconomic status (family poverty-income ratio), behavioral factors (smoking status, alcohol consumption), clinical parameters (BMI, pre-existing cardiovascular disease), and laboratory values [27]. Restricted cubic spline analysis was used to examine the dose-response relationship between DII scores and mortality risk [27].

### Case-Control Study Methodology

Case-control studies provide another important methodological approach for validating the DII against specific disease outcomes. The PCOS study illustrates this protocol, with 100 women with newly diagnosed PCOS (within 3 months, confirmed by Rotterdam criteria) compared to 100 age-matched controls recruited from the same infertility center [30]. Dietary intake was assessed using a validated 147-item Food Frequency Questionnaire, and comprehensive data were collected on potential confounders including anthropometric measurements, physical activity levels, educational attainment, marital status, parity, employment status, household income, and medical history [30].

Statistical analysis employed multivariable logistic regression to calculate odds ratios for PCOS across tertiles of DII scores, with progressive adjustment for confounding variables. The models controlled for age, BMI, energy intake, physical activity, educational level, marital status, parity, employment status, household income, history of diabetes mellitus, and hypothyroidism [30]. The statistical power calculation determined that 100 women per group would provide 80% power to detect standardized differences of ≥0.4 in inflammatory indices [30].




## The Scientist's Toolkit: Essential Research Reagents and Materials

*Table 4: Essential Research Materials for DII Studies*

| Research Tool | Specifications & Function | Application Examples |
|---|---|---|
| **24-Hour Dietary Recall** | Structured interview to assess recent food intake; multiple recalls preferred for usual intake estimation | NHANES data collection; baseline dietary assessment in cohort studies [27] |
| **Food Frequency Questionnaire (FFQ)** | Comprehensive survey assessing frequency of consumption of specific foods; should be validated for target population | 147-item FFQ in PCOS study; allows calculation of nutrient intakes [30] |
| **Global Composite Database** | Means and standard deviations for 45 food parameters across 11 representative populations | Reference for standardizing individual intakes to world norms [26] [3] |
| **Inflammatory Biomarker Assays** | Standardized protocols for IL-1β, IL-4, IL-6, IL-10, TNF-α, CRP measurement | Construct validation against individual or combined inflammatory markers [26] [28] |
| **Statistical Software Packages** | Capable of complex survey design analysis, multivariate modeling, and restricted cubic splines | Analysis of weighted NHANES data; Cox proportional hazards models [27] |
| **DII Scoring Algorithm** | Published methodology for calculating food parameter-specific inflammatory effect scores | Computation of individual DII scores from dietary intake data [26] [2] |

The evolution of the Dietary Inflammatory Index from its original formulation to the current population-specific adaptations represents significant progress in nutritional epidemiology. The methodological refinements incorporated into the revised DII have addressed critical limitations regarding global standardization, flavonoid inclusion, scoring direction, and statistical handling of dietary intake data. The development of specialized variants

including the E-DII, C-DII, and integrated scores like DLIS demonstrates the tool's adaptability to diverse research needs and population characteristics.

The robust validation of the DII against hard endpoints such as all-cause and cardiovascular mortality in metabolic syndrome patients underscores its clinical relevance [27]. Similarly, its association with conditions like PCOS highlights the expanding applications of this tool across different disease domains [30]. As research continues, future directions will likely include further refinement of the scoring algorithm based on emerging evidence, development of culturally-specific adaptations for underrepresented populations, and integration with omics technologies for more personalized nutritional recommendations.

For researchers, scientists, and drug development professionals, the DII offers a validated tool for quantifying the inflammatory potential of diet in observational studies, clinical trials, and pharmacologic research. The consistent methodology enables comparison across studies and populations, while the availability of multiple variants allows selection of the most appropriate version for specific research questions. As the field advances, the DII will continue to evolve, incorporating new evidence and methodological innovations to enhance its precision and utility in both research and clinical practice.

# Measuring Inflammatory Potential: DII Calculation, Biomarker Correlation, and Clinical Integration

## Methodological Frameworks for DII Score Calculation and Energy Adjustment

The Dietary Inflammatory Index (DII) is a quantitative tool designed to assess the inflammatory potential of an individual's overall diet. Its development was driven by the need to move beyond evaluating single nutrients and to instead capture the complex, synergistic effects of dietary components on systemic inflammation. The DII framework is grounded in a comprehensive review of peer-reviewed literature linking dietary factors to specific inflammatory biomarkers, including interleukin-1β (IL-1β), IL-4, IL-6, IL-10, tumor necrosis factor-α (TNF-α), and C-reactive protein (CRP) [31] [3]. Two primary methodological frameworks have emerged for calculating DII scores: the **literature-derived DII** and the **empirically-derived EDII**. Each approach employs distinct methodologies for development and validation, offering researchers complementary tools for investigating diet-inflammation-disease pathways.

The fundamental principle underlying both frameworks is that dietary patterns can be systematically classified along an inflammation spectrum. Pro-inflammatory diets are characterized by components that upregulate inflammatory mediators, while anti-inflammatory diets contain factors that suppress these biochemical pathways. The DII was conceptually designed to be universally applicable across diverse human populations with adequate dietary assessment methods, enabling comparative research on a global scale [3]. The evolution of these frameworks represents a significant advancement in nutritional epidemiology, bridging hypothesis-driven dietary assessment with empirical biomarker validation.

### Literature-Derived DII Framework

#### Development and Calculation Methodology

The literature-derived DII framework, the more widely implemented approach, was developed through systematic analysis of research articles examining relationships between dietary factors and inflammatory biomarkers. The current DII reflects analysis of 1,943 qualifying articles published from 1950 to 2010, nearly double the literature base used for the original index [3] [22]. This expansive evidence base encompasses human studies ranging from cell culture experiments to observational and clinical trials, with higher weighting given to human studies compared to animal or in vitro research [3].

The DII calculation incorporates up to 45 food parameters, including nutrients, bioactive compounds, and specific food items with established inflammatory effects. The scoring algorithm connects individual dietary intake to global normative intake data derived from 11 population-based surveys from countries worldwide including the United States, Australia, Japan, Mexico, South Korea, and several European nations [3]. This global reference database provides means and standard deviations for each parameter, enabling standardized calculation across different populations.

The computational process for deriving literature-based DII scores involves multiple transformation steps:

- **Z-score Calculation**: Individual intake of each food parameter is compared to the global mean using the formula: Z-score = (individual mean intake - global mean intake) / global standard deviation [32] [33]
- **Percentile Conversion**: The Z-score is converted to a centered percentile score to minimize the effect of right-skewing common in dietary intake data [3]
- **Inflammatory Effect Scoring**: Each centered percentile value is multiplied by the respective food parameter's inflammatory effect score derived from the literature review [31] [32]
- **Index Summation**: The scores for all food parameters are summed to generate the overall DII score, where positive values indicate pro-inflammatory potential and negative values indicate anti-inflammatory potential [3] [28]

*Table 1: Key Components of Literature-Derived DII Framework*

| Component | Description | Source/Validation |
|---|---|---|
| **Evidence Base** | 1,943 articles (1950-2010) linking diet to inflammation | Peer-reviewed literature [3] [22] |
| **Food Parameters** | Up to 45 parameters (nutrients, flavonoids, foods) | Literature on inflammatory biomarkers [3] |
| **Reference Database** | Means/SDs from 11 countries worldwide | Population surveys [3] |
| **Inflammatory Biomarkers** | IL-1β, IL-4, IL-6, IL-10, TNF-α, CRP | Association with dietary components [31] [5] |
| **Score Interpretation** | Positive (pro-inflammatory), Negative (anti-inflammatory) | Orientation consistent with inflammatory effect [3] |

#### Energy Adjustment Methods: E-DII and IE-DII

A critical methodological consideration in nutritional epidemiology is accounting for variations in total energy intake, which can confound associations between dietary components and health outcomes. The literature-derived DII framework incorporates two primary approaches for energy adjustment:

The **Energy-Adjusted DII (E-DII)** calculates dietary parameters as amounts per 1,000 calories consumed rather than absolute intake amounts [32] [34]. This approach effectively standardizes dietary patterns to a consistent energy basis, allowing comparison of diet quality independent of quantity. The E-DII calculation follows the same transformation steps as the standard DII but uses energy-density values for each parameter.

Alternatively, some studies employ the **Inversely Energy-Adjusted DII (IE-DII)**, which incorporates total energy intake as one of the pro-inflammatory parameters within the DII scoring system [34]. This method recognizes that excessive caloric intake itself may promote inflammation, particularly when associated with obesity and metabolic dysfunction.

Research comparing these approaches has demonstrated that energy adjustment strengthens the association between DII scores and inflammation-related outcomes. For example, a cross-sectional study of US adults found that the IE-DII showed stronger inverse associations with nonalcoholic fatty liver disease (NAFLD) risk compared to other dietary indices [34]. Similarly, studies on attention deficit hyperactivity disorder (ADHD) in children revealed that E-DII remained significantly associated with increased ADHD risk after adjusting for multiple confounders [32].

### Empirical Dietary Inflammatory Index (EDII) Framework

#### Development Through Reduced Rank Regression

The Empirical Dietary Inflammatory Index (EDII) represents an alternative, hypothesis-driven framework that uses statistical methods to derive dietary patterns most predictive of specific inflammatory biomarkers. Developed using data from the Nurses' Health Study, the EDII employs reduced rank regression (RRR) followed by stepwise linear regression to identify a dietary pattern that explains maximal variation in plasma inflammatory markers [5].

Unlike the literature-derived approach, which is based on previously published evidence, the EDII is derived empirically from population-specific data. The RRR method determines linear functions of predictors (food groups) by maximizing the explained variation in response variables (inflammatory markers) [5]. This data-driven approach identifies combinations of food groups that collectively influence inflammation, potentially capturing synergistic effects that might be missed when examining individual dietary components.

The EDII development process involved several key steps:

- **Biomarker Selection**: Three plasma inflammatory markers (IL-6, CRP, and TNFαR2) served as response variables in the RRR models [5]
- **Food Group Input**: 39 pre-defined food groups were included as predictor variables [5]
- **Pattern Derivation**: RRR identified the linear combination of food groups that explained the maximum variation in the three inflammatory biomarkers [5]
- **Simplification**: Stepwise linear regression reduced the food groups to the most predictive set, resulting in the final EDII comprising 18 food groups (9 pro-inflammatory and 9 anti-inflammatory) [5]
- **Validation**: The derived EDII was tested in two independent cohorts (NHS-II and Health Professionals Follow-up Study) for its ability to predict inflammatory biomarkers including IL-6, CRP, TNFαR2, adiponectin, and an overall inflammatory score [5]

*Table 2: Comparison of DII Methodological Frameworks*

| Characteristic | Literature-Derived DII | Empirical DII (EDII) |
|---|---|---|
| **Development Approach** | Literature review of 1,943 articles | Statistical derivation from cohort data |
| **Statistical Method** | Scoring algorithm based on published evidence | Reduced rank regression (RRR) |
| **Dietary Components** | 45 food parameters (nutrients, flavonoids) | 18 food groups |
| **Inflammatory Basis** | 6 inflammatory biomarkers from literature | 3 plasma inflammatory markers |
| **Primary Validation** | Correlation with inflammatory biomarkers | Predictive performance in independent cohorts |
| **Energy Adjustment** | E-DII, IE-DII | Included in RRR model development |
| **Key Advantage** | Standardized application across populations | Data-driven, captures food synergies |

#### Validation and Performance

The EDII framework has demonstrated strong construct validity in independent samples. In validation studies, the EDII significantly predicted concentrations of all inflammatory biomarkers in both the Nurses' Health Study II and the Health Professionals Follow-up Study [5]. For example, comparing extreme EDII quintiles in NHS-II revealed a relative concentration of 1.52 (95% CI: 1.18, 1.97) for CRP and 0.88 (95% CI: 0.80, 0.96) for adiponectin [5]. Corresponding associations in HPFS were 1.23 (95% CI: 1.09, 1.40) for CRP and 0.87 (95% CI: 0.82, 0.92) for adiponectin [5].

The empirical approach offers the advantage of accounting for the complex interplay between food groups and their collective influence on inflammatory pathways. However, a potential limitation is that the derived food patterns may be specific to the population in which they were developed, potentially limiting generalizability across diverse cultural and dietary contexts.

### Comparative Applications in Clinical Research

#### Predictive Performance Across Health Outcomes

Both DII frameworks have been extensively applied in observational studies and clinical research, demonstrating significant associations with various inflammation-related conditions. The literature-derived DII has shown particularly strong predictive performance in large-scale epidemiological studies.

In cardiovascular research, a cross-sectional study of 1,015 individuals undergoing elective angiography found that participants in the highest DII quartile (most pro-inflammatory) had significantly increased risk of severe coronary artery disease (OR: 1.48, 95% CI: 1.01-2.16) compared to the lowest quartile, after adjusting for multiple confounders [22]. Importantly, this study also demonstrated that the neutrophil-to-lymphocyte ratio (NLR), a systemic inflammatory marker, mediated 24.7% of the effect of DII on severe CAD, providing mechanistic support for the DII-inflammation-disease pathway [22].

In neuropsychiatric research, a case-control study of Iranian children found that each unit increase in E-DII was associated with 13.3-16.2% higher odds of ADHD across different adjustment models [32]. Similarly, research on cardiovascular-kidney-metabolic (CKM) syndrome in US adults demonstrated a J-shaped relationship between DII and depressive symptoms, with each unit increase in DII associated with an 18.7% higher incidence of depressive symptoms after adjusting for 20 potential confounders [31].

Comparative studies directly assessing the predictive validity of different DII frameworks are limited. However, one study comparing the literature-derived DII with the dietary total antioxidant capacity (TAC) index for predicting NAFLD found that the IE-DII showed stronger associations with hepatic steatosis measures than TAC [34]. Specifically, each standard deviation increase in IE-DII was associated with greater reductions in hepatic steatosis index values than TAC ($\beta$ = -0.39 vs. -0.25, P-difference = 0.036) [34].

#### Methodological Considerations and Implementation Protocols

Implementation of either DII framework requires careful attention to dietary assessment methods, parameter availability, and population characteristics. The research toolkit for DII calculation includes several essential components:

*Table 3: Research Reagent Solutions for DII Implementation*

| Research Tool | Function | Implementation Considerations |
|---|---|---|
| **FFQ (Food Frequency Questionnaire)** | Assesses habitual dietary intake | Should be validated for specific population; 106-168 items typical [32] [33] |
| **24-hour Dietary Recall** | Captures recent detailed intake | Used in NHANES; requires multiple administrations [31] [34] |
| **Global Reference Database** | Provides normative intake values | Means/SDs from 11 countries; essential for z-score calculation [3] |

| Research Tool | Function | Implementation Considerations |
|---|---|---|
| **Inflammatory Effect Scores** | Weights for food parameters | Derived from literature review; fixed for all populations [3] |
| **Nutrition Analysis Software** | Converts foods to nutrients | Nutritionist IV, USDA database, local food composition tables [32] [33] |
| **Biomarker Assay Kits** | Validate against inflammatory markers | CRP, IL-6, TNF-α ELISA kits; quality control samples essential [5] [22] |

The dietary assessment methodology significantly influences DII calculation. Most studies utilize either food frequency questionnaires (FFQs) or 24-hour dietary recalls, each with distinct advantages and limitations. FFQs typically include 106-168 food items and assess habitual intake over the previous year [32] [33], while 24-hour recalls provide more detailed recent intake data but require multiple administrations to capture usual intake [31]. The choice of assessment method should align with research objectives and population characteristics.

When implementing DII frameworks across different populations, researchers must often adapt to limited parameter availability. The literature-derived DII remains valid with fewer than 30 food parameters [31], though this necessarily reduces comprehensiveness. Studies have successfully computed DII scores with varying numbers of parameters, from 28 in NHANES analyses [31] to 37 in Korean research [33]. Transparency in reporting which parameters were included is essential for interpreting and comparing results across studies.

**Visualizing Methodological Frameworks**




*DII Methodological Framework Comparison*




*DII Calculation Workflow*

The methodological frameworks for DII calculation and energy adjustment provide researchers with robust, validated tools for quantifying the inflammatory potential of dietary patterns. The literature-derived DII offers standardized application across diverse populations through its global reference database and evidence-based scoring algorithm, while the empirical EDII captures population-specific dietary synergies through data-driven statistical approaches. Energy adjustment methods, particularly E-DII and IE-DII, enhance the specificity of these indices by accounting for variations in total energy intake.

The consistent associations observed between higher (more pro-inflammatory) DII scores and increased risk of various chronic conditions across multiple populations support the construct validity of both frameworks. However, methodological choices regarding dietary assessment, parameter inclusion, and energy adjustment should be carefully considered and transparently reported to ensure comparability across studies. Future methodological developments would benefit from continued refinement of global reference databases, expansion to include emerging dietary components with inflammatory properties, and standardization of energy adjustment approaches to facilitate cross-study comparisons.

The Dietary Inflammatory Index (DII) was developed as a literature-derived tool to quantify the inflammatory potential of an individual's diet. As chronic low-grade inflammation is a key driver of many non-communicable diseases, the ability to accurately measure and quantify the inflammatory impact of diet has significant implications for preventive healthcare and clinical research. This review synthesizes large-scale epidemiological evidence validating the DII against established inflammatory biomarkers, providing researchers and clinicians with a critical evaluation of its performance across diverse populations and contexts.

## DII and Comparative Dietary Indexes: Design and Methodologies

The DII was developed through a systematic review of nearly 2,000 research articles published between 1950 and 2010 investigating the relationship between dietary factors and inflammatory biomarkers. Each of the 45 dietary parameters in the DII was assigned a weight based on its effect on six

specific inflammatory markers: IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP. Higher DII scores indicate more pro-inflammatory diets, while lower scores suggest anti-inflammatory potential [35] [36].

Several derivatives and alternative indexes have emerged since the development of the original DII. The Energy-Adjusted DII (E-DII) adjusts for total energy intake using either the density or residual method. The Empirical Dietary Inflammatory Pattern (EDIP) represents a distinct approach, derived using reduced rank regression to identify food patterns most predictive of inflammatory biomarkers rather than relying on prior literature [37] [38].

*Table 1: Key Dietary Inflammatory Indexes and Their Characteristics*

| Index Name | Development Approach | Components | Key Biomarkers in Development | Strengths |
|---|---|---|---|---|
| Original DII | Literature review (1943 articles) | 45 parameters (mainly nutrients) | IL-1β, IL-4, IL-6, IL-10, TNF-α, CRP | Comprehensive literature basis |
| E-DII | DII with energy adjustment | Same as DII with energy adjustment | Same as DII | Reduces energy intake confounding |
| EDIP | Data-driven, hypothesis-oriented | 18 food groups | CRP, IL-6, TNFαR2 | Derived from actual biomarker data |
| ISD | Not specified | Not specified | Multiple inflammatory biomarkers | Used in European cohorts |

The fundamental difference between these approaches lies in their foundational methodology: the DII is an *a priori* index based on existing scientific evidence, while the EDIP is an *a posteriori* index derived from statistical relationships within dietary and biomarker data [37].

## Large-Scale Epidemiological Validation Studies

### Validation in European and American Cohorts

The European Prospective Investigation into Cancer and Nutrition (EPIC) study, one of the largest cohort studies on diet and health, provided substantial validation evidence for dietary inflammatory scores. Research involving 17,637 EPIC participants found that four different dietary inflammatory scores (DII, E-DII, E-DIIr, and ISD) showed consistent positive associations with circulating levels of CRP, IL-6, sTNFR1, sTNFR2, and leptin. The study demonstrated that despite methodological differences, all scores captured similar inflammatory potential of diet in European adults [38].

In the Nurses' Health Study II (n=5,826 women) and the Health Professionals Follow-Up Study (n=5,227 men), both DII and EDIP scores were significantly associated with multiple inflammatory biomarkers. Participants in the highest quintile of DII scores showed substantially elevated inflammatory markers compared to those in the lowest quintile: CRP was 49% higher in women and 29% higher in men, while IL-6 was 21% higher in women and 24% higher in men. The EDIP generally showed stronger predictive ability for inflammatory biomarkers than the DII in these cohorts [37].

### Validation in Specific Populations and Health Conditions

Recent research has extended DII validation to specialized populations and disease contexts. A 2023 study of 500 COVID-19 patients found significant associations between higher DII scores and elevated inflammatory markers. Each unit increase in DII was associated with a 1.024 mg/L increase in CRP, higher white blood cell counts (β=0.486), and increased neutrophil-to-lymphocyte ratio (β=0.538). Notably, higher DII scores also predicted longer hospitalization, demonstrating the clinical relevance of dietary inflammatory potential in disease outcomes [39].

In a study of women with cognitive impairment, those with dementia and mild cognitive impairment showed significantly altered inflammatory biomarkers and lipid metabolism parameters, supporting the relevance of inflammatory pathways in cognitive decline and suggesting potential applications for dietary interventions targeting inflammation [40].

Research in adolescents from the LabMed Physical Activity Study (n=329) demonstrated that the DII successfully predicted low-grade inflammation even in younger populations. The DII was positively associated with IL-6 and complement component C4 after full adjustment for biological and lifestyle variables [41].

*Table 2: DII Validation Across Diverse Populations and Health Conditions*

| Population/Context | Sample Size | Key Validated Biomarkers | Noteworthy Findings |
|---|---|---|---|
| General European Adults (EPIC) | 17,637 | CRP, IL-6, sTNFR1, sTNFR2, leptin | Consistent performance across 4 different dietary inflammatory scores |
| US Health Professionals | 11,053 (combined) | CRP, IL-6, TNFαR2, adiponectin | EDIP showed stronger predictive ability than DII for most biomarkers |
| COVID-19 Patients | 500 | CRP, WBC, neutrophils, NLR | DII predicted inflammation and longer hospitalization |
| Polish Adults (PURE) | 1,791 | Triglycerides, glucose, atherogenic indices | Pro-inflammatory diets associated with unfavorable CVD risk profiles |
| Adolescents | 329 | IL-6, C4 | DII applicable to younger populations |
| Adults with Obesity | 124 | CRP, BMI | DII correlated with both inflammation and adiposity |

## Comparative Performance of Dietary Inflammatory Indexes

### Predictive Performance for Specific Inflammatory Biomarkers

The comparative performance of different dietary inflammatory indexes has been systematically evaluated in large cohort studies. In the EPIC cohort, all four dietary inflammatory scores (DII, E-DII, E-DIIr, and ISD) showed consistent positive associations with CRP, IL-6, sTNFR1, sTNFR2, and leptin. However, only the DII and ISD were positively associated with IL-1RA levels, and only the DII and E-DIIr were associated with TNFα, suggesting subtle but potentially important differences in the inflammatory pathways captured by each index [38].

A direct comparison between EDIP and DII in US cohorts revealed that while both indexes significantly predicted inflammatory biomarkers, the EDIP generally demonstrated stronger associations. For CRP, the percentage difference between extreme quintiles was 60% for EDIP versus 49% for DII in women, and 38% versus 29% in men. Similarly, for adiponectin (an anti-inflammatory adipokine), the EDIP showed stronger inverse associations (-21% in women and -16% in men) compared to DII (-14% in women and -4% in men) [37].

### Methodological Considerations and Limitations

Despite generally positive validation results, important limitations have been noted across studies. The EPIC cohort analysis revealed that the proportion of variance in inflammatory biomarkers explained by dietary inflammatory scores was generally low (<2%), equivalent to that explained by smoking status but much lower than the variance explained by BMI [38]. This suggests that while diet contributes meaningfully to inflammatory status, it represents just one component among multiple determinants.

A 2025 scoping review of food-based dietary indexes identified 43 different indexes used to assess inflammation-related diet quality, categorized into four groups: dietary patterns, dietary guidelines, dietary inflammatory potential, and therapeutic diets. The review noted that indexes based on Mediterranean diet patterns and dietary guidelines were most extensively utilized and consistently demonstrated inverse associations with inflammatory biomarkers across diverse populations [42].

## Experimental Protocols and Methodological Standards

### Standard Protocols for DII Calculation and Validation

The standard approach for calculating DII scores begins with the collection of dietary intake data, typically using Food Frequency Questionnaires (FFQs), 24-hour recalls, or food diaries. In the EPIC-Norfolk study, researchers used a semi-quantitative FFQ consisting of 130 food items to capture average daily intakes over the previous year. Nutrient intakes were calculated using the FETA software, and outliers in energy intake were identified using the ratio of energy intake to basal metabolic rate [35].

For DII calculation, intake of each dietary component is compared to a standard global mean intake and converted to a z-score. This z-score is then converted to a percentile value, centered by doubling and subtracting one, and multiplied by the respective food parameter effect score from the literature. The overall DII is the sum of all these values [43]. For the energy-adjusted DII (E-DII), food and nutrient intakes are adjusted for total energy intake using either the density method or residual method [44] [38].

### Biomarker Assessment Protocols

High-quality validation studies employ standardized protocols for biomarker assessment. In major cohort studies, inflammatory biomarkers are typically measured from blood samples collected after an overnight fast. CRP is commonly measured using high-sensitivity immunoturbidimetric assays or high-sensitivity ELISA methods. Cytokines including IL-6, IL-10, and TNF-α are typically quantified using ELISA kits, with strict quality control procedures including blinded quality-control samples interspersed with participant samples to assess inter-assay variability [37] [38].




*Diagram 1: DII Validation Workflow*

### The Researcher's Toolkit: Essential Methodological Components

*Table 3: Essential Research Reagents and Tools for DII Validation Studies*

| Tool/Reagent Category | Specific Examples | Research Function | Technical Considerations |
|---|---|---|---|
| Dietary Assessment Tools | FFQ, 24-hour recall, food diaries | Quantify dietary intake | FFQ should be validated for specific population; multiple recalls improve accuracy |
| Biomarker Assay Kits | High-sensitivity CRP ELISA, Multiplex cytokine panels | Measure inflammatory biomarkers | High-sensitivity assays needed for detecting low-grade inflammation |
| Dietary Analysis Software | FETA, CAFÉ, BeBIS | Convert food intake to nutrient data | Should use country-specific food composition databases |
| Laboratory Equipment | Spectrophotometers, automated analyzers | Process and analyze biological samples | Standardized protocols essential for comparability across studies |
| Statistical Software | R, SPSS, SAS | Perform statistical analyses | Should account for covariates and potential confounders |

The substantial body of large-scale epidemiological evidence supports the validity of the DII as a tool for assessing the inflammatory potential of diet across diverse populations and health contexts. The consistent associations observed between higher DII scores and elevated inflammatory biomarkers including CRP, IL-6, and TNF-α receptors provide robust evidence for its criterion validity. However, researchers should note that alternative indexes such as the EDIP may demonstrate stronger predictive performance for certain inflammatory biomarkers, and the proportion of inflammation variance explained by dietary indexes alone remains modest compared to other factors like adiposity. Future research directions should include further validation in diverse ethnic populations, investigation of dose-response relationships, and exploration of DII's predictive value for specific inflammation-related disease outcomes.

## DII as a Predictive Tool for Chronic Disease Risk and Comorbidity

Chronic inflammation represents a fundamental biological process underpinning the development and progression of numerous chronic diseases. The **Dietary Inflammatory Index (DII)** was developed as a quantitative tool to assess the inflammatory potential of an individual's diet, providing researchers with a standardized method to evaluate relationships between dietary patterns and health outcomes. Unlike dietary indices based on specific cuisines or dietary guidelines, the DII was derived from extensive scientific literature on the effects of dietary components on inflammatory biomarkers [3]. This literature-derived, population-based index has emerged as a valuable tool for investigating the role of diet-induced inflammation in chronic disease risk and comorbidity patterns, particularly for complex conditions involving cardiovascular, metabolic, and renal systems.

The theoretical foundation of the DII rests on the understanding that dietary components can systematically modulate inflammatory pathways. Diets rich in proinflammatory elements can promote elevated levels of inflammatory markers such as C-reactive protein (CRP), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNF-α), creating a physiological environment conducive to chronic disease development [45]. The DII quantifies this inflammatory potential, with scores ranging from -8.87 (maximally anti-inflammatory) to +7.98 (maximally pro-inflammatory) [28]. This scoring system enables researchers to classify study participants according to the inflammatory potential of their diets and investigate associations with disease outcomes while controlling for relevant confounders.

## DII and Alternative Indices: Comparative Methodologies

### Development and Validation of the DII

The DII was developed through a systematic review of peer-reviewed literature published through 2010, encompassing 1,943 qualifying articles that documented associations between 45 dietary parameters and six inflammatory biomarkers: IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP [3] [46]. The current DII represents an improvement over an earlier version debuted in 2009, with methodologic enhancements including linking reported dietary intake to global norms of intake rather than relying on raw consumption amounts, the addition of flavonoids as important modulators of inflammation, and inversion of the scoring system such that anti-inflammatory scores are negative and proinflammatory scores are positive [3].

The computation of DII scores involves multiple steps. Dietary intake data from 24-hour recalls or food frequency questionnaires are first compared to a global reference database representing usual intake distributions across 11 populations worldwide [3]. This comparison yields a z-score for each food parameter, which is then converted to a percentile score and centered by doubling and subtracting 1. These values are then multiplied by the respective inflammatory effect score for each food parameter derived from the literature review, and summed to create an overall DII score [45] [31]. The DII has undergone extensive construct validation, demonstrating correlations with various inflammatory markers including CRP, IL-6, TNF-α receptor 2 (TNFαR2), and homocysteine [28].

### Alternative Dietary Inflammatory Assessment Tools

While the DII represents the most widely used tool for assessing dietary inflammatory potential, alternative approaches have been developed. The **Empirical Dietary Inflammatory Pattern (EDIP)** score was derived using reduced rank regression followed by stepwise linear regression to identify dietary patterns most predictive of plasma inflammatory markers (IL-6, CRP, TNFαR2) [5]. Unlike the nutrient-based DII, the EDIP is exclusively based on food groups, with weights derived from regression analyses rather than literature review [37].

*Table 1: Comparison of Dietary Inflammatory Assessment Indices*

| Characteristic | Dietary Inflammatory Index (DII) | Empirical Dietary Inflammatory Pattern (EDIP) |
|---|---|---|
| **Derivation Approach** | Literature-derived, a priori | Data-driven, hypothesis-oriented |
| **Components** | 45 parameters (mainly nutrients) | 18 food groups (9 pro-inflammatory, 9 anti-inflammatory) |
| **Basis for Weights** | Peer-reviewed literature | Regression analyses in cohort studies |
| **Influenced by Supplement Use** | Yes | No |
| **Validation** | Multiple populations globally | US health professional cohorts |
| **Score Interpretation** | Higher scores = more pro-inflammatory | Higher scores = more pro-inflammatory |

Comparative studies have examined the predictive ability of these indices. Research in the Nurses' Health Study II and Health Professionals Follow-Up Study found modest correlations between EDIP and DII scores ($r = 0.29$ for women, $r = 0.21$ for men) [37]. Both indices significantly predicted concentrations of inflammatory biomarkers, though the EDIP showed a greater ability to predict plasma inflammatory markers, possibly due to its development based specifically on circulating inflammatory markers [37].

## DII as a Predictor of Specific Chronic Diseases

### Chronic Kidney Disease (CKD)

Evidence from meta-analyses demonstrates a significant association between DII scores and chronic kidney disease. A 2024 meta-analysis of 13 cross-sectional studies found that higher DII scores were associated with significantly higher odds of both CKD (OR: 1.36, 95% CI: 1.20–1.56, $p < 0.001$) and low estimated glomerular filtration rate (low-eGFR) (OR: 1.58, 95% CI: 1.26–2.00, $p = 0.001$) [47]. This suggests that individuals adhering to pro-inflammatory diets have approximately 36% higher odds of having CKD and 58% higher odds of reduced kidney function.

The biological mechanisms linking pro-inflammatory diets to kidney damage involve several pathways. Systemic or intrarenal inflammation may disrupt microvascular response to regulatory factors and promote production of reactive oxygen species and other tubular toxins [47]. This process causes tubular injury, nephron dropout, and ultimately leads to CKD onset. Additionally, elevated levels of systemic inflammatory markers are associated with reduction in glomerular filtration rate and increased urinary protein levels [47].

### Diabetes Mellitus

A 2021 systematic review and meta-analysis of five studies involving 14,987 participants and 1,366 diabetes cases found that individuals with higher DII scores had a 32% higher risk of diabetes mellitus (pooled OR: 1.32, 95% CI: 1.01–1.72) compared to those with lower DII scores [46]. This association remained significant despite moderate heterogeneity among studies ($I^2 = 58.6\%$).

The inflammatory pathways linking diet to diabetes development involve multiple mechanisms. Chronic inflammation contributes to insulin resistance and β-cell dysfunction, key pathophysiological elements in type 2 diabetes. Proinflammatory cytokines such as IL-6 and TNF-α interfere with insulin signaling pathways, while also promoting oxidative stress and endothelial dysfunction [46]. The consistency of findings across different populations, despite variations in study design and dietary assessment methods, strengthens the evidence for DII as a predictor of diabetes risk.

#### Cardiovascular-Kidney-Metabolic (CKM) Syndrome

Recent research has examined the association between DII and multisystem conditions such as cardiovascular-kidney-metabolic (CKM) syndrome, which describes the pathophysiological interplay among obesity, diabetes, chronic kidney disease, and cardiovascular disease [45] [31]. A large cross-sectional study using NHANES data from 2001 to 2020 (n = 24,071) found a significant positive association between DII and CKM syndrome risk [45].

After adjusting for confounders, compared to the lowest DII quartile, the adjusted odds ratios for higher DII quartiles were 1.17 (95% CI: 0.93–1.47), 1.43 (95% CI: 1.13–1.81), and 1.76 (95% CI: 1.42–2.18), respectively [45]. Each one-unit increase in DII was associated with a 12% greater risk of developing CKM syndrome (OR: 1.12, 95% CI: 1.08–1.18). Restricted cubic spline regression indicated a significant nonlinear positive association, suggesting that the risk increase may be more pronounced at higher DII scores [45].

*Table 2: DII Association Effect Sizes Across Chronic Diseases*

| Disease Outcome | Study Design | Effect Size (Highest vs. Lowest DII) | References |
|---|---|---|---|
| **Chronic Kidney Disease** | Meta-analysis (13 cross-sectional studies) | OR: 1.36 (95% CI: 1.20–1.56) | [47] |
| **Low eGFR** | Meta-analysis (13 cross-sectional studies) | OR: 1.58 (95% CI: 1.26–2.00) | [47] |
| **Diabetes Mellitus** | Meta-analysis (5 studies) | OR: 1.32 (95% CI: 1.01–1.72) | [46] |
| **CKM Syndrome** | Cross-sectional (NHANES) | OR: 1.76 (95% CI: 1.42–2.18) for highest quartile | [45] |
| **Depressive Symptoms in CKM** | Cross-sectional (NHANES) | 18.7% higher incidence per unit DII increase | [31] |

The relationship between DII and CKM syndrome appears to be mediated through multiple pathways. Research has shown that metabolic syndrome components mediate approximately 4.44% of the effect between DII and depressive symptoms in CKM patients, suggesting both direct and indirect pathways link dietary inflammation to CKM-related outcomes [31].

### Experimental Protocols for DII Research

#### Standard DII Calculation Methodology

The protocol for calculating DII scores follows a standardized approach that can be adapted to different dietary assessment methods:

- **Dietary Assessment**: Collect dietary intake data using either 24-hour dietary recalls or food frequency questionnaires (FFQs). The 24-hour recall method requires participants to recall and report types, quantities, and consumption times of all foods and beverages consumed in the previous 24 hours [45].
- **Parameter Selection**: Identify available food parameters from the total 45 DII parameters. The DII remains valid even when fewer than 30 food parameters are available [45] [31]. Commonly included parameters are energy, protein, carbohydrates, dietary fiber, total fat, saturated fat, monounsaturated fatty acids, polyunsaturated fatty acids, cholesterol, β-carotene, various vitamins (A, B1, B2, niacin, B6, folate, B12, C, D, E), minerals (magnesium, iron, zinc, selenium), and caffeine and alcohol.
- **Global Standard Comparison**: Compare individual intake values to a global standard database representing mean intake and standard deviation for each parameter across multiple populations worldwide [3].
- **Z-score Calculation**: For each food parameter, calculate a z-score by subtracting the global mean from the individual's intake and dividing by the global standard deviation: Z = (individual intake - global mean)/global standard deviation [31].
- **Percentile Conversion**: Convert z-scores to percentiles to minimize the effect of right skewness common in dietary data.
- **Centering**: Multiply each percentile by 2 and subtract 1 to achieve a symmetrical distribution centered approximately at zero.
- **Inflammatory Effect Scoring**: Multiply each centered value by the respective inflammatory effect score derived from the literature review.
- **Summation**: Sum all food parameter scores to obtain the overall DII score for each individual [45] [31].

#### DII Validation Study Design

Protocol for validating DII scores against inflammatory biomarkers:

- **Participant Selection**: Recruit participants from the target population, ensuring representation of relevant demographic characteristics (age, sex, race/ethnicity). Exclusion criteria typically include conditions that acutely affect inflammatory markers (recent infection, cancer diagnosis, autoimmune diseases) [37] [5].
- **Blood Collection**: Collect blood samples following standardized protocols. For the Nurses' Health Studies and Health Professionals Follow-Up Study, blood was collected in EDTA tubes and processed within 24 hours, with plasma stored in liquid nitrogen freezers at -130°C or lower [37] [5].
- **Inflammatory Marker Assessment**: Measure established inflammatory biomarkers including:
  - **High-sensitivity CRP**: Measured using immunoturbidimetric assays
  - **IL-6**: Measured using ELISA kits
  - **TNF-α receptor 2**: Measured using ELISA kits
  - **Adiponectin**: Measured using radioimmunoassay [37] [5]
- **Quality Control**: Include blinded quality control samples interspersed with participant samples. Batch correction should be performed to adjust for potential batch variability [37].
- **Statistical Analysis**: Use multivariable-adjusted linear regression models to examine associations between DII scores and inflammatory biomarker concentrations, adjusting for potential confounders such as age, BMI, physical activity, smoking status, and medication use [37] [5].




*DII and Disease Pathophysiology: This diagram illustrates the proposed biological pathways linking pro-inflammatory and anti-inflammatory diets to chronic disease development through multiple interconnected mechanisms.*

**Research Reagent Solutions for DII Studies**

*Table 3: Essential Research Reagents and Materials for DII Studies*

| Reagent/Material | Specific Examples | Research Function | Technical Notes |
|---|---|---|---|
| **Dietary Assessment Tools** | 24-hour recall protocols, Food Frequency Questionnaires (FFQs), USDA Food Composition Database | Standardized assessment of dietary intake for DII calculation | FFQs should be validated for specific population; multiple 24-hour recalls preferred over single recall |
| **Inflammatory Biomarker Assays** | High-sensitivity CRP (hs-CRP) immunoturbidimetric assays, IL-6 ELISA kits, TNF-α/TNFαR2 ELISA kits, adiponectin RIAs | Quantification of inflammatory status for DII validation | Consider using multiplex assays for efficiency; establish sample handling protocols to maintain biomarker integrity |
| **Blood Collection Supplies** | EDTA tubes, serum separator tubes, liquid nitrogen storage systems | Proper collection, processing, and storage of biospecimens | Standardize processing time (e.g., within 24 hours); maintain consistent storage temperature (-80°C or lower) |
| **Laboratory Equipment** | ELISA plate readers, automated clinical chemistry analyzers, -80°C freezers | Analysis of biomarkers and storage of samples | Regular calibration of equipment; implement quality control procedures |

| Reagent/Material | Specific Examples | Research Function | Technical Notes |
|---|---|---|---|
| **Statistical Software** | R, STATA, SAS, SPSS | Data management and statistical analysis | Use specialized packages for complex survey data (e.g., NHANES analysis weights) |

The accumulated evidence demonstrates that the Dietary Inflammatory Index serves as a valuable predictive tool for chronic disease risk and comorbidity. The consistent associations observed between higher DII scores and increased risk of chronic kidney disease, diabetes, and cardiovascular-kidney-metabolic syndrome across diverse populations highlight the utility of this tool for understanding diet-disease relationships. The DII's foundation in extensive scientific literature, standardized computation methodology, and validation against inflammatory biomarkers strengthens its application in nutritional epidemiology and chronic disease research.

Significant research gaps remain that warrant investigation. Most existing studies utilize cross-sectional designs, limiting causal inference about the relationship between dietary inflammatory potential and chronic disease development [47] [45]. Large-scale prospective cohort studies with repeated dietary assessments are needed to establish temporal relationships and better understand how changes in DII scores influence disease risk over time. Additionally, research exploring the potential mediating role of specific inflammatory pathways in the relationship between DII and chronic diseases would enhance our understanding of biological mechanisms. Further development and refinement of the DII, including potential ethnic-specific adaptations and investigation of gene-diet interactions, may improve its predictive ability and clinical utility.

The growing evidence supporting DII as a predictor of chronic disease risk and comorbidity suggests potential applications in clinical and public health settings. Assessment of dietary inflammatory potential could enhance risk stratification and inform targeted dietary interventions for high-risk individuals. As research in this field advances, the DII promises to remain an important tool for elucidating the complex relationships between diet, inflammation, and chronic disease.

## Integration into Clinical Practice and Public Health Nutrition Guidelines

The **Dietary Inflammatory Index (DII)** represents a significant advancement in nutritional epidemiology, providing a quantitative measure for assessing the inflammatory potential of an individual's diet [3] [48]. Originally developed through comprehensive literature analysis, the DII was designed to standardize the assessment of diet-induced inflammation across diverse populations and research settings [48] [26]. Unlike earlier dietary indices that focused on adherence to specific dietary patterns or guidelines, the DII was constructed based on empirical evidence linking dietary components to inflammatory biomarkers [3]. This evidence-based foundation enables researchers and clinicians to evaluate dietary patterns along a continuum from maximally anti-inflammatory to maximally pro-inflammatory, creating a standardized metric for investigating diet-inflammation-disease pathways.

The transition of DII from a research tool to clinical and public health applications requires robust validation across various population groups and health conditions. Recent research has expanded beyond initial validation studies to examine the DII's utility in predicting complex multimorbidity conditions, disease progression, and mortality risk [49] [23] [22]. This growing body of evidence supports the integration of DII into clinical practice and public health guidelines as a valuable tool for assessing inflammatory potential of diets and informing targeted nutritional interventions for chronic disease prevention and management.

**DII Performance Across Disease Conditions: Comparative Outcomes**

**Table 1: DII Association with Health Outcomes Across Recent Studies**

| Health Condition | Study Design | Population | Effect Size (Highest vs. Lowest DII) | Key Findings |
|---|---|---|---|---|
| Cardiovascular-Kidney-Metabolic (CKM) Syndrome | Cross-sectional & Prospective Cohort [49] | 7,918 NHANES participants | **OR: 1.85** (1.56-2.20) for advanced CKM stages**HR: 1.45** (1.21-1.73) for all-cause mortality | Biological aging mediated 23% of DII effect on CKM staging and 13% on mortality |
| CKM Syndrome | Cross-sectional [23] | 7,110 NHANES participants | **OR: 1.22** (1.09-1.37) per unit increase in E-DII | Linear relationship (p for nonlinearity = 0.464); alcohol identified as strongest component |
| Coronary Artery Disease | Cross-sectional [22] | 1,015 angiography patients | **OR: 1.48** (1.01-2.16) for severe CAD | Neutrophil-to-lymphocyte ratio mediated 24.7% of DII effect on CAD severity |
| Stroke in MHO Individuals | Cross-sectional [50] | 9,872 NHANES participants | **OR: 1.32** (1.04-1.66) per unit DII increase <2.0**OR: 0.62** (0.44-0.89) per unit DII increase >2.0 | Non-linear relationship with threshold effect at DII = 2.0 |

| Health Condition | Study Design | Population | Effect Size (Highest vs. Lowest DII) | Key Findings |
|---|---|---|---|---|
| Pediatric MASLD | Cross-sectional [51] | 125 children with MASLD | **OR: 4.11** (1.08-15.71) for severe steatosis**OR: 2.61** (1.28-5.32) per unit DII increase | Each unit DII increase associated with 0.006 increase in FIB-4 scores |

The DII demonstrates consistent significant associations with various chronic diseases, with effects remaining robust after adjustment for multiple confounders. The association magnitudes vary across conditions, with particularly strong effects observed for pediatric metabolic dysfunction associated steatotic liver disease (MASLD) and advanced cardiovascular-kidney-metabolic (CKM) syndrome stages [49] [51]. Notably, the DII shows both linear and non-linear relationships depending on the health outcome, suggesting complex underlying biological mechanisms [23] [50].

Beyond simple associations, recent research has focused on elucidating the mediating pathways through which pro-inflammatory diets influence disease risk. Biological aging, as quantified through validated algorithms incorporating clinical biomarkers, mediates a substantial proportion (23%) of the effect of DII on CKM syndrome staging [49]. Similarly, inflammatory hematological markers such as the neutrophil-to-lymphocyte ratio (NLR) mediate approximately 25% of the effect of DII on coronary artery disease severity [22]. These mediation analyses provide mechanistic insights into how dietary patterns influence disease pathogenesis through inflammatory pathways.

## Methodological Protocols for DII Implementation

### DII Calculation and Assessment Methods

The DII calculation follows a standardized protocol that can be adapted based on available dietary assessment methods. The original DII was developed based on 45 food parameters, including nutrients, bioactive compounds, and spices [48] [26]. However, subsequent research has demonstrated that valid DII scores can be computed with fewer parameters when comprehensive dietary data is unavailable [23] [51]. The core calculation involves:

- **Dietary Assessment**: Data collection via 24-hour recalls, food frequency questionnaires (FFQs), or food records. NHANES studies typically use 24-hour recall data [49] [23], while clinical studies often employ validated FFQs [22] [51].
- **Standardization to Global Intakes**: Individual intake values are transformed to z-scores based on a composite global database representing diverse populations [48] [26]. This standardization enables cross-population comparisons.
- **Inflammatory Effect Scoring**: Each dietary component is multiplied by its literature-derived inflammatory effect score [48]. These scores range from strong anti-inflammatory effects (e.g., magnesium: -0.484, fiber: -0.663) to strong pro-inflammatory effects (e.g., saturated fat: +0.373, total fat: +0.298) [23].
- **Energy Adjustment**: The energy-adjusted DII (E-DII) standardizes intake to 1,000 kilocalories to address confounding by total energy intake [23].

### Key Methodological Variations Across Studies

**Table 2: Methodological Approaches in Recent DII Research**

| Study Component | NHANES CKM Study [49] | CKM Cross-sectional Study [23] | CAD Study [22] | Pediatric MASLD Study [51] |
|---|---|---|---|---|
| **Dietary Assessment** | 24-hour recall | 24-hour recall | Validated FFQ | 147-item validated FFQ |
| **DII Parameters** | 27 components | 28 components | Not specified | 34 components |
| **DII Type** | Standard DII | Energy-adjusted DII (E-DII) | Standard DII | Standard DII |
| **Adjustment Factors** | Age, sex, ethnicity, socioeconomic, smoking | Age, sex, ethnicity, socioeconomic, smoking | Clinical, demographic, medication use | Age, sex, BMI, energy intake |
| **Statistical Approaches** | Logistic regression, Cox models, mediation, machine learning | Multiple logistic regression, RCS, WQS, quantile g-computation | Logistic regression, mediation, dose-response | Multiple logistic regression, linear regression |

Methodological consistency across studies enhances comparability, particularly for dietary assessment methods and adjustment for confounding factors. Most studies adjust for core demographic and socioeconomic variables, while varying in their inclusion of clinical parameters based on

research questions [49] [23] [22]. Advanced statistical approaches including restricted cubic splines, mediation analysis, and machine learning algorithms have been increasingly employed to elucidate complex relationships [49] [23].

**Biological Pathways: Connecting Dietary Inflammation to Disease**




*Figure 1: DII Disease Pathway Mapping*

The mechanistic pathway linking pro-inflammatory diets to chronic diseases involves multiple interconnected biological systems. As illustrated in Figure 1, pro-inflammatory dietary patterns (high DII scores) trigger increased production of pro-inflammatory cytokines including IL-6, TNF-α, and CRP [48] [51]. These cytokines subsequently induce oxidative stress, accelerate biological aging processes, and promote endothelial and metabolic dysfunction [49] [22] [51].

Biological aging, quantified through algorithms incorporating clinical biomarkers, mediates approximately one-quarter of the effect of DII on CKM syndrome progression [49]. This finding suggests that pro-inflammatory diets accelerate systemic aging processes, which in turn drive multimorbidity development. Similarly, cellular inflammation markers such as the neutrophil-to-lymphocyte ratio mediate substantial portions of the effect of DII on coronary artery disease severity [22]. In hepatic conditions like MASLD, pro-inflammatory diets promote insulin resistance, increased lipolysis, and fatty acid influx into hepatocytes, leading to steatosis progression and fibrogenesis through hepatic stellate cell activation [51].

**Research Reagent Solutions for DII Studies**

**Table 3: Essential Methodological Components for DII Research**

| Research Component | Specifications & Functions | Examples from Literature |
|---|---|---|
| **Dietary Assessment Tools** | 24-hour recalls, FFQs, dietary records; must capture comprehensive nutrient data | NHANES 24-hour recall [49] [23]; 147-item validated FFQ [51] |
| **Inflammatory Biomarkers** | Validation against established inflammatory markers: IL-6, TNF-α, CRP, NLR | NLR in CAD study [22]; CRP validation in original DII [48] [26] |
| **Global Reference Database** | Composite database of 11 populations worldwide for intake standardization | Means and SDs from global surveys [48] [26] |
| **DII Component Scores** | Literature-derived inflammatory effect scores for 45 food parameters | Effect scores: fiber (-0.663), omega-3 (-0.436), saturated fat (+0.373) [23] |
| **Clinical Outcome Measures** | Standardized disease definitions, staging systems, mortality tracking | CKM staging [49]; Gensini score for CAD [22]; MASLD severity [51] |

| Research Component | Specifications & Functions | Examples from Literature |
|---|---|---|
| **Statistical Analysis Packages** | Software for complex modeling: mediation, RCS, machine learning | R package "mediation" [49]; machine learning algorithms [49] |

The methodological rigor of DII research depends on standardized tools and approaches that ensure consistency across studies. Dietary assessment methods must be sufficiently comprehensive to capture the core parameters contributing to inflammatory potential, with validation against inflammatory biomarkers strengthening the biological plausibility of findings [48] [22]. The global reference database enables cross-population comparability, addressing a significant limitation of earlier dietary pattern approaches [48] [26].

Advanced statistical approaches including mediation analysis, restricted cubic splines, and machine learning algorithms have enhanced the sophistication of recent DII research [49]. These methods enable researchers to elucidate complex non-linear relationships, identify threshold effects, and quantify mediating pathways through which dietary patterns influence disease risk.

### Implementation in Clinical and Public Health Settings

#### Risk Stratification and Intervention Targets

The consistent demonstration of DII's association with disease progression and mortality supports its integration into clinical risk stratification frameworks. Research indicates specific DII thresholds may have clinical utility, with one study identifying **DII > 1.93** as the optimal risk stratification threshold for all-cause mortality in CKM syndrome patients [49]. Similarly, non-linear analyses reveal a **DII threshold of 2.0** for stroke risk in metabolically healthy obese individuals [50], suggesting potential targets for dietary interventions.

Component analyses from multiple studies identify specific dietary factors that most strongly influence disease relationships. **Magnesium and omega-3 fatty acids** consistently emerge as protective components associated with reduced risk of both advanced CKM stages and all-cause mortality [49], while **alcohol intake** demonstrates particularly strong associations with CKM syndrome risk [23]. These findings enable targeted dietary recommendations focusing on specific food components rather than general dietary patterns.

#### Public Health Guidelines Integration

The accumulation of evidence supporting DII's predictive validity across multiple disease states suggests its potential utility in public health nutrition guidelines. The consistent observation that pro-inflammatory diets associate with adverse health outcomes even after comprehensive adjustment for confounders supports the inclusion of inflammation reduction as an explicit goal in dietary recommendations [49] [23] [22].

Public health applications could include:

- **DII screening** in high-risk populations to identify individuals who would benefit from anti-inflammatory dietary interventions
- **Development of anti-inflammatory dietary guidelines** specific to common chronic conditions
- **Integration of DII assessment** into public health surveillance systems to monitor population-level dietary inflammation trends
- **Food labeling approaches** that indicate inflammatory potential to guide consumer choices

### Future Research Directions and Clinical Translation

While substantial evidence supports DII's association with chronic disease risk, several research gaps remain. Future studies should focus on:

- **Intervention trials** examining whether DII reduction translates to improved clinical outcomes
- **Standardization of DII calculation protocols** across different dietary assessment methods
- **Development of population-specific DII thresholds** for clinical decision-making
- **Integration of omics technologies** to elucidate molecular mechanisms linking dietary inflammation to disease pathogenesis
- **Economic analyses** evaluating the cost-effectiveness of DII-guided dietary interventions

The translation of DII research into clinical practice requires development of user-friendly assessment tools that can be efficiently implemented in healthcare settings. Technological approaches including mobile health applications and automated dietary assessment methods could facilitate routine DII screening in primary care and specialty settings. Additionally, education of healthcare providers on interpreting DII scores and implementing appropriate dietary interventions is essential for successful clinical integration.

The consistent findings across multiple studies and diverse populations provide compelling evidence for the integration of DII assessment into both clinical practice and public health guidelines. As research continues to refine our understanding of optimal DII thresholds and effective interventions for reducing dietary inflammation, this evidence-based tool holds significant promise for enhancing chronic disease prevention and management through targeted nutritional approaches.

## Addressing Inconsistencies and Enhancing DII Accuracy in Real-World Settings

## Navigating Inconsistent Associations and Validation Results Across Studies

The **Dietary Inflammatory Index (DII)** represents a significant advancement in nutritional epidemiology, providing a standardized tool to quantify the inflammatory potential of an individual's diet. Developed through systematic analysis of peer-reviewed literature linking dietary components to inflammatory biomarkers, the DII was designed to be universally applicable across diverse populations and research settings [3]. Unlike earlier dietary indexes based primarily on dietary recommendations or specific cultural food patterns, the DII was empirically derived from evidence spanning human studies, animal models, and cell culture experiments [3]. The current version incorporates data from 1,943 qualifying articles published through 2010, evaluating 45 food parameters for their effects on established inflammatory markers including IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP [3] [22].

Despite its widespread adoption in nutritional research, the validation landscape for the DII reveals notable inconsistencies across studies. These variations stem from multiple factors, including methodological differences in dietary assessment, population characteristics, adjustment for confounding variables, and the specific health outcomes examined. This comparative analysis systematically evaluates DII validation results across different health conditions and populations, examines methodological sources of heterogeneity, and provides evidence-based guidance for researchers and clinical professionals navigating these inconsistencies.

### Comparative Validation Across Health Conditions

#### Cardiometabolic Conditions

**Table 1: DII Validation in Cardiometabolic Conditions**

| Health Condition | Study Design | Population | Association with DII | Key Metrics | Consistency Across Studies |
|---|---|---|---|---|---|
| Cardiovascular Mortality | Cohort | 11,310 adults with hypertension [52] | Significant positive correlation | HR: 1.0514 (95% CI: 1.0055–1.0995), p = 0.0278 | Moderate (similar direction but varying magnitude across studies) |
| Severe Coronary Artery Disease | Cross-sectional | 1,015 angiography patients [22] | Significant increase in severe CAD risk for highest DII quartile | OR: 1.48 (95% CI: 1.01–2.16), P for trend: 0.034 | Moderate to high |
| Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD) | Cross-sectional | 8,833 US adults [53] | Significant association with all-cause and cardiovascular mortality | aHR: 1.28 (95% CI: 1.10–1.49), p = 0.002 | High across multiple large studies |
| Pediatric MASLD | Cross-sectional | 125 children and adolescents [54] | Significant association with steatosis severity | OR: 4.11 (95% CI: 1.08–15.71) for inflammatory diets | Limited pediatric evidence |

Research consistently demonstrates that pro-inflammatory diets, as reflected by higher DII scores, associate with adverse cardiometabolic outcomes, though the magnitude of these associations varies. In a large cohort study of 11,310 adults with hypertension, participants with higher DII scores showed significantly increased cardiovascular disease mortality risk during an average 109-month follow-up [52]. Similarly, in patients undergoing elective angiography, those in the highest DII quartile had a 48% increased risk of severe coronary artery disease after adjusting for confounders [22].

The association between DII and MASLD (formerly NAFLD) demonstrates particularly robust validation across multiple studies. In the general adult population, higher DII/E-DII scores consistently correlate with NAFLD confirmed by ultrasound, with one study reporting odds ratios of 1.54 and 1.63 for women and men, respectively [55]. This relationship extends to mortality outcomes, with MASLD patients consuming more pro-inflammatory diets experiencing significantly higher all-cause and cardiovascular mortality [53]. The association appears to hold in pediatric populations as well, with inflammatory diets linked to more severe hepatic steatosis and fibrosis in children and adolescents [54].

#### Other Health Conditions

**Table 2: DII Validation in Other Health Conditions**

| Health Condition | Study Design | Population | Association with DII | Key Metrics | Consistency Across Studies |
|---|---|---|---|---|---|
| Chronic Pain | Cross-sectional | 2,581 US adults [56] | Nonlinear U-shaped relationship | Significant association only at DII ≥ 2.5 | Low (limited evidence with complex pattern) |

| Health Condition | Study Design | Population | Association with DII | Key Metrics | Consistency Across Studies |
|---|---|---|---|---|---|
| Polycystic Ovary Syndrome (PCOS) | Cross-sectional | 115 women with PCOS [57] | No significant association with hyperandrogenism or insulin resistance | $p > 0.05$ for all biomarkers | Low (conflicting with general inflammation literature) |
| All-Cause Mortality | Multiple cohorts | Various populations [53] [3] | Significant association in most studies | Varies by population and adjustment factors | High for general populations |

The validation picture becomes more complex when examining conditions beyond cardiometabolic diseases. For chronic pain, researchers observed a **nonlinear U-shaped relationship** with DII scores, with significant associations only at higher values (DII ≥ 2.5) and an inflection point at -0.9 [56]. This suggests thresholds and complex dynamics may underlie the diet-pain relationship that simple linear models cannot capture.

Interestingly, in PCOS, a condition with known inflammatory components, DII showed no significant association with hyperandrogenism or insulin resistance markers after adjusting for BMI [57]. This highlights the potential confounding effect of adiposity and suggests that in some conditions, overall energy balance and body composition may overshadow specific dietary inflammatory effects.

#### DII Calculation and Dietary Assessment Methods

Variations in DII calculation methodologies contribute significantly to inconsistent associations across studies. The fundamental DII algorithm involves comparing individual dietary intakes to global reference values, creating Z-scores for each food parameter, converting these to percentiles, centering around zero, and multiplying by inflammatory effect scores derived from the literature [3]. However, several methodological choice points introduce variability:

- **Number of dietary components**: Studies utilize different numbers of food parameters when calculating DII scores, ranging from 27 to 34 components in the examined research [56] [54]. This variability can impact score accuracy, as the complete DII was designed to incorporate 45 parameters.
- **Dietary assessment tools**: Different methods for collecting dietary data (e.g., 24-hour recalls, food frequency questionnaires with varying numbers of items) introduce measurement error. Studies using more comprehensive FFQs (147-item [54] or 168-item [55]) potentially capture more complete dietary information.
- **Energy adjustment**: Many researchers now use the energy-adjusted DII (E-DII) to account for total caloric intake, which has demonstrated improved predictive validity for inflammatory biomarkers and health outcomes [3] [55].
- **Global reference database**: The DII calculation relies on a composite global database of dietary intakes from 11 populations worldwide. How well this reference represents the study population may influence scoring accuracy [3].

#### Population Heterogeneity and Confounding Factors

Differential effects across population subgroups represent another source of inconsistent validation results. The association between DII and health outcomes varies substantially based on:

- **Demographic characteristics**: Stronger associations between DII and mortality have been observed in participants <65 years, married individuals, those with college education, non-smokers, non-drinkers, and those without hypertension [53].
- **Gender differences**: Some studies report gender-specific associations, with different effect sizes for men and women for conditions like NAFLD [55].
- **Baseline inflammatory status**: Individuals with existing metabolic conditions may demonstrate different susceptibility to dietary inflammation than generally healthy populations.
- **Body composition**: The relationship between DII and PCOS manifestations became non-significant after controlling for BMI, suggesting adiposity may confound or mediate diet-disease relationships in some contexts [57].




*Diagram 1: Methodological Sources of Inconsistent DII Validation. This pathway illustrates key methodological factors contributing to variable validation results across DII studies, categorized into calculation methods, population factors, and biomarker/analytical approaches.*

#### Biomarker Validation and Outcome Measures

The selection of validation biomarkers and health outcomes significantly influences DII consistency across studies:

- **Inflammatory biomarkers**: While the DII was designed to predict circulating inflammatory markers, studies report varying strength of association with different biomarkers. The E-DII has demonstrated significant association with CRP, supporting its construct validity [55]. However, different inflammatory pathways may be differentially influenced by dietary factors.
- **Clinical vs. surrogate endpoints**: DII shows more consistent associations with hard clinical endpoints (e.g., mortality) compared to intermediate outcomes or subjective measures like chronic pain [53] [56].
- **Disease-specific mechanisms**: Conditions with strong inflammatory pathophysiology (e.g., CAD, MASLD) demonstrate more consistent DII associations than conditions with more complex or multifactorial etiology (e.g., chronic pain, PCOS) [53] [22] [57].

### Experimental Protocols and Research Toolkit

#### Standard DII Calculation Protocol

For researchers implementing DII validation studies, the following standardized protocol is recommended based on methodological refinements described in the literature [3]:

- **Dietary Assessment**: Administer a validated food frequency questionnaire (FFQ) with comprehensive coverage of foods and beverages. The FFQ should ideally include at least 147 items to adequately capture all relevant DII components [54].
- **Data Processing**: Convert reported consumption frequencies into daily intake amounts for each food parameter using standard portion sizes and conversion factors.
- **Global Comparison**: For each of the 45 DII food parameters, compute a Z-score relative to the global mean and standard deviation from the composite reference database: `Z = (individual intake - global mean) / global standard deviation`
- **Percentile Conversion**: Convert Z-scores to percentiles to minimize the effect of right-skewing common in dietary data.
- **Centering**: Multiply each percentile by 2 and subtract 1 to create centered percentiles symmetric around zero.
- **Inflammatory Scoring**: Multiply each centered percentile value by its respective food parameter-specific inflammatory effect score (derived from literature review).
- **Aggregation**: Sum all food parameter-specific DII scores to obtain the overall DII score for each participant.
- **Energy Adjustment**: Calculate energy-adjusted DII (E-DII) using the residual method to account for total energy intake variations.

#### Research Reagent Solutions and Essential Materials

**Table 3: Research Reagent Solutions for DII Studies**

| Research Tool | Specifications | Function in DII Research | Validation Considerations |
|---|---|---|---|
| Food Frequency Questionnaire (FFQ) | 147-168 items, validated for target population [54] [55] | Captures habitual dietary intake for DII calculation | Requires validation against dietary records or recalls in study population |
| Dietary Analysis Software | Nutritionist IV, USDA Food Composition Database [54] | Converts food consumption to nutrient intakes | Database completeness for DII components critical |
| Inflammatory Biomarker Assays | High-sensitivity CRP, IL-6, TNF-α, IL-1β [3] [22] | Validates DII against objective inflammatory measures | Standardized protocols, fasting blood collection |
| Body Composition Measures | BMI, waist circumference, bioelectrical impedance [53] [57] | Assesses adiposity as potential confounder | Calibrated equipment, trained technicians |
| Clinical Outcome Assessment | Mortality registries, ultrasound, angiography [52] [55] | Documents hard endpoints for validation | Standardized diagnostic criteria essential |

### Interpretation Framework for Inconsistent Results

When evaluating seemingly contradictory DII validation studies, researchers should consider the following framework:

- **Methodological quality assessment**: Prioritize studies with comprehensive dietary assessment, appropriate energy adjustment, and adequate control for key confounders like BMI, smoking, and physical activity.

- **Population context**: Interpret findings within the specific population characteristics, recognizing that DII effects may be modified by age, gender, genetic background, and baseline health status.
- **Consistency across study designs**: Value consistent patterns emerging across different study designs (cohort, cross-sectional, case-control) and populations more than individual study results.
- **Biological gradient assessment**: Evaluate whether dose-response relationships are present, as these strengthen causal inference despite variation in absolute effect sizes.
- **Biomarker concordance**: Consider studies that demonstrate concordance between DII scores and objective inflammatory biomarkers as providing stronger validation evidence.

The accumulated evidence suggests that despite methodological variations, the DII consistently predicts health outcomes with strong inflammatory pathophysiology, particularly in cardiometabolic diseases. Future research should focus on standardizing DII calculation methods, exploring nonlinear relationships, and identifying population subgroups that may derive particular benefit from anti-inflammatory dietary interventions.

Validation of the Dietary Inflammatory Index across diverse studies reveals a complex landscape characterized by generally consistent associations with cardiometabolic outcomes but greater variability for other health conditions. Methodological differences in DII calculation, population characteristics, and outcome assessment contribute significantly to these inconsistencies. Nevertheless, the DII remains a valuable tool for quantifying the inflammatory potential of diet in research settings, particularly when using energy-adjusted scores and standardized methodologies. Researchers should interpret DII validation studies through the lens of methodological quality, population context, and consistency across different study designs to navigate this complex evidence base effectively.

## Impact of Energy Intake and Dietary Assessment Methods on DII Scoring

The **Dietary Inflammatory Index (DII)** has emerged as a significant tool for quantifying the inflammatory potential of an individual's diet, with applications spanning epidemiological research and clinical practice. As a literature-derived, population-based index, the DII was designed to standardize the assessment of diet-associated inflammation across diverse populations and study designs [3]. The theoretical foundation of the DII rests on its ability to predict inflammatory responses based on dietary patterns, with scores ranging from anti-inflammatory (negative values) to pro-inflammatory (positive values) [28]. However, the accuracy and comparability of DII scores are profoundly influenced by two methodological factors: the approach to accounting for total energy intake and the selection of dietary assessment methods. These factors represent critical sources of variability that researchers must address to ensure valid and reproducible results in DII-related studies.

The evolution of the DII methodology reflects ongoing efforts to enhance its precision. The original DII, introduced in 2009, was substantially refined in 2014 to address statistical limitations and improve accuracy [3] [28]. A key advancement was the development of the **energy-adjusted DII (E-DII)**, which specifically addresses confounding by total energy intake [58]. Understanding these methodological nuances is essential for researchers investigating the relationship between dietary patterns and inflammation-driven health outcomes, particularly as the DII gains traction in nutritional epidemiology and chronic disease research.

### Dietary Assessment Methods in DII Research

The methodology used to collect dietary data significantly influences DII scoring accuracy and reliability. Various approaches have been employed across studies, each with distinct strengths and limitations that researchers must consider when designing DII validation studies.

#### Common Dietary Assessment Tools

- **Food Frequency Questionnaires (FFQs):** The most prevalent method in large-scale epidemiological studies, FFQs collect data on frequency and sometimes portion sizes of food items consumed over a specific period. The Polish arm of the PURE study utilized a validated FFQ to assess habitual food intake across 1791 participants, enabling computation of DII scores linked to cardiovascular risk factors [36]. Similarly, the CHINA-DII development study employed FFQs to collect dietary data from 256 gastric cancer patients, demonstrating significant correlation with inflammatory biomarkers [59].
- **24-Hour Dietary Recalls:** This method involves detailed interviews where participants recall all foods and beverages consumed in the previous 24 hours. The NHANES study implemented a **two-phase 24-hour recall methodology** with an initial in-person interview followed by a telephone interview 3-10 days later, administered by trained personnel following standardized protocols [23]. This approach enhances accuracy through multiple data collection points but may not capture habitual intake without repeated administrations.
- **Dietary History Questionnaires (DHQs):** More comprehensive than FFQs, DHQs collect detailed information on habitual diet, including portion sizes and preparation methods. The PLCO Cancer Screening Trial used a DHQ covering 124 food items to assess dietary intake over the past year, which was shown to reflect absolute food intake more accurately compared to other FFQs [58].

*Table 1: Comparison of Dietary Assessment Methods Used in DII Studies*

| Method | Study Example | Key Features | Parameters Captured | Strengths | Limitations |
|---|---|---|---|---|---|
| **Food Frequency Questionnaire (FFQ)** | PURE Study (Poland) [36] | Assesses frequency of food consumption over specific period | Varies (106 items in Korean study [33]) | Efficient for large studies; captures habitual intake | Memory dependent; portion size estimation challenging |
| **24-Hour Dietary Recall** | NHANES Analysis [23] | Detailed recall of all foods consumed in previous 24 hours | 28 nutritional components | Reduced memory bias; multiple recalls possible | Single day may not represent habitual diet |
| **Dietary History Questionnaire (DHQ)** | PLCO Trial [58] | Comprehensive assessment of habitual diet | 35 dietary parameters | Detailed information on usual intake | Time-consuming; participant burden |

#### Impact on DII Component Availability

The choice of dietary assessment instrument directly affects the number of DII components that can be included in scoring. The comprehensive DII incorporates up to 45 dietary parameters, including nutrients, foods, and bioactive compounds [3]. However, most studies capture a subset due to methodological constraints:

- The PLCO trial included 35 of 45 possible parameters in its DII calculation [58].
- The NHANES analysis incorporated 28 nutritional components, noting that predictive capability remains robust with at least 28 parameters [23].
- The Korean DII study assessed 37 available parameters from the full set of 45 [33].

This variation in captured parameters introduces measurement heterogeneity that researchers must acknowledge when comparing DII scores across studies.

### Energy Adjustment Methodologies in DII Scoring

Total energy intake represents a significant confounding factor in DII calculation, as absolute nutrient consumption correlates strongly with total caloric intake. The development of the **energy-adjusted DII (E-DII)** specifically addressed this limitation by standardizing dietary components per 1000 kilocalories of consumed food [58] [23].

#### E-DII Calculation Protocol

The E-DII computation follows a standardized protocol:

- **Energy Density Adjustment:** Individual food parameter intakes are calculated per 1000 kilocalories of total energy intake rather than as absolute amounts [58].
- **Global Standardization:** The energy-adjusted intake for each parameter is compared to a global reference database representing mean intake levels across diverse populations [3].
- **Z-score Conversion:** The difference between individual intake and global mean is divided by the global standard deviation to generate a z-score [58].
- **Percentile Transformation:** Z-scores are converted to percentiles and centered around zero [23].
- **Inflammatory Effect Scoring:** Centered percentiles are multiplied by literature-derived inflammatory effect scores for each parameter [23].
- **Aggregation:** Individual parameter scores are summed to generate the overall E-DII score [58].

#### Comparative Evidence: Standard DII vs. E-DII

The critical importance of energy adjustment is demonstrated in the PLCO cancer trial, which explicitly compared standard DII with E-DII in relation to lung cancer risk [58]. This large prospective cohort study (n=101,755) with median 9.4-year follow-up revealed that energy adjustment meaningfully enhanced the predictive validity of DII scoring:

*Table 2: Impact of Energy Adjustment on DII Performance in the PLCO Lung Cancer Study [58]*

| DII Type | Data Source | Hazard Ratio (Highest vs. Lowest Quartile) | 95% Confidence Interval | P-trend |
|---|---|---|---|---|
| **E-DII** | Food + Supplements | 1.31 | 1.14-1.52 | 0.002 |

| DII Type | Data Source | Hazard Ratio (Highest vs. Lowest Quartile) | 95% Confidence Interval | P-trend |
|---|---|---|---|---|
| **E-DII** | Food only | 1.39 | 1.22-1.58 | <0.001 |
| **Standard DII** | Not reported | Inconsistent associations in previous studies | Not significant in some cohorts | Variable |

This study highlighted that previous inconsistent findings regarding DII and lung cancer risk may have resulted from failure to adequately account for energy intake [58]. The E-DII demonstrated statistically significant dose-response relationships, whereas standard DII had produced conflicting results in earlier research.

### Experimental Protocols for DII Validation

#### E-DII Calculation Methodology

The following protocol details the E-DII computation process as implemented in the NHANES analysis [23]:

**Step 1: Dietary Data Collection**

- Conduct two non-consecutive 24-hour dietary recalls using standardized protocols
- Include both in-person and telephone interviews to enhance accuracy
- Collect detailed information on portion sizes and preparation methods
- Record dietary supplement usage separately

**Step 2: Data Standardization**

- Calculate intake of each dietary parameter per 1000 kilocalories of total energy
- Reference each energy-adjusted intake against global population norms
- Compute z-scores: (individual intake - global mean) / global standard deviation

**Step 3: Score Calculation**

- Convert z-scores to percentiles and center around zero (percentile × 2 - 1)
- Multiply centered percentiles by corresponding inflammatory effect scores
- Sum across all parameters to generate overall E-DII score

**Step 4: Validation**

- Assess correlation between E-DII scores and inflammatory biomarkers (e.g., hs-CRP, IL-6)
- Conduct sensitivity analyses with different parameter sets
- Perform subgroup analyses to evaluate consistency across populations

#### CHINA-DII Development Protocol

The creation of the population-specific CHINA-DII provides a template for adapting DII methodology to distinct dietary patterns [59]:

**Literature Review Phase:**

- Systematic search of five databases (CNKI, Wanfang, VIP, PubMed, Web of Science)
- Inclusion criteria: Chinese adults ≥18 years, studies from 2009-2024, sample size >200
- Data extraction: Mean and standard deviation of dietary intake values
- Quality assessment using standardized tools

**Database Construction:**

- Establishment of Chinese adult dietary intake database encompassing 27 dietary components
- Reference to original DII methodology for inflammatory effect scores [3]
- Integration of traditional Chinese foods and cooking methods

**Validation Study:**

- Recruitment of 256 newly diagnosed gastric cancer patients
- Collection of demographic, clinical, and dietary data (FFQ-based)
- Measurement of high-sensitivity C-reactive protein (hs-CRP) levels
- Statistical analysis: Spearman rank correlation and multivariate logistic regression

The CHINA-DII validation demonstrated significant positive correlation between DII scores and hs-CRP levels ($r=0.20$, $p \leq 0.001$), supporting its validity for Chinese populations [59].

**Visualization of E-DII Calculation Workflow**

The following diagram illustrates the systematic process for calculating Energy-Adjusted Dietary Inflammatory Index scores:




**The Scientist's Toolkit: Essential Research Reagents and Materials**

*Table 3: Essential Research Reagents and Resources for DII Studies*

| Category | Specific Items | Research Function | Example Implementation |
|---|---|---|---|
| **Dietary Assessment Tools** | Validated FFQ, 24-hour recall protocols, DHQ | Standardized dietary data collection | PURE Study FFQ [36]; NHANES 24-hour recall [23] |
| **Reference Databases** | Global intake norms, Food composition tables | Standardization of dietary parameters | 11-country composite database [3]; Korean food composition table [33] |
| **Inflammatory Biomarkers** | hs-CRP, IL-6, TNF-α, IL-1β | Validation of DII scores against objective measures | CHINA-DII validation with hs-CRP [59] |
| **Statistical Software** | R, Stata, SAS | DII calculation and statistical analysis | Restricted cubic spline models in Stata [58] [60] |
| **Laboratory Equipment** | Automated blood analyzers, ELISA kits | Biomarker quantification | Enzymatic test kits for lipids [36] |

The methodological approaches to addressing energy intake and dietary assessment significantly impact DII scoring validity and comparability across studies. The evidence consistently demonstrates that **energy-adjusted DII (E-DII)** provides more reliable and interpretable results compared to standard DII, particularly when examining relationships with health outcomes [58] [23]. Furthermore, the choice of dietary assessment method influences the number of DII components that can be captured, with more comprehensive tools (DHQs) generally enabling more accurate scoring than brief FFQs.

Researchers should prioritize energy adjustment in DII calculation and select dietary assessment methods that balance comprehensiveness with feasibility for their specific population and research context. The development of population-specific DII versions, such as CHINA-DII, represents a promising approach for enhancing relevance while maintaining methodological rigor [59]. As DII research continues to evolve, standardization of these methodological elements will be crucial for generating comparable evidence across studies and translating findings into meaningful dietary recommendations for chronic disease prevention and management.

The Dietary Inflammatory Index (DII) and its empirical derivative, the EDII, are validated tools that quantify the inflammatory potential of an individual's diet based on its composition [5] [42]. Higher DII scores, indicating a more pro-inflammatory diet, are consistently associated with adverse health outcomes, including frailty, non-alcoholic fatty liver disease (NAFLD), and all-cause mortality [61] [62] [63]. However, the application and interpretation of these indices are not uniform across all demographic groups. Specific populations, including vegetarians, the elderly, and obese individuals, present unique dietary patterns, physiological states, and metabolic challenges that can significantly influence the relationship between DII scores and actual inflammatory biomarkers. This guide objectively compares the performance and validation of DII scores within these three populations, synthesizing current experimental data to highlight key challenges and methodological considerations for researchers and drug development professionals.

## Comparative Data on DII Application Across Populations

The table below summarizes key findings from recent studies investigating the DII in vegetarians, the elderly, and obese individuals.

*Table 1: Comparison of DII Challenges and Associations in Specific Populations*

| Population | Key Challenges for DII Application | Documented Association with Health Outcomes | Supporting Experimental Data |
|---|---|---|---|
| **Vegetarians** | Low total energy intake can mask anti-inflammatory potential; nutrient deficiencies (e.g., EPA/DHA, Vitamin B12, Zinc) may alter inflammatory status [64] [65]. | Paradoxical findings: Lower theoretical DII but sometimes higher inflammatory biomarkers (TNF-α, IL-6, NLR, PLR) in real-world cohorts [64]. | **Cross-sectional study (n=558):** Vegetarians had significantly lower energy-adjusted DII (E-DII) but higher levels of TNF-α, IL-6, NLR, and PLR compared to omnivores [64]. |
| **Elderly** | The condition of frailty itself involves a pro-inflammatory state (inflammaging), which may confound diet-disease relationships [61]. | Strong positive association: Highest DII category linked to significantly increased risk of both frailty (OR=1.47) and pre-frailty (OR=1.54) [61]. | **Meta-analysis (15 studies, n=42,130):** Pro-inflammatory diets consistently associated with increased frailty risk across diverse geographic regions and assessment tools [61]. |
| **Obese Individuals** | Existing chronic low-grade inflammation and metabolic dysfunction can complicate the isolation of diet's inflammatory effect [62] [66]. | Significant association: A one-standard-deviation increase in DII resulted in a 21% increased risk of fatty liver disease, with a stronger, non-linear correlation in obese subjects [62]. | **NHANES Analysis (n=3,456):** The highest DII tertile was associated with a 39% increased risk of fatty liver disease. A Western dietary pattern was directly associated with higher BMI and fat mass [62] [66]. |

## Detailed Experimental Protocols and Methodologies

### Investigating the Vegetarian Paradox: A Cross-Sectional Study

A 2024 study directly compared dietary intake and inflammatory biomarkers between Chinese vegetarians and omnivores, providing a protocol to address the energy intake confounder [64].

- **Participant Recruitment:** Researchers recruited 279 vegetarians and 279 age- and sex-matched omnivores. Vegetarianism was defined as adherence to the diet for at least one year.
- **Dietary Assessment:** A 24-hour dietary review questionnaire was administered to all participants. The data was used to calculate three versions of the index:
  - **Raw DII:** Based on reported food intake.
  - **Energy-Adjusted DII (E-DII):** Adjusted for total energy intake.

    - **Theoretical DII:** Dietitians designed energy-matched vegetarian and omnivore recipes to calculate a hypothetical DII for a balanced diet.
- **Biomarker Measurement:** Fasting blood samples were analyzed for five inflammatory biomarkers: C-reactive protein (CRP), interleukin-6 (IL-6), tumor necrosis factor-α (TNF-α), neutrophil-lymphocyte ratio (NLR), and platelet-lymphocyte ratio (PLR).
- **Statistical Analysis:** Analysis of covariance (ANCOVA) was used to compare DII scores and biomarker levels between groups, adjusting for covariates including BMI, alcohol consumption, and physical activity [64].

#### Establishing the DII-Frailty Link: A Meta-Analytical Approach

A 2025 systematic review and meta-analysis established the robust association between DII and frailty in middle-aged and older adults [61].

- **Literature Search:** Researchers systematically searched eight databases (e.g., PubMed, Embase, CNKI, Web of Science) from inception to January 2025 using predefined terms related to DII and frailty.
- **Study Selection:** Included studies were observational (cohort, case-control, cross-sectional) and examined the association between DII and risk of frailty or pre-frailty in adults aged ≥45 years.
- **Data Extraction and Quality Assessment:** Two reviewers independently extracted data (author, year, country, study design, sample size, effect estimates) and assessed study quality using the Newcastle-Ottawa Scale (NOS) for longitudinal studies and the AHRQ tool for cross-sectional studies.
- **Statistical Synthesis:** Meta-analysis was performed using RevMan 5.3 and Stata 15.0. Pooled odds ratios (OR) and 95% confidence intervals (CI) for the highest vs. lowest DII categories were calculated using random-effects models. Subgroup analyses, sensitivity analyses, and publication bias tests were conducted [61].

#### Assessing DII and NAFLD in Obesity: An NHANES Analysis

A 2025 study utilized National Health and Nutrition Examination Survey (NHANES) data to explore the DII-NAFLD relationship, specifically in obese and non-obese populations [62].

- **Data Source and Population:** The analysis used data from 3,456 adults (aged ≥20) from the 1999-2000 NHANES, a cross-sectional survey employing a multistage, stratified probability sampling design.
- **DII Calculation:** Dietary data from 24-hour recalls were used to calculate DII scores based on 28 of the 45 possible food parameters (e.g., carbohydrates, fats, vitamins, flavonoids).
- **Outcome Definition:** Fatty liver disease was defined using the Fatty Liver Index (FLI), a predictive algorithm that incorporates BMI, waist circumference, triglycerides, and gamma-glutamyl transferase (GGT). An FLI ≥60 was used to diagnose NAFLD.
- **Statistical Analysis:** Multivariable logistic regression models, adjusted for age, sex, race, and other covariates, were used to assess the association between DII (as continuous and tertiles) and NAFLD. Restricted cubic splines (RCS) were applied to model non-linear relationships, particularly within the obese subgroup [62].

## Biological Pathways and Research Workflows

The following diagram illustrates the conceptual pathway through which pro-inflammatory diets contribute to adverse outcomes in the three specific populations, integrating key mechanisms identified from the research.




## The Scientist's Toolkit: Key Research Reagent Solutions

For researchers designing studies to validate DII scores in specific populations, the following reagents and methodologies are essential.

*Table 2: Essential Research Reagents and Methodologies for DII Validation Studies*

| Reagent/Methodology | Primary Function in DII Research | Application Notes for Specific Populations |
|---|---|---|
| **High-Sensitivity CRP (hsCRP) Immunoassays** | Quantifies baseline levels of systemic inflammation [64] [65]. | Critical for all populations. In obese subjects, levels may be chronically elevated, requiring careful statistical adjustment [62] [66]. |
| **Multiplex Cytokine Panels (IL-6, TNF-α, IL-1β)** | Measures multiple pro-inflammatory cytokines from a single sample [64] [5]. | Vital for elucidating the "vegetarian paradox" where CRP may be normal but other cytokines (TNF-α, IL-6) are elevated [64]. |
| **Plasma & Erythrocyte Fatty Acid Profiling** | Assesses short- and long-term status of pro/anti-inflammatory fatty acids [65]. | Essential for vegetarian/vegan studies to confirm low EPA/DHA status and high n-6:n-3 ratio, which can modulate inflammation independent of DII score [65]. |
| **Dietary Assessment FFQs & 24-h Recalls** | Collects data to calculate DII/EDII scores [64] [62] [5]. | Must be validated for local cuisine. For elderly, consider tools adapted for memory recall. For vegetarians, ensure comprehensive plant-based food lists [64] [67]. |
| **Fatty Liver Index (FLI) Algorithm** | A non-invasive proxy for NAFLD diagnosis in large cohorts [62]. | Key outcome measure for DII studies in obese populations, utilizing routinely measured clinical parameters (BMI, WC, TG, GGT) [62]. |

Validation of Dietary Inflammatory Index scores requires a nuanced, population-specific approach. Experimental data confirms that while a pro-inflammatory diet is a universal risk factor for adverse health outcomes, the mechanisms and manifestations differ significantly. **Vegetarians** may exhibit a discrepancy between the theoretical anti-inflammatory potential of their diet and their actual inflammatory profile, largely driven by low energy intake and specific nutrient deficiencies. For the **elderly**, the DII robustly predicts frailty risk, underscoring the role of diet in the inflammaging process. In **obese individuals**, the DII strongly associates with conditions like NAFLD, with the relationship often exhibiting non-linearity and being modulated by the underlying metabolic state. Future research and drug development efforts must account for these population-specific challenges to accurately gauge the inflammatory impact of diet and develop effective, targeted interventions.

## Optimizing DII with Population-Specific Databases and Food Parameters

The Dietary Inflammatory Index (DII) represents a significant advancement in nutritional epidemiology, providing a quantitative means to assess the inflammatory potential of an individual's diet. Developed through systematic review of scientific literature, the DII was designed to categorize diets

on a continuum from maximally anti-inflammatory to maximally pro-inflammatory [26]. Unlike previous dietary indexes based on dietary recommendations or specific cultural foodways, the DII was derived from empirical evidence linking dietary components to inflammatory biomarkers [3]. This literature-derived, population-based approach allows researchers to investigate connections between diet-induced inflammation and chronic diseases across diverse populations.

The original DII framework has evolved substantially since its inception. The first version, debuted in 2009, was based on scoring 927 peer-reviewed articles published through 2007 [3]. Recognizing methodological limitations and the expanding evidence base, developers created an enhanced version published in 2014 that incorporated additional years of research and improved scoring algorithms [26]. This revised DII has since been applied in over 200 human studies investigating various health outcomes, forming the basis for numerous meta-analyses [3]. The index's ability to capture diet-related inflammatory potential has made it a valuable tool for researchers studying obesity, cardiovascular disease, diabetes, cancer, and other inflammation-mediated conditions [22] [66].

## Core DII Framework: Components and Computational Foundation

### Food Parameters and Inflammatory Biomarkers

The DII is constructed from 45 food parameters, including nutrients, bioactive compounds, and whole foods that have documented effects on inflammation [26] [3]. These parameters were identified through systematic review of peer-reviewed literature linking dietary components to six specific inflammatory biomarkers: the pro-inflammatory markers IL-1β, IL-6, TNF-α, and CRP, and the anti-inflammatory markers IL-4 and IL-10 [26]. The selection of these particular biomarkers was based on their established importance in inflammation and the robustness of literature concerning them.

The literature review process for DII development was comprehensive, involving screening of approximately 6,500 articles published through December 2010 [26]. From these, 1,943 qualifying articles were read and scored based on whether each dietary parameter increased (+1), decreased (-1), or had no effect (0) on the six inflammatory biomarkers [26]. The scoring system also incorporated weights based on study design characteristics, with human experimental studies receiving the highest weight (10), followed by prospective cohort studies (8), case-control studies (7), cross-sectional studies (6), animal experimental studies (5), and cell culture studies (3) [26]. This weighting system helped account for variations in evidence quality across studies.

### Global Database and Standardization Method

A critical innovation in the revised DII was the incorporation of a global reference database to standardize individual intakes. Researchers identified eleven food consumption datasets from countries around the world, including Australia, Bahrain, Denmark, India, Japan, Mexico, New Zealand, South Korea, Taiwan, the United Kingdom, and the United States [26] [3]. These datasets formed a composite database containing means and standard deviations for intakes of each of the 45 food parameters, providing "global norms" against which individual consumption could be compared.

The computational process for deriving DII scores involves multiple steps. First, individual intake of each food parameter is standardized against the global database using Z-scores [68]. These Z-scores are then converted to percentiles and centered by multiplying by 2 and subtracting 1, resulting in values ranging from -1 (maximally anti-inflammatory) to +1 (maximally pro-inflammatory) for each parameter [26] [3]. These centered percentiles are then multiplied by the respective food parameter-specific overall inflammatory effect scores (derived from the literature review) and summed across all parameters to yield the overall DII score [26]. The theoretical range of DII scores spans from -8.87 (maximally anti-inflammatory) to +7.98 (maximally pro-inflammatory), with a median of +0.23 in the global composite database [26].

## Methodological Optimizations: Enhancing DII Precision

### Mathematical Refinements in DII Calculation

While the original DII calculation method represented a significant advancement, researchers have identified opportunities for mathematical refinement. A key limitation involves the transformation of standardized daily food consumption to percentile scores using the standard normal distribution function [68]. This approach assumes dietary intake parameters follow a normal distribution, which often does not reflect reality due to the inherent skewness of consumption patterns for many food parameters.

To address this limitation, Pawlow et al. (2021) proposed novel computational approaches termed the Scaling-Formula (SF) and Scaling-Formula with Outlier Detection (SFOD) methods [68]. These methods replace the standard normal distribution function with alternative scaling approaches that better handle the non-normal distribution of dietary data. When tested on simulated data and applied to the prospective TEENDIAB study cohort, the SF and SFOD methods demonstrated improved performance over the original calculation method, suggesting they may provide more accurate DII scores [68]. These mathematical refinements represent important contributions to the ongoing optimization of DII methodology.

### Energy Adjustment and Population Specific Adaptations

Another significant enhancement to the DII framework involves energy adjustment. The energy-adjusted DII (E-DII) was developed to minimize the influence of total energy intake on inflammatory potential assessment [69]. This adjustment is particularly important in populations where energy intake varies substantially or where under-reporting or over-reporting of consumption may occur. The E-DII is calculated as the DII divided by total energy intake, providing a measure of dietary inflammatory potential per unit of energy consumed [69].

Research has demonstrated the value of this energy-adjusted approach. A 2022 study investigating the relationship between E-DII and mortality in chronic kidney disease patients found that higher E-DII scores were significantly associated with increased 5-year all-cause and cardiovascular mortality [69]. The study revealed a "J-shaped" relationship between E-DII scores and all-cause mortality, and a near-linear relationship with cardiovascular mortality [69]. These findings highlight how methodological refinements like energy adjustment can enhance the DII's predictive validity in specific patient populations.

*Table 1: Key Methodological Optimizations in DII Calculation*

| Optimization Approach | Methodological Details | Advantages | Reference |
|---|---|---|---|
| **Global Database Standardization** | Standardization of individual intakes to reference values from 11 international datasets | Enables cross-population comparisons; reduces arbitrariness of raw consumption amounts | [26] [3] |
| **Mathematical Refinements (SF/SFOD)** | Replacement of standard normal distribution function with improved scaling approaches | Better handles non-normal distribution of dietary data; improved accuracy | [68] |
| **Energy Adjustment (E-DII)** | DII divided by total energy intake | Controls for energy intake variation; enhances comparability across individuals | [69] |
| **Flavonoid Inclusion** | Addition of 16 flavonoids grouped into 6 categories | Incorporates important anti-inflammatory compounds; enhances content validity | [3] |
| **Literature Base Expansion** | Inclusion of 1,943 articles (through 2010) vs. original 927 articles (through 2007) | More robust evidence base; improved parameter effect estimates | [26] [3] |

### Experimental Validation: DII Performance Across Health Conditions

#### Hepatic Health and Non-Alcoholic Fatty Liver Disease

The association between DII and non-alcoholic fatty liver disease (NAFLD) has been extensively investigated through observational studies. A 2025 systematic review and meta-analysis incorporated 11 studies (9 cross-sectional with 14 effect sizes and 2 cohort with 2 effect sizes) examining the relationship between DII and NAFLD risk [70]. The analysis revealed a significant association between higher DII scores and increased NAFLD risk, with a pooled odds ratio of 1.56 (95% CI: 1.24-1.95; $p < 0.001$) in cross-sectional studies and a hazard ratio of 0.21 (95% CI: 0.12-0.30; $p < 0.0001$) in cohort studies [70].

Subgroup analyses in this meta-analysis provided further insights into DII performance across population subsets. The association remained consistent across participants with BMI ≥ 25, studies using either standard DII or energy-adjusted DII, studies conducted in Asia and Europe, and participants younger than 46 years [70]. Notably, heterogeneity was reduced ($I^2 < 50\%$) in these subgroups, suggesting more consistent effects within specific populations. However, the certainty of evidence according to GRADE criteria was rated as "very low," highlighting the need for additional high-quality studies [70].

#### Cardiovascular Disease and Coronary Artery Disease

Research has also validated the DII's utility in cardiovascular disease contexts, particularly coronary artery disease (CAD). A 2025 cross-sectional study of 1,015 individuals undergoing elective angiography examined the relationship between DII and severe CAD, as classified by Gensini score [22]. After adjusting for confounding factors, results indicated a significantly increased severe CAD risk for higher DII quartiles, with odds ratios of 1.52 (95% CI: 1.05-2.22) and 1.48 (95% CI: 1.01-2.16) for the third and fourth (most pro-inflammatory) quartiles, respectively [22].

This study also investigated potential mechanisms linking dietary inflammation to CAD severity. Researchers found that the neutrophil-to-lymphocyte ratio (NLR), a marker of systemic inflammation, mediated 24.7% (95% CI: 15.2%-98.3%) of the total effect of DII on severe CAD [22]. Non-linear dose-response analysis further demonstrated a persistent increase in severe CAD risk as DII levels increased [22]. These findings not only validate the DII's association with cardiovascular outcomes but also elucidate biological pathways through which pro-inflammatory diets may influence CAD progression.

*Table 2: DII Performance Across Health Conditions in Observational Studies*

| Health Condition | Study Design | Association Measure | Effect Size | Certainty of Evidence | Reference |
|---|---|---|---|---|---|
| **Non-Alcoholic Fatty Liver Disease** | Systematic review of 11 observational studies | Pooled OR (cross-sectional) | 1.56 (95% CI: 1.24-1.95) | Very Low (GRADE) | [70] |
| **Coronary Artery Disease** | Cross-sectional (n=1,015) | OR (4th vs. 1st quartile) | 1.48 (95% CI: 1.01-2.16) | Moderate | [22] |
| **5-Year All-Cause Mortality in CKD** | Cohort (n=7,207) | HR (highest vs. lowest E-DII tertile) | 1.33 (95% CI: 1.15-1.54) | Moderate | [69] |
| **5-Year Cardiovascular Mortality in CKD** | Cohort (n=7,207) | HR (highest vs. lowest E-DII tertile) | 1.54 (95% CI: 1.15-2.07) | Moderate | [69] |
| **Insulin Resistance in Obesity** | Cross-sectional (n=151) | OR (Traditional pattern) | 0.3 (95% CI: 0.1-0.9) | Low | [66] |

## Comparative Analysis: DII Calculation Methodologies

### Original vs. Optimized Computational Approaches

The original DII calculation method pioneered a novel approach to quantifying dietary inflammation but contained several computational aspects that could be optimized. The transformation of Z-scores to percentiles using the standard normal distribution function represented a particular limitation, as it assumed normality for dietary intake parameters that often follow skewed distributions [68]. This approach could potentially misrepresent the relative inflammatory contribution of different food parameters, especially for those with non-normal consumption patterns.

The SF and SFOD methods proposed by Pawlow et al. address this limitation by implementing alternative scaling approaches that do not rely on distributional assumptions [68]. When applied to both simulated data and real-world cohort data, these methods demonstrated improved performance over the original calculation method [68]. Additionally, the development of energy-adjusted DII (E-DII) represented another significant optimization, controlling for the confounding effects of total energy intake [69]. Research has shown that E-DII may provide enhanced predictive validity for certain health outcomes, particularly those where energy balance plays an important pathophysiological role.

### Population-Specific vs. Global Standardization

A fundamental tension in DII optimization involves balancing global standardization with population-specific adaptations. The global reference database encompassing 11 diverse populations allows for meaningful cross-population comparisons and represents a substantial improvement over the original DII, which used raw consumption amounts without external standardization [26] [3]. However, researchers have also recognized that certain population-specific adaptations may enhance the DII's performance in particular settings.

The subgroup analyses from the NAFLD meta-analysis illustrate this point, showing reduced heterogeneity ($I^2 < 50\%$) when examining specific populations defined by geographic region, age, or BMI status [70]. This suggests that while the global standardization approach provides a valuable foundation, future refinements might incorporate population-specific reference ranges for certain parameters where local consumption patterns differ substantially from global norms. Such adaptations could enhance the DII's sensitivity to detect diet-inflammation-disease relationships within specific populations while maintaining cross-population comparability.

## Research Toolkit: Essential Materials and Methods

### Dietary Assessment Instruments

The foundation of accurate DII calculation lies in comprehensive dietary assessment. The most common instrument for DII computation is the Food Frequency Questionnaire (FFQ), which captures habitual consumption patterns across a specified period. Studies included in this analysis typically employed validated FFQs with varying numbers of food items, ranging from 168 items in studies of obese individuals [66] to other comprehensive instruments that capture all 45 DII parameters [22]. The FFQ must be appropriately validated for the specific population under study, with consideration for local food customs and availability.

Alternative dietary assessment methods include 24-hour dietary recalls, which provide detailed information about recent intake but may not capture habitual patterns unless administered multiple times [69]. Some studies have utilized dietary history interviews or food diaries. Regardless of the specific instrument, key requirements for DII calculation include comprehensiveness (covering all DII parameters of interest), appropriate quantification of portion sizes, and validation against biomarkers or other objective measures when possible.

### Computational Tools and Statistical Approaches

Calculating DII scores requires specialized computational approaches that implement the multi-step standardization and scoring algorithm. While no single software package dominates DII research, studies commonly use statistical programming environments like R or specialized nutritional analysis software that can be customized to implement the DII algorithm [68]. The computation involves several distinct phases: data cleaning and processing, standardization against global reference values, transformation to centered percentiles, application of food parameter-specific effect scores, and summation across parameters.

Statistical analysis of DII-disease relationships typically employs multivariable regression models (logistic, linear, or Cox proportional hazards depending on the outcome) with comprehensive adjustment for potential confounders [70] [69] [22]. Common adjustments include age, sex, BMI, physical activity, smoking status, energy intake, and pre-existing health conditions. Mediation analysis may be employed to investigate potential biological pathways linking dietary inflammation to health outcomes [22]. More advanced approaches include non-linear dose-response analysis and stratified analyses to examine effect modification by population characteristics.




*DII Calculation and Validation Workflow*

*Table 3: Essential Research Reagents and Tools for DII Studies*

| Tool Category | Specific Examples | Application in DII Research | Technical Considerations |
|---|---|---|---|
| **Dietary Assessment** | 168-item FFQ [66], 24-hour dietary recall [69] | Capture consumption of all DII parameters | Requires validation for specific population; portion size estimation critical |
| **Biomarker Analysis** | CRP, IL-6, TNF-α [26], NLR [22] | Validate DII against inflammatory markers; mediate DII-disease pathways | Standardized protocols; consideration of biological variability |
| **Global Reference Database** | 11-population composite [26] [3] | Standardize individual intakes to global norms | Missing parameters imputed from other datasets or literature |
| **Statistical Software** | R, SAS, STATA, SPSS | Implement DII algorithm; statistical analysis of associations | Custom programming required for DII calculation |
| **Effect Score Library** | 45-parameter effect scores [26] | Weight each parameter's inflammatory contribution | Derived from 1,943 articles; weighted by study quality |

**Future Directions: Advancing DII Research and Application**

The ongoing optimization of DII methodology continues to open new research avenues. Future directions include further refinement of mathematical computation approaches, particularly regarding handling of non-normal distributions and missing data [68]. Additionally, as nutritional science evolves, periodic updates to the underlying literature base and effect scores will be necessary to incorporate new evidence about diet-inflammation relationships [3]. The development of population-specific reference ranges while maintaining global comparability represents another promising direction for enhancing the DII's precision and applicability across diverse settings.

Beyond methodological refinements, important substantive questions remain about how dietary inflammation interacts with other environmental and genetic factors to influence disease risk. Future research should investigate effect modification by factors such as microbiome composition, genetic polymorphisms in inflammatory pathways, and non-dietary environmental exposures [22]. Additionally, more prospective studies and randomized trials are needed to establish causal relationships between DII scores and health outcomes and to determine whether dietary interventions targeting DII reduction actually improve inflammatory markers and clinical endpoints [70] [69].

As the DII continues to be optimized and validated across populations and health conditions, it holds promise as both a research tool and potential clinical assessment metric. The ongoing methodological refinements summarized in this review—including mathematical improvements, energy adjustment, and population-specific considerations—collectively enhance the DII's precision and practical utility for researchers and clinicians seeking to quantify and address the inflammatory potential of diet.

## Comparative Analysis of Dietary Inflammatory Indices and Validation Across Populations

In nutritional epidemiology, the accurate assessment of diet's impact on health outcomes requires robust, validated tools. The **Dietary Inflammatory Index (DII)** and its derivatives—the **Energy-Adjusted DII (E-DII)** and **Empirical DII (EDII)**—represent structured approaches to quantify the inflammatory potential of an individual's diet [71] [72]. In parallel, the **Healthy Eating Index-2015 (HEI-2015)** serves as a benchmark for diet quality based on adherence to the Dietary Guidelines for Americans [73] [74]. Understanding the comparative performance, methodological foundations, and applications of these indices is crucial for researchers investigating diet-disease relationships, particularly within the context of inflammatory pathways. This guide provides an objective comparison of these indices, supported by experimental data and detailed methodologies from recent scientific investigations.

### Index Conceptual Frameworks and Design Principles

The dietary indices compared in this guide are founded on distinct conceptual frameworks and are designed for specific assessment purposes.

*Table 1: Core Characteristics of Dietary Assessment Indices*

| Index Name | Primary Purpose | Theoretical Foundation | Score Interpretation | Key Components |
|---|---|---|---|---|
| **DII** | Assess inflammatory potential of diet | Literature linking 45 food parameters to 6 inflammatory biomarkers [71] | Higher scores = more pro-inflammatory | Nutrients and food components affecting inflammatory biomarkers |
| **E-DII** | Assess inflammatory potential, energy-adjusted | Same as DII, but parameters expressed per 1000 kcal [72] | Higher scores = more pro-inflammatory | Energy-adjusted versions of DII parameters |
| **HEI-2015** | Measure diet quality against guidelines | Dietary Guidelines for Americans 2015-2020 [74] | Higher scores = better diet quality | 13 components: fruits, vegetables, whole grains, dairy, etc. |

The **DII** is a literature-derived index developed through systematic review of nearly 2,000 research articles examining associations between dietary parameters and inflammatory biomarkers [71]. It was designed to categorize an individual's diet on a continuum from maximally anti-inflammatory to maximally pro-inflammatory. The **E-DII** applies the same underlying algorithm but with all dietary parameters energy-adjusted, enabling better comparison across individuals with different caloric intakes [72]. In contrast, the **HEI-2015** evaluates compliance with federal dietary recommendations through 13 dietary components that reflect different food groups and dietary recommendations [74].




*Index Calculation and Validation Pathways: This diagram illustrates the distinct methodological pathways for calculating DII, E-DII, and HEI-2015 scores from dietary data and their relationship to different validation endpoints.*

## Methodological Protocols and Computational Approaches

### DII and E-DII Calculation Methodology

The computational protocol for DII and E-DII follows a standardized multi-step process derived from the original development work by Shivappa et al. [71]:

- **Dietary Assessment**: Dietary intake data is collected typically through 24-hour recalls or Food Frequency Questionnaires (FFQs). For NHANES-based studies, two 24-hour dietary recalls are used, with the first conducted in-person at a Mobile Examination Center and the second via telephone 3-10 days later [71] [75].
- **Parameter Standardization**: Each dietary parameter is standardized to a global reference database by calculating a z-score: `z = (actual intake - global mean)/global standard deviation`.
- **Centering to Percentiles**: The z-score is converted to a centered percentile score to minimize the effect of right skewing: `centered percentile = (2 * cumulative distribution function(z) - 1) * 1 + 0)`.
- **Inflammatory Effect Multiplication**: The centered percentile score is multiplied by the respective food parameter's inflammatory effect score (derived from literature review) to obtain the food parameter-specific DII score.
- **Aggregation**: All food parameter-specific DII scores are summed to create the overall DII score for each participant [71].

The E-DII follows identical procedures except that the reference database is energy-adjusted, with each parameter expressed per 1000 kilocalories [72]. The DII calculation formula can be summarized as: $\mathbf{DII = b_1 * n_1 + b_2 * n_2 \ldots + b_n * n_n}$, where b refers to the literature-derived inflammatory effect score for each food parameter and n refers to the food parameter-specific percentiles [71].

### HEI-2015 Calculation Methodology

The HEI-2015 computational protocol follows these key steps:

- **Dietary Data Processing**: Food intake data from 24-hour recalls are processed through the USDA Food Patterns Equivalents Database to convert consumed foods into equivalent amounts of dietary components [74].
- **Component Scoring**: Thirteen dietary components are scored on a density basis per 1000 calories (except fatty acid ratio):
  - **Adequacy Components** (higher intake yields higher score): Total fruits (0-5 points), whole fruits (0-5), total vegetables (0-5), greens and beans (0-5), whole grains (0-10), dairy (0-10), total protein foods (0-5), seafood and plant proteins (0-5), fatty acids ratio (0-10)
  - **Moderation Components** (lower intake yields higher score): Refined grains (0-10), sodium (0-10), added sugars (0-10), saturated fats (0-10) [74]
- **Score Aggregation**: All component scores are summed to create a total score ranging from 0 to 100, with higher scores indicating better diet quality [74].

## Comparative Performance Analysis

### Statistical Association with Health Outcomes

Recent studies provide direct comparative data on the performance of these indices in predicting various health conditions.

*Table 2: Performance Comparison Across Health Conditions*

| Health Condition | Index | Association Strength | Population | Study Reference |
|---|---|---|---|---|
| **Hypertension** | DII | Significant positive association (OR increment with DII increase) [71] | 45,023 US adults (NHANES) | [71] |
| **Rheumatoid Arthritis** | DII | OR: 2.99 (95% CI: 1.08-8.24) for highest vs. lowest tertile [76] | 150 Iranian adults | [76] |
| **Rheumatoid Arthritis** | HEI-2015 | OR: 0.33 (95% CI: 0.12-0.87) for highest vs. lowest tertile [76] | 150 Iranian adults | [76] |
| **Frailty** | E-DII | HR: 1.05 (95% CI: 1.01-1.08) per 1-point increase [77] | 15,249 US adults (NHANES) | [77] |
| **Frailty** | HEI-2015 | HR: 0.93 (95% CI: 0.89-0.97) per 10-point increase [77] | 15,249 US adults (NHANES) | [77] |
| **Mortality** | E-DII | HR: 1.05 (95% CI: 1.01-1.08) per 1-point increase [78] | 15,249 US adults (NHANES) | [78] |
| **Mortality** | HEI-2015 | HR: 0.93 (95% CI: 0.89-0.97) per 10-point increase [78] | 15,249 US adults (NHANES) | [78] |
| **Depression** | HEI-2015 | OR: 0.66 (95% CI: 0.50-0.87) for Q4 vs. Q1 [75] | 11,091 US adults (NHANES) | [75] |
| **Depression** | DII | No significant association [75] | 11,091 US adults (NHANES) | [75] |

### Biomarker Correlation and Validation

Validation against inflammatory biomarkers provides critical evidence for the physiological relevance of these indices:

- **HEI-2015 and Inflammatory Biomarkers**: In a cross-sectional study of 1,989 middle-to-older aged adults, higher HEI-2015 scores were significantly associated with lower CRP concentrations ($\beta = -0.06$, $p < 0.05$) and interleukin-6 (IL-6) levels ($\beta = -0.04$, $p < 0.05$) after adjusting for age and sex. These associations persisted for CRP and white blood cell counts in fully adjusted models [74].
- **DII and Inflammatory Biomarkers**: The DII has been validated against multiple inflammatory markers including CRP, TNF-α, IL-1β, IL-6, and IL-10 in previous studies [71]. A study on overweight and obese Iranian women found significant associations between E-DII and quality of life measures, though not with hs-CRP levels in this particular population [79].
- **Comparative Variance Explanation**: Analysis of NHANES 2015-2018 data demonstrated that E-DII explains approximately 52% of the variance in HEI-2015 scores in both adjusted and unadjusted models, whereas standard DII explains only 17-26% of the variance [72]. This suggests E-DII has stronger alignment with diet quality as measured by HEI-2015.

## Inter-index Correlations and Comparative Strengths

The relationship between inflammatory indices and diet quality indices reveals important insights for researchers:

- **DII/HEI-2015 Inverse Relationship**: A consistent inverse association exists between DII and HEI-2015 scores, with lower DII scores (anti-inflammatory) associated with higher HEI-2015 scores (better diet quality) [73]. This relationship was particularly strong for E-DII, where the difference between HEI-2015 scores for the first versus fourth quartile of E-DII scores was -25.90, significantly larger than the comparable difference for standard DII (-15.64) [72].
- **Energy Adjustment Impact**: The energy adjustment in E-DII appears to enhance the index's performance. In one analysis, HEI-2015 scores were significantly higher in the first E-DII quartile (most anti-inflammatory) compared to the first DII quartile, suggesting E-DII better discriminates dietary patterns with genuinely lower inflammatory potential [72].

- **Gender-Specific Patterns**: The DII-HEI relationship shows gender variations, with males in the "Low Inflammation-Low Energy" group having significantly higher HEI scores (63.58) compared to the "High Inflammation-High Energy" group (38.99, $\beta = 24.51$, $p < 0.0001$). For females, those in the "Low Inflammation-Moderate Energy" group had significantly higher HEI scores (65.08) compared to the "High Inflammation-High Energy" group (38.83, $\beta = 26.95$, $p < 0.0001$) [73].

## Research Reagent Solutions and Essential Materials

*Table 3: Essential Research Materials for Dietary Index Implementation*

| Research Reagent/Material | Function in Dietary Assessment | Application Context | Specification Requirements |
|---|---|---|---|
| **24-Hour Dietary Recall Protocol** | Collect detailed dietary intake data | Primary data collection for all indices | Standardized interview protocol, multiple passes, estimation aids |
| **Food Frequency Questionnaire (FFQ)** | Assess habitual dietary intake over time | DII/E-DII calculation in observational studies | Validated for population, 100+ food items, portion size images |
| **USDA Food Patterns Equivalents Database** | Convert foods to dietary components for HEI-2015 | HEI-2015 calculation | Current version compatible with dietary data structure |
| **Global Nutrient Database** | Standardize dietary parameters for DII calculation | DII/E-DII reference comparison | Representative mean and standard deviation for 45 parameters |
| **Inflammatory Biomarker Assays** | Validate index against physiological measures | DII validation studies | CRP, IL-6, TNF-α, IL-1β, IL-4, IL-10 measurements |
| **Nutrition Analysis Software** | Process dietary data and calculate nutrient intake | All indices | NUTRITIONIST IV, FFQ Software, ASA24-based systems |

The comparative analysis of DII, E-DII, and HEI-2015 reveals distinct strengths and applications for each index. The **E-DII** demonstrates superior performance in explaining variance in diet quality and appears more sensitive to energy-adjusted dietary patterns, making it particularly valuable for studies where caloric intake varies substantially across participants. The **standard DII** remains a validated tool for assessing inflammatory potential, with demonstrated associations across multiple health conditions. The **HEI-2015** provides the strongest measure of overall diet quality according to established dietary guidelines and shows consistent associations with health outcomes, though it may not specifically capture inflammatory pathways.

For researchers designing studies on diet-inflammatory disease relationships, using E-DII alongside HEI-2015 provides complementary information on both inflammatory potential and overall diet quality. The consistent inverse relationship between these indices confirms that better diet quality generally translates to lower inflammatory potential, while also highlighting that they capture distinct aspects of the diet-health relationship. Future research should continue to validate these indices against expanded biomarker panels and across diverse populations to further refine their utility in nutritional science and chronic disease prevention.

In nutritional epidemiology, dietary indices have emerged as essential tools for quantifying the relationship between diet and health, particularly concerning chronic inflammation. Chronic low-grade inflammation is a known precursor to numerous diseases, including cardiovascular disease, type 2 diabetes, osteoporosis, and cancer [6]. To assess the inflammatory potential of diet, researchers have developed several indices, each with distinct methodologies and applications. The **Empirical Dietary Inflammatory Index (EDII)** and the **Dietary Inflammatory Index (DII)** were pioneering in this field, but limitations remained, including nutrient-focused approaches and reliance on single inflammatory biomarkers [80] [81].

The **Empirical Anti-inflammatory Diet Index (eADI)** represents a novel advancement designed to overcome these limitations. Developed through rigorous statistical methodology and validated against multiple inflammatory biomarkers, the eADI aims to provide a more robust, food-based tool for predicting diet-related inflammation [6]. This guide provides a comprehensive comparison of the eADI against established indices, detailing its experimental validation, performance metrics, and practical applications for researchers and healthcare professionals engaged in nutritional science and chronic disease prevention.

**Table 1: Key Characteristics of Major Dietary Inflammatory Indices**

| Index Name | Abbreviation | Primary Focus | Component Basis | Number of Components | Key Biomarkers for Development | Scoring Interpretation |
|---|---|---|---|---|---|---|
| Empirical Anti-inflammatory Diet Index | eADI-17 | Anti-inflammatory potential | Food groups | 17 (11 anti-, 6 pro-inflammatory) | hsCRP, IL-6, TNF-R1, TNF-R2 | Higher score = stronger anti-inflammatory effect |

| Index Name | Abbreviation | Primary Focus | Component Basis | Number of Components | Key Biomarkers for Development | Scoring Interpretation |
|---|---|---|---|---|---|---|
| Empirical Dietary Inflammatory Index | EDII | Inflammatory potential | Food groups | 28 (12 anti-, 16 pro-inflammatory) | CRP, IL-6, TNF-α | Higher score = more pro-inflammatory |
| Dietary Inflammatory Index | DII | Inflammatory potential | Nutrients/foods | 45 (maximum) | Literature review of 6 cytokines | Higher score = more pro-inflammatory |
| Healthy Eating Index-2010 | HEI-2010 | Adherence to dietary guidelines | Food groups/nutrients | 13 | USDA Dietary Guidelines | Higher score = healthier diet |
| Alternative Healthy Eating Index-2010 | AHEI-2010 | Chronic disease prevention | Food groups/nutrients | 11 | Foods/nutrients predictive of chronic disease | Higher score = healthier diet |

The **Empirical Dietary Inflammatory Index (EDII)** was developed as a food-based alternative to nutrient-focused indices, incorporating 28 food groups (12 anti-inflammatory and 16 pro-inflammatory) based on their associations with inflammatory biomarkers including CRP, IL-6, and TNF-α [80]. The **Dietary Inflammatory Index (DII)** takes a different approach, scoring individuals based on up to 45 food parameters with pro- or anti-inflammatory properties determined through extensive literature review [2]. While these indices have proven valuable in predicting various health outcomes, limitations persist, including the EDII's development on a limited biomarker panel and the DII's complex calculation method [6].

The **Healthy Eating Index (HEI)** and **Alternative Healthy Eating Index (AHEI)** represent complementary approaches focused on overall diet quality rather than inflammation specifically. The HEI measures adherence to the Dietary Guidelines for Americans, while the AHEI incorporates additional food and nutrients predictive of chronic disease risk [82]. Research has demonstrated that both indices are associated with reduced chronic disease risk, with the AHEI sometimes showing stronger predictive ability for conditions like coronary heart disease and diabetes [83] [82].

## Experimental Validation of the eADI

### Study Design and Population

The development and validation of the eADI followed a rigorous cross-sectional design using data from the Cohort of Swedish Men-Clinical (COSM-CS) [6]. The study population consisted of **4,432 men** with a mean age of 74 ± 6 years, a demographic particularly relevant for inflammation research given the established relationship between aging and elevated inflammatory markers (inflammaging). Participants were randomly divided into two groups: a **Discovery group** (n = 2,216) for initial index development and a **Replication group** (n = 2,216) for validation purposes [6].

Key exclusion criteria included hsCRP levels >20 mg/L (suggestive of acute infection or intensive inflammatory processes), missing data on inflammatory biomarkers or education status, and implausible energy intake values. This selective approach ensured the study focused specifically on low-grade chronic inflammation rather than acute inflammatory responses [6].

### Biomarker Assessment

A key innovation in the eADI development was the use of multiple inflammatory biomarkers to capture different aspects of the immune response:

- **High-sensitivity C-reactive protein (hsCRP)**: Measured using an Architect Ci8200 analyzer with high-sensitivity latex enhanced immunonephelometric assay
- **Interleukin-6 (IL-6)**: Determined using panels from Olink Proteomics
- **Tumor necrosis factor receptor 1 (TNF-R1)**: Measured via Olink Proteomics panels
- **Tumor necrosis factor receptor 2 (TNF-R2)**: Assessed using Olink Proteomics methodology

All biomarker concentrations were log2-transformed for analysis, with IL-6, TNF-R1, and TNF-R2 presented as Normalized Protein Expression (NPX) values in arbitrary units on a log2 scale [6]. This multi-biomarker approach addressed a significant limitation of previous indices that relied on single biomarkers, particularly CRP alone.

### Dietary Assessment and Food Group Selection

Dietary intake was assessed using a **145-item food frequency questionnaire (FFQ)** that asked participants to indicate their consumption frequency across eight predefined categories ranging from "never/seldom" to "≥3 times per day" [6]. The FFQ had been previously validated in a

subset of 248 men using 14 repeated 24-hour recall interviews, demonstrating strong correlation coefficients for energy-adjusted macro- and micronutrient intake (0.65 and 0.62, respectively) [6].

The food selection process for the eADI employed sophisticated statistical approaches:

- **10-fold feature selection with filtering** based on Lasso regression to identify food groups most strongly correlated with the inflammatory biomarkers
- Food groups were categorized as either anti-inflammatory or pro-inflammatory based on their association direction with the biomarkers
- The final eADI-17 included **17 food groups**: 11 with anti-inflammatory potential and 6 with pro-inflammatory potential [6]

#### Index Scoring System

The eADI scoring system was designed for practical application:

- Consumption levels for each food group were categorized into tertiles
- For anti-inflammatory foods, the highest consumption tertile received 1 point, middle tertile 0.5 points, and lowest tertile 0 points
- For pro-inflammatory foods, the scoring was reversed: lowest consumption received 1 point, middle tertile 0.5 points, and highest tertile 0 points
- All points were summed to create a total eADI score ranging from 0-17, with higher scores indicating stronger anti-inflammatory potential [6]

This straightforward scoring system enhances the index's utility in both research and clinical settings for personalized nutrition advice.

### Performance Comparison of Dietary Indices

#### Association with Inflammatory Biomarkers

**Table 2: Comparative Performance of Dietary Indices Against Inflammatory Biomarkers**

| Dietary Index | hsCRP Association | IL-6 Association | TNF-R1 Association | TNF-R2 Association | Other Reported Associations |
|---|---|---|---|---|---|
| **eADI-17** | -0.17 (Spearman correlation)12% lower per 4.5-point increment | -0.23 (Spearman correlation)6% lower per 4.5-point increment | -0.28 (Spearman correlation)8% lower per 4.5-point increment | -0.26 (Spearman correlation)9% lower per 4.5-point increment | Robust across discovery and replication cohorts |
| **EDII** | Positive association(pro-inflammatory diet) | Positive association(pro-inflammatory diet) | Not specifically reported | Not specifically reported | Associated with anxiety disorders (OR: 2.09 for top vs bottom category) [80] |
| **DII** | Variable across studies | Variable across studies | Not specifically reported | Not specifically reported | Associated with oxidative stress (lipoperoxidation) [81] |
| **HEI-2010** | Limited direct association | Limited direct association | Not specifically reported | Not specifically reported | Associated with Faecalibacterium prausnitzii levels [81] |
| **AHEI-2010** | Limited direct association | Limited direct association | Not specifically reported | Not specifically reported | Associated with fecal SCFAs [81] |

The eADI-17 demonstrates consistent, significant inverse correlations with all four inflammatory biomarkers assessed in the validation study [6]. In the Replication group, each 4.5-point increment (2 standard deviations) in eADI-17 score was associated with substantially lower concentrations of all measured biomarkers: 12% lower for hsCRP, 6% lower for IL-6, 8% lower for TNF-R1, and 9% lower for TNF-R2 [6]. The robustness of these findings was confirmed by nearly identical results in both the Discovery and Replication groups.

Comparative studies of other indices show more variable performance. The EDII has demonstrated significant associations with inflammatory biomarkers and mental health outcomes, with one study reporting that individuals in the highest EDII category had 2.09 times greater odds of anxiety disorders compared to those in the lowest category [80]. The DII has shown associations with oxidative stress parameters, while the HEI and AHEI appear more strongly linked to gut microbiota composition and metabolic health than direct inflammatory markers [81].

#### Predictive Validity for Health Outcomes

While the eADI is newly developed and prospective health outcome data are limited, other indices have established predictive validity:

- The AHEI-2010 has demonstrated stronger associations with chronic disease risk reduction compared to the HEI-2010, particularly for coronary heart disease and diabetes [83]
- In a study comparing HEI-2010 and AHEI-2010 in relation to diabetes status, adults with type 2 diabetes showed higher scores on both indices compared to adults with prediabetes and without diabetes, though neither index was clearly superior in predicting diabetes status [82]
- The DII has been associated with metabolic syndrome risk and related conditions across multiple studies, though results have sometimes been inconsistent [2]

The multi-biomarker foundation of the eADI suggests potential for strong predictive validity regarding inflammation-related health outcomes, though prospective studies are needed to confirm this potential.

## Research Reagent Solutions Toolkit

**Table 3: Essential Research Materials and Methods for Dietary Index Validation**

| Reagent/Instrument | Specification/Model | Primary Function | Key Applications in Validation |
|---|---|---|---|
| **High-Sensitivity CRP Assay** | Architect Ci8200 analyzer with 6K2601 immunonephelometric assay (Abbott Laboratories) | Quantification of hsCRP with high sensitivity | Assessment of low-grade inflammation; correlation with dietary patterns |
| **Multiplex Inflammation Panel** | Olink Proteomics CVD II and CVD III panels | Simultaneous measurement of multiple inflammatory biomarkers | Detection of IL-6, TNF-R1, TNF-R2 using NPX technology |
| **Food Frequency Questionnaire** | 145-item FFQ (COSM-CS version) | Comprehensive dietary assessment | Estimation of habitual intake of food groups for index calculation |
| **Biobanking Storage System** | -80°C freezer systems | Preservation of biological samples | Maintenance of sample integrity for batch analysis of biomarkers |
| **Dietary Analysis Software** | Nutritionist IV software (adapted for local foods) | Nutrient calculation from FFQ data | Energy adjustment and food group quantification |
| **Statistical Software** | R, SPSS, or similar packages | Advanced statistical analysis | Performance of Lasso regression, correlation analysis, and linear regression |

## Experimental Workflow for Index Development and Validation

The development and validation of dietary indices like the eADI follows a systematic process that can be visualized as follows:




*Diagram 1: Experimental workflow for eADI development and validation*

## Methodological Considerations and Limitations

The validation of any dietary index requires careful consideration of methodological limitations. For the eADI, several aspects deserve attention:

The **cross-sectional design** of the validation study prevents establishment of causal relationships between dietary patterns and inflammatory status [6]. While the robust discovery-replication approach strengthens the findings, prospective studies are needed to confirm predictive validity for health outcomes.

The **population homogeneity** (exclusively older Swedish men) limits generalizability to other demographic groups, including women, younger populations, and diverse ethnicities [6]. The authors explicitly note the need for validation in more diverse populations.

The **dietary assessment method** (FFQ) introduces potential measurement error, though the instrument was previously validated and demonstrated reasonable accuracy for nutrient estimation [6].

Despite these limitations, the eADI represents a significant methodological advancement through its **multi-biomarker approach** and **food-based scoring system**, addressing key limitations of previous indices that relied on single biomarkers or nutrient-based approaches [6].

The Empirical Anti-inflammatory Diet Index (eADI) represents a validated, robust tool for assessing the inflammatory potential of diet using a food-based approach. Its development against multiple inflammatory biomarkers addresses a significant limitation of previous indices and enhances its biological plausibility. The eADI demonstrates consistent, significant associations with key inflammatory markers including hsCRP, IL-6, TNF-R1, and TNF-R2, with higher scores predicting substantially lower concentrations of all measured biomarkers.

For researchers and healthcare professionals, the eADI offers a practical tool with clear scoring criteria suitable for both research and clinical applications. Its food-group basis facilitates translation into dietary recommendations, potentially refining future dietary guidelines and enhancing personalized nutrition approaches for inflammation reduction [6].

While the eADI shows promise, further validation in diverse populations and prospective studies examining hard health endpoints will strengthen its utility. As research in nutritional immunology advances, the eADI provides a valuable addition to the methodological toolkit for quantifying diet-inflammation relationships and developing evidence-based dietary interventions for chronic disease prevention and management.

The Dietary Inflammatory Index (DII) was developed to quantify the inflammatory potential of an individual's overall diet, moving beyond the limitations of analyzing single foods or nutrients in isolation [59]. However, a significant limitation of the original DII is that its reference database was primarily constructed using dietary intake data from Western populations [59]. This poses a challenge for its application in non-Western contexts like China, where dietary cultures, food composition, and eating habits differ substantially. To address this lack of population-specific tools, researchers developed the China Dietary Inflammatory Index (CHINA-DII), an index tailored specifically to reflect the dietary patterns and inflammatory impacts relevant to Chinese adults [59]. This guide objectively compares the CHINA-DII with other DII variants and details its development, validation, and application, providing a resource for researchers and drug development professionals working in nutritional epidemiology and chronic disease prevention.

## Comparative Analysis of Dietary Inflammatory Indices

The development of the CHINA-DII represents an evolution in the field of dietary inflammation assessment. The table below compares its key features with those of the original DII and its standard application in Chinese research.

*Table 1: Comparison of the Original DII and the CHINA-DII*

| Feature | Original DII (Shivappa et al., 2014) | CHINA-DII (2025) | Standard DII in Chinese Studies |
|---|---|---|---|
| **Development Basis** | 45 dietary components; global intake data from 11 countries [59]. | 27 dietary components; intake data from Chinese adult populations [59]. | Applies the original DII algorithm to dietary data from Chinese cohorts [84]. |
| **Target Population** | Designed for global application, with a Western-centric reference database [59]. | Specifically developed and validated for Chinese adults [59]. | Used to study Chinese populations but not tailored to their specific dietary patterns. |
| **Key Validated Associations** | Associated with CVD (e.g., MI in men) [85], inflammatory biomarkers (hs-CRP, IL-6) [85]. | Positively associated with hs-CRP and gastric cancer risk in Chinese cohorts [59] [86]. | Associated with hypertension and geographic disparities in inflammation among Chinese middle-aged/elderly [84]. |
| **Primary Utility** | Assessing dietary inflammatory potential across diverse, international populations. | Evaluating dietary inflammation and its disease links within the context of Chinese dietary culture. | Investigating diet-inflammation-disease pathways in China using a standardized, international tool. |

The CHINA-DII was developed to directly address the geographic and cultural dietary gap. A nationwide spatial analysis of DII scores in China revealed significant disparities, with higher (more pro-inflammatory) scores concentrated in the northwestern region and lower (more anti-inflammatory) scores in the southeastern region [84]. This geographic pattern underscores the need for a population-specific tool that can accurately capture such local variations in dietary inflammatory potential.

## Development and Validation of the CHINA-DII

### Development Protocol

The construction of the CHINA-DII followed a rigorous, multi-stage process to ensure its validity and relevance for the target population.

*Table 2: Key Stages in the CHINA-DII Development Protocol*

| Stage | Description | Outcome |
|---|---|---|
| **1. Literature Search & Screening** | A systematic search was conducted in five databases (CNKI, Wanfang, VIP, PubMed, Web of Science) for studies published between 2009 and 2024 | Identification of 33 eligible studies that met the inclusion criteria [59]. |

| Stage | Description | Outcome |
|---|---|---|
| | reporting dietary intake in Chinese adults [59]. | |
| **2. Database Establishment** | Dietary data from the included literature were extracted and synthesized. | Creation of a standardized dietary intake database for Chinese adults, encompassing 27 dietary components [59]. |
| **3. Index Calculation** | The CHINA-DII score was calculated for each individual based on their dietary data, following the statistical methodology of the original DII but using the newly established Chinese reference intake database [59]. | Each individual receives a CHINA-DII score, where a higher score indicates a more pro-inflammatory diet [59]. |

**Validation against Inflammatory Biomarkers**

The validity of the CHINA-DII was assessed by examining its relationship with high-sensitivity C-reactive protein (hs-CRP), a well-established blood biomarker of systemic inflammation.

- **Study Population**: The validation study recruited 256 newly diagnosed gastric cancer patients from a university hospital in China [59].
- **Methods**: Demographic, clinical, and dietary data (collected via a food frequency questionnaire) were obtained from participants. The CHINA-DII score was calculated for each participant, and its correlation with hs-CRP levels was analyzed using Spearman rank correlation and multivariate logistic regression [59].
- **Key Findings**:
  - The average CHINA-DII score was -1.91, and the mean hs-CRP level was 3.68 mg/L [59].
  - A statistically significant positive correlation was found between CHINA-DII scores and hs-CRP levels ($r = 0.20$, $p \leq 0.001$) [59].
  - Individuals with higher CHINA-DII scores had a 1.90-fold increased risk of having elevated hs-CRP (≥3 mg/L) compared to those with lower scores (OR = 1.90; 95% CI: 1.01–3.55). For each one-standard-deviation increase in the CHINA-DII score, the risk of elevated hs-CRP increased by 1.50 times (OR = 1.50, 95% CI: 1.10–2.06) [59].

This robust correlation with a objective inflammatory biomarker confirms that the CHINA-DII is a valid tool for reflecting the inflammatory impact of diet in Chinese adults.




*Figure 1: Workflow for the development and biomarker validation of the CHINA-DII.*

## Application in Disease Association Studies

The CHINA-DII has been applied in epidemiological research to investigate the link between pro-inflammatory diets and chronic diseases, particularly gastric cancer (GC), in the Chinese population.

### Association with Gastric Cancer Risk

A case-control study conducted in Southeastern China exemplifies the application of the CHINA-DII in etiological research.

- **Study Design**: A 1:1 sex-matched case-control study involving 336 newly diagnosed GC cases and 336 healthy controls from Fujian Province [86].
- **Dietary Assessment**: Dietary intake was collected using a structured, semi-quantitative Food Frequency Questionnaire (FFQ) covering 78 food items [86].
- **Key Findings**:
  - Higher CHINA-DII scores (indicating a more pro-inflammatory diet) were significantly associated with an increased risk of gastric cancer (OR = 1.45, 95% CI: 1.05–1.99) [86].
  - Each one-standard-deviation increase in the CHINA-DII score was associated with a 1.26-fold increase in GC risk (OR = 1.26, 95% CI: 1.07–1.48) [86].
  - The study also found that higher intakes of the anti-inflammatory nutrients vitamin C and vitamin D were significantly associated with a lower GC risk [86].

These findings highlight the value of the CHINA-DII as a tool for identifying dietary patterns that contribute to cancer risk in specific populations.

### Subgroup Analysis Reveals Vulnerable Populations

The gastric cancer study further performed subgroup analyses, which revealed that the association between a pro-inflammatory diet and disease risk was not uniform across the population. The effect was more pronounced in certain subgroups, suggesting they may be more susceptible to the adverse effects of dietary inflammation [86].

*Table 3: Subgroup Analysis of CHINA-DII and Gastric Cancer Risk*

| Subgroup | Adjusted Odds Ratio (OR) | 95% Confidence Interval (CI) |
|---|---|---|
| **Age ≤ 55 years** | 2.44 | 1.51 – 3.96 |
| **Married population** | 1.41 | 1.01 – 1.96 |
| **Non-smokers** | 1.70 | 1.14 – 2.54 |
| **High perceived daily stress** | 2.82 | 1.67 – 4.75 |

The data indicates that younger individuals, non-smokers, and those experiencing high daily stress exhibit a stronger association between dietary inflammation and gastric cancer risk. This underscores the potential for the CHINA-DII to help identify high-risk populations who would benefit most from targeted dietary interventions.




*Figure 2: The conceptual pathway linking a pro-inflammatory diet to chronic disease risk, as measured by the CHINA-DII, and the role of effect modifiers.*

## The Scientist's Toolkit: Key Research Reagents and Materials

Successfully conducting research with the CHINA-DII requires a specific set of methodological tools and reagents. The following table details the essential components used in the featured validation and application studies.

*Table 4: Essential Research Reagents and Materials for CHINA-DII Studies*

| Item | Specification / Function | Application in CHINA-DII Research |
|---|---|---|
| **Food Frequency Questionnaire (FFQ)** | A validated, semi-quantitative questionnaire designed for Chinese dietary patterns, often containing ~78 food items across 13 categories [86]. | To collect detailed data on habitual dietary intake over the past year, which is the foundation for calculating the CHINA-DII score [59] [86]. |
| **Chinese Food Composition Table** | A comprehensive database providing nutrient profiles for over 2,000 foods and ingredients commonly consumed in China [84]. | To convert reported food consumption frequencies and portion sizes from the FFQ into daily intakes of the specific nutrients required for CHINA-DII calculation [84]. |
| **High-Sensitivity C-Reactive Protein (hs-CRP) Assay** | An immunoassay kit for the quantitative measurement of low levels of CRP in serum or plasma, with high precision. | To measure systemic inflammation as an objective biomarker for validating the CHINA-DII score [59]. |
| **Statistical Analysis Software (e.g., R, SAS, SPSS)** | Software packages capable of performing Spearman rank correlation, multivariate logistic regression, and other advanced statistical models. | To analyze the relationship between CHINA-DII scores, inflammatory biomarkers, and disease outcomes, while adjusting for potential confounders like age, sex, and BMI [59] [86]. |

The development and validation of the CHINA-DII mark a significant advancement in the field of nutritional epidemiology in China. By building a dietary index based on local consumption data and validating it against robust inflammatory biomarkers and hard clinical endpoints like gastric cancer, researchers have created a tool with demonstrated validity and applicability for the Chinese population. The evidence shows that the CHINA-DII is not only a significant predictor of systemic inflammation but is also positively associated with the risk of major chronic diseases. Its ability to identify vulnerable subgroups, such as younger adults and individuals under high stress, provides a powerful target for public health initiatives aimed at reducing the burden of inflammation-related diseases through precision nutrition. For the research community, the CHINA-DII offers a superior, population-specific instrument for investigating the intricate links between diet, inflammation, and health in China.

This comparative guide provides a systematic analysis of the Dietary Inflammatory Index (DII) and Healthy Eating Index-2015 (HEI-2015) as tools for assessing diet's impact on inflammatory markers. While both indices evaluate dietary patterns, they approach assessment from distinct perspectives: the DII specifically quantifies diet's inflammatory potential based on established literature, whereas HEI-2015 measures adherence to U.S. dietary guidelines. Evidence from large observational studies and systematic reviews demonstrates that both tools independently predict inflammatory biomarker levels, with pro-inflammatory DII scores and lower HEI-2015 scores associated with elevated CRP, IL-6, white blood cell count, and other inflammatory indicators. Crucially, joint effects analysis reveals that high-quality diets can counteract pro-inflammatory diets, whereas anti-inflammatory diets alone cannot compensate for poor overall diet quality. This guide synthesizes experimental data, methodological protocols, and mechanistic pathways to inform research applications in nutritional epidemiology and chronic disease prevention.

### Dietary Inflammatory Index (DII): Concept and Development

The Dietary Inflammatory Index (DII) is a literature-derived, population-based scoring system designed specifically to quantify the inflammatory potential of an individual's diet [3] [1]. Developed through a comprehensive review of peer-reviewed research articles published between 1950 and 2010, the DII links dietary components to six established inflammatory biomarkers: IL-1β, IL-4, IL-6, IL-10, TNF-α, and CRP [3] [1]. The index was created to address a significant gap in nutritional epidemiology—the lack of a standardized, evidence-based tool specifically focused on diet-induced inflammation. The theoretical bounds of the DII range from -8.87 (maximally anti-inflammatory) to +7.98 (maximally pro-inflammatory), with scores below zero indicating anti-inflammatory diets and scores above zero indicating pro-inflammatory diets [28].

The DII represents a paradigm shift from traditional dietary indices because it is not based on dietary guidelines or specific cultural foodways, but rather on empirical evidence linking dietary factors to inflammatory responses [3]. This evidence base has grown exponentially, with the original DII leveraging 1,943 qualifying articles that documented associations between dietary parameters and inflammatory markers [3]. The DII has undergone significant methodological refinements since its initial conception, including enhanced scoring algorithms that link reported dietary intake to global norms of intake from 11 populations worldwide, addressing previous limitations related to right-skewing of dietary data and arbitrary use of raw consumption amounts [3].

### Healthy Eating Index-2015 (HEI-2015): Concept and Development

The Healthy Eating Index-2015 (HEI-2015) is a diet quality index developed by the National Cancer Institute (NCI) and the U.S. Department of Agriculture (USDA) to assess alignment with the 2015-2020 Dietary Guidelines for Americans [87]. Unlike the hypothesis-driven DII, the HEI-2015 represents a normative approach to dietary assessment, measuring how well an individual's diet conforms to recommended eating patterns. The HEI-2015 consists of 13 components, including 9 adequacy components (total fruits, whole fruits, total vegetables, greens and beans, whole grains, dairy, total protein foods, seafood and plant proteins, and fatty acids) and 4 moderation components (refined grains, sodium, added sugars, and saturated fats) [87].

Each component is scored based on established standards, with the total score ranging from 0 to 100, where higher scores indicate better diet quality and closer adherence to dietary guidelines [87]. The HEI-2015 evaluates density-based standards (e.g., per 1,000 calories or as a percentage of calories) to allow for comparison across different energy intake levels. The adequacy components are scored such that higher intake yields higher scores, while the moderation components are reverse-scored, with lower intake receiving higher scores [87]. This structure captures the multidimensional nature of diet quality, addressing both sufficient consumption of beneficial food groups and limited consumption of components associated with negative health outcomes.

## Methodological Protocols

### DII Calculation Protocol

The calculation of DII scores follows a standardized protocol that can be applied across different populations and dietary assessment methods [3]. The methodological workflow involves multiple steps:

#### Step 1: Dietary Data Collection

- Collect dietary intake data using validated methods (24-hour recalls, food frequency questionnaires, or food records)
- Identify and quantify intake of up to 45 food parameters that constitute the DII, including:
  - Macronutrients: carbohydrates, protein, fat, fiber
  - Fatty acids: saturated, monounsaturated, polyunsaturated, omega-3, omega-6, trans-fat
  - Vitamins: A, B6, B12, C, D, E, thiamin, riboflavin, niacin
  - Minerals: iron, magnesium, zinc, selenium
  - Bioactive compounds: flavonoids, beta-carotene
  - Spices and food components: garlic, ginger, onions, caffeine, tea [3] [21]

#### Step 2: Standardization Against Global Intake Database

- Link reported dietary intakes to a global comparative database comprising means and standard deviations from 11 populations worldwide (including the United States, United Kingdom, Bahrain, Mexico, Australia, South Korea, Taiwan, India, New Zealand, Japan, and Denmark)
- For each food parameter, calculate a z-score by subtracting the global mean from the actual intake and dividing by the global standard deviation [3]

**Step 3: Conversion to Percentiles and Centering**

- Convert z-scores to percentiles to minimize the effect of right-skewing common in dietary data
- Center the distribution by multiplying percentiles by 2 and subtracting 1, creating values from -1 (minimally anti-inflammatory) to +1 (maximally pro-inflammatory) [3]

**Step 4: Apply Inflammatory Effect Scores and Summation**

- Multiply each centered percentile value by its corresponding inflammatory effect score derived from the literature review
- Sum across all food parameters to obtain the overall DII score [3]

**Energy Adjustment Protocol:**

- To account for variations in energy intake, researchers often calculate energy-adjusted DII (E-DII) by expressing dietary parameters per 1,000 calories consumed [21]




**HEI-2015 Calculation Protocol**

The HEI-2015 calculation follows a distinct protocol focused on assessing adherence to dietary guidelines:

**Step 1: Dietary Data Collection and Processing**

- Collect dietary intake data using 24-hour recalls, food records, or food frequency questionnaires

- Process foods and beverages using the USDA Food Patterns Equivalent Database (FPED) to convert consumed items into equivalent amounts of HEI-2015 components [87] [75]

**Step 2: Component Calculation**

- For each of the 13 components, calculate intake amounts following standardized definitions:
  - Adequacy components: calculate cup or ounce equivalents per 1,000 calories
  - Moderation components: calculate as percent of calories or ounce equivalents per 1,000 calories [87]

**Step 3: Component Scoring**

- Score each component based on established standards:
  - Adequacy components: higher intake receives higher scores, up to a maximum
  - Moderation components: lower intake receives higher scores, up to a maximum
- Apply proportional scoring for intermediate values between minimum (0) and maximum scores [87]

**Step 4: Total Score Calculation**

- Sum scores across all 13 components to obtain the total HEI-2015 score (range: 0-100) [87]

**Implementation Considerations:**

- The calculation can be performed using the "Dietaryindex" package in R or SAS software provided by the National Cancer Institute
- When multiple dietary recalls are available, average component scores across days for more stable estimates [87]

**Inflammatory Biomarker Assessment Protocols**

The validation of both DII and HEI-2015 relies on established protocols for measuring inflammatory biomarkers:

**Blood Collection and Processing:**

- Collect venous blood samples following standardized phlebotomy procedures
- Process samples within 2 hours of collection through centrifugation
- Store plasma or serum at -80°C until analysis [88] [87]

**Biomarker Assay Methods:**

- **High-sensitivity C-reactive protein (hs-CRP):** Assessed using immunoturbidimetric assays on automated clinical chemistry analyzers
- **Interleukins (IL-6, IL-1β, IL-4, IL-10):** Measured using enzyme-linked immunosorbent assays (ELISA) or multiplex immunoassays
- **Tumor Necrosis Factor-alpha (TNF-α):** Quantified via ELISA or electrochemiluminescence immunoassays
- **White blood cell (WBC) count and differential:** Analyzed using automated hematology analyzers with flow cytometry [88] [87] [4]

**Quality Control:**

- Implement standard calibration procedures using certified reference materials
- Include internal quality control samples with known concentrations in each assay batch
- For multi-center studies, standardize procedures across sites and implement cross-validation [87]

## Comparative Performance Data

### Independent Associations with Inflammatory Markers

*Table 1: Independent Associations of DII and HEI-2015 with Inflammatory Biomarkers*

| Biomarker | DII Association | HEI-2015 Association | Population | Source |
|---|---|---|---|---|
| **C-reactive Protein (CRP)** | Significant positive association ($P < 0.001$) | Significant inverse association ($P < 0.001$) | 80 PCOS patients vs. 80 controls [88] | PCOS Case-Control Study |
| **White Blood Cell (WBC) Count** | Significant positive association ($\beta = 0.08$, $P < 0.001$) | Significant inverse association ($\beta = -0.10$, $P < 0.001$) | 19,110 U.S. adults [87] | NHANES 2009-2018 |
| **Neutrophil Count** | Significant positive association ($\beta = 0.07$, $P < 0.001$) | Significant inverse association ($\beta = -0.09$, $P < 0.001$) | 19,110 U.S. adults [87] | NHANES 2009-2018 |

| Biomarker | DII Association | HEI-2015 Association | Population | Source |
|---|---|---|---|---|
| **Neutrophil-to-Lymphocyte Ratio (NLR)** | Significant positive association (β = 0.05, P < 0.001) | Significant inverse association (β = -0.06, P < 0.001) | 19,110 U.S. adults [87] | NHANES 2009-2018 |
| **Systemic Immune-Inflammation Index (SII)** | Significant positive association (β = 0.06, P < 0.001) | Significant inverse association (β = -0.07, P < 0.001) | 19,110 U.S. adults [87] | NHANES 2009-2018 |
| **Cardiovascular Disease Mortality** | Positive association (HR: 1.0514, 95% CI: 1.0055-1.0995) | Inverse association (HR: 0.9404, 95% CI: 0.8846-0.9998) | 11,310 hypertensive adults [52] | NHANES 1999-2016 |

The data consistently demonstrate that DII and HEI-2015 show opposing yet complementary relationships with inflammatory markers. Higher DII scores (pro-inflammatory diets) are consistently associated with elevated levels of multiple inflammatory biomarkers, while higher HEI-2015 scores (better diet quality) are associated with lower inflammatory biomarker levels across diverse populations [88] [87] [52]. The associations remain significant after adjustment for potential confounders including age, sex, BMI, physical activity, and socioeconomic status.

**Joint Effects Analysis**

*Table 2: Joint Effects of DII and HEI-2015 on Inflammatory Markers*

| Dietary Pattern | DII Characterization | HEI-2015 Characterization | WBC Association | Neu Association | NLR Association | SII Association |
|---|---|---|---|---|---|---|
| **Pattern 1** | Pro-inflammatory | Low quality | Reference | Reference | Reference | Reference |
| **Pattern 2** | Anti-inflammatory | Low quality | No significant reduction | No significant reduction | No significant reduction | No significant reduction |
| **Pattern 3** | Pro-inflammatory | High quality | Significant reduction | Significant reduction | Significant reduction | Significant reduction |
| **Pattern 4** | Anti-inflammatory | High quality | Significant reduction | Significant reduction | Significant reduction | Significant reduction |

Data derived from NHANES 2009-2018 analysis of 19,110 participants [87]

The joint effects analysis reveals crucial insights about the interplay between diet quality and inflammatory potential. Participants were categorized into four dietary patterns based on combined DII and HEI-2015 scores. The results demonstrate that high-quality diets (Patterns 3 and 4) significantly reduce inflammatory markers regardless of DII classification, whereas anti-inflammatory diets alone (Pattern 2) cannot compensate for poor overall diet quality [87]. This suggests that overall diet quality exerts a stronger influence on inflammatory status than the specific inflammatory potential of the diet.

**Key Component Contributions**

*Table 3: Key Dietary Components Contributing to Inflammatory Potential*

| Index | Most Influential Anti-inflammatory Components | Most Influential Pro-inflammatory Components |
|---|---|---|
| **DII** | Fiber, seafood and plant proteins, omega-3 fatty acids, flavonoids, vitamins A, C, D, E | Saturated fats, trans fats, carbohydrates, iron, energy intake |
| **HEI-2015** | Whole fruits, seafood and plant proteins, whole grains, greens and beans | Added sugars, saturated fats, refined grains, sodium |

Component analysis based on weighted quantile sum (WQS) regression and SHapley Additive exPlanations (SHAP) analysis [87] [75]

The WQS analysis of DII components revealed that dietary fiber, seafood and plant proteins, and alcohol intake were the primary contributors to inflammatory marker levels [87]. For HEI-2015, SHAP analysis identified added sugars, whole fruits, and saturated fats as the most influential components affecting depression risk (a condition with established inflammatory pathways) [75]. This component-level analysis provides insights for targeted dietary interventions.

## Biological Pathways and Mechanisms




The biological pathways linking dietary patterns to systemic inflammation involve multiple interconnected mechanisms. Pro-inflammatory diets (high DII scores) and poor-quality diets (low HEI-2015 scores) activate the NF-κB pathway, a master regulator of inflammation that increases production of inflammatory cytokines including IL-6, TNF-α, and IL-1β [1] [89]. These cytokines stimulate hepatic production of acute-phase proteins such as CRP and fibrinogen while also activating circulating leukocytes [89].

Parallel pathways involve oxidative stress generation from pro-oxidant dietary components and gut microbiota-mediated inflammation. Poor-quality diets disrupt gut microbial homeostasis, leading to increased intestinal permeability and endotoxemia, which further activates inflammatory pathways [75]. The resulting chronic low-grade systemic inflammation contributes to the pathogenesis of numerous chronic diseases, including cardiovascular diseases, diabetes, cancer, and depression [1] [89].

Anti-inflammatory diets (low DII scores) and high-quality diets (high HEI-2015 scores) counteract these processes through multiple mechanisms: (1) providing antioxidants that reduce oxidative stress; (2) supplying fiber and polyphenols that support beneficial gut microbiota; (3) offering omega-3 fatty acids that serve as precursors to anti-inflammatory resolvins and protectins; and (4) containing bioactive compounds that directly inhibit NF-κB activation and inflammatory cytokine production [87] [3] [89].

## The Researcher's Toolkit

### Essential Research Reagents and Materials

*Table 4: Essential Research Materials for Dietary Inflammation Studies*

| Category | Item | Specification/Function | Example Applications |
|---|---|---|---|
| **Dietary Assessment** | Automated 24-hour recall system | Standardized dietary data collection (e.g., USDA Automated Multiple-Pass Method) | NHANES dietary data collection [87] |
| | Food frequency questionnaire (FFQ) | Semi-quantitative assessment of habitual dietary intake | Epidemiologic cohort studies [88] |

| Category | Item | Specification/Function | Example Applications |
|---|---|---|---|
| | Food composition database | Nutrient calculation reference (e.g., FNDDS, NDB) | Nutrient analysis for DII calculation [75] |
| **Laboratory Analysis** | High-sensitivity CRP assay | Quantification of low-grade inflammation (detection limit <0.1 mg/L) | Inflammation biomarker measurement [88] [4] |
| | Multiplex cytokine array | Simultaneous measurement of multiple cytokines (IL-6, TNF-α, IL-1β, IL-10) | Comprehensive inflammatory profiling [4] [89] |
| | Hematology analyzer | Complete blood count with differential (WBC, neutrophils, lymphocytes) | Cellular inflammation assessment [87] |
| **Data Analysis** | DII calculation algorithm | Standardized computation of dietary inflammatory scores | DII score calculation [3] |
| | HEI-2015 scoring algorithm | Standardized diet quality assessment | HEI-2015 score calculation [87] |
| | Statistical software packages | Advanced statistical analysis (R, SAS, Stata) | Joint effects analysis, confounding adjustment [87] |

**Methodological Considerations for Researchers**

**Dietary Assessment Selection:**

- For DII calculation, 24-hour recalls provide more current intake data but FFQs better capture habitual intake
- Multiple dietary assessments (≥2) per participant improve reliability for both indices
- Consider using the "Dietaryindex" package in R for standardized calculation of both DII and HEI-2015 [87]

**Biomarker Selection and Measurement:**

- Include multiple inflammatory biomarkers representing different pathways (acute phase proteins, cytokines, cellular markers)
- Standardize collection, processing, and storage protocols to minimize pre-analytical variability
- Consider batch analysis with quality controls to reduce assay variability [87] [89]

**Confounding Control:**

- Adjust for established covariates: age, sex, BMI, physical activity, smoking, socioeconomic status
- Consider additional potential confounders: medication use, existing health conditions, genetic factors
- Use directed acyclic graphs (DAGs) to identify minimal sufficient adjustment sets [88] [87]

**Joint Effects Analysis Approach:**

- Categorize participants into joint exposure groups based on both DII and HEI-2015
- Include interaction terms in regression models to test for effect modification
- Consider using weighted quantile sum regression to identify component importance [87]

This comparative analysis demonstrates that both DII and HEI-2015 provide valuable, complementary information about diet's relationship with inflammatory status. The DII specifically captures dietary inflammatory potential based on established literature, while HEI-2015 assesses overall diet quality relative to dietary guidelines. The joint effects analysis reveals their interdependent nature: high-quality diets can counteract pro-inflammatory diets, but anti-inflammatory diets alone cannot compensate for poor overall diet quality [87].

For researchers investigating diet-inflammation relationships, employing both indices provides a more comprehensive assessment than either index alone. The consistent associations observed between these indices and inflammatory biomarkers across diverse populations support their construct validity and utility in nutritional epidemiology [88] [87] [52]. Future research should focus on longitudinal studies with repeated dietary assessments, investigation of these relationships in diverse populations, and intervention studies testing whether improvements in these indices directly reduce inflammation and chronic disease risk.

## Conclusion

The validation of Dietary Inflammatory Index scores establishes them as robust, scientifically-grounded tools for quantifying the inflammatory potential of diet, with significant implications for biomedical research and clinical practice. Consistent positive correlations with inflammatory biomarkers like CRP, IL-6, and TNF-α confirm their biological relevance. Key advancements include the development of population-specific indices and empirical derivatives like the eADI, which enhance predictive accuracy. Future directions should focus on standardizing validation protocols

across diverse global populations, longitudinal studies to establish causal relationships with hard clinical endpoints, and integrating these indices into personalized nutrition and pharmaceutical interventions to modulate chronic inflammation effectively.

## References


- 1. sciencedirect.com/topics/nursing-and-health-professions/ dietary ... [https://www.sciencedirect.com/topics/nursing-and-health-professions/dietary-inflammatory-index]
- 2. Scientific basis of dietary inflammatory index (DII): A dietary tool to metabolic syndrome risk - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S2667268525000415]
- 3. Perspective: The Dietary (DII)—Lessons Learned... Inflammatory Index [https://pmc.ncbi.nlm.nih.gov/articles/PMC6416047/]
- 4. Development and Validation of an Empirical Dietary Inflammatory Index - PubMed [https://pubmed.ncbi.nlm.nih.gov/27358416/]
- 5. Development and Validation of an Empirical Dietary Inflammatory Index - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC4958288/]
- 6. Development of empirical anti-inflammatory diet index: a cross-sectional study | Nutrition Journal | Full Text [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-025-01165-x]
- 7. Development and Validation of the China Dietary Inflammatory Index (CHINA-DII) [https://www.mdpi.com/2072-6643/17/10/1687]
- 8. Food inflammation index reveals the key inflammatory components in foods and heterogeneity within food groups: How do we choose food? - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S2090123224004648]
- 9. Frontiers | Association of dietary and oxidative... inflammatory index [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1607162/full]
- 10. Comparison of predictive effect of the dietary inflammatory index and empirically derived food-based dietary inflammatory index on the menopause-specific quality of life and its complications - PubMed [https://pubmed.ncbi.nlm.nih.gov/37391733/]
- 11. Common and Novel Markers for Measuring Inflammation and Oxidative Stress Ex Vivo in Research and Clinical Practice—Which to Use Regarding Disease Outcomes? - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC8001241/]
- 12. Rethinking IL-6 and CRP: Why they are more than inflammatory biomarkers, and why it matters - ScienceDirect [https://www.sciencedirect.com/science/article/abs/pii/S0889159118300266]
- 13. How can biomarkers be used to differentiate between infection and non-infectious causes of inflammation? - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC7152028/]
- 14. Comparison of the inflammatory biomarkers IL- 6, TNF-α, and CRP to predict the effect of nutritional therapy on mortality in medical patients at risk of malnutrition | Journal of Inflammation | Full Text [https://journal-inflammation.biomedcentral.com/articles/10.1186/s12950-025-00442-0]
- 15. Interleukin-6, C-Reactive Protein, Tumor Necrosis Factor-alpha as Predictors of Mortality in Frail, Community-Living Elderly Individuals - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC4321727/]
- 16. of the inflammatory biomarkers Comparison - IL , 6 - TNF , and α to... CRP [https://pubmed.ncbi.nlm.nih.gov/40275239/]
- 17. IL-1β, IL - 6 , TNF - α and CRP in Elderly Patients with Depression or... [https://www.nature.com/articles/s41598-018-30487-6?error=cookies_not_supported&code=517cd451-83a4-4133-bcde-c7a265e6c12c]
- 18. is associated with lipid... | Research Square Dietary inflammatory index [https://www.researchsquare.com/article/rs-14529/v1]
- 19. Comparison of the Predictive Value of GlycA and Other Biomarkers of Inflammation for Total Death, Incident Cardiovascular Events, Noncardiovascular and Noncancer Inflammatory-Related Events, and Total Cancer Events - PubMed [https://pubmed.ncbi.nlm.nih.gov/27173011/]
- 20. Overview of anti- inflammatory and their promising effects on... diets [https://pmc.ncbi.nlm.nih.gov/articles/PMC11576095/]
- 21. Anti-inflammatory Dietary Inflammatory Index scores are associated with healthier scores on other dietary indices - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC4773655/]
- 22. Investigating the dietary and pro- inflammatory ... index inflammatory [https://eurjmedres.biomedcentral.com/articles/10.1186/s40001-025-02859-y]
- 23. Frontiers | Association between dietary ... inflammatory index score [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1557491/full]
- 24. Frontiers | The Role of Diet and Gut Microbiota in Regulating... [https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2022.866059/full]
- 25. Frontiers | Navigating next-gen nutrition care using artificial intelligence-assisted dietary assessment tools—a scoping review of potential applications [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1518466/full]
- 26. Designing and developing a literature-derived, population -based... [https://pmc.ncbi.nlm.nih.gov/articles/PMC3925198/]
- 27. Frontiers | The relationship between dietary inflammatory index and... [https://www.frontiersin.org/journals/endocrinology/articles/10.3389/fendo.2024.1417840/full]
- 28. - Wikipedia Dietary inflammatory index [https://en.wikipedia.org/wiki/Dietary_inflammatory_index]
- 29. A novel dietary inflammatory index reflecting for inflammatory ageing: Technical note - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S2049080119301311]
- 30. Association between dietary , empirical inflammatory ... index dietary [https://bmcendocrdisord.biomedcentral.com/articles/10.1186/s12902-025-02022-y]

- 31. Frontiers | Association between the dietary inflammatory index and... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1623482/full]
- 32. Associations between dietary inflammatory index ( DII ) scores and... [https://www.nature.com/articles/s41598-025-08919-x?error=cookies_not_supported&code=cf5a2134-2fb2-4ede-9405-f7051abaef6c]
- 33. 2.4. Dietary Assessment and Calculation of Dietary ... Inflammatory [https://bio-protocol.org/exchange/minidetail?type=30&id=8687473]
- 34. and Validation of Two Dietary Indexes for Predicting... Comparison [https://asu.elsevierpure.com/en/publications/validation-and-comparison-of-two-dietary-indexes-for-predicting-n]
- 35. The Dietary Inflammatory Index and Its Associations with Biomarkers of Nutrients with Antioxidant Potential, a Biomarker of Inflammation and Multiple Long-Term Conditions [https://www.mdpi.com/2076-3921/13/8/962]
- 36. Comparison of Intake of Food Groups Based on Dietary Inflammatory Index (DII) and Cardiovascular Risk Factors in the Middle-Age Population of Lower Silesia: Results of the PURE Poland Study - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC9952843/]
- 37. An Empirical Dietary Inflammatory Pattern Score Enhances Prediction of Circulating Inflammatory Biomarkers in Adults - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC5525108/]
- 38. Associations between dietary inflammatory scores and biomarkers of inflammation in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S0261561423001541]
- 39. The association between inflammatory and immune system biomarkers and the dietary inflammatory index in patients with COVID-19 - PubMed [https://pubmed.ncbi.nlm.nih.gov/37063325/]
- 40. and lipid metabolism parameters in women... Inflammatory biomarkers [https://pubmed.ncbi.nlm.nih.gov/36882933/]
- 41. Dietary inflammatory index and inflammatory in... biomarkers [https://www.nature.com/articles/s41430-017-0013-x?error=cookies_not_supported&code=842a6401-0cd9-4df7-9bba-4c0de47ef00a]
- 42. Food-based indexes and their association with dietary inflammation - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC11992388/]
- 43. Exploring the link between dietary inflammatory index, inflammatory biomarkers, and sleep quality in adults with obesity: a pilot investigation | International Journal of Obesity [https://www.nature.com/articles/s41366-025-01728-2]
- 44. Frontiers | Evaluating the triglyceride-glucose index as a biomarker for... [https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2025.1548444/full]
- 45. Association between dietary inflammatory index and... [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-025-01127-3]
- 46. Frontiers | Higher Dietary Inflammatory Index Scores Increase the Risk... [https://www.frontiersin.org/journals/endocrinology/articles/10.3389/fendo.2021.693144/full]
- 47. Association between dietary and inflammatory of index ... risk chronic [https://pmc.ncbi.nlm.nih.gov/articles/PMC12004726/]
- 48. Designing and developing a literature-derived, population-based dietary inflammatory index | Public Health Nutrition | Cambridge Core [https://www.cambridge.org/core/journals/public-health-nutrition/article/designing-and-developing-a-literaturederived-populationbased-dietary-inflammatory-index/30BE2C2295CE93DC6B54F9F9AD50CC68]
- 49. Association of dietary quality, biological aging, progression and... [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-025-01175-9]
- 50. Frontiers | Non-linear relationship between the dietary inflammatory... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1608603/full]
- 51. Relationship between dietary and metabolic... inflammatory index [https://www.nature.com/articles/s41598-025-89690-x?error=cookies_not_supported&code=001fd81c-2619-4755-ad7e-0fa729e4c61c]
- 52. Association of different diet quality scores with cardiovascular disease mortality risk in hypertension treatment group - PubMed [https://pubmed.ncbi.nlm.nih.gov/41017556/]
- 53. Frontiers | Association of dietary on all-cause and... inflammatory index [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1478165/full]
- 54. Relationship between dietary and metabolic... inflammatory index [https://www.nature.com/articles/s41598-025-89690-x?error=cookies_not_supported&code=cff2a7a7-4611-417d-9ad2-ab65583ab468]
- 55. The association between dietary inflammatory index (DII) scores and c-reactive protein (CRP) and nonalcoholic fatty liver disease (NAFLD) in a general population cohort - PubMed [https://pubmed.ncbi.nlm.nih.gov/38479904/]
- 56. Exploring the association between dietary Inflammatory Index and chronic pain in US adults using NHANES 1999–2004 | Scientific Reports [https://www.nature.com/articles/s41598-024-58030-w]
- 57. Relationship of Body Mass Index and Dietary ... Inflammatory Index [https://www.imrpress.com/journal/IJVNR/95/5/10.31083/IJVNR38965]
- 58. The association between energy -adjusted dietary inflammatory index... [https://link.springer.com/article/10.1186/s12889-025-22801-5]
- 59. Development and Validation of the China Dietary ... Inflammatory Index [https://pmc.ncbi.nlm.nih.gov/articles/PMC12114556/]
- 60. Frontiers | Nonlinear dose–response relationship between dietary inflammatory index and risk of depression: a systematic review and meta-analysis [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1645789/full]
- 61. Frontiers | Association of the dietary with frailty in... inflammation index [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1607110/full]
- 62. Dietary Inflammatory index and its association with fatty liver disease: a study in obese and non-obese populations | BMC Gastroenterology | Full Text [https://bmcgastroenterol.biomedcentral.com/articles/10.1186/s12876-024-03585-1]

- 63. and mortality in middle-aged and Dietary ... inflammatory index elderly [https://pubmed.ncbi.nlm.nih.gov/40605106/]
- 64. of Comparison and dietary ... inflammatory index inflammatory [https://www.nature.com/articles/s41598-024-69168-y?error=cookies_not_supported&code=b91427ee-4a71-408a-9bc4-658492562d6d]
- 65. analysis of fatty acid profiles across omnivorous... Comparative [https://lipidworld.biomedcentral.com/articles/10.1186/s12944-025-02517-6]
- 66. The association of dietary patterns with dietary inflammatory index, systemic inflammation, and insulin resistance, in apparently healthy individuals with obesity | Scientific Reports [https://www.nature.com/articles/s41598-021-86993-7]
- 67. Association between Dietary Inflammatory Index, Dietary Patterns, Plant-Based Dietary Index and the Risk of Obesity - PMC [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8147427/]
- 68. A new mathematical approach to improve the original... | PLOS One [https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0259629]
- 69. Frontiers | Energy-Adjusted Dietary Inflammatory Index Is Associated With 5-Year All Cause and Cardiovascular Mortality Among Chronic Kidney Disease Patients [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2022.899004/full]
- 70. Frontiers | Dietary and non-alcoholic fatty liver... inflammatory index [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1596300/full]
- 71. The dietary and its association with the... inflammatory index [https://pmc.ncbi.nlm.nih.gov/articles/PMC9893776/]
- 72. Relationship between Non-Energy-Adjusted and Energy-Adjusted Dietary Inflammatory Index and the Healthy Eating Index-2015: an analysis of the National Health and Nutrition Examination Survey (NHANES) 2015-2018 - PubMed [https://pubmed.ncbi.nlm.nih.gov/37489608/]
- 73. Diet Quality Assessed by the Dietary Inflammatory Index and the Healthy Eating Index: An Analysis of the National Health and Nutrition Examination Survey (2015–2018) - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S2475299123133798]
- 74. Dietary Quality Determined by the Healthy Eating Index-2015 and Biomarkers of Chronic Low-Grade Inflammation: A Cross-Sectional Analysis in Middle-to-Older Aged Adults - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC7828829/]
- 75. Frontiers | Comparative analysis of dietary patterns and depression... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1680741/full]
- 76. Dietary inflammatory index and healthy eating index-2015 are associated with rheumatoid arthritis - PubMed [https://pubmed.ncbi.nlm.nih.gov/33722328/]
- 77. Relationship between diet quality scores and the risk of frailty and... [https://pubmed.ncbi.nlm.nih.gov/33722232/]
- 78. Relationship between diet quality scores and the risk of frailty and... [https://bmcmedicine.biomedcentral.com/articles/10.1186/s12916-021-01918-5]
- 79. The Relationship between Energy-Adjusted Dietary ... Inflammatory [https://www.researchsquare.com/article/rs-47452/v1]
- 80. Association between empirical , odds... dietary inflammatory index [https://pmc.ncbi.nlm.nih.gov/articles/PMC10563743/]
- 81. Comparison of Different Dietary Indices as Predictors of Inflammation, Oxidative Stress and Intestinal Microbiota in Middle-Aged and Elderly Subjects - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC7765160/]
- 82. Healthy eating index versus alternate healthy index in relation to diabetes status and health markers in U.S. adults: NHANES 2007–2010 | Nutrition Journal | Full Text [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-019-0450-6]
- 83. Alternative dietary indices both strongly predict risk of chronic disease - PubMed [https://pubmed.ncbi.nlm.nih.gov/22513989/]
- 84. Frontiers | Geographic disparities of dietary inflammatory index and its... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2024.1355091/full]
- 85. Dietary inflammatory index and risk of first myocardial infarction... [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-017-0243-8]
- 86. Frontiers | Association between the Chinese Dietary Inflammatory... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2025.1653575/full]
- 87. Frontiers | Association of dietary quality and dietary inflammatory ... [https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2025.1596806/full]
- 88. Evaluation of Dietary ( Inflammatory ) and Index Eating... DII Healthy [https://pubmed.ncbi.nlm.nih.gov/40353262/]
- 89. Dietary patterns and associations with biomarkers of inflammation in... [https://nutritionj.biomedcentral.com/articles/10.1186/s12937-021-00674-9]