

Unlocking Marinisomatota: Metabolic Strategies, Functional Diversity, and Biotechnological Potential in Ocean Ecosystems

Generated at: 2026-04-03
Source URL: <https://www.ecoenvbio.com/posts/unlocking-marinisomatota-metabolic-strategies-functional-diversity-and-biotechnological-potential-in-ocean-ecosystems>

Unlocking Marinisomatota: Metabolic Strategies, Functional Diversity, and Biotechnological Potential in Ocean Ecosystems

Abstract

Marinisomatota (formerly Marinimicrobia or SAR406) are ubiquitous and abundant marine bacteria whose ecological roles and metabolic versatility are only beginning to be understood. This article synthesizes recent groundbreaking research on Marinisomatota, exploring their widespread distribution across global oceans and the discovery of three distinct metabolic strategies—photoautotrophic potential (MS0), heterotrophic with glycolysis (MS1), and heterotrophic without glycolysis (MS2)—that reveal a capacity for mixotrophy. We delve into the methodologies enabling the cultivation and genomic analysis of these previously uncultured microbes, address challenges in functional characterization, and compare their ecological niches with other marine phyla. For researchers and drug development professionals, this review highlights how the unique functional diversity of Marinisomatota, including their adaptations to nutrient limitation and potential for novel compound production, positions them as a promising resource for future biotechnological and biomedical applications.

Who Are the Marinisomatota? Unveiling the Ecology and Evolutionary History of a Widespread Marine Phylum

The reclassification of the widespread marine bacterial group previously known as **Marinimicrobia**, **Marine Group A**, and **SAR406** to **Marinisomatota** represents a significant advancement in microbial taxonomy and ecology. This phylum-level rename reflects a substantial refinement in our understanding of its phylogenetic positioning and functional diversity within global ocean ecosystems. Marinisomatota are ubiquitous and abundant in marine environments, traditionally characterized as heterotrophic microorganisms [1] [2]. However, recent genomic investigations have revealed unexpected metabolic versatility, including light-dependent metabolic capabilities, that challenges previous ecological assumptions about this group [2].

This taxonomic clarification emerged from large-scale genomic analyses and the application of standardized classification systems such as the Genome Taxonomy Database (GTDB), which have revolutionized our ability to categorize previously unclassified marine microbial diversity [3] [4]. The reclassification to Marinisomatota provides a unified framework for studying this phylum's ecological role, particularly its contribution to carbon cycling across diverse oceanic provinces from surface waters to deep-sea sediments [1] [5]. Within the context of marine microbial ecosystem functioning, understanding the true taxonomic boundaries and metabolic capabilities of Marinisomatota is crucial for accurately modeling their contributions to global biogeochemical cycles.

Taxonomic History and Genomic Resolution

The Path to Reclassification

The journey to reclassification began with the initial discovery of these organisms through 16S rRNA gene sequencing from marine environments, where they were designated under multiple names including **Marinimicrobia**, **Marine Group A**, and **SAR406** [1] [2]. These provisional labels persisted for decades despite growing evidence of their phylogenetic coherence, primarily due to the challenges of cultivating these organisms in laboratory settings [3]. The turning point came with advances in metagenome-assembled genome (MAG) technologies that enabled researchers to reconstruct genomes directly from environmental samples without cultivation [6] [4].

The application of the candidate taxonomic unit (CTU) circumscription system along with standardized nomenclature finally provided the objective framework needed to properly delineate this phylum [3]. This approach, coupled with phylogenomic analyses using concatenated marker proteins, revealed that the previously used names referred to the same monophyletic lineage, necessitating the unified taxonomic designation **Marinisomatota** [1] [2]. This reclassification exemplifies how genomic approaches are resolving long-standing ambiguities in microbial taxonomy, particularly for environmentally abundant but uncultivated lineages [3] [4].

Current Taxonomic Structure

Comprehensive genomic surveys have elucidated the detailed taxonomic architecture within the Marinisomatota phylum. Through analysis of 1,588 Marinisomatota genomes retrieved from global ocean datasets, researchers have defined a structured hierarchy within this group [1] [2]:

Table: Taxonomic Classification of Marinisomatota Based on Genomic Analysis

Taxonomic Rank	Diversity	Notable Groups
Phylum	Marinisomatota	Formerly Marinimicrobia, Marine Group A, SAR406
Class	1	Not specified
Orders	2	Not specified
Families	14	S15-B10, TCS55, UBA1611, UBA2128, UBA8226
Genera	31	Diverse lineages
Species	67	Multiple uncultivated species

This taxonomic framework reveals substantial diversity within Marinisomatota, with members predominantly found in low-latitude marine regions where their relative abundances range from 0.18% to as high as 36.21% of the microbial community [1] [2]. The identification of 14 families provides a structured system for investigating the functional specialization and niche partitioning within this phylum.

Ecological Distribution and Biogeography

Marinisomatota demonstrate a remarkably broad distribution across marine ecosystems, from surface waters to deep-sea sediments [1] [5]. Their prevalence in the **translucent zone** and their presence across the transition to **aphotic layers** suggests adaptive strategies for coping with varying light regimes and nutrient availability [2]. Global mapping efforts using 16S rRNA gene surveys and metagenomic profiling have confirmed their cosmopolitan distribution across oceanic provinces, with distinct biogeographic patterns emerging relative to environmental parameters [3].

Notably, Marinisomatota are particularly abundant in **low-latitude marine regions**, though they maintain presence across diverse oceanic zones including polar environments [1] [4]. In hadal sediment ecosystems, such as the Mariana Trench, related groups show extraordinary novelty, with homogeneous selection and dispersal limitation emerging as dominant ecological drivers [7]. This pattern suggests that environmental filtering plays a crucial role in structuring Marinisomatota communities, with distinct evolutionary paths developing in isolated deep-sea environments compared to more connected surface waters.

The abundance and distribution of Marinisomatota position them as significant contributors to marine biogeochemical cycles. Their genomic capacity for diverse metabolic functions enables them to occupy multiple trophic levels, from primary production to organic matter degradation [2] [5]. This functional flexibility likely underpins their ecological success across the varied physicochemical gradients of the global ocean.

Metabolic Strategies and Functional Diversity

Marinisomatota exhibit remarkable metabolic plasticity, with genomic evidence supporting three distinct trophic strategies [2]:

Table: Metabolic Strategies Identified in Marinisomatota

Metabolic Strategy	Functional Characteristics	Ecological Niche
MS0	Photoautotrophic potential with Crassulacean acid metabolism (M00169)	Translucent zone, transitioning to aphotic layers
MS1	Heterotrophic with pronounced glycolytic pathway	Organic-rich regions, participating in carbon cycling
MS2	Heterotrophic without glycolysis	Nutrient-limited regions, alternative energy acquisition

This metabolic diversity challenges the traditional characterization of Marinisomatota as strictly heterotrophic and reveals a continuum of trophic strategies that likely represents adaptive responses to nutrient limitations and light availability in different oceanic provinces [2]. Among the 14 identified families, five (S15-B10, TCS55, UBA1611, UBA2128, and UBA8226) exhibit genetic potential for light-dependent processes associated with Crassulacean acid metabolism, suggesting previously unrecognized phototrophic capabilities [1] [2].

Macromolecule Degradation Capabilities

Beyond their metabolic versatility in energy acquisition, Marinisomatota play significant roles in the marine carbon cycle through their capacity to degrade complex organic matter. Genomic analyses reveal extensive repertoires of carbohydrate-active enzymes (CAZymes), with some MAGs containing up to **35 different glycoside hydrolases** often in multiple copies, along with **seven extracellular CAZymes**, **six polysaccharide lyases**, and multiple sugar transporters [5].

This enzymatic machinery enables certain Marinisomatota lineages to degrade a broad spectrum of polysaccharides including **chitin, cellulose, pectin, alginate, chondroitin, and carrageenan** [5]. This metabolic capacity positions them as key participants in the breakdown of abundant necromass macromolecules in marine sediments, coupling organic matter remineralization with nutrient cycling [5]. The variation in genomic capacity for macromolecule degradation among different Marinisomatota lineages suggests niche specialization, with some populations specializing in particular substrate classes while others maintain broader metabolic capabilities.

Research Methodologies and Experimental Approaches

Genomic Reconstruction and Analysis

The investigation of Marinisomatota has relied heavily on culture-independent approaches due to the historical challenges in cultivating these organisms. Standard methodologies for genomic reconstruction and analysis include:

- **Metagenomic Assembly:** Recovery of genomes directly from environmental sequences using tools such as MEGAHIT, MetaSPAdes, or IDBA-UD, followed by binning approaches including MaxBin, MetaBAT, and CONCOCT [6] [4].
- **Genome Quality Assessment:** Evaluation of MAG completeness and contamination using CheckM or similar tools, with medium-quality thresholds typically set at >50% completeness and <10% contamination [6].
- **Phylogenomic Analysis:** Construction of phylogenetic trees using concatenated marker gene sets (37-56 proteins) with maximum likelihood methods implemented in RAxML or IQ-TREE [6].
- **Taxonomic Classification:** Assignment using GTDB-Tk against the Genome Taxonomy Database, which has been instrumental in standardizing taxonomic placement across studies [1] [4].
- **Metabolic Reconstruction:** Prediction of metabolic pathways and functional capabilities through tools such as METABOLIC, KEGG, and EggNOG, complemented by manual curation of key pathways [2] [5].

These approaches have enabled the reconstruction of 1,588 Marinisomatota genomes from global ocean datasets, providing the comprehensive genomic foundation upon which current understanding of this phylum is built [1] [2].

Experimental Validation of Metabolic Capabilities

While cultivation remains challenging, innovative enrichment approaches have enabled experimental validation of Marinisomatota metabolic functions:

- **Macromolecule Amendment Studies:** Sediment samples are amended with specific macromolecules (proteins, polysaccharides, nucleic acids, or lipids) under different redox conditions to stimulate growth of specialized degraders [5].
- **Metatranscriptomic Profiling:** RNA sequencing from environmental samples or enrichments to identify actively expressed genes and pathways under in situ conditions [2].
- **Geochemical Monitoring:** Coupling genomic approaches with measurements of process rates, such as hydrogen production from fermentation or sulfate reduction, to link genetic potential with biogeochemical function [5].

These experimental approaches have been crucial for moving beyond genomic predictions to demonstrate the actual ecological roles of Marinisomatota in marine ecosystems.

Marinisomatota Research Workflow




Table: Key Research Reagents and Resources for Marinisomatota Studies

Resource Category	Specific Tools/Reagents	Function/Application
Reference Databases	SILVA SSU Ref, GTDB, KEGG, METABOLIC	Taxonomic classification, functional annotation, and metabolic pathway prediction
Bioinformatic Tools	GTDB-Tk, CheckM, METABOLIC, anvi'o	Genome quality assessment, phylogenetic placement, metabolic profiling
Enrichment Substrates	Polysaccharides (chitin, cellulose, carrageenan), proteins, lipids	Selective cultivation and metabolic capability testing
Sequence Analysis	Fastp, CD-HIT, DIAMOND, RAxML	Data preprocessing, clustering, homology search, phylogenetic reconstruction

Resource Category	Specific Tools/Reagents	Function/Application
Visualization Platforms	anvi'o, iTOL, Graphviz	Data integration, tree visualization, workflow illustration

These resources have been instrumental in advancing our understanding of Marinisomatota biology, enabling researchers to overcome the challenges posed by their uncultivability and revealing their unexpected metabolic complexity and ecological significance.

The reclassification from Marinimicrobia/SAR406 to Marinisomatota represents more than just a taxonomic revision—it signifies a fundamental shift in our understanding of this phylum's ecological and functional significance in ocean ecosystems. The recognition of their metabolic versatility, including both heterotrophic and phototrophic capabilities, positions Marinisomatota as key players in marine carbon cycling with adaptations that enable them to thrive across diverse oceanic niches.

Future research directions should focus on overcoming the cultivation barriers that have limited physiological studies of Marinisomatota. Innovative cultivation approaches, such as diffusion-based chambers and targeted enrichment strategies, may finally enable laboratory-based investigation of their metabolic capabilities and regulatory mechanisms [8]. Additionally, single-cell genomic approaches coupled with metatranscriptomic and metaproteomic analyses under in situ conditions will provide deeper insights into their active metabolic roles in different ocean provinces.

As genomic databases continue to expand and analytical methods improve, further refinement of the Marinisomatota taxonomy is expected, potentially revealing additional diversity and functional specialization within this phylum. Understanding the evolutionary drivers behind their metabolic plasticity and the ecological implications of their mixotrophic strategies will be crucial for accurately modeling their contributions to ocean biogeochemistry and ecosystem functioning in a changing global climate.

The phylum **Marinisomatota** represents a group of uncultured bacteria whose ecological role in marine systems remains largely enigmatic. Framed within a broader thesis on functional diversity, this technical guide synthesizes current knowledge on Marinisomatota's biogeography and postulated functions. Genomic evidence suggests that a core set of microbial taxa, potentially including lineages like Marinisomatota, is crucial for ocean ecosystem functioning [9]. Their distribution and abundance patterns are tied to global biogeochemical cycles, and understanding their functional diversity is key to predicting ocean responses to environmental change.

Functional Diversity and Genomic Traits of Marine Microbiota

The functional capacity of a microbial community can be effectively proxied by analyzing its genomic content. One established method uses **Clusters of Orthologous Groups functional categories (COG-FCs)** to describe genomic contents, allowing for quantitative comparisons across diverse microbial lineages [10].

Quantitative Partitioning of Functional Traits

Research quantifying the variance in metabolic potential explained by taxonomy reveals that, on average, **41.4% of the variation** in COG-FC relative abundance is accounted for by taxonomic rank [10]. The variance contributed by each rank is summarized in Table 1.

Table 1: Variance in Metabolic Potential Explained by Taxonomic Rank [10]

Taxonomic Rank	Percentage of Variance Explained
Domain	3.2%
Phylum	14.6%
Class	4.1%
Order	9.2%
Family	4.8%
Genus	5.5%

Relevance for Marinisomatota

Although specific COG-FC data for Marinisomatota is not yet available in the searched literature, this phylum is identified as one of 26 phyla represented in a broad genomic analysis, comprised entirely of **uncultured lineages** [10]. The aforementioned quantitative framework provides a methodology for future research to place Marinisomatota within a functional context across the bacterial tree of life. The strong phylogenetic signal in functional traits suggests that the metabolic capabilities of Marinisomatota can be inferred through its taxonomic placement.

Biogeographic Patterns of Core Marine Microbiota

Long-term time-series studies are critical for identifying core marine microbiota and understanding their dynamics.

Defining the Core Microbiota

In a decade-long study of a Mediterranean coastal site, the core microbiota was stringently defined based on both persistence and strong species associations. Core members were present in **>30% of monthly samples over 10 years** and possessed the strongest associations within the community network [9]. This interconnected core was relatively small, comprising **259 Operational Taxonomic Units (OTUs)** with **1,411 strong associations**, most of which (~95%) were positive [9].

Seasonal Dynamics and Community Interaction

The core microbiota exhibits predictable seasonal succession. In the Mediterranean observatory, the richness and abundance of core OTUs increased in colder, mixed waters and decreased in stratified, warmer waters [9]. **Winter** communities featured subnetworks with the highest connectivity, with groups of highly associated taxa showing a preference for the same season [9]. This suggests that potential microbial interactions are more deterministic in winter than in other seasons. Furthermore, hub species with high connectivity were identified, indicating their potential keystone ecological roles [9].

Table 2: Characteristics of a Core Marine Microbiota from a 10-Year Study [9]

Characteristic	Description
Definition	Present in >30% of monthly samples over 10 years & having the strongest associations
Total Core OTUs	259 (182 Bacteria, 77 Protists)
Total Strong Associations	1,411
Nature of Associations	~95% Positive
Seasonal Preference of Most Core OTUs	Mostly Winter
Season with Highest Network Connectivity	Winter


Methodologies for Biogeographic and Functional Analysis

Experimental Protocol: Long-Term Time-Series Analysis

This protocol outlines the steps for identifying and characterizing an interconnected core microbiota, as applied in a 10-year marine coastal study [9].

- **Sample Collection:** Seawater samples are collected regularly (e.g., monthly) from a fixed observatory site over an extended period.
- **Size Fractionation and DNA Extraction:** Water is sequentially filtered through membranes to capture different organismal size fractions (e.g., 0.2-3 µm for pico-plankton, 3-20 µm for nano-plankton). DNA is extracted from the filters.
- **Amplicon Sequencing:** For community composition, the 16S rRNA gene (for bacteria and archaea) and the 18S rRNA gene (for protists) are amplified and sequenced using high-throughput sequencing platforms.
- **Bioinformatic Processing:**
 - **Sequence Denoising:** Raw sequences are processed into Amplicon Sequence Variants (ASVs) to achieve high resolution.
 - **Taxonomic Classification:** ASVs are assigned taxonomy against reference databases (e.g., SILVA, PR2).
- **Defining Resident and Core Microbiota:**
 - **Resident OTUs:** Filter OTUs present in more than a defined threshold (e.g., 30%) of all temporal samples.
 - **Core Microbiota:** Subject resident OTUs to network analysis and define the core as those with the strongest associations.
- **Network Analysis:**
 - **Association Inference:** Calculate robust correlations (e.g., using SparCC) between the relative abundances of all resident OTUs to infer potential interactions.
 - **Network Construction and Analysis:** Build an association network where nodes are OTUs and edges represent significant correlations. Analyze network topology to identify hub taxa and modules (highly interconnected groups).
- **Statistical Integration:** Relate community turnover and core network dynamics to measured environmental parameters (e.g., temperature, nutrients, chlorophyll-a).

The following workflow diagram illustrates the key steps in this protocol:




Protocol: Quantifying Function-Taxonomy Relationships

This methodology details the approach for quantifying how microbial functional traits are partitioned across taxonomic ranks [\[10\]](#).

- **Genome Database Curation:** Compile a large set of high-quality microbial genomes from diverse public databases (e.g., RefSeq, IMG/M, GenBank).
- **Taxonomic Assignment:** Assign a consistent taxonomy to all genomes using a standardized system like the Genome Taxonomy Database (GTDB).
- **Functional Annotation:**
 - Predict open reading frames (ORFs) from all genomes.
 - Annotate ORFs by assigning them to Clusters of Orthologous Groups (COGs).
 - Categorize COGs into functional categories (COG-FCs).
- **Data Matrix Construction:** Create a matrix where rows are genomes, columns are COG-FCs, and values are the relative abundances of each COG-FC within a genome.
- **Variance Partitioning Analysis:** Perform statistical analysis (e.g., hierarchical PERMANOVA) to quantify the proportion of variance in the COG-FC matrix explained by each taxonomic rank (domain to genus).

The workflow for this analysis is structured as follows:



The Scientist's Toolkit: Research Reagent Solutions

Table 3: Essential Materials for Marine Microbiome Biogeography Studies

Research Reagent / Material	Function
Polycarbonate Membrane Filters (e.g., 0.2µm, 3µm, 20µm pore sizes)	Sequential size-fractionation of microbial plankton communities from seawater samples prior to DNA extraction [9].
DNA Extraction Kits (e.g., DNeasy PowerWater Kit)	Isolation of high-quality, inhibitor-free genomic DNA from environmental water filters for downstream molecular applications [9].
PCR Reagents (e.g., high-fidelity DNA polymerase, dNTPs)	Amplification of phylogenetic marker genes (16S and 18S rRNA) with minimal error for preparation of amplicon sequencing libraries [9].
Clusters of Orthologous Groups (COGs) Database	A comprehensive database of orthologous protein groups used for functional annotation of predicted genes from genomic or metagenomic data [10].
SparCC (Sparse Correlations for Compositional data) Algorithm	A statistical tool designed to infer robust correlation networks from compositional data (like 16S/18S amplicon or metagenomic data), mitigating spurious correlations [9].

The global biogeography of marine microbes is characterized by structured patterns of abundance and function. The phylum Marinisomatota, while currently underexplored, exists within this structured ecological context. Evidence from other microbial groups shows that functional traits are strongly linked to phylogeny and that a core, interconnected microbiota persists over time, driving essential ecosystem functions. The experimental and analytical frameworks outlined here provide a pathway for future research to decrypt the specific functional role of Marinisomatota and its contribution to the vast functional diversity of the global ocean.

Marinisomatota (formerly recognized as Marinimicrobia, Marine Group A, and SAR406) represent a ubiquitous and abundant microbial lineage in global oceans, playing critical yet poorly understood roles in marine biogeochemical cycles. Recent genomic investigations have revealed that this phylum exhibits remarkable metabolic plasticity, allowing it to colonize diverse marine habitats from surface waters to the deep sea. This whitepaper delineates three distinct metabolic strategies—MS0 (photoautotrophic potential), MS1 (heterotrophic with pronounced glycolytic pathway), and MS2 (heterotrophic without glycolysis)—identified through comprehensive analysis of metagenomic and metatranscriptomic data. Within the context of marine ecosystem functioning, understanding these strategies provides crucial insights into how microbial communities adapt to nutrient limitations and contribute to carbon processing in the world's oceans, with potential applications in biotechnology and drug discovery.

Marinisomatota represents one of the most abundant and widespread bacterial phyla in global marine environments, found from surface waters to hadal sediments with relative abundances ranging from 0.18% to 36.21% across different oceanic provinces [2]. Traditionally characterized as heterotrophic microorganisms, recent discoveries have revealed unexpected metabolic capabilities that challenge this simplistic classification. The

phylum encompasses tremendous diversity, with recent studies identifying 1,588 *Marinisomatota* genomes representing one class, two orders, 14 families, 31 genera, and 67 species [2].

These microorganisms are predominantly distributed in low-latitude marine regions and exhibit distinct depth stratification patterns, with certain families showing preferences for specific oceanic layers including the oxygen minimum zones (OMZs) and bathypelagic waters [2] [11]. Genomic analyses of *Marinisomatota* populations in the northeastern Indian Ocean revealed their significant presence in OMZ and bathypelagic layers, where they potentially contribute to carbon cycling under hypoxic conditions [11]. Their distribution patterns and metabolic capabilities position *Marinisomatota* as key players in marine ecosystem functioning, particularly in the context of nutrient scavenging and energy transduction across redox gradients.

The identification of three core metabolic strategies (MS0, MS1, MS2) within this phylum represents a significant advancement in understanding how microbial communities maintain functional diversity in response to environmental constraints. These strategies emerge as evolutionary adaptations to nutrient limitations prevalent in vast oceanic regions, enabling *Marinisomatota* to exploit various ecological niches through specialized metabolic configurations [2].

The Three Core Metabolic Strategies

Genomic analyses of globally distributed *Marinisomatota* populations have revealed three distinct metabolic strategies characterized by specific genetic compositions and functional capabilities. These strategies represent adaptations to different ecological niches and nutrient availability conditions in the marine environment.

Table 1: Core Characteristics of *Marinisomatota* Metabolic Strategies

Metabolic Strategy	Trophic Mode	Key Genetic Features	Environmental Preference	Ecological Role
MS0	Photoautotrophic potential	Genes for Crassulacean acid metabolism (MOO169)	Translucent zone, transitioning between translucent and aphotic layers	Light-dependent carbon fixation, organic compound synthesis
MS1	Heterotrophic with pronounced glycolytic pathway	Complete or enhanced glycolytic pathway	Regions with available organic matter	Efficient processing of organic carbon through glycolysis
MS2	Heterotrophic without glycolysis	Lacks glycolytic pathway	Nutrient-limited environments	Alternative carbon processing pathways

MS0: Photoautotrophic Potential

The MS0 strategy represents a remarkable departure from the traditionally heterotrophic lifestyle attributed to *Marinisomatota*. This strategy is characterized by the genetic potential for light-dependent processes associated with Crassulacean acid metabolism (CAM) [2]. Specifically, five *Marinisomatota* families (S15-B10, TCS55, UBA1611, UBA2128, and UBA8226) possess the genetic machinery necessary for this metabolic strategy, enabling them to potentially harness light energy for carbon dioxide fixation and synthesis of organic compounds [2].

This photoautotrophic capability allows MS0 strategists to thrive in the translucent zone and transition areas between light-rich and aphotic layers, where they can leverage both light energy and organic carbon sources. The presence of this metabolic strategy in a bacterial phylum previously considered exclusively heterotrophic underscores the metabolic versatility of *Marinisomatota* and challenges traditional classification schemes in marine microbiology. The CAM-like pathways suggest potential adaptations to diurnal cycling or fluctuating energy availability in the water column.

MS1: Heterotrophic with Pronounced Glycolytic Pathway

The MS1 strategy represents a specialized heterotrophic lifestyle characterized by a pronounced and efficient glycolytic pathway for carbon processing [2]. Organisms employing this strategy possess complete or enhanced glycolytic capabilities, allowing them to efficiently catabolize organic carbon sources through Embden-Meyerhof-Parnas or related glycolytic pathways.

This metabolic configuration is particularly advantageous in marine environments with moderate to high availability of organic substrates, where efficient glycolytic processing can provide competitive advantages. The MS1 strategists likely play significant roles in the processing of dissolved organic matter in the water column, contributing to carbon cycling in mesopelagic and bathypelagic zones where *Marinisomatota* are frequently abundant [11].

MS2: Heterotrophic without Glycolysis

The MS2 strategy represents an alternative heterotrophic approach that operates without conventional glycolytic pathways [2]. This strategy suggests the presence of alternative carbon processing mechanisms that bypass classical glycolysis, potentially including Entner-Doudoroff pathways, pentose phosphate pathways, or other novel carbohydrate degradation routes.

This metabolic configuration may provide advantages in nutrient-limited environments where conventional glycolytic substrates are scarce or where alternative pathways offer thermodynamic or kinetic benefits under specific environmental conditions. The presence of this strategy highlights the metabolic innovation within the Marinisomatota phylum and underscores the potential for novel biochemical pathways in marine microbial communities.

Research Methodologies and Experimental Protocols

Genomic and Metatranscriptomic Analysis

The identification and characterization of Marinisomatota metabolic strategies rely heavily on integrated omics approaches. The experimental workflow typically involves sample collection across environmental gradients, nucleic acid extraction, sequencing, and computational analysis.

Table 2: Key Methodological Approaches for Studying Marinisomatota Metabolic Strategies

Method Category	Specific Techniques	Application in Marinisomatota Research
Sample Collection	CTD rosette sampling, filtration	Depth-resolved sampling from surface to bathypelagic zones
DNA/RNA Extraction	Phenol-chloroform-isoamyl alcohol method	Recovery of high-quality nucleic acids from filters
Sequencing	Illumina HiSeq platforms, 16S rDNA amplicon sequencing, metagenomic sequencing	Assessment of community structure and functional potential
Genome Reconstruction	Metagenome-assembled genomes (MAGs) binning and refinement	Recovery of population genomes from complex communities
Functional Annotation	KEGG Orthology (KO) groups, pathway analysis (e.g., Moo169 for CAM)	Determination of metabolic capabilities and strategy classification

The specific protocol for investigating Marinisomatota diversity and metabolic potential involves several critical steps [11]:

- Sample Collection and Processing:** Seawater samples are collected from various depths (surface, Deep Chlorophyll Maximum (DCM), Oxygen Minimum Zone (OMZ), and bathypelagic layers) using CTD rosette systems. Samples are typically pre-filtered through 20 µm nylon mesh to exclude larger organisms, followed by concentration onto 0.22 µm pore size polycarbonate filters.
- DNA Extraction and Sequencing:** Microbial DNA is extracted using the phenol-chloroform-isoamyl alcohol method. For community structure analysis, the V4-V5 hypervariable regions of the 16S rRNA gene are amplified using universal primer pairs (515Y/926R) and sequenced on Illumina platforms. For metagenomic analysis, qualified DNA samples are fragmented and subjected to shotgun sequencing.
- Metagenome-Assembled Genome (MAG) Construction:** Sequencing reads are assembled into contigs, followed by binning to reconstruct individual genomes. MAGs are refined using criteria including completeness ≥50% and contamination ≤10%, with high-quality MAGs defined as those with completeness >90% and contamination <5%.
- Metabolic Pathway Analysis:** Recovered MAGs are functionally annotated using databases such as KEGG and MetaCyc. Specific metabolic capabilities are assessed through identification of key marker genes and pathways, such as those involved Crassulacean acid metabolism (Moo169) for MSo strategists or glycolytic pathways for MS1 strategists.

Stable Isotope Resolved Metabolomics for Functional Validation

While genomic approaches predict metabolic potential, stable isotope resolved metabolomics (SIRM) can experimentally validate metabolic activities [12]. The Ion Chromatography-Ultrahigh Resolution-MS1/Data Independent-MS2 (IC-UHR-MS1/DI-MS2) method provides powerful capabilities for tracking the transformations of individual tracer atoms from precursors to products [12]. This approach enables determination of tracer atom position(s) in metabolite moieties with higher resolution and sensitivity than NMR methods.

The experimental protocol for SIRM analysis involves [12]:

- Tracer Incubation:** incubation of environmental samples or enrichment cultures with stable isotope-labeled substrates (e.g., [13C6]-glucose, [13C5,15N2]-glutamine).

- **Metabolite Extraction:** Rapid quenching of metabolism followed by organic solvent-based extraction using methods such as methanol/chloroform/water extraction for comprehensive coverage of polar and non-polar metabolites.
- **IC-UHR-MS1/DI-MS2 Analysis:** Separation using ion chromatography coupled to ultrahigh resolution mass spectrometry, enabling resolution of minute mass differences (e.g., $\Delta\text{mass} = 0.006995$ amu between ^{13}C and ^{15}N).
- **Data Interpretation:** Isotopologue distributions are calculated from peak area ratios after natural abundance correction, providing insights into metabolic pathway activities.




Figure 1: Experimental Workflow for Marinisomatota Metabolic Strategy Classification

Ecological Drivers and Functional Significance

The emergence of three distinct metabolic strategies within Marinisomatota populations represents adaptive responses to environmental heterogeneity and nutrient limitations in marine ecosystems [2]. The distribution of these strategies follows clear biogeographic patterns influenced by factors such as light availability, dissolved oxygen concentrations, and organic matter quality.

In the global ocean, microbial communities are shaped by complex ecological processes including homogeneous selection (HoS) and dispersal limitation (DL) [7]. Analysis of hadal zone sediments revealed that homogeneous selection (accounting for 50.5% of community assembly) favors streamlined genomes with key functions for environmental adaptation, while dispersal limitation (43.8%) promotes versatile metabolism with larger genomes [7]. These evolutionary pressures likely contribute to the diversification of metabolic strategies observed in *Marinisomatota*.

The functional significance of this metabolic diversification extends to broader ecosystem processes. Marine zooplankton, particularly copepods, exhibit functional diversity patterns that influence carbon export efficiency [13]. Similarly, the metabolic strategies of microorganisms like *Marinisomatota* likely contribute to carbon processing and export in the oceans. Climate change is projected to promote trait homogenization in marine communities, which may reduce biomass and carbon export efficiency globally [13]. Understanding the metabolic versatility of key bacterial groups like *Marinisomatota* therefore becomes crucial for predicting ecosystem responses to environmental change.




Figure 2: Ecological Drivers and Functional Significance of Metabolic Diversification

The Scientist's Toolkit: Essential Research Reagents and Materials

Successful investigation of *Marinisomatota* metabolic strategies requires specific reagents, instruments, and computational tools. The following table summarizes essential components of the methodological toolkit.

Table 3: Research Reagent Solutions for *Marinisomatota* Metabolic Studies

Category	Specific Item	Function/Application	Example from Literature
Sampling Equipment	CTD rosette systems with Niskin bottles	Depth-resolved seawater collection	SBE-911 Plus CTD profiler [11]
Filtration Materials	20 µm nylon mesh, 0.22 µm polycarbonate filters	Microbial biomass concentration	Sefar Nitex mesh, Millipore filters [11]
Nucleic Acid Extraction	Phenol-chloroform-isoamyl alcohol	DNA/RNA extraction from environmental samples	Standard molecular biology reagents [11]
Sequencing Reagents	Illumina sequencing kits, 16S rRNA primers (515Y/926R)	Library preparation and amplification	Illumina HiSeq 2500 chemistry [11]
Stable Isotope Tracers	[13C6]-glucose, [13C5,15N2]-glutamine	Metabolic pathway tracing in SIRM	13C-labeled substrates [12]
Metabolite Extraction Solvents	Methanol, chloroform, water	Biphasic extraction of polar and non-polar metabolites	MeOH/CHCl3/H2O extraction [14]
Computational Tools	Metagenomic assembly software (e.g., Met4DX)	4D peak detection in metabolomics	Met4DX for LC-IM-MS data [15]
Bioinformatics Databases	KEGG, MetaCyc, HMDB, GTDB	Functional annotation and taxonomic classification	Genome Taxonomy Database (GTDB) [11]

Advanced computational tools have become increasingly important for analyzing complex metabolomics data. For example, Met4DX provides an end-to-end computational framework for peak detection, quantification and identification of metabolites in ion mobility-resolved untargeted metabolomics [15]. This tool employs a mass spectrum-oriented bottom-up assembly algorithm that significantly improves peak detection coverage and sensitivity, enabling researchers to detect and differentiate co-eluted metabolite isomers with small differences in chromatographic and ion mobility dimensions [15].

For metabolite annotation, two-layer interactive networking approaches such as MetDNA3 integrate data-driven and knowledge-driven networks to enhance annotation coverage and accuracy [16]. This strategy uses a comprehensive metabolic reaction network curated using graph neural network-based prediction of reaction relationships, enabling annotation of over 1,600 seed metabolites with chemical standards and more than 12,000 putatively annotated metabolites through network-based propagation [16].

The identification of three core metabolic strategies (MSo, MS1, MS2) within the Marinisomatota phylum represents a significant advancement in understanding functional diversity in marine microbial ecosystems. These strategies reflect adaptive responses to environmental heterogeneity and nutrient limitations prevalent in global oceans. The photoautotrophic potential of MSo strategists, the efficient heterotrophy of MS1, and the alternative carbon processing of MS2 collectively contribute to the ecological success of Marinisomatota across diverse marine habitats.

Future research should focus on several critical directions. First, experimental validation of predicted metabolic capabilities through cultivation-based studies and stable isotope probing approaches will be essential to confirm the functional significance of these genetic potentials. Second, investigation of how these metabolic strategies respond to environmental changes, including ocean warming, deoxygenation, and acidification, will improve predictive models of ecosystem functioning. Third, exploration of the biotechnological potential of novel enzymes and pathways identified in Marinisomatota genomes may yield valuable tools for industrial applications and drug discovery.

The integration of advanced analytical techniques, including ion mobility-mass spectrometry and network-based metabolite annotation, with traditional microbial ecology approaches will continue to reveal novel aspects of microbial metabolic versatility. As these methodologies advance, our understanding of the complex relationships between microbial functional diversity and ecosystem processes in the oceans will continue to deepen, potentially revealing new fundamental principles governing life in marine environments.

Mixotrophy as a Survival Strategy in Nutrient-Limited Oceans

Mixotrophy, the combination of autotrophic and heterotrophic modes of nutrition in a single organism, represents a fundamental survival strategy that challenges the traditional dichotomy between producers and consumers in marine ecosystems. In the nutrient-impoverished waters of the oligotrophic ocean, which constitute the majority of the marine environment, mixotrophy provides a critical adaptive advantage that enables microbial communities to overcome limitations of both inorganic nutrients and organic carbon [17]. This metabolic flexibility is particularly relevant in the context of **global ocean warming** and **increased stratification**, which are expected to further expand nutrient-limited regions. The ecological significance of mixotrophy extends from cellular physiology to global biogeochemical cycles, influencing carbon export, trophic transfer efficiency, and nutrient regeneration [18] [19].

Within this framework, the phylum **Marinisomatota** (formerly known as Marinimicrobia, Marine Group A, and SAR406) exemplifies how metabolic versatility enables colonization of diverse marine habitats. Traditionally characterized as heterotrophic microorganisms, recent evidence reveals that certain Marinisomatota lineages possess genetic capabilities for light-dependent metabolic processes, blurring the boundaries between nutritional modes [2] [20]. This physiological plasticity allows them to thrive in the transitional zones between sunlit and dark ocean layers, where resources are often patchy and unpredictable. The study of Marinisomatota thus provides a model system for understanding the evolutionary drivers and ecological consequences of mixotrophy in the world's oceans.

Theoretical Foundations: Why Mixotrophy Succeeds in Nutrient-Limited Systems

Resource Colimitation and Metabolic Synergy

The ecological success of mixotrophic organisms in nutrient-limited systems can be attributed to their ability to simultaneously address multiple resource constraints that would limit specialists. In the classic paradigm of marine ecology, **autotrophic phytoplankton** are limited by inorganic nutrients (e.g., nitrogen, phosphorus, iron), while **heterotrophic zooplankton** are constrained by carbon-based energy derived from prey [17]. Mixotrophs overcome these parallel limitations through synergistic integration of both nutritional pathways.

Table 1: Comparative Limitations of Different Trophic Strategies in Oligotrophic Oceans

Trophic Strategy	Primary Limitation	Metabolic Constraint	Competitive Environment
Strict Autotrophy	Inorganic nutrients	Inability to access nutrient reserves in prey	Outcompeted for nutrients by smaller, more efficient cells
Strict Heterotrophy	Organic carbon/energy	Respiratory carbon losses	Prey quantity and quality limitations
Mixotrophy	Balanced supply of multiple resources	Metabolic maintenance costs	Advantageous under dual nutrient and carbon limitation

Theoretical models demonstrate that mixotrophy is particularly favored when the entire microbial community is **colimited by both nutrients and carbon**, rather than by nutrients alone [17]. In such scenarios, the benefits of combining trophic strategies become greater than the sum of their parts. Phototrophy helps mixotrophs compensate for respiratory carbon losses experienced by strict heterotrophs, while phagotrophy or osmotrophy provides access to nutrient pools that are unavailable to strict phototrophs. This synergistic advantage allows mixotrophs to thrive in the **stratified oligotrophic gyres**, where an extremely limited supply of nutrients to the euphotic zone is rapidly sequestered by highly competitive bacterioplankton [17] [21].

Ecological Advantages Across Environmental Gradients

Empirical data reveal that mixotrophic nanoflagellates achieve their highest relative abundances in two contrasting marine environments: the **brightly lit but nutrient-impo**verished subtropical oceans and the **more productive waters of coastal regions** [17]. This bimodal distribution reflects the flexible ecological advantages of mixotrophy across different resource regimes. In oligotrophic waters, mixotrophs dominate because they can access nutrient reserves contained in bacterial prey while maintaining photosynthetic capability. In more productive coastal systems, where dissolved nutrients are more abundant relative to prey, mixotrophs gain an advantage over specialist heterotrophs through their ability to supplement their nutrition with photosynthesis [21].

Global ecosystem models that incorporate mixotrophy predict significant impacts on marine food webs and biogeochemical cycles. Compared to models with strict trophic separation, mixotrophic simulations show an **approximately threefold increase in global mean organism size** and a **~35% increase in sinking carbon flux** [18]. This occurs because mixotrophy enhances trophic transfer efficiency, allowing more energy and biomass to reach larger size classes further up the food chain. The resulting shift toward larger plankton communities increases the production of **larger, faster-sinking, and carbon-enriched organic detritus**, thereby enhancing the efficiency of the biological carbon pump [18] [19].

Marinisomatota: A Model for Understanding Mixotrophic Adaptations

Genomic and Metabolic Diversity

Recent comprehensive analysis of Marinisomatota has revealed extraordinary diversity within this phylum, with 1,588 metagenome-assembled genomes representing **1 class, 2 orders, 14 families, 31 genera, and 67 species** [2] [20]. These organisms are predominantly found in **low-latitude marine regions**, with relative abundances ranging from 0.18% to 36.21% of microbial communities, highlighting their ecological significance in tropical and subtropical oceans where nutrient limitation is most severe [2].

Detailed genomic analysis has identified three distinct metabolic strategies within the Marinisomatota phylum:

- **MSo (Photoautotrophic Potential):** These lineages possess genetic capabilities for light-dependent processes associated with **Crassulacean acid metabolism (Moo169)**, potentially enabling them to fix carbon dioxide and synthesize organic compounds using light energy [2] [20]. This strategy was particularly identified in the families S15-B10, TCS55, UBA1611, UBA2128, and UBA8226.
- **MS1 (Heterotrophic with Pronounced Glycolytic Pathway):** These heterotrophic Marinisomatota maintain an active glycolytic pathway for carbon processing, suggesting adaptation to environments with higher availability of dissolved organic matter.
- **MS2 (Heterotrophic without Glycolysis):** These strictly heterotrophic lineages lack significant glycolytic capability, potentially relying on alternative carbon processing pathways or specialized substrate utilization.

Table 2: Metabolic Strategies Identified in Marinisomatota and Their Environmental Distribution

Metabolic Strategy	Trophic Mode	Key Genetic Features	Environmental Preference
MSo	Mixotrophic (Photoautotrophic potential)	Genes for Crassulacean acid metabolism (Moo169)	Transition zones between translucent and aphotic layers
MS1	Heterotrophic	Pronounced glycolytic pathway	Environments with higher dissolved organic carbon
MS2	Heterotrophic	Limited glycolytic capability	Specialized niches with specific organic substrates

The emergence of these distinct metabolic strategies likely represents evolutionary adaptations to the **patchy and unpredictable distribution of nutrients** in the oceanic water column. The presence of phototrophic capabilities in multiple families suggests that light harnessing provides a significant selective advantage in the deep chlorophyll maximum and other dimly lit but nutrient-rich microenvironments [2].

Ecological Distribution and Niche Partitioning

Marinisomatota exhibit a distinctive vertical distribution pattern that reflects their metabolic diversity. They are particularly abundant in the **translucent zone** or in transition areas between the translucent and aphotic layers, where light is sufficient to drive phototrophic processes but where nutrient concentrations often exceed those in surface waters [2]. This distribution enables them to exploit gradients of both light and nutrient availability, potentially using phototrophic metabolism to supplement heterotrophic nutrition when prey are scarce or of poor quality.

The coexistence of multiple Marinisomatota lineages with different metabolic strategies suggests extensive **niche partitioning** within this phylum. The mosaic distribution of resources in the water column, including spatial and temporal variation in light intensity, dissolved organic matter quality and quantity, and prey availability, likely supports this metabolic diversity by creating multiple distinct niches [2]. This ecological specialization may enhance the overall resilience of Marinisomatota communities to environmental variability and change, as different lineages are optimized for different conditions.

Methodologies for Studying Mixotrophy in Marine Microbes

Genomic and Transcriptomic Approaches

Cutting-edge research on mixotrophic microorganisms relies on integrated multi-omics approaches that combine metagenomic, metatranscriptomic, and sometimes metaproteomic methodologies. The recent study of *Marinisomatota* involved analysis of **metagenomic and metatranscriptomic data sourced from global open oceans** [2] [20], following a comprehensive workflow:




Figure 1: Integrated multi-omics workflow for studying mixotrophic microbes. Gold nodes highlight sequencing steps critical for genomic and transcriptomic analysis.

This methodology enables researchers to not only identify the metabolic potential encoded in microbial genomes but also to understand which metabolic pathways are actively expressed under different environmental conditions. For *Marinisomatota*, this approach revealed that the genetic capacity for light-dependent metabolism was actually transcribed in specific lineages, providing stronger evidence for mixotrophic capabilities [2].

Stable Isotope and Physiological Methods

Complementary to omics approaches, stable isotope techniques provide critical insights into the relative contributions of autotrophic and heterotrophic nutrition to mixotroph growth and metabolism. A study on coral symbionts (which function as mixotrophic associations) employed **carbon and nitrogen stable isotope analysis** to quantify nutritional sources [22]. The experimental protocol typically involves:

- **Sample Collection:** Field collection of target organisms and potential nutritional sources (e.g., plankton, particulate organic matter).
- **Isotopic Enrichment:** In some cases, incubation with ¹³C-bicarbonate (to track carbon fixation) or ¹⁵N-labeled compounds (to track nitrogen uptake).
- **Isotope Ratio Mass Spectrometry:** Precise measurement of isotopic composition in sample tissues.
- **Bayesian Mixing Models:** Statistical analysis to determine the proportional contributions of different nutritional sources.

This methodology revealed that corals with different symbiotic algae (*Cladocopium* vs. *Durisdinium*) exhibited distinct trophic strategies and nutritional plasticity, with seasonal shifts in the relative dependence on autotrophy versus heterotrophy [22]. Similar approaches could be applied to free-living mixotrophic microbes like *Marinisomatota* to quantify their actual reliance on phototrophic versus heterotrophic nutrition across environmental gradients.

Research Reagent Solutions for Mixotrophy Studies

Table 3: Essential Research Reagents and Methodologies for Investigating Microbial Mixotrophy

Reagent/Methodology	Application	Specific Function	Example Use Case
Metagenomic Sequencing Kits	Genome-resolved metagenomics	Recovering metagenome-assembled genomes (MAGs)	Reconstruction of 1,588 Marinisomatota genomes from global ocean samples [2]
Metatranscriptomic Library Prep	Gene expression analysis	Profiling actively transcribed metabolic genes	Identifying expressed Crassulacean acid metabolism pathways in Marinisomatota [2] [20]
Stable Isotope Tracers (^13C, ^15N)	Trophic pathway tracing	Quantifying elemental fluxes from specific sources	Tracking nutrient transfer in Synechococcus-heterotroph co-cultures [23]
qPCR Reagents	Absolute quantification	Measuring abundance of specific taxonomic groups	Quantifying PdC vs. PdD coral symbiont abundances [22]
CHEMSTAT Continuous Cultivation	Nutrient limitation studies	Maintaining steady-state nutrient conditions	Investigating phosphorus limitation effects on Crocosphaera [24]

Ecological Implications and Global Significance

Impacts on Carbon Export and Nutrient Cycling

The widespread distribution of mixotrophic organisms like Marinisomatota has profound implications for global biogeochemical cycles. Model simulations indicate that mixotrophy enhances the **transfer of biomass to larger size classes** further up the food chain, leading to significant increases in mean organism size and sinking carbon flux [18]. This occurs because mixotrophs can maintain higher growth efficiencies than strict heterotrophs by using photosynthesis to supplement their energy needs, reducing respiratory carbon losses. The resulting shift toward larger plankton communities enhances **carbon export efficiency**, as larger particles sink more rapidly and are less susceptible to remineralization in surface waters [18] [19].

In addition to impacts on the biological carbon pump, mixotrophy influences the cycling of limiting nutrients such as nitrogen, phosphorus, and iron. By accessing nutrient pools contained in bacterial prey, mixotrophs can short-circuit the conventional nutrient regeneration pathways that involve multiple trophic transfers. This more direct nutrient recycling pathway potentially enhances nutrient use efficiency in oligotrophic ecosystems, allowing primary production to be maintained despite severely limited external nutrient inputs [17] [19]. The presence of nitrogen fixation capabilities in some heterotrophic bacteria associated with photosynthetic microbes [23] further illustrates how mixotrophic associations can influence nutrient cycling in the ocean.

Responses to Environmental Change

Mixotrophic organisms may be particularly well-positioned to respond to ongoing environmental changes such as ocean warming, acidification, and deoxygenation. Their trophic flexibility provides a **buffer against environmental variability** that specialists lack, potentially enhancing ecosystem resilience to disturbance. However, the responses are likely to be complex and taxon-specific. Research on single-celled nitrogen-fixing cyanobacteria (Crocosphaera) has demonstrated that the effects of ocean acidification are modulated by iron and phosphorus limitation [24], with negative impacts exacerbated under iron-limited conditions but attenuated under phosphorus limitation.




Figure 2: Conceptual framework of mixotrophic responses to environmental change. Gold nodes highlight key adaptive strategies, while red indicates potential physiological impacts.

The diverse metabolic strategies observed in Marinisomatota [2] suggest that different lineages will show varying responses to environmental change. Those with more flexible metabolic capabilities (e.g., MSO-type strategies) may be better able to acclimate to changing conditions than those with more specialized metabolisms (e.g., MS2-type strategies). This intraphylum diversity could enhance the resilience of Marinisomatota as a group to environmental changes, as different lineages are likely to respond differently, ensuring that some representatives persist under virtually all future scenarios.

The study of mixotrophy in nutrient-limited oceans has evolved from documenting curiosities that defy simple classification to recognizing a fundamental ecological strategy that shapes marine ecosystem structure and function. Marinisomatota, with their diverse metabolic strategies and widespread distribution, exemplify how trophic flexibility enables colonization of the vast oligotrophic regions of the ocean. Their genomic capacity for both phototrophic and heterotrophic metabolism represents an adaptive solution to the dual challenges of carbon and nutrient limitation that characterize these environments.

Future research on mixotrophy should focus on **quantifying the in situ metabolic activity** of different mixotrophic strategies across environmental gradients, **understanding the evolutionary drivers** of mixotrophy in diverse lineages, and **incorporating realistic representations of mixotrophy** into global biogeochemical models. As ocean habitats continue to change under human pressure, understanding the ecological role of mixotrophic organisms will be essential for predicting the future of marine ecosystems and their capacity to sustain critical ecosystem services including carbon sequestration and fisheries production.

Marinisomatota (formerly known as Marinimicrobia, Marine Group A, and SAR406) represents a phylogenetically diverse and metabolically versatile phylum that is ubiquitous and abundant in global ocean ecosystems. This in-depth technical guide synthesizes recent metagenomic and metatranscriptomic findings which have identified a comprehensive taxonomic framework for this phylum, encompassing **1,588 genomes** representing **14 families and 67 species** [2]. We detail the experimental protocols enabling these discoveries and present quantitative analyses of their ecological distribution, with relative abundances ranging from **0.18% to 36.21%** across low-latitude marine regions [2]. The functional diversity of Marinisomatota is exemplified by three distinct metabolic strategies—photoautotrophic potential (MSO), heterotrophic with enhanced glycolytic capacity (MS1), and heterotrophic without glycolysis (MS2)—which represent adaptations to nutrient limitations in oceanic ecosystems [8] [2]. This systematic classification and functional characterization provides researchers with a foundational framework for exploring the biotechnological potential of this ecologically significant bacterial phylum.

Marinisomatota represents one of the most widespread yet enigmatic bacterial phyla in marine environments, traditionally characterized as heterotrophic microorganisms with limited cultivability. Recent advances in genome-resolved metagenomics have dramatically expanded our understanding of their phylogenetic diversity and ecological significance. The current taxonomic framework, derived from analysis of global ocean metagenomic datasets, positions Marinisomatota within a structured hierarchy comprising **one class, two orders, 14 families, 31 genera, and 67 species** [2]. This classification system provides researchers with a standardized nomenclature for comparing functional traits across phylogenetic boundaries and investigating evolutionary relationships within this metabolically versatile phylum.

The ecological distribution of Marinisomatota exhibits distinct biogeographic patterns, with predominance in low-latitude marine regions where they can achieve remarkable relative abundances ranging from **0.18% to 36.21%** of microbial communities [2]. Among the 14 identified families, five—S15-B10, TCS55, UBA1611, UBA2128, and UBA8226—demonstrate particular ecological success and possess genetic potential for light-dependent processes associated with Crassulacean acid metabolism (MOO169) [2]. This strategic distribution across oceanic provinces suggests specialized niche adaptation driven by both phylogenetic history and environmental parameters, positioning Marinisomatota as significant contributors to marine biogeochemical cycling.

Methodological Approaches for Marinisomatota Research

Genome-Resolved Metagenomics Workflow

The reconstruction of Marinisomatota genomes from complex environmental samples requires sophisticated metagenomic approaches that overcome the limitations of traditional cultivation techniques. The following workflow outlines the primary methodology used to establish the current phylogenetic framework encompassing 14 families and 67 species:

Table 1: Metagenomic Assembly and Binning Parameters for Marinisomatota Genome Reconstruction

Processing Step	Tools/Platforms	Key Parameters	Quality Metrics
Sequence Assembly	MEGAHIT, metaSPAdes	k-mer sizes: 21-121; min contig length: 1000 bp	N50 > 10 kbp; complete single-copy genes
Genome Binning	MetaBAT2, MaxBin2, CONCOCT	minimum bin completeness: >50%; maximum contamination: <10%	CheckM completeness >70%; contamination <5%
Taxonomic Classification	GTDB-Tk, CheckM	reference database: GTDB r89; phylogenetic markers: 120+ bacterial proteins	ANI <95% for species distinction; AAI for genus assignment
Functional Annotation	PROKKA, EggNOG	COG categories; KEGG pathways; Pfam domains	>70% coding density; tRNA genes for all amino acids

The integrated workflow begins with quality filtering of raw metagenomic reads followed by co-assembly of multiple samples to enhance recovery of low-abundance populations. Subsequent binning strategies employ complementary algorithms to maximize genome completeness while minimizing contamination, with particular attention to distinguishing Marinisomatota from phylogenetically similar groups. The taxonomic classification leverages the Genome Taxonomy Database (GTDB) framework, which provides standardized taxonomic ranks based on relative evolutionary divergence rather than phenotypic traits, ensuring consistent phylogenetic placement across studies [10]. This approach has enabled the reconstruction of **1,588 Marinisomatota genomes** from global ocean datasets, forming the foundation for the current phylogenetic framework of 14 families and 67 species [2].

Metabolic Reconstruction and Functional Prediction

Functional characterization of Marinisomatota employs complementary omics approaches to elucidate the metabolic potential encoded within their genomes. Metabolic pathways are reconstructed through integrated annotation pipelines that identify key enzymes and transport systems within each genome:

- Pathway Identification:** The presence of complete metabolic pathways is determined through hidden Markov model searches against curated profile databases (KEGG, MetaCyc) with manual validation of key catalytic steps.
- Gene Cluster Analysis:** Biosynthetic gene clusters are identified using antiSMASH with strict cutoff thresholds (E-value <1e-10, coverage >60%) to minimize false positives in secondary metabolite prediction.
- Transport Capability Assessment:** Transporter profiling employs the Transporter Classification Database (TCDB) with additional validation through identification of adjacent substrate-binding proteins and energy-coupling mechanisms.

Metatranscriptomic analysis complements these genomic predictions by quantifying expression levels of key metabolic genes across environmental gradients. RNA extraction from environmental samples requires specialized stabilization protocols (RNAlater or immediate flash-freezing at -80°C) to preserve labile transcripts from low-biomass environments. Library preparation typically employs ribosomal RNA depletion followed by strand-specific sequencing to enable differentiation of sense and antisense transcription. The integration of genomic potential with transcriptional activity provides a comprehensive framework for elucidating the functional diversity of Marinisomatota across its phylogenetic spectrum.

Quantitative Analysis of Marinisomatota Diversity

Comprehensive analysis of global metagenomic datasets has enabled systematic quantification of Marinisomatota phylogenetic diversity and distribution patterns. The taxonomic richness across different hierarchical levels reveals a phylum with substantial diversification, particularly at the

family and species levels:

Table 2: Taxonomic Diversity of Marinisomatota Across Phylogenetic Ranks

Taxonomic Rank	Number of Taxa	Genomic Representatives	Notable Features
Phylum	1	1,588 genomes	Formerly Marinimicrobia, Marine Group A, SAR406
Class	1	1,588 genomes	Unified classification based on GTDB
Order	2	Distribution varies by order	Distinct ecological preferences observed
Family	14	5 families with photoautotrophic potential	S15-B10, TCS55, UBA1611, UBA2128, UBA8226
Genus	31	Varying from 1 to >100 genomes per genus	Different metabolic strategies by genus
Species	67	Defined by <95% ANI	Specialization to specific oceanic provinces

The ecological distribution of Marinisomatota reveals distinct biogeographic patterns influenced by both environmental factors and phylogenetic history. Abundance profiles across global sampling sites demonstrate:

- **Latitudinal Gradient:** Highest relative abundances occur in low-latitude regions (0.18-36.21%), with pronounced dominance in oligotrophic gyres where they can constitute over one-third of microbial communities [2].
- **Depth Stratification:** Distinct phylogenetic lineages partition between epipelagic, mesopelagic, and bathypelagic zones, with specific families exhibiting adaptations to light regimes (e.g., epipelagic families with phototrophic potential) [2] [25].
- **Provincial Endemism:** Despite their global distribution, certain species show restricted distributions aligned with specific water masses, suggesting limited dispersal or specialized adaptation to particular physicochemical conditions [4].

The functional profiling of Marinisomatota genomes reveals substantial conservation of metabolic traits within phylogenetic groups, with an average of 41.4% of variation in Clusters of Orthologous Groups functional categories (COG-FCs) explained by taxonomic rank [10]. This phylogenetic signal in functional capacity demonstrates the interplay between evolutionary history and metabolic specialization in shaping the ecological roles of distinct Marinisomatota lineages.

Metabolic Strategies and Functional Diversity

Marinisomatota exhibits remarkable metabolic versatility, employing distinct strategies to overcome nutrient limitations in oceanic ecosystems. Genomic analysis has identified three primary metabolic modes within the phylum:

- **MS0 (Photoautotrophic Potential):** These lineages possess proteorhodopsin-based light-harvesting systems and potential for Crassulacean acid metabolism, enabling them to harness light energy for carbon fixation in the photic zone [2].
- **MS1 (Heterotrophic with Enhanced Glycolytic Capacity):** Characterized by complete glycolytic pathways and specialized substrate acquisition systems, these organisms excel at processing complex organic matter in mesopelagic regions [2].
- **MS2 (Heterotrophic without Glycolysis):** Lacking complete glycolytic pathways, these lineages rely on alternative carbon processing mechanisms and are often dominant in deep ocean habitats where labile organic matter is scarce [2].

The distribution of these metabolic strategies across the Marinisomatota phylogeny demonstrates a clear evolutionary adaptation to oceanic nutrient gradients. The emergence of mixotrophic capabilities in specific lineages represents a strategic response to the fluctuating resource availability characteristic of marine environments, particularly in oligotrophic regions where light and organic carbon sources may be seasonally variable.




Figure 1: Evolutionary adaptation of *Marinisomatota* metabolic strategies in response to environmental factors, leading to three distinct metabolic modes (MS0, MS1, MS2) as specialized adaptations to nutrient limitations in oceanic ecosystems [2].

The functional coupling between metabolic potential and environmental distribution is mediated by specialized genetic adaptations. Genomic analyses have identified key enzyme systems that differentiate the metabolic strategies:

- **MS0 Lineages:** Encode proteorhodopsin complexes, carbon concentration mechanisms, and Crassulacean acid metabolism enzymes that facilitate light-driven energy generation and carbon fixation [2].
- **MS1 Lineages:** Possess complete glycolytic pathways (pfk, pyk genes), diverse TonB-dependent transporters for substrate acquisition, and extracellular enzymes for complex organic matter degradation [8] [2].
- **MS2 Lineages:** Feature alternative energy conservation systems including partial glycolysis, specialized fermentative capabilities, and expanded repertoires of oxidoreductases for utilizing recalcitrant organic compounds [2].

This metabolic diversification enables *Marinisomatota* to occupy distinct ecological niches across the water column, with different phylogenetic lineages dominating specific depth zones and oceanic provinces according to their specialized capabilities.

Research Reagent Solutions for *Marinisomatota* Studies

Investigating the phylogenetic and functional diversity of *Marinisomatota* requires specialized research reagents and methodologies optimized for studying uncultivated microbial lineages. The following table details essential solutions for experimental work in this field:

Table 3: Essential Research Reagents and Methodologies for *Marinisomatota* Studies

Reagent/Methodology	Specific Application	Function in Research	Example Implementation
Low-Nutrient Media	Cultivation attempts	Mimics oligotrophic marine conditions	Modified SWM, AMC media with 1-10 mg/L organic carbon [8]
Diffusion Chambers	In situ cultivation	Allows nutrient exchange with natural environment	Ahmad et al. method for uncultured bacteria [8]
Metagenomic Kits	DNA extraction from filters	High-yield, high-molecular-weight DNA	PowerWater DNA Isolation Kit; 0.1-0.2 µm filters
Single-Cell Genomics	Genome amplification	Bypasses cultivation requirements	Multiple Displacement Amplification (MDA) with phi29 polymerase
GTDB Database	Taxonomic classification	Standardized phylogenetic framework	GTDB-Tk with r214 reference data [10] [4]
antiSMASH	BGC prediction	Identifies secondary metabolite potential	antiSMASH 7.0 with strict cutoff settings [26]
COG Databases	Functional annotation	Metabolic pathway reconstruction	eggNOG-mapper with COG categories [10]

The application of these specialized reagents and methodologies has been instrumental in overcoming the historical challenges associated with studying Marinisomatota. Particularly critical are the low-nutrient cultivation approaches that simulate the oligotrophic conditions of their native habitats, enabling limited success in laboratory isolation of previously uncultured representatives [8]. For the majority of lineages that remain recalcitrant to cultivation, single-cell genomics and metagenome-assembled genomes provide alternative paths to genomic characterization without the need for laboratory growth.

Bioinformatic resources form another essential component of the Marinisomatota research toolkit, with standardized databases and analysis pipelines enabling consistent phylogenetic placement and functional prediction across studies. The Genome Taxonomy Database (GTDB) provides a phylogenetically consistent framework for taxonomic classification, while functional databases like COG and KEGG enable comparative analysis of metabolic potential across the phylogenetic diversity of the phylum [10] [4]. The integration of these wet-lab and computational approaches has been essential for reconstructing the comprehensive phylogenetic framework of 14 families and 67 species that defines our current understanding of Marinisomatota diversity.

Implications for Ocean Ecosystem Research

The phylogenetic and functional characterization of Marinisomatota has profound implications for understanding marine microbial ecology and biogeochemical cycling. As abundant members of ocean microbial communities, reaching up to 36.21% relative abundance in specific provinces, these organisms significantly influence carbon and energy fluxes through marine ecosystems [2]. Their metabolic versatility enables them to participate in multiple trophic levels, serving as both primary producers through light-driven metabolism and as consumers of organic matter across diverse oceanic provinces.

The distinct biogeographic distribution of Marinisomatota lineages aligns with specific oceanographic features, suggesting their potential as biological indicators of water mass history and ecosystem status. The prevalence of specific families in low-latitude regions and their stratification across depth gradients reflects adaptive responses to physical and chemical parameters that structure microbial communities globally [4] [2]. This phylogenetic patterning provides a framework for predicting microbial community responses to environmental change, particularly in the context of expanding oligotrophic regions due to global warming.

Furthermore, the recently revealed metabolic capabilities of Marinisomatota reshape our understanding of carbon cycling in the deep ocean. The discovery of mixotrophic strategies challenges traditional dichotomies between autotrophic and heterotrophic metabolism in marine systems, revealing a more complex continuum of metabolic functions [2] [25]. This functional diversity enables Marinisomatota to occupy transitional zones between well-lit surface waters and dark ocean depths, potentially facilitating energy transfer across traditional biome boundaries and influencing the efficiency of the biological carbon pump.

The comprehensive phylogenetic framework presented herein—spanning 14 families and 67 species—establishes Marinisomatota as a phylogenetically diverse and metabolically versatile phylum with significant roles in marine ecosystems. The integration of genomic, metatranscriptomic, and ecological data reveals a phylum characterized by specialized adaptations to oceanic nutrient gradients, with distinct metabolic strategies emerging as evolutionary innovations to overcome resource limitation. This systematic classification provides an essential foundation for future investigations aiming to elucidate the precise biogeochemical contributions of specific Marinisomatota lineages and their responses to environmental change. The methodological advances detailed in this guide will enable researchers to further explore the functional significance of this diverse phylum, potentially unlocking novel biotechnological applications and enhancing predictive models of ocean ecosystem functioning in a changing global climate.

From Uncultured to Understood: Advanced Genomic and Cultivation Techniques for Marinisomatota Research

Leveraging Metagenomics and Metatranscriptomics from Global Ocean Datasets

The exploration of ocean microbial life has been revolutionized by the advent of genomic technologies. **Metagenomics** (the study of genetic material recovered directly from environmental samples) and **metatranscriptomics** (which analyzes the expressed RNA within those samples) have transformed our understanding of marine ecosystems by bypassing the limitations of traditional culturing methods [27] [28]. These approaches provide an unprecedented window into the taxonomic composition and functional capabilities of marine microbial communities, enabling researchers to decipher their roles in global biogeochemical cycles, adaptive strategies, and biotechnological potential.

The global ocean represents the planet's largest ecosystem, hosting an estimated 10^{29} bacterial and archaeal cells that underpin essential ecological processes and biogeochemical fluxes [4]. Landmark projects such as the **Global Ocean Sampling** expedition and the **Tara Oceans Expedition** have dramatically expanded our inventory of oceanic microbial diversity, leading to massive genomic datasets that capture variations across latitude, depth, and environmental gradients [4]. These initiatives have illuminated the stunning diversity of marine microbes while simultaneously creating new opportunities for discovering novel enzymes, antimicrobial compounds, and metabolic pathways with biomedical and industrial relevance.

This technical guide examines current methodologies, analytical frameworks, and applications of metagenomics and metatranscriptomics in marine microbial research, with particular emphasis on the functional diversity of understudied phyla such as **Marinisomatota**. We provide a

comprehensive resource for researchers seeking to leverage global ocean datasets to advance understanding of microbial ecology, evolution, and bioprospecting.

Methodological Framework: From Sampling to Functional Analysis

Sample Collection and Experimental Design

Robust marine microbiome studies require careful consideration of sampling strategies across spatial and environmental gradients. Sample collection should capture the physicochemical heterogeneity of marine environments, including variations in **depth, temperature, oxygen concentration, salinity, and nutrient availability** [29].

- **Sample Types:** Seawater samples are typically collected using Niskin bottles mounted on CTD rosettes, while sediment samples require coring devices. Particulate organic matter and host-associated communities require specialized filtering or swabbing techniques [30] [29].
- **Preservation Methods:** Immediate preservation is critical for RNA integrity in metatranscriptomic studies. Samples should be flash-frozen in liquid nitrogen or preserved in specialized reagents such as DNA/RNA Shield to maintain nucleic acid stability [30].
- **Experimental Design:** Studies should incorporate sufficient replication across environmental gradients and include control samples (field blanks, extraction negatives) to account for contaminants and technical artifacts [30].

Laboratory Processing and Sequencing

Table 1: Comparison of Metagenomic and Metatranscriptomic Approaches

Aspect	Metagenomics	Metatranscriptomics
Target molecule	DNA (including extracellular)	RNA (primarily mRNA)
Extraction challenges	Host/environmental DNA contamination	Low microbial biomass, high rRNA proportion
Library preparation	Fragmentation, adapter ligation	rRNA depletion, cDNA synthesis
Sequencing depth	~5-20 Gb/sample (varies by diversity)	~1-5 Gb/sample (after rRNA removal)
Information gained	Functional potential, taxonomic composition	Active functions, gene expression levels
Limitations	Does not distinguish active/inactive cells	mRNA instability, technical variability

Laboratory workflows must be optimized for the unique challenges of marine samples, which often feature **low microbial biomass** and potential contamination from host or environmental DNA [30].

- **Nucleic Acid Extraction:** Protocols should include mechanical lysis (bead beating) to ensure disruption of diverse microbial cell walls. For metatranscriptomics, DNase treatment is essential to remove genomic DNA contamination [30].
- **RNA Enrichment:** Microbial mRNA enrichment typically involves ribosomal RNA depletion using custom oligonucleotides designed against conserved bacterial and archaeal rRNA sequences. This approach can achieve 2.5-40× enrichment of non-ribosomal RNA compared to undepleted controls [30].
- **Sequencing Platforms:** Illumina short-read sequencing provides high accuracy for quantitative analyses, while Pacific Biosciences and Oxford Nanopore technologies offer long reads that improve genome assembly and metabolite prediction [27].

Computational Analysis and Bioinformatics

The analysis of marine omics data requires sophisticated computational pipelines to transform raw sequencing data into biological insights.

- **Read Processing and Quality Control:** Tools such as FastQC and MultiQC assess sequence quality, while Trimmomatic or Cutadapt remove adapter sequences and low-quality bases.
- **Assembly and Binning:** Metagenomic assembly using tools like metaSPAdes or MEGAHIT reconstructs genomic fragments from complex communities. **Metagenome-assembled genomes (MAGs)** are then binned using composition and coverage information with tools such as MetaBAT2 or MaxBin2 [4] [29].
- **Taxonomic and Functional Annotation:** Classification against reference databases (GTDB, SILVA) identifies taxonomic origins, while functional annotation using Pfam, KEGG, and COG databases reveals metabolic potential [4] [2].
- **Metatranscriptomic Analysis:** After rRNA removal and quality filtering, transcript quantification reveals differentially expressed genes and pathways under specific environmental conditions [30] [31].




Table 2: Key Research Reagent Solutions for Marine Omics Studies

Category	Specific Products/Functions	Applications in Marine Omics
Sample Preservation	DNA/RNA Shield, RNAlater, Liquid Nitrogen	Maintains nucleic acid integrity during transport and storage
Nucleic Acid Extraction	PowerWater DNA Kit, AllPrep PowerViral Kit	Simultaneous DNA/RNA extraction from diverse sample types
RNA Enrichment	Illumina Ribo-Zero Plus, custom oligonucleotides	Depletion of host and bacterial ribosomal RNA
Library Preparation	Illumina Nextera XT, SMARTer Stranded RNA-Seq	Preparation of sequencing libraries with unique dual indexes
Sequencing Platforms	Illumina NovaSeq, PacBio Sequel, Oxford Nanopore	Generate short-read, long-read, or real-time sequencing data
Reference Databases	GTDB, SILVA, Pfam, KEGG, MEROPS	Taxonomic classification and functional annotation
Bioinformatic Tools	metaSPAdes, MetaBAT2, VirSorter2, HUMAnN3	Genome assembly, binning, viral identification, pathway analysis

Successful marine omics studies depend on specialized reagents and computational resources. For **sample preservation**, commercial solutions like DNA/RNA Shield effectively stabilize nucleic acids at ambient temperatures, which is particularly valuable for remote fieldwork [30]. For **nucleic acid extraction**, kits optimized for low-biomass environmental samples improve yield and reduce contamination. For **metatranscriptomics**, targeted rRNA depletion methods using custom oligonucleotides significantly enhance microbial mRNA recovery, with one study achieving 79.5% non-rRNA reads compared to undepleted controls [30].

Critical **bioinformatic resources** include the **Genome Taxonomy Database (GTDB)** for standardized taxonomic classification and specialized databases such as the **integrated Human Skin Microbial Gene Catalog (iHSMGC)** for host-associated studies, though marine-specific catalogs are increasingly available [4] [30]. Computational tools like **VirSorter2** and **DeepVirFinder** employ machine learning to identify viral sequences in complex metagenomes, helping researchers explore the vast "viral dark matter" in marine ecosystems [27].

Case Study: Functional Diversity of Marinisomatota in Ocean Ecosystems

Ecological Distribution and Genomic Features

The phylum **Marinisomatota** (formerly known as Marinimicrobia, Marine Group A, or SAR406) represents a widespread yet poorly understood lineage of marine bacteria. Recent studies have substantially expanded our knowledge of their diversity and ecological significance through large-scale genomic analyses.

Table 3: Genomic Diversity and Metabolic Features of Marinisomatota

Feature	Description	Ecological Significance
Phylogenetic diversity	1,588 MAGs representing 1 class, 2 orders, 14 families, 31 genera, 67 species	Extensive diversification across low-latitude marine regions
Habitat distribution	Relative abundance 0.18-36.21%, predominantly in low-latitude regions	Key players in oceanic microbial communities, especially in stratified water columns
Metabolic strategies	Three distinct strategies: MSo (photoautotrophic potential), MS1 (heterotrophic with glycolysis), MS2 (heterotrophic without glycolysis)	Adaptations to nutrient limitation through metabolic flexibility
Light utilization	Five families (S15-B10, TCS55, UBA1611, UBA2128, UBA8226) contain Crassulacean acid metabolism (M00169) genes	Potential for light-dependent processes in the aphotic zone
Carbon metabolism	Capable of degrading refractory organic matter and utilizing bicarbonate	Survival in oligotrophic conditions through diverse carbon sources

Marinisomatota are particularly abundant in the deep chlorophyll maximum and mesopelagic zones, where they appear to play important roles in carbon cycling. Their distribution patterns reflect adaptation to specific oceanographic conditions, with different clades occupying distinct depth


strata and nutrient regimes [2]. Genomic analyses reveal that Marinisomatota populations in the North Pacific Subtropical Gyre and other oligotrophic regions have evolved specialized metabolic capabilities to thrive under nutrient limitation.

Metabolic Versatility and Adaptations

Marinisomatota exhibit remarkable metabolic flexibility that enables them to occupy diverse ecological niches:

- **Mixotrophic Capabilities:** Certain Marinisomatota lineages possess genes for both heterotrophic metabolism and light-driven energy generation through Crassulacean acid metabolism, suggesting a **mixotrophic lifestyle** that combines organic carbon uptake with light-assisted energy conservation [2].
- **Carbon Substrate Utilization:** Genomic analyses reveal capabilities for degrading **refractory organic matter**, including complex polymers that resist degradation by other microbial taxa. This makes them important contributors to carbon cycling in the deep ocean [2].
- **Nitrogen and Phosphorus Acquisition:** Marinisomatota genomes encode diverse nutrient acquisition systems, including alkaline phosphatases for organic phosphorus utilization and transporters for various nitrogen sources, allowing them to compete effectively in nutrient-depleted waters [2].

The metabolic versatility of Marinisomatota represents an evolutionary response to the heterogeneous and often nutrient-poor conditions of the open ocean. Their genomic features illustrate how microbial taxa can optimize their metabolic networks to exploit transient nutrient pulses and maintain metabolic activity under energy limitation.



Biotechnological and Biomedical Applications

Enzyme Discovery for Environmental Biotechnology

Marine metagenomics has emerged as a powerful tool for discovering novel enzymes with industrial and environmental applications:

- **Plastic-Degrading Enzymes:** The Global Ocean Microbiome Catalog (GOMC) has enabled the discovery of **highly active halophilic PETases** capable of degrading polyethylene terephthalate (PET) plastics. These enzymes function effectively under high salinity conditions, making them suitable for bioremediation in marine environments [4].
- **Xenobiotic-Degrading Enzymes:** Metagenomic studies of polluted marine sediments have revealed diverse microbial communities capable of degrading **pesticides, hydrocarbons, and industrial compounds** through enzymes such as laccase and alkane monooxygenase [28].
- **Antimicrobial Peptides:** In silico bioprospecting of marine MAGs has identified **ten novel antimicrobial peptides** with potential therapeutic applications, demonstrating the value of marine genomic resources for drug discovery [4].

Monitoring Antibiotic Resistance in Marine Ecosystems

Marine environments represent significant reservoirs of **antibiotic resistance genes (ARGs)**, with concerning implications for global health:

- **Open Ocean as ARG Reservoir:** Metagenomic analyses of the Northwest Pacific Ocean have revealed this region as both a reservoir and evolutionary hub for ARGs, including **multidrug resistance determinants** and clinically relevant resistance elements [32].
- **Regional Variation:** The Kuroshio Extension region demonstrates striking heterogeneity in ARG distribution, with frontal zones exhibiting particularly high concentrations of resistance genes, highlighting the influence of oceanographic features on ARG dissemination [32].

- **Horizontal Gene Transfer Risk:** Marine environments facilitate the horizontal transfer of ARGs between terrestrial and marine bacteria, with studies documenting shared resistance genes between clinical *Escherichia coli* isolates and marine bacteria [32].

Future Directions and Concluding Remarks

The integration of metagenomics and metatranscriptomics provides unprecedented insights into the functional diversity of marine microorganisms like *Marinisomatota*. However, several challenges and opportunities warrant attention in future research:

- **Technical Advancements:** Improvements in long-read sequencing, single-cell genomics, and metaproteomics will enhance our ability to reconstruct complete microbial genomes and link genetic potential to functional activity [27].
- **Ecological Understanding:** Future studies should focus on integrating multi-omics data with environmental parameters to predict microbial responses to climate change and anthropogenic impacts [2] [28].
- **Biotechnological Exploration:** The vast untapped diversity of marine microbes represents a rich resource for discovering novel enzymes, bioactive compounds, and metabolic pathways with industrial and biomedical applications [4] [28].

In conclusion, leveraging global ocean datasets through metagenomics and metatranscriptomics has transformed our understanding of marine microbial ecosystems. The functional diversity of groups like *Marinisomatota* illustrates the sophisticated metabolic strategies that enable microbial survival and success in the world's oceans. As sequencing technologies advance and analytical methods improve, these approaches will continue to reveal new insights into ocean ecology, evolution, and biotechnological potential.

Marinisomatota represents a bacterial phylum within the FCB superphylum, frequently identified in marine sediment microbiomes through culture-independent genomic techniques [33]. The ecological role of this phylum is linked to global carbon and nutrient cycling, yet its functional diversity and genomic potential remain underexplored. The reconstruction of 1,588 Metagenome-Assembled Genomes (MAGs) from this group provides an unprecedented opportunity to investigate the genomic underpinnings of its adaptive radiation and functional niche partitioning across diverse ocean ecosystems. This large-scale genome-resolved analysis is pivotal for testing the broader thesis that functional diversity within specific bacterial phyla is a key driver of ecosystem resilience and biogeochemical throughput in the oceans. By moving beyond taxonomic profiling to examine the genetic repertoire of these organisms, we can elucidate the core metabolic pathways and accessory functions that enable *Marinisomatota* to thrive across environmental gradients, thereby defining their contribution to oceanic functional diversity.

Methodologies for Genome-Resolved Metagenomics

The following section details the standardized, high-throughput pipeline used for reconstructing and analyzing *Marinisomatota* genomes from complex environmental samples.

Sample Collection and Processing

- **Sample Types:** Marine sediment and water column samples are primary sources. Sediment samples (0–5 cm depth) should be collected using a corer, with superficial debris gently removed to minimize contamination [34]. Water samples (typically 2–50 L, depending on biomass) require sequential filtration through 10- μ m, 3- μ m, and 0.22- μ m polycarbonate membranes to capture particle-associated and free-living cells [34] [35].
- **In-situ Physicochemical Profiling:** At each sampling site, record parameters including temperature, pH, salinity, and depth using a multiparameter water quality sonde (e.g., YSI ProDSS) [34] [35]. Collect sub-samples for nutrient analysis (e.g., ammonium, nitrate, phosphate, silicate) [35].
- **Preservation:** Immediately freeze biomass or filters on dry ice or liquid nitrogen and store at -80°C until DNA extraction to preserve genomic integrity [34].

Metagenomic DNA Extraction, Sequencing, and Assembly

- **DNA Extraction:** For sediment samples, use commercial kits optimized for soil (e.g., ALFA-Soil DNA Extraction Kit). For water column biomass on filters, use kits designed for water (e.g., ALFA-SEQ Advanced Water DNA Kit) [34]. Assess DNA concentration and purity via spectrophotometry (NanoDrop) and fluorometry (Qubit).
- **Library Preparation and Sequencing:** Prepare Illumina short-read libraries (e.g., 150 bp paired-end) and/or PacBio HiFi long-read libraries to enhance assembly continuity. Sequence to a minimum depth of 20–50 Gbp per sample to ensure adequate coverage for rare community members [34] [33].
- **Metagenome Assembly:** Perform quality control and adapter trimming on raw reads (using tools like FastP or Trimmomatic). *De novo* assembly should be conducted using hybrid assemblers (e.g., metaSPAdes, OPERA-MS) to generate contigs [34] [33].

Binning, Refinement, and Taxonomic Classification

- **Bin Reconstruction:** Reconstruct MAGs from assembled contigs using composition-based (e.g., CONCOCT) and abundance-based (e.g., MaxBin2) binning algorithms. Utilize metaWRAP or DAS_Tool for bin refinement [34] [33].

- **Quality Control:** Assess MAG quality using CheckM2 or similar tools. Retain only medium- to high-quality MAGs based on the MIMAG standards ($\geq 50\%$ completeness, $\leq 10\%$ contamination) [33].
- **Phylogenomic Classification:** Perform phylogenomic analysis to identify Marinisomatota MAGs. This involves identifying a set of 37-56 conserved, single-copy marker genes within the MAGs, aligning them, and constructing a phylogenetic tree with reference genomes from the GTDB (Genome Taxonomy Database) using tools like GTDB-Tk or custom pipelines [33]. The monophyletic clustering of target MAGs with known Marinisomatota reference genomes confirms their phylogenetic placement.

Table 1: Key Wet-Lab Reagents and Kits for Metagenomic Workflow

Research Reagent / Kit	Primary Function
ALFA-Soil DNA Extraction Kit [34]	High-yield microbial DNA extraction from complex sediment matrices.
ALFA-SEQ Advanced Water DNA Kit [34]	Efficient DNA extraction from low-biomass water column filters.
Illumina DNA Prep Kit	Preparation of sequencing libraries for Illumina short-read platforms.
PacBio SMRTbell Prep Kit	Preparation of libraries for long-read, high-fidelity (HiFi) sequencing.
Polycarbonate Membrane Filters (0.22-3µm) [34]	Size-fractionated collection of microbial cells from water samples.

Genomic and Functional Annotation

- **Metabolic Pathway Reconstruction:** Annotate MAGs using automated pipelines (e.g., PROKKA, DRAM) followed by manual curation. Identify key metabolic pathways (e.g., carbon fixation, sulfur respiration, nitrogen cycling) using KEGG and MetaCyc databases [34] [33].
- **Identification of Novelty:** To detect novel protein families (NPFs), compare all predicted protein sequences from MAGs against public databases (eggNOG, Pfam, RefSeq) using tools like HMMER and Diamond. Proteins with no significant homology constitute NPFs [33].
- **Calculation of Functional Diversity Metrics:** Map functional traits (e.g., KEGG orthologs, CAZymes) to calculate metrics such as functional richness (number of unique functions) and functional divergence (variance in abundance of traits) within the Marinisomatota population [36].

The following workflow diagram summarizes the core computational and laboratory processes for reconstructing and analyzing Marinisomatota MAGs.




Diagram 1: Workflow for Marinisomatota MAG Reconstruction and Analysis

Key Genomic Findings and Functional Diversity

The analysis of 1,588 Marinisomatota MAGs reveals extensive taxonomic novelty and functional versatility, positioning this phylum as a significant player in marine biogeochemistry.

Phylogenetic Diversity and Global Distribution

Phylogenomic analyses confirm that Marinisomatota forms a monophyletic lineage within the FCB superphylum [33]. The 1,588 MAGs likely encompass multiple novel classes, orders, and families, as a significant proportion of genomes from similar studies are phylogenetically distinct from previously described bacterial phyla [33]. Comparison of 16S rRNA genes from these MAGs with public databases demonstrates that Marinisomatota has a **global distribution**, with high sequence homology (>95%) to sequences retrieved from diverse habitats including coastal waters, deep-sea sediments, and hypersaline ponds [33]. This wide distribution suggests a high degree of physiological plasticity and niche adaptation.

Metabolic Versatility and Ecosystem Functioning

Metabolic reconstruction of the MAGs reveals a remarkable capacity for diverse energy generation and nutrient cycling processes, as summarized in Table 2.

Table 2: Key Metabolic Functions in Marinisomatota MAGs

Metabolic Process	Key Genes/Pathways Identified	Inferred Ecological Role
Carbon Cycling	Wood-Ljungdahl pathway; diverse Glycoside Hydrolases (GHs) for polysaccharide degradation [33]	Anaerobic degradation of complex organic matter; potential for mixotrophy and carbon fixation.
Sulfur Respiration	Dissimilatory sulfite reductase (dsrABC) [33]	Reduction of sulfur compounds for energy generation, coupling organic matter remineralization to the sulfur cycle.

Metabolic Process	Key Genes/Pathways Identified	Inferred Ecological Role
Nitrogen Metabolism	Genes for nitrate reduction (narGHI) and putative ammonification [33]	Involvement in the later stages of denitrification, contributing to nitrogen turnover in sediments.
Novel Genetic Repertoire	High proportion (~9% on average) of Novel Protein Families (NPFs) [33]	Potential for uncharacterized biochemical pathways, possibly linked to adaptation and specialized metabolisms.

The functional diversity of a community, including the range of traits expressed by groups like Marinisomatota, is a critical component of ecosystem functioning. Studies on marine copepods have established that functional diversity metrics (e.g., functional richness, divergence) have emergent covariance with key ecosystem processes like **primary production, zooplankton biomass, and carbon export efficiency** [36]. This suggests that the specific traits carried by Marinisomatota, even if they are not the most abundant taxa, could disproportionately influence these broader ecosystem functions, aligning with the mass ratio hypothesis [36].

The following diagram illustrates the integrated role of Marinisomatota in marine biogeochemical cycles, based on genomic evidence.




Diagram 2: Marinisomatota's Role in Biogeochemical Cycling

The Scientist's Toolkit: Key Reagents and Computational Tools

Table 3: Essential Resources for Genome-Resolved Metagenomic Analysis

Tool / Resource Name	Category	Primary Function in Analysis
metaSPAdes [33]	Computational Tool	De novo metagenomic assembly from sequencing reads into contigs.
CheckM2	Computational Tool	Assesses quality (completeness/contamination) of reconstructed MAGs.
GTDB-Tk [33]	Computational Tool	Provides standardized taxonomic classification of MAGs against the Genome Taxonomy Database.
PROKKA / DRAM [33]	Computational Tool	Rapid annotation of MAGs and metabolism-centric annotation, respectively.
HMMER	Computational Tool	Profile hidden Markov model searches for identifying novel protein families (NPFs).
Polycarbonate Filter Membranes (0.22µm) [34]	Laboratory Consumable	Collection of microbial biomass from water samples for DNA extraction.
ALFA-Soil DNA Kit [34]	Laboratory Consumable	Extraction of high-quality metagenomic DNA from complex sediment samples.
Illumina & PacBio Sequencers	Instrumentation	Generation of short-read and long-read sequence data, respectively.

The large-scale reconstruction of 1,588 Marinisomatota MAGs establishes this phylum as a widespread, metabolically versatile, and functionally diverse component of marine ecosystems. Its genomic toolkit for anaerobic carbon degradation, sulfur respiration, and nitrogen cycling, combined with a significant reservoir of novel protein families, underscores a profound capacity to influence oceanic biogeochemical fluxes. This study validates the thesis that functional diversity within specific bacterial lineages is a critical determinant of ecosystem functioning. The presented genomic

resource and methodological framework pave the way for future research to culture these organisms, experimentally validate predicted metabolic pathways, and quantitatively model their contribution to carbon and nutrient cycling in a changing global ocean.

The vast majority of marine microorganisms, essential players in global biogeochemical cycles, have historically resisted cultivation in laboratory settings, earning them the designation "Microbial Dark Matter" (MDM) [37]. Current estimates suggest that >99% of marine microorganisms remain uncultured and uncharacterized, creating a significant gap in our understanding of oceanic ecosystems and their functional diversity [8]. This challenge is particularly acute for the phylum **Marinisomatota** (formerly Marinimicrobia, Marine Group A, and SAR406), which are ubiquitous and abundant in marine environments yet poorly characterized due to their resistance to traditional cultivation techniques [8] [38].

The inability to cultivate these microorganisms in pure culture has profound implications for understanding the functional diversity of Marinisomatota in ocean ecosystems. Without pure cultures, detailed microbiological studies—including precise metabolic characterization, validation of genomic predictions, and exploration of biotechnological potential—remain challenging [8]. This gap underscores the critical need for innovative cultivation strategies that can bridge the divide between culture-independent genomic surveys and traditional microbiological experimentation.

The Hurdle: Why Most Marine Microbes Resist Laboratory Cultivation

The pervasive uncultivability of marine microorganisms stems from multiple factors that traditional cultivation methods fail to address. The **nutrient-rich media** typically used in laboratories create artificially high substrate concentrations that inhibit oligotrophic specialists adapted to nutrient-scarce marine environments [37]. Furthermore, laboratory conditions cannot replicate the **complex ecological interactions** and syntrophic relationships that microbial communities depend on in their natural habitats, particularly for members of the Candidate Phyla Radiation (CPR) and DPANN archaea that may have symbiotic lifestyles [37].

For Marinisomatota specifically, their metabolic versatility presents unique cultivation challenges. Recent genomic analyses have revealed that Marinisomatota members exhibit **three distinct metabolic modes**: MSo (photoautotrophic potential), MS1 (heterotrophic with enhanced glycolytic capacity), and MS2 (heterotrophic without glycolysis) [38]. This metabolic plasticity suggests they may transition between trophic strategies in response to fluctuating environmental conditions, a behavior difficult to replicate under static laboratory conditions [8] [38]. Additionally, their adaptation to specific oceanic niches—with distinct subpopulations dominating in different depth zones and water masses—indicates specialized environmental requirements that must be understood and replicated for successful cultivation [38].

Innovative Cultivation Methodologies

Diffusion-Based Cultivation Systems

Diffusion-based cultivation methods represent a paradigm shift from traditional axenic culture techniques by maintaining microorganisms in a more natural chemical environment while allowing controlled nutrient exchange. Ahmad et al. (2025) developed an innovative **diffusion-based integrative cultivation method** using modified low-nutrient media to efficiently isolate previously uncultured bacteria from marine sediments [8].

Table 1: Key Components of Diffusion-Based Cultivation Chambers


Component	Function	Specification/Composition
Semi-permeable Membrane	Allows diffusion of nutrients and signaling molecules while protecting cells from direct exposure to high nutrient concentrations	Typically 0.03-0.2 µm pore size
Low-Nutrient Media	Recreates oligotrophic conditions similar to natural marine environments	Modified with various carbon sources, mimicking natural concentrations
Inoculum Source	Provides diverse microbial community from environmental samples	Marine sediments, water samples, or microbial mats
Diffusion Chamber Design	Creates interface between natural environment and laboratory conditions	Can be deployed in situ or in laboratory simulators

This approach enabled the successful cultivation of species from rarely cultured phyla, including **Verrucomicrobiota** and **Balneolota**, significantly outperforming traditional cultivation methods [8]. Remarkably, the application of this novel technique yielded **196 isolates**, of which **115 represented previously uncultured taxa**, achieving an exceptional novelty ratio of **58%** [8].

Advanced Cultivation Workflows

The successful cultivation of fastidious marine microorganisms requires integrated workflows that combine genomic insights with tailored cultivation conditions. The following diagram illustrates a comprehensive experimental workflow for cultivating previously uncultured marine microorganisms,

incorporating both in situ and laboratory-based approaches:



This workflow begins with comprehensive **metagenomic analysis** of the source environment to identify taxonomic composition and predict metabolic capabilities of target organisms [4] [37]. For Marinisomatota, this might reveal their potential for mixotrophic adaptations, informing media design [38]. The cultivation phase then employs either **diffusion chamber systems** or **in situ incubation devices** that maintain environmental conditions while allowing microbial growth monitoring [8] [39]. Successful cultures are subsequently transferred to laboratory conditions for isolation and purification, followed by functional validation of metabolic capabilities.

Niche-Mimicking Cultivation Strategies

Cultivation success rates have improved significantly through approaches that specifically replicate the chemical and physical conditions of target microorganisms' native habitats. For example, studies of nitrite-oxidizing bacteria (NOB) in the Mariana Trench revealed clear **niche partitioning** between slope (6000-10,000 m) and bottom (>10,000 m) sediments, with distinct adaptations to pressure, nutrient availability, and oxidative stress [40]. Slope-dominant NOB possessed expanded genetic arsenals for antioxidation and osmoprotection, suggesting these functions should be supported in cultivation attempts [40].

For Marinisomatota, their distribution patterns across global oceans provide crucial insights for cultivation strategies. These organisms are predominantly found in **low-latitude marine regions**, with relative abundances ranging from **0.18% to 36.21%** across different oceanic provinces [38]. Their prevalence in specific depth zones and water masses indicates specialized environmental requirements that must be replicated for successful cultivation.

The Scientist's Toolkit: Essential Reagents and Materials

Table 2: Research Reagent Solutions for Cultivating Marine Microorganisms

Reagent/Material	Function	Application Example
Modified Low-Nutrient Media	Recreates oligotrophic conditions of marine environments	Isolation of Verrucomicrobiota and Balneolota from marine sediments [8]
Semi-permeable Membranes	Enables chemical exchange while protecting cells	Diffusion chambers for in situ cultivation [8]
Multiple In Situ Nucleic Acid Collections (MISNAC)	Preserves in situ gene expression profiles during sampling	Seasonal studies of deep-sea microbial communities [39]

Reagent/Material	Function	Application Example
GuHCl-based Lysis Buffer	Efficient nucleic acid extraction from recalcitrant cells	Metagenomic analysis of hypersaline microbial mats [37]
RNAlater Stabilization Solution	Preserves RNA integrity during sample transport	Metatranscriptomic studies of Mariana Trench sediments [40]
Casein Starch Agar (SCA)	Selective isolation of marine Actinomycetota	Drug discovery from marine sediments [41]

Marinisomatota: A Case Study in Connecting Cultivation with Functional Diversity

The relationship between cultivation success and understanding functional diversity is exemplified by recent advances in Marinisomatota research. Prior to innovative cultivation approaches, this phylum was known primarily from metagenomic studies indicating its ubiquity in marine environments [38]. The reconstruction of **1,588 Marinisomatota genomes** from global ocean datasets revealed astonishing metabolic diversity, leading to the identification of three distinct metabolic groups: MS0 (photoautotrophic potential), MS1 (heterotrophic with enhanced glycolytic capacity), and MS2 (heterotrophic without glycolysis) [38].

This genomic insight fundamentally changed cultivation strategies for these organisms. Instead of applying uniform culture conditions, researchers could now design targeted approaches based on predicted metabolic capabilities. For putative MS0 members, cultivation attempts could incorporate **light exposure** and limited organic carbon sources to select for photoautotrophic populations. For MS1 and MS2 types, varying the availability of organic carbon substrates and monitoring glycolytic activity would help distinguish these functional groups [38].

The functional diversity of Marinisomatota has significant implications for ocean ecosystem research. Their metabolic flexibility suggests they play important roles in **carbon cycling** across different oceanic provinces, potentially transitioning between autotrophic and heterotrophic strategies in response to nutrient availability and light conditions [38]. This adaptability may explain their widespread distribution and persistence in various marine environments, from surface waters to the deep sea [38].

Innovative cultivation methods are progressively overcoming the challenge of the "uncultured majority" in marine microbiology. Approaches such as diffusion-based cultivation, niche-mimicking conditions, and genomics-guided media design have demonstrated remarkable success in isolating previously uncultured taxa, achieving novelty rates as high as 58% [8]. These advances are particularly crucial for understanding the functional diversity of ecologically significant groups like Marinisomatota, whose metabolic versatility appears fundamental to their success in global oceans [38].

Future cultivation efforts will likely incorporate increasingly sophisticated multi-omics data to design targeted isolation strategies. As metagenomic and metatranscriptomic datasets continue to expand, particularly for understudied extreme environments like hadal zones and hypersaline systems, they will provide unprecedented insights into the metabolic requirements and interactions of uncultured microorganisms [37] [40] [39]. This knowledge, combined with advanced cultivation devices that better simulate natural environmental conditions, promises to further narrow the gap between molecular surveys and cultured representatives, ultimately transforming our understanding of microbial life in ocean ecosystems.

The **global ocean** represents a vast and relatively untapped reservoir of biological and biochemical diversity, serving as a critical source for discovering novel **bioactive compounds** with significant potential in **biotechnological and biomedical applications** [42] [43]. Marine microorganisms survive in diverse and often extreme environments, leading to the evolution of **specialized metabolic pathways** that produce unique secondary metabolites and enzymes not found in terrestrial organisms [43]. The past two decades have witnessed remarkable advances in sequencing technologies, enabling researchers to access the genetic potential of marine microorganisms without the limitations of traditional cultivation methods [4]. Among the myriad of marine microbes, the phylum **Marinisomatota** has emerged as a particularly promising target for bioprospecting efforts due to its extensive **functional diversity** in ocean ecosystems and its genetic potential for producing novel bioactive compounds [37].

In silico bioprospecting represents a paradigm shift in the discovery of bioactive compounds, leveraging computational approaches to mine genomic and metagenomic datasets for genes encoding potentially valuable enzymes, antimicrobial peptides, and other therapeutic molecules [4] [44]. This approach allows researchers to efficiently screen thousands of genomes or metagenome-assembled genomes (MAGs) for target sequences before engaging in costly laboratory experiments and synthesis [45]. The integration of **high-throughput sequencing** with advanced bioinformatics tools has dramatically accelerated the discovery pipeline, enabling the identification of novel CRISPR-Cas systems, antimicrobial peptides, and plastic-degrading enzymes from marine microbial genomes [4].

Computational Framework for Marine Bioprospecting

In silico bioprospecting relies on the availability of comprehensive genomic resources and specialized computational tools. The establishment of **global genome catalogues** has been instrumental in providing the raw data necessary for large-scale mining efforts. The **Global Ocean Microbiome Catalogue (GOMC)**, for instance, integrates 43,191 bacterial and archaeal genomes from publicly available marine metagenomes,

encompassing 138 distinct phyla and significantly expanding the known diversity of marine microbiomes [4]. Similarly, databases such as **MAGdb** provide curated collections of high-quality metagenome-assembled genomes (HMAGs), with 99,672 genomes meeting strict quality standards (>90% completeness and <5% contamination) [46].

Table 1: Key Genomic Resources for Marine In Silico Bioprospecting

Resource Name	Genome Count	Key Features	Relevance to Marine Bioprospecting
Global Ocean Microbiome Catalogue (GOMC)	43,191 MAGs	138 phyla; 3,470 genera; 20,295 potentially novel species	Unified catalogue from diverse marine ecosystems; Enables discovery of novel CRISPR systems, antimicrobial peptides, and enzymes
MAGdb	99,672 HMAGs	>90% completeness, <5% contamination; Covers clinical, environmental, animal categories	High-quality reference genomes; Standardized quality metrics facilitate reliable comparative genomics
Marine Microbial Dark Matter Databases	364 MAGs (Solar Lake study)	~30% classified as microbial dark matter; Includes uncultured lineages	Access to previously unexplored genetic diversity from extreme environments

The functional annotation of genomes typically employs classification systems such as **Clusters of Orthologous Groups (COGs)**, which categorize proteins into functional groups based on sequence homology [10]. Quantitative analyses have demonstrated that taxonomic rank explains approximately 41.4% of the variation in COG functional category relative abundance, with phylum-level classification accounting for 14.6% of this variance [10]. This taxonomic coherence in functional potential provides a valuable framework for guiding bioprospecting efforts toward specific microbial lineages with heightened probabilities of harboring desired metabolic capabilities.

Core Methodological Workflow

The in silico bioprospecting pipeline involves multiple interconnected steps, from data acquisition to functional validation. The following diagram illustrates a generalized workflow for genome mining and target identification:




Diagram 1: In Silico Bioprospecting Workflow. The process begins with sample collection and progresses through sequencing, assembly, and computational analysis before experimental validation.

Genome Recovery and Quality Assessment

The initial phase involves processing raw sequencing data to reconstruct microbial genomes. From 237.02 Tb of sequence data from 24,395 publicly available marine metagenomes, researchers have successfully reconstructed 43,191 medium- to high-quality MAGs with average completeness of 82.33% and minimal contamination (1.79%) [4]. **Quality assessment** is critical at this stage, with tools like **CheckM** commonly used to evaluate completeness and contamination based on the presence of single-copy marker genes [46]. High-quality MAGs should meet the **MIMAG standards** (Minimum Information about a Metagenome-Assembled Genome) with >90% completeness and <5% contamination to ensure reliable downstream analyses [46].

Functional Annotation and Target Identification

Following genome recovery, **functional annotation** identifies genes encoding potentially valuable bioactive compounds. This process typically involves:

- **Gene calling** using tools like Prodigal or MetaGeneMark
- **Homology searches** against curated databases (e.g., COG, Pfam, TIGRFAM)
- **Domain identification** to detect specific protein families of interest
- **Pathway reconstruction** to identify complete biosynthetic gene clusters

Advanced approaches incorporate **machine learning algorithms** to predict novel antimicrobial peptides or enzyme families based on conserved sequence features and structural motifs [4]. For instance, deep learning-based bioinformatics applied to marine genomes has led to the discovery of ten novel antimicrobial peptides and three enzymes capable of degrading polyethylene terephthalate (PET) [4].

Structural Prediction and Molecular Docking

For candidate bioactive compounds, **structural prediction** provides insights into function and potential applications. Tools like **I-TASSER** predict three-dimensional protein structures from amino acid sequences, while **PharmMapper** facilitates reverse docking to identify potential molecular targets [44] [45]. **Molecular docking** simulations using platforms like **PatchDock** evaluate the binding compatibility between predicted structures and target receptors, generating scores based on three-dimensional conformational compatibility and intermolecular forces [44]. These computational assessments help prioritize candidates for subsequent experimental validation.

Marinisomatota: A Case Study in Functional Diversity

Ecological Significance and Genomic Features

The phylum **Marinisomatota** represents a fascinating group of marine bacteria with considerable **biotechnological potential**. Recent studies of hypersaline microbial mats in Solar Lake, Egypt, recovered multiple Marinisomatota MAGs, revealing their **genetic capacity for polysaccharide degradation** [37]. This functional capability positions Marinisomatota as key players in carbon cycling within marine ecosystems and suggests potential applications in biofuel production and bioremediation.

Marinisomatota genomes demonstrate **adaptations to extreme environments**, including mechanisms for coping with high salinity, UV radiation, and fluctuating nutrient conditions [37]. These adaptations often involve the production of specialized enzymes (extremozymes) and secondary metabolites that stabilize cellular structures and protect against environmental stressors. The genetic machinery underlying these adaptations represents a valuable resource for bioprospecting efforts aimed at discovering novel bioactives with industrial applications.

Table 2: Functional Capabilities of Marinisomatota and Related Marine Microbes

Microbial Group	Habitat	Key Functional Capabilities	Bioprospecting Relevance
Marinisomatota	Hypersaline microbial mats	Polysaccharide degradation; Carbon cycling	Biofuel production; Bioremediation enzymes
Candidate Lokiarchaeota	Hypersaline microbial mats	Mixotrophic lifestyle; Carbon fixation	Novel metabolic pathways; Evolutionary insights
Candidate Heimdallarchaeota	Hypersaline microbial mats	Mixotrophic lifestyle; Methanol utilization	Specialty chemical production
Myxococcota	Hypersaline microbial mats	Complete photosynthetic gene cluster	Bioenergy applications; Light-harvesting pigments
Planctomycetota	Anoxic marine basins	Extremely large genomes (up to 18.4 Mb); Diverse metabolic pathways	Novel gene discovery; Complex natural products

In Silico Bioprospecting for Antimicrobial Compounds

Marinisomatota and other marine microbes represent promising sources for novel **antimicrobial peptides** (AMPs), which are increasingly important in addressing the global crisis of **multidrug-resistant pathogens** [45]. Computational approaches have been successfully applied to discover AMPs from diverse marine sources, including spider venoms and marine microorganisms [44] [45].

The process for AMP discovery involves:

- **Peptide characterization** using tools like **Heliquest** to determine physicochemical parameters (net charge, hydrophobicity, hydrophobic moment, molecular weight)
- **Target identification** through reverse docking with **PharmMapper**
- **Molecular docking** using **PatchDock** to evaluate binding affinity to microbial targets
- **Dynamics simulations** with **GROMACS** to assess interaction stability

This approach has identified potential targets for known antimicrobial peptides, including outer membrane proteins F (1MPF) and A (1QJP), flavoprotein fumarate reductase (1D4E), and ATP-dependent Holliday junction DNA helicase (1IN4) [44]. Similarly, studies on the peptide Oligoventin have identified enoyl-ACP reductase and thymidylate synthase ThyX as potential molecular targets [45].

The following diagram illustrates the molecular docking process for antimicrobial peptide discovery:




Diagram 2: Antimicrobial Peptide Discovery Pipeline. Computational workflow for identifying and validating antimicrobial peptides from sequence data to candidate selection.

Experimental Protocols and Methodologies

Metagenomic Assembly and Binning Protocol

Sample Preparation and Sequencing:

- Collect marine samples (water, sediment, or microbial mats) using sterile containers
- Extract DNA using commercial kits (e.g., DNeasy PowerSoil Kit) with modifications for difficult samples
- Perform quality control on extracted DNA using fluorometric quantification and gel electrophoresis
- Prepare sequencing libraries with appropriate insert sizes for short-read (Illumina) or long-read (PacBio, Nanopore) platforms
- Sequence using high-throughput platforms (e.g., Illumina NovaSeq) with sufficient depth (~70 million reads per sample for complex communities) [37]

Data Processing and Genome Reconstruction:

- **Quality Filtering:** Remove adapters and low-quality sequences using Fastp (vo.23.2) or similar tools
- **Metagenomic Assembly:** Assemble quality-filtered reads using metaSPAdes or MEGAHIT with optimized parameters for complex communities
- **Binning:** Recover MAGs using multiple binning tools (e.g., MetaBAT2, MaxBin2, CONCOCT) followed by integration and refinement with metaWRAP [46]
- **Quality Assessment:** Evaluate MAG quality based on completeness and contamination using CheckM with default parameters
- **Taxonomic Classification:** Assign taxonomy to MAGs using GTDB-Tk against the Genome Taxonomy Database

Molecular Docking and Dynamics Protocol

Target Identification and Preparation:

- Identify potential targets through reverse docking using PharmMapper with default settings
- Retrieve protein structures from PDB database (<https://www.rcsb.org/>)
- Prepare receptor files by removing water molecules and adding hydrogen atoms using UCSF Chimera
- Optimize structures through energy minimization (1000 steps steepest descent, 10 steps conjugate gradient) [45]

Ligand Preparation:

- Predict peptide three-dimensional structure using I-TASSER server
- Perform energy minimization with UCSF Chimera (steepest descent step size 0.02Å)
- Save structures in .MOL2 and .PDB formats for docking analyses

Molecular Docking:

- Perform docking using PatchDock server with Clustering RMSD 4.0 and Default Complex Type
- Evaluate results based on geometric shape complementarity scores
- Analyze hydrogen bonds and hydrophobic interactions using UCSF Chimera FindhBond tool (relax constraints: 2Å and 20 degrees; maximum distance 4Å)
- Validate binding sites using I-TASSER server for ligand site prediction

Molecular Dynamics Simulations:

- Conduct simulations using GROMACS (v.2022.4)
- Set simulation parameters: temperature 300K, pressure 1 bar, simulation time 10ns [44]
- Analyze trajectory files for complex stability and interaction persistence

Research Reagent Solutions for In Silico Bioprospecting

Table 3: Essential Computational Tools and Databases for Marine Bioprospecting

Tool/Resource	Category	Function	Access
GTDB-Tk	Taxonomic Classification	Standardized taxonomic assignment of microbial genomes	https://ecogenomics.github.io/GTDBTk/
CheckM	Quality Assessment	Evaluate completeness and contamination of MAGs	https://github.com/Ecogenomics/CheckM
metaWRAP	Binning Pipeline	Integrate and refine bins from multiple tools	https://github.com/bxlab/metaWRAP
I-TASSER	Structure Prediction	Predict 3D protein structures from amino acid sequences	https://zhanggroup.org/I-TASSER/
PharmMapper	Target Identification	Reverse docking for potential target identification	http://www.lilab-ecust.cn/pharmmapper/
PatchDock	Molecular Docking	Evaluate receptor-ligand binding compatibility	https://bioinfo3d.cs.tau.ac.il/PatchDock/
GROMACS	Dynamics Simulation	Molecular dynamics simulations of biomolecules	https://www.gromacs.org/
UCSF Chimera	Visualization	Molecular visualization and analysis	https://www.cgl.ucsf.edu/chimera/
Heliquest	Peptide Analysis	Determine physicochemical parameters of peptides	https://heliquest.ipmc.cnrs.fr/

In silico bioprospecting represents a powerful approach for unlocking the **functional potential** of marine microbial diversity, with **Marinisomatota** serving as an exemplary case of the promising bioactive resources awaiting discovery in ocean ecosystems. The integration of

high-throughput sequencing, advanced bioinformatics, and computational modeling has created an efficient pipeline for identifying novel enzymes, antimicrobial peptides, and other valuable compounds from marine genomes before laboratory synthesis and validation [4] [42].

Future advances in in silico bioprospecting will likely focus on improving **algorithm accuracy** for predicting protein structure and function, expanding **reference databases** with diverse marine genomes, and developing **integrated platforms** that combine multi-omics data for comprehensive functional assessment [46]. As these computational methods continue to evolve, they will dramatically accelerate the discovery of novel bioactives from marine sources, contributing to drug development, industrial biotechnology, and environmental sustainability while highlighting the invaluable functional diversity of marine microorganisms like Marinisomatota in ocean ecosystem research.

The oceans represent the planet's most extensive ecosystem, within which marine microorganisms act as the fundamental engines of global biogeochemical cycles. The direct linkage between the genetic potential of these microorganisms and the large-scale cycling of carbon and nutrients represents a frontier in marine microbial ecology. This connection is epitomized by the **functional diversity** found within specific microbial clades, which directly influences ecosystem-level processes. The advent of **genome-resolved metagenomics** has revolutionized this field by enabling researchers to move beyond mere taxonomic identification to a detailed understanding of the metabolic capabilities encoded within microbial genomes. By analyzing these genomic blueprints, scientists can now decipher the specific roles microorganisms play in carbon fixation, organic matter degradation, and nutrient transformation across diverse marine environments.

This technical guide explores the critical link between genetic potential and ecosystem function, with a specific focus on the phylum **Marinisomatota** (formerly known as Marinimicrobia, Marine Group A, and SAR406) as a model system. These organisms exemplify how genomic diversity translates to functional diversity in marine ecosystems. Through detailed analysis of their metabolic strategies, distribution patterns, and enzymatic capabilities, we can establish a mechanistic understanding of how microbial communities drive the oceanic carbon pump and nutrient regeneration processes that sustain life on Earth.

Quantitative Profiling of Marinisomatota Diversity and Distribution

Comprehensive analysis of Marinisomatota across global ocean basins has revealed remarkable patterns of abundance, diversity, and distribution. These findings provide critical baseline data for understanding their potential contribution to carbon and nutrient cycling on a global scale.

Table 1: Global Distribution and Abundance of Marinisomatota

Parameter	Value/Range	Significance
Latitudinal Preference	Low-latitude marine regions	Highest abundance in tropical and subtropical waters
Relative Abundance	0.18% to 36.21%	Can dominate microbial communities in specific niches
Genomic Diversity	1,588 genomes recovered	Represents substantial phylogenetic diversity
Taxonomic Resolution	1 class, 2 orders, 14 families, 31 genera, 67 species	Extensive radiation within the phylum
Notable Families	S15-B10, TCS55, UBA1611, UBA2128, UBA8226	Families with photoautotrophic potential

Table 2: Metabolic Strategies Identified in Marinisomatota


Metabolic Strategy	Trophic Mode	Key Characteristics	Ecological Niche
MSo	Photoautotrophic potential	Capacity for light-dependent carbon fixation via Crassulacean acid metabolism (Moo169)	Translucent zone or transition between translucent and aphotic layers
MS1	Heterotrophic	Pronounced glycolytic pathway for organic carbon degradation	Various depths, depending on organic matter availability
MS2	Heterotrophic	Lacks glycolysis; alternative pathways for carbon processing	Environments with specific organic substrate availability

The emergence of these distinct metabolic strategies likely represents an evolutionary response to nutrient limitations within the ocean, allowing Marinisomatota to occupy diverse ecological niches and maximize resource utilization in different marine environments [2] [1]. The coexistence of multiple trophic modes within a single phylum exemplifies the metabolic plasticity that enables microbial taxa to significantly influence both carbon fixation and carbon degradation pathways within oceanic ecosystems.

Methodologies for Linking Genetic Potential to Ecosystem Function

Genome-Resolved Metagenomics Workflow

Establishing robust connections between genetic potential and ecosystem function requires sophisticated methodological approaches centered on genome-resolved metagenomics. The following protocol outlines the key steps for processing marine samples from collection to functional annotation:



Sample Collection and Processing:

- Collect marine samples (water, sediment, or microbial mats) using appropriate methods (e.g., Niskin bottles for water, corers for sediments)
- Immediately preserve samples at -80°C or in DNA/RNA stabilization buffers to prevent degradation
- For metatranscriptomic studies, add RNase inhibitors and process samples rapidly to preserve RNA integrity

DNA Extraction and Sequencing:

- Extract high-molecular-weight DNA using kits specifically designed for environmental samples (accounting for diverse cell wall types and potential inhibitors)
- Assess DNA quality and quantity using fluorometric methods (e.g., Qubit) and fragment analyzers
- Prepare sequencing libraries using appropriate kits for Illumina, PacBio, or Oxford Nanopore platforms
- Sequence to sufficient depth (typically 20-50 million reads per sample for complex marine communities)

Bioinformatic Processing:

- Perform quality control using Fastp [\[1\]](#) or similar tools to remove adapter sequences and low-quality reads
- Assemble quality-filtered reads into contigs using metaSPAdes, MEGAHIT, or similar metagenomic assemblers
- Bin contigs into metagenome-assembled genomes (MAGs) using composition and coverage information with tools like MaxBin, MetaBAT, or CONCOCT
- Assess genome quality using CheckM or similar tools, reporting completeness and contamination estimates according to established standards [\[4\]](#)

Functional Annotation:

- Annotate protein-coding genes using Prokka or similar pipelines
- Perform functional assignment using KEGG, COG, Pfam, and TIGRFAM databases
- Identify specific metabolic pathways of interest (e.g., carbon fixation, sulfur oxidation, nitrogen metabolism) using specialized tools like METABOLIC or HMMER searches against custom profile HMMs
- For *Marinisomatota* specifically, screen for genes associated with Crassulacean acid metabolism (MOO169) and glycolytic pathways [\[2\]](#) [\[1\]](#)

Experimental Validation of Predicted Functions

While genomic predictions provide valuable insights, experimental validation is essential to confirm the hypothesized ecological functions. Key validation approaches include:

Metatranscriptomic Analysis:

- Extract RNA from the same environment using RNA stabilization protocols
- Remove rRNA and prepare mRNA sequencing libraries
- Sequence and map reads to reference MAGs to identify actively transcribed genes
- Compare expression levels of key metabolic genes across different environmental conditions

Heterologous Expression:

- Clone putative functional genes into expression vectors
- Express in model bacterial systems (e.g., *E. coli*)
- Purify and characterize enzyme activity in vitro
- For example, putative PETases discovered in marine microbiomes have been validated through heterologous expression and confirmation of plastic-degrading activity [\[4\]](#)


Isotopic Tracer Experiments:

- Incubate environmental samples with stable isotope-labeled substrates (e.g., ¹³C-bicarbonate, ¹⁵N-nitrate)
- Track incorporation into biomass or specific metabolic products
- Combine with single-cell sorting or density gradient centrifugation to link activity to specific phylogenetic groups

Carbon Cycling Pathways in Marine Microbiomes

Genomic Blueprints of Carbon Transformation

The genetic potential for carbon cycling represents one of the most critical links between microbial communities and ecosystem function. Different microbial groups possess distinct genetic repertoires that determine their roles in the marine carbon cycle, from CO₂ fixation to complex carbon degradation.



Marinisomatota as Model Mixotrophs in the Carbon Cycle

Marinisomatota represent a particularly interesting case study in microbial contributions to carbon cycling due to their metabolic plasticity. Genomic analysis has revealed that different members of this phylum employ distinct metabolic strategies:

Photoautotrophic Potential (MS0):

- Several Marinisomatota families (S15-B10, TCS55, UBA1611, UBA2128, and UBA8226) encode proteins associated with Crassulacean acid metabolism (M00169) [2] [1]
- This light-dependent carbon fixation pathway allows these organisms to function as primary producers in the translucent zone
- The capacity to fix inorganic carbon positions these organisms as significant contributors to carbon uptake in oligotrophic waters

Heterotrophic Carbon Processing (MS1 and MS2):

- MS1 genotypes possess complete glycolytic pathways for efficient processing of organic carbon
- MS2 genotypes lack glycolysis but utilize alternative pathways for carbon metabolism
- Certain Marinisomatota MAGs show genetic potential for polysaccharide degradation, contributing to the breakdown of complex organic matter [47] [48]
- This functional diversity enables different Marinisomatota populations to participate in various stages of organic carbon remineralization

The coexistence of autotrophic and heterotrophic strategies within a single phylum exemplifies how genomic diversity within microbial groups creates functional redundancy and resilience in marine carbon cycling processes.

Nutrient Cycling and Interconnections with Carbon Flow

Integrated Nutrient Cycling Pathways

The cycling of carbon in marine ecosystems is intrinsically linked to the transformation of essential nutrients, particularly nitrogen, phosphorus, and sulfur. Microorganisms mediate these interconnected cycles through specialized metabolic pathways encoded in their genomes.

Table 3: Microbial Genes Involved in Nutrient Cycling Processes

Nutrient Cycle	Key Metabolic Process	Representative Genes	Microbial Groups
Nitrogen Cycle	Nitrogen Fixation	<i>nifH</i> , <i>nifD</i> , <i>nifK</i>	Cyanobacteria, KSB1 bacteria [47] [48]
	Denitrification	<i>nirS</i> , <i>nirK</i> , <i>nosZ</i>	Marinisomatota, other heterotrophic bacteria [47]
Sulfur Cycle	Thiosulfate Oxidation	SOX gene complex (<i>soxA</i> , <i>soxB</i> , <i>soxX</i> , <i>soxY</i> , <i>soxZ</i>)	Asgardarchaeota, Coatesbacteria [47] [48]
	Sulfate Reduction	<i>dsrA</i> , <i>dsrB</i>	Various sulfate-reducing bacteria and archaea
Phosphorus Cycle	Organic P Hydrolysis	Alkaline phosphatase (<i>phoA</i> , <i>phoX</i>)	Diverse marine bacteria and archaea

Genomic Evidence for Coupled Biogeochemical Processes

Microbial genomes frequently reveal the genetic potential for coupled biogeochemical transformations that link multiple nutrient cycles:

Marinisomatota in Carbon-Nitrogen Coupling:

- Genomic evidence indicates that some Marinisomatota can degrade nitrogen-containing organic compounds
- This simultaneous processing of carbon and nitrogen structures positions these organisms at the interface of multiple biogeochemical cycles

Microbial Dark Matter in Extreme Environments:

- In hypersaline microbial mats like Solar Lake, previously uncharacterized microbial dark matter (including Asgardarchaeota and candidate phyla) have been found to encode genes for thiosulfate oxidation (SOX complex), nitrogen fixation, and denitrification [47] [48]
- These findings demonstrate that uncultured microbial groups play significant roles in connecting carbon, nitrogen, and sulfur cycles in specialized ecosystems

Hadal Zone Adaptations:

- In deep-sea trench sediments, microbial communities exhibit genomic adaptations for utilizing aromatic compounds (carbon cycling) under oligotrophic conditions while simultaneously maintaining antioxidant systems for high-pressure adaptation [7]
- This coupling of carbon metabolism with stress response mechanisms illustrates how environmental constraints shape functional genetic potential

The interconnection between nutrient cycles at the genomic level creates a network of biogeochemical processes that ultimately controls ecosystem-level carbon and nutrient fluxes in the ocean.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 4: Essential Research Reagents for Marine Microbial Genomics

Category	Specific Product/Kit	Application	Key Features
Sample Preservation	RNAlater, DNA/RNA Shield	Nucleic acid stabilization	Preserves integrity during storage and transport
DNA Extraction	DNeasy PowerSoil Pro Kit	Environmental DNA extraction	Effective with low biomass and inhibitor removal
RNA Extraction	RNeasy PowerMicrobiome Kit	Environmental RNA extraction	Maintains RNA integrity, removes co-purified inhibitors
Library Preparation	Illumina DNA Prep, Nextera XT	Sequencing library construction	Compatible with low-input samples, dual index barcoding
Long-read Sequencing	PacBio SMRTbell, Oxford Nanopore	Complete genome assembly	Resolves repetitive regions, complete genomes
Functional Annotation	KofamKOALA, eggNOG-mapper	Metabolic pathway prediction	KEGG orthology assignment, hidden Markov models
Genome Binning	MetaBAT2, MaxBin 2.0	MAG reconstruction	Uses sequence composition and coverage patterns

Category	Specific Product/Kit	Application	Key Features
Taxonomic Classification	GTDB-Tk	Genome-based taxonomy	Standardized taxonomy based on bacterial and archaeal genomes
Metabolic Analysis	METABOLIC, HMMER	Pathway profiling	Custom HMM profiles for key metabolic functions

The direct linkage between genetic potential and ecosystem function represents a paradigm shift in our understanding of marine biogeochemical cycles. Through genome-resolved metagenomics, we can now identify the specific microbial actors responsible for carbon fixation, organic matter degradation, and nutrient transformations that collectively regulate ocean fertility and carbon sequestration. The functional diversity within groups like Marinisomatota—spanning photoautotrophic, heterotrophic, and mixotrophic strategies—exemplifies how metabolic plasticity at the genomic level translates to ecological resilience at the ecosystem level.

Future research in this field will increasingly focus on quantifying the expression of genetic potential through metatranscriptomic and metaproteomic approaches, coupled with stable isotope probing to directly link taxonomy to function. As global sequencing efforts continue to expand [4], and as analytical methods become more sophisticated, we will develop increasingly predictive models of how marine ecosystems respond to environmental change based on the genetic makeup of their microbial communities. This knowledge is not only fundamental to understanding ocean biogeochemistry but also critical for projecting how climate change will alter the ocean's biological carbon pump and nutrient cycles that support marine life.

Navigating Research Challenges: Common Pitfalls in Characterizing Marinisomatota Function and Metabolism

Addressing Primer Biases and Detection Gaps in Community Profiling

The accurate profiling of microbial communities, essential for understanding ecosystems from the human gut to the global ocean, is fundamentally constrained by methodological choices in molecular amplification. **Primer bias**—the distortion of microbial community representation due to mismatches in primer binding sites—presents a significant challenge in molecular ecology, potentially leading to the underestimation of key taxa and flawed ecological inferences [49] [50]. This technical guide addresses these biases within the specific context of studying the functional diversity of **Marinisomatota** (formerly Marinimicrobia) in ocean ecosystems. Marinisomatota are ubiquitous and abundant in marine environments, traditionally characterized as heterotrophic microorganisms but now recognized for their metabolic versatility, including the capacity for light-dependent processes in some lineages [2]. Their ecological roles are shaped by diverse metabolic strategies, yet their comprehensive profiling is susceptible to the primer-related detection gaps discussed herein. This guide provides researchers with a structured framework to identify, quantify, and mitigate these biases through rigorous experimental design and data analysis, thereby enabling more accurate characterization of complex microbial communities.

Quantitative Evidence of Primer Bias Across Ecosystems

The impact of primer selection is quantifiable and varies significantly across different genetic markers and ecosystems. The following summaries highlight key findings from recent studies.

Table 1: Documented Primer Bias Across Genetic Markers and Ecosystems

Genetic Marker	Ecosystem	Documented Bias	Performance Summary
ITS1 & ITS2 [51]	Human Gut Mycobiome	Enhanced coverage of fungal taxa compared to 18S rRNA.	ITS2 yielded 183 OTUs, ITS1 158 OTUs, versus only 58 OTUs for 18S.
ITS2 (Plant) [49]	Large Mammalian Herbivore Diets	Consistent underrepresentation of graminoids.	>40% of graminoid species failed to amplify <i>in vitro</i> ; relative abundance was underestimated.
COI [50]	Marine Metazoan Biodiversity	Highly variable taxon recovery based on primer set.	Mismatches, especially near the 3' end, significantly reduce PCR efficacy and biodiversity estimates.
Multi-Marker Combination [51]	Human Gut Mycobiome	Outperformed single-marker approaches.	The combination of ITS1–ITS2–18S produced the highest fungal richness.

The consistency of primer bias across diverse systems underscores its fundamental nature. In fungal profiling, the choice between ITS1, ITS2, and 18S rRNA markers leads to dramatically different outcomes. One study found that **ITS2 primers yielded 183 operational taxonomic units (OTUs)**, ITS1 yielded 158 OTUs, while 18S primers detected only 58 OTUs from the same samples, highlighting a stark difference in taxonomic coverage [51]. Similarly, in plant DNA metabarcoding of herbivore diets, both tested ITS2 primer pairs (**ITS-S2F/ITS4** and **UniPlant F/R**)

consistently underrepresented graminoids, with over 40% of graminoid species failing to amplify during *in vitro* tests [49]. This bias can lead to incorrect ecological conclusions, such as overestimating diet overlap among herbivores.


The root of this bias often lies in **primer-template mismatches**. Research on the Cytochrome c Oxidase Subunit I (COI) gene for marine metazoan biodiversity found that the number and position of these mismatches are critical. Exceeding three mismatches in a single primer, or three in one primer and two in the other, can inhibit PCR entirely. Mismatches within the last 5 bases of the primer's 3' end are particularly detrimental to amplification efficiency [50]. Given the ubiquity of these issues, a single primer set is often insufficient for accurate biodiversity assessment, arguing for a multi-marker approach [51] [52].

Experimental Protocols for Bias Assessment and Mitigation

To ensure the accuracy of community profiling data, researchers must implement protocols to actively identify and account for primer biases. The following workflows provide a pathway to achieve this.

Core Workflow for Bias Evaluation

The following diagram outlines the critical stages for a robust evaluation of primer bias in any profiling study.



Detailed Experimental Procedures

1. In Silico Primer Evaluation

- **Objective:** Computationally predict primer coverage and specificity before wet-lab work.
- **Procedure:**
 - **Database Curation:** Compile a relevant, high-quality reference database. For marine microbes, this may involve downloading complete genomes from NCBI RefSeq and filtering for marine taxa using databases like WoRMS (World Register of Marine Species) [50].
 - **Mismatch Analysis:** Extract the target gene region (e.g., 16S, 18S, ITS, COI) from the reference database. Use bioinformatics scripts (e.g., in R or Python) to align primer sequences to these targets and identify the number and position of mismatches. Primers with >3 mismatches, especially in the 3' region, should be flagged [50].
 - **Coverage Estimation:** Calculate the percentage of target taxa in your reference database that would be amplified with a minimal number of mismatches.

2. Mock Community Construction & Validation

- **Objective:** Create a defined microbial community with known composition to serve as a ground truth for testing.
- **Procedure:**
 - **Design:** Create mock communities with different abundance distributions (e.g., even, dominant for a specific taxon). A study on plant primers used four types: equal, grass-dominant, forb-dominant, and tree/shrub-dominant [49].
 - **Pooling Strategies:** Employ two pooling approaches to isolate PCR bias from other effects [49]:
 - **MC-A (Pool after PCR):** Amplify each specimen's DNA separately with unique index primers, then pool the resulting amplicons in known concentrations. This controls for post-amplification steps.
 - **MC-B (Pool before PCR):** Pool the genomic DNA of all specimens in known concentrations before a single PCR amplification. This directly tests the bias introduced during amplification.

- **Validation:** Use alternative quantification methods (e.g., qPCR, fluorometry) to confirm the initial DNA concentrations in the mock community.

3. Wet-Lab Primer Testing & Amplification

- **Objective:** Empirically determine the amplification efficiency of different primer sets.
- **Procedure:**
 - **Standardized PCR:** Amplify the mock communities (both MC-A and MC-B) with each candidate primer set using the same thermocycling conditions, polymerase, and number of PCR cycles.
 - **Replication:** Perform a minimum of three technical replicates per primer set to account for stochastic PCR effects.
 - **Control:** Include a negative control (no template) to detect contamination.

4. Sequencing & Bioinformatic Processing

- **Objective:** Generate and process sequencing data to determine observed community composition.
- **Procedure:**
 - **Sequencing:** Sequence the amplicons on a platform like Illumina MiSeq or DNBSEQ.
 - **Bioinformatics:** Process raw reads using standardized pipelines (e.g., QIIME 2, USEARCH) for quality filtering, denoising, and clustering into OTUs or ASVs (Amplicon Sequence Variants) [51].
 - **Taxonomic Assignment:** Classify sequences against reference databases (e.g., SILVA for 16S/18S, UNITE for ITS) [51].

5. Bias Quantification & Decision

- **Objective:** Quantify the discrepancy between expected and observed compositions to guide primer selection.
- **Procedure:**
 - **Statistical Comparison:** Compare the expected composition of the mock community to the observed sequencing results. Calculate metrics like **Fold-Change** for each taxon: $(\text{Observed Relative Abundance} / \text{Expected Relative Abundance})$.
 - **Amplification Efficiency:** For a taxon, a fold-change of 1 indicates perfect efficiency; <1 indicates under-representation; >1 indicates over-representation.
 - **Decision:** Select the primer set that shows the highest fidelity to the expected mock community structure or the smallest bias against your taxa of interest (e.g., Marinisomatota).

A Case Study: From Bias to Functional Insights in Marinisomatota

Applying these bias-aware methods is crucial for accurate profiling of specific groups like **Marinisomatota**. These bacteria are abundant in global oceans and exhibit remarkable metabolic versatility, with recent studies identifying three distinct strategies: photoautotrophic potential (MS0), heterotrophic with pronounced glycolysis (MS1), and heterotrophic without glycolysis (MS2) [2]. Primer bias in 16S rRNA surveys could easily lead to the underrepresentation of certain Marinisomatota lineages, distorting our understanding of their ecological roles.

Once communities are accurately profiled, tools like **METABOLIC** can be used to infer functional potential from genomic data. METABOLIC is a scalable software that annotates microbial genomes, validates protein motifs, analyzes metabolic pathways, and calculates contributions to biogeochemical cycles [52]. It can process genomes from isolates, metagenome-assembled genomes (MAGs), or single-cell genomes.

For a community-scale analysis, METABOLIC can incorporate metagenomic read mappings to determine genome abundance. It then reconstructs **functional networks** based on "metabolic handoffs"—where sequential metabolic transformations are distributed among different organisms [52]. This allows researchers to visualize the interconnected roles of community members, including Marinisomatota, in driving processes like carbon and sulfur cycling. METABOLIC produces outputs such as heatmaps, functional network diagrams, and Sankey diagrams illustrating microbial contributions to biogeochemical processes, providing a systems-level view of the community's functional potential [52].

Table 2: Research Reagent Solutions for Primer Bias and Functional Analysis

Reagent / Tool	Function / Application	Specifications & Considerations
Mock Communities [49]	Ground truth for evaluating primer bias.	Should include taxa and abundance distributions relevant to the study ecosystem (e.g., marine samples).
DNBSEQ / Illumina Systems [51]	High-throughput amplicon sequencing.	Platform choice affects read length and cost; ensure compatibility with amplicon size.
SILVA & UNITE Databases [51]	Taxonomic classification of 16S/18S and ITS sequences.	Requires regular updating to ensure comprehensive taxonomic coverage.
METABOLIC Software [52]	Functional profiling of genomes/MAGs; biogeochemical cycle inference.	Integrates KEGG, Pfam, TIGRfam, and custom HMMs for comprehensive annotation.

Reagent / Tool	Function / Application	Specifications & Considerations
GTDB (Genome Taxonomy Database) [29]	Standardized microbial taxonomy based on genomes.	Essential for consistent taxonomic naming of novel groups like Marinisomatota.
KOfam, TIGRfam, Pfam HMMs [52]	Hidden Markov Model databases for protein family annotation.	Used by METABOLIC with manually curated cutoffs to minimize false positives.

Primer bias is an inherent and significant challenge in molecular community profiling that cannot be ignored. As demonstrated across diverse ecosystems, the choice of genetic marker and primer set directly and measurably impacts the perceived diversity, composition, and dynamics of microbial communities. A commitment to rigorous, bias-aware methodologies—combining *in silico* checks, mock community validation, and multi-marker approaches—is fundamental for generating reliable data. For the study of complex groups like Marinisomatota, moving beyond biased amplicon sequencing to **genome-resolved metagenomics** and functional tools like METABOLIC is the path forward. This integrated strategy allows researchers to not only identify which microorganisms are present but also to accurately delineate their metabolic strategies and interconnected roles in global biogeochemical cycles, turning robust profiling data into meaningful ecological insight.

Resolving Functional Annotations for Hypothetical and Novel Genes

In the evolving field of marine microbial genomics, a significant portion of sequenced genes remains functionally uncharacterized. These hypothetical proteins (HPs)—predicted from genomic sequences but lacking experimental validation of their biological functions—represent a substantial knowledge gap in understanding ocean ecosystems. In marine bacteria like the widespread **Marinisomatota** phylum (formerly known as Marinimicrobia, Marine Group A, and SAR406), hypothetical genes can constitute a considerable proportion of genomic content [2]. Resolving their functions is critical for advancing our understanding of marine microbial ecology, biogeochemical cycling, and biotechnological applications.

The challenge is particularly acute in marine systems where microbial diversity vastly exceeds current cultivation capabilities. As of 2024, global ocean microbiome catalogs encompass over 43,000 metagenome-assembled genomes (MAGs) spanning 138 phyla, with a striking 43% of bacterial and archaeal genomes representing potentially novel taxa [4]. Within this reservoir of genetic novelty, hypothetical genes represent both a challenge and an opportunity—they may encode novel metabolic pathways, unique adaptations to marine environments, or biotechnologically valuable enzymes, but their characterization requires sophisticated computational and experimental approaches.

This technical guide provides a comprehensive framework for resolving functional annotations of hypothetical and novel genes, with specific emphasis on methodologies relevant to **Marinisomatota** research. We integrate cutting-edge computational predictions with experimental validation strategies, contextualized within the unique constraints and opportunities of marine microbiome studies.

Computational Prediction and Prioritization Pipelines

Structural and Functional Annotation Workflows

Initial functional annotation requires integrated bioinformatics pipelines that combine multiple prediction algorithms. The core workflow begins with genome quality assessment and proceeds through sequential layers of analysis (Figure 1). For Marinisomatota genomes, which are primarily obtained through metagenomic assembly, quality verification using tools like CheckM is essential to ensure reliable downstream analyses [53].

Table 1: Key Bioinformatics Tools for Functional Annotation

Tool Category	Specific Tools	Primary Function	Considerations for Marine Microbes
Genome Quality Assessment	CheckM	Assess completeness and contamination	Crucial for metagenome-assembled genomes
Structural Annotation	PROKKA, RAST	Gene calling and initial function assignment	Performance varies with atypical GC content
Domain Identification	InterProScan, Pfam	Protein family classification	Identifies conserved domains in novel proteins
Transmembrane Prediction	TMHMM	Helical transmembrane regions	Important for transporter identification
Subcellular Localization	PSORTb, Cell-PLoc	Cellular compartment prediction	Informed by gram-status (often negative for marine bacteria)
Tertiary Structure Prediction	AlphaFold2, ColabFold	3D protein structure modeling	Reveals functional insights when sequence similarity is low

Advanced structural annotation leverages both ab initio gene prediction and homology-based methods. For *Marinisomatota*, with their distinctive genomic features including potential mixotrophic capabilities, specialized attention should be paid to metabolic pathway genes that may be underrepresented in reference databases [2]. Subsequent functional annotation employs integrated pipelines that combine results from multiple databases including **Pfam**, **TIGRFAM**, **COG**, and **GO** through platforms like **InterProScan** [53].

Prioritization Strategies for Hypothetical Proteins

With typically dozens to hundreds of hypothetical proteins per genome, strategic prioritization is essential for efficient research. The following criteria help identify HPs with high probability of functional significance:

- **Essentiality:** Determine whether genes are essential using database comparisons (e.g., DEG) or genomic context [53].
- **Conservation:** Identify HPs with orthologs across multiple *Marinisomatota* lineages or related phyla.
- **Genomic Context:** Analyze operonic structures and gene neighborhoods for guilt-by-association predictions [54].
- **Expression Evidence:** Prioritize HPs with supporting transcriptomic or proteomic data.
- **Structuredness:** Assess predicted secondary structure and disorder; well-structured proteins often have more definable functions.

For *Marinisomatota* research, particular attention should be paid to HPs located near genes with established roles in marine nutrient cycling, light-harvesting, or stress response, as these may represent novel components of known adaptive strategies [2].

Advanced Functional Prediction Methodologies

Genomic Language Models and Semantic Design

A transformative approach in functional prediction leverages genomic language models like **Evo**, which learns semantic relationships across prokaryotic genes by training on vast genomic datasets [54]. These models employ a "guilt-by-association" principle at scale, recognizing that functionally related genes often cluster together in prokaryotic genomes.

The semantic design methodology (Figure 2) enables function-guided discovery through several applications:

- **Contextual Autocomplete:** Partial gene sequences can be completed with functionally consistent extensions, useful for fragmentary metagenomic data.
- **Operon Prediction:** Generating missing components of multi-gene systems based on genomic neighbors.
- **Novel Enzyme Discovery:** Designing sequences for desired catalytic activities based on contextual prompts.

For *Marinisomatota* research, this approach could generate hypotheses about the functions of hypothetical genes located in defense islands, photosynthetic gene clusters, or nutrient utilization operons [54]. The generated sequences can be tested for structural stability and functional potential before experimental validation.

Structure-Based Functional Inference

When sequence similarity offers limited functional insights, tertiary structure predictions can reveal unexpected relationships. Tools like **AlphaFold2** and **ColabFold** enable accurate protein structure prediction even for novel sequences [53]. Subsequent structural comparison against databases like **PDB** and **FoldSeek** can identify distant homologs with established functions.

For *Marinisomatota* hypothetical proteins, structural analysis might reveal:

- Novel catalytic triads in putative enzymes
- DNA-binding motifs in regulatory proteins
- Transmembrane domains in potential transporters
- Protein-protein interaction interfaces

Particularly valuable is the identification of structural motifs associated with marine-relevant functions such as light-sensing domains, osmoregulatory components, or heavy metal binding sites [55].

Experimental Validation Frameworks

Molecular Cloning and Heterologous Expression

Functional characterization typically requires recombinant protein production. For marine microbes like *Marinisomatota*, this presents special challenges due to potential codon usage biases, requirement for specific ion concentrations, or unusual temperature optima.

Table 2: Experimental Approaches for Functional Validation

Method Category	Specific Techniques	Information Gained	Marine-Specific Considerations
Heterologous Expression	E. coli, Vibrio systems	Protein production for biochemical assays	Codon optimization often needed for high GC genes
Phenotypic Screening	Growth complementation	Reveals role in metabolic pathways	Marine-simulating media conditions
Protein Interaction	Yeast two-hybrid, AP-MS	Identifies interaction partners	May require marine-like salt conditions
Enzyme Assays	Spectrophotometry, HPLC	Catalytic activity measurement	Temperature and pH reflecting native habitat
Localization Studies	Fluorescence tagging	Subcellular distribution	Membrane integrity in osmotic stress

Essential steps in cloning *Marinisomatota* genes include:

- **Codon Optimization:** Adapt rare codon usage to expression host
- **Vector Selection:** Choose appropriate promoters and tags
- **Expression Screening:** Test different temperatures and inducers
- **Solubility Enhancement:** Use fusion tags or specialized strains

Successful examples include the expression of novel polyethylene terephthalate-degrading enzymes from marine microbes, where activity was confirmed *in vitro* after computational prediction [4].

Functional Assays for Marine-Relevant Activities

Hypothesized functions should be tested through targeted biochemical assays:

- **Enzymatic Activities:** For predicted enzymes, establish assays with putative substrates including marine-specific compounds like dimethylsulfoniopropionate (DMS), methylphosphonate, or algal polysaccharides.
- **Metabolic Complementarity:** Introduce genes into mutant strains lacking specific metabolic capabilities to test functional rescue.
- **Binding Studies:** Use surface plasmon resonance or microscale thermophoresis to test interactions with potential ligands.
- **Cellular Phenotypes:** Express genes in model organisms to observe morphological or growth changes.

For *Marinisomatota*, particular attention should be paid to testing predicted functions related to their documented metabolic versatility, including potential involvement in light-driven metabolism, organic matter degradation, or specialized nutrient acquisition [2] [56].

Case Studies in Marine Microbial Genomics

Functional Prediction in *Marinisomatota*

Recent research on *Marinisomatota* illustrates the power of integrated approaches. Xiang et al. (2025) reconstructed 1,588 *Marinisomatota* genomes from global ocean datasets, identifying three distinct metabolic strategies: MSo (photoautotrophic potential), MS1 (heterotrophic with enhanced glycolytic capacity), and MS2 (heterotrophic without glycolysis) [2]. This classification was enabled by:

- **Comparative Genomics:** Identifying genes and pathways differentially present across metabolic types
- **Phylogenomic Analysis:** Mapping metabolic traits onto evolutionary relationships
- **Metatranscriptomics:** Validating expression of key genes in environmental samples

The study revealed that certain *Marinisomatota* families encode proteins for Crassulacean acid metabolism (MOO169), explaining their ability to transition between photic and aphotic zones [2]. This discovery emerged from persistent investigation of previously hypothetical genes adjacent to known metabolic markers.

Mercury Biotransformation Genes in Deep-Sea Sediments

A comprehensive study of mercury cycling genes in deep-sea sediments demonstrates the value of targeted functional screening. By analyzing 101 sediment layers from the South China Sea to the Mariana Trench, researchers identified novel Hg-methylating microorganisms affiliated with Desulfobacterota, Spirochaetota, and Zixibacteria [55]. The methodology included:

- **Hidden Markov Model Searches:** Using custom models to identify hgcAB and mer genes
- **Genomic Context Analysis:** Correlating gene presence with geochemical measurements
- **Phylogenetic Placement:** Determining evolutionary relationships of novel genes
- **Adaptation Signature Detection:** Identifying deep-sea-specific sequence features

This approach expanded the diversity of known Hg-transforming taxa and revealed unique ecophysiological adaptations, providing a template for investigating other biogeochemical cycles [55].

The Scientist's Toolkit: Research Reagent Solutions

Table 3: Essential Research Reagents for Functional Genomics

Reagent Category	Specific Examples	Function in Workflow	Implementation Notes
Cloning Systems	pET vectors, Gateway	Recombinant protein expression	Codon-optimized for GC-rich marine genes
Expression Hosts	E.BL21, Vibrio natriegens	Heterologous protein production	Marine-adapted hosts for difficult proteins
Annotation Software	rTOOLS, Pharokka	Phage genome annotation	Superior functional annotation performance [57]
Protein Purification	His-tag resins, SEC	Isolation of recombinant proteins	May require specialized buffers mimicking marine conditions
Activity Assays	PNPP, ONPG substrates	Enzymatic activity detection	Marine-relevant substrates enhance ecological relevance
Antibiotic Resources	Custom resistance markers	Selection in cloning workflows	Consider marine bacterial susceptibility profiles

Resolving the functions of hypothetical and novel genes in Marinisomatota and other marine microbes requires increasingly sophisticated integration of computational prediction and experimental validation. As genomic language models advance and structural prediction becomes more accurate, the marine microbiology field is poised to transition from cataloging genetic diversity to understanding functional significance.

The unique adaptations of Marinisomatota to oceanic environments—from their metabolic flexibility to their distribution across ocean depths—make them particularly rewarding targets for functional genomics. By applying the methodologies outlined in this guide, researchers can accelerate the discovery of novel biological mechanisms with potential applications in biotechnology, climate science, and drug development.

Future directions will likely include high-throughput functional screening, single-cell omics, and machine learning approaches that integrate multiple data types to predict gene function with increasing accuracy. As these tools mature, the vast repository of hypothetical genes in marine microbes will transform from a catalog of mysteries into a roadmap for biological discovery.




Figure 1. Workflow for functional annotation of hypothetical proteins from Marinisomatota and other marine microbes. The pipeline integrates computational predictions with experimental validation, with specific considerations for marine microbial genomics.




Figure 2. Semantic design approach using genomic language models for functional discovery. This methodology leverages the genomic context of known genes to generate novel sequences with related functions, applicable to understanding *Marinisomatota* metabolic diversity.

Differentiating True Autotrophy from Heterotrophic Capabilities

Within marine microbial ecology, accurately distinguishing obligate autotrophs from metabolically versatile organisms is fundamental for understanding carbon cycling and energy flow in ocean ecosystems. This differentiation is particularly critical for phyla like **Marinisomatota** (formerly SAR406), which are abundant in global oceans and exhibit complex metabolic strategies that blur traditional trophic classifications [1]. Relying solely on phylogenetic markers or the presence of a few key genes can lead to misclassification, as the genetic potential for autotrophy does not necessarily equate to its ecological implementation. This technical guide provides researchers with a structured framework, combining genomic, experimental, and analytical approaches to conclusively determine an organism's trophic mode, with a specific focus on its application within the context of *Marinisomatota* research.

Defining True Autotrophy and Key Differentiators

True autotrophy is defined as the ability of an organism to use inorganic carbon (CO_2 or HCO_3^-) as its sole carbon source for biomass synthesis, deriving energy from light (photoautotrophy) or the oxidation of inorganic chemicals (chemoautotrophy). The core differentiator from heterotrophy is the *assimilation of inorganic carbon* versus the *assimilation of organic carbon compounds*.

The challenge in differentiation arises from widespread **mixotrophy** (the capability of both autotrophic and heterotrophic metabolism) and the presence of heterotrophic contaminants in cultivation experiments. Key physiological and genomic differentiators are summarized in Table 1.

Table 1: Key Differentiators Between True Autotrophy and Heterotrophic Capabilities

Feature	True Autotrophy	Heterotrophy
Primary Carbon Source	Inorganic carbon (CO_2 , HCO_3^-)	Organic carbon compounds
Energy Source	Light (photo-) or inorganic chemical oxidation (chemo-)	Oxidation of organic compounds
Essential Metabolic Pathways	Complete carbon fixation pathways (e.g., Calvin Cycle, rTCA, 3-HP/4-HB)	Transporters and pathways for organic carbon degradation (e.g., glycolysis)

Feature	True Autotrophy	Heterotrophy
Stable Isotope Probing (SIP)	> ¹³ C-bicarbonate/ ¹⁴ CO ₂ into biomass in the absence of organic carbon	> ¹³ C- or ¹⁴ C-labeled organic compounds into biomass
Growth Validation	Growth in minimal medium with CO ₂ as sole carbon source	Requires one or more organic carbon sources for growth

For Marinisomatota, recent metagenomic studies have revealed a phylum in metabolic transition. While traditionally characterized as heterotrophic, certain members possess genes for **proteorhodopsin-based light harvesting** and Crassulacean acid metabolism (CAM)-like carbon fixation pathways [1]. This does not necessarily indicate full photoautotrophy but suggests a **mixotrophic strategy** where light energy is potentially used to enhance survival under nutrient limitation, complicating clear-cut classification.

Methodological Approaches for Differentiation

A multi-faceted approach is required to unambiguously assign a trophic mode. The following methodologies should be employed in concert.

Genomic and Metagenomic Analysis

Genome-resolved metagenomics provides the foundational blueprint for an organism's metabolic potential.

- **Target:** Identify key autotrophy marker genes.
- **Protocol:**
 - **Genome Assembly & Curation:** Recover high-quality metagenome-assembled genomes (MAGs) or isolate genomes. Standards suggest >80% completeness and <5% contamination for reliable metabolic inference [4].
 - **Pathway Identification:** Use tools like METASCAPE, KEGG, or MetaCyc to screen for complete carbon fixation pathways. The presence of a full pathway (e.g., all genes of the Calvin-Benson-Bassham cycle) is a minimum prerequisite.
 - **Contextual Gene Analysis:** Assess the genomic context. The co-localization of carbon fixation genes with light-harvesting genes (e.g., proteorhodopsin) or inorganic electron transport chains (e.g., for Fe²⁺, S⁰ oxidation) strengthens the case for autotrophy.
 - **Expression Profiling:** Conduct metatranscriptomic or metaproteomic analysis to confirm the expression of key autotrophic genes *in situ* [1]. The mere presence of genes is insufficient.

Stable Isotope Probing (SIP)

SIP provides direct evidence of inorganic carbon assimilation into biomass.


- **Target:** Quantify the incorporation of ¹³C-labeled bicarbonate into biomass.
- **Protocol:**
 - **Incubation:** Incubate a microbial sample (enrichment culture, isolated cells, or filtered biomass) with ¹³C-labeled sodium bicarbonate (e.g., 1-5 mM final concentration) in a minimal medium. **Crucially, the medium must contain no organic carbon sources** to rule out heterotrophic uptake of minerogenic CO₂.
 - **Control:** Run a parallel incubation with ¹²C-bicarbonate.
 - **Incubation Conditions:** Maintain conditions (temperature, light, pressure) that are environmentally relevant to the source organism.
 - **Biomass Recovery & Analysis:** After incubation, collect cells by filtration. Analyze the isotopic enrichment of bulk biomass or specific biomarkers (like phospholipid-derived fatty acids, DNA/RNA) using **Isotope-Ratio Mass Spectrometry (IRMS)** or **Nanoscale Secondary Ion Mass Spectrometry (NanoSIMS)** for single-cell resolution [58].

Cultivation-Based Assays

Pure culture or highly enriched culture studies remain the gold standard for validating autotrophy.

- **Target:** Demonstrate growth with CO₂ as the sole carbon source.
- **Protocol:**
 - **Medium Design:** Prepare a strictly inorganic mineral medium. A carbon-free base is supplemented with bicarbonate (e.g., 10-30 mM NaHCO₃) as the sole carbon source. The gas headspace should be air or a CO₂-enriched mix (e.g., 5% CO₂).
 - **Growth Monitoring:** Track growth over multiple serial transfers into fresh inorganic medium using cell counts (e.g., flow cytometry [59]), protein content, or optical density. A consistent increase in biomass confirms autotrophic potential.
 - **Control for Contamination:** Include controls with added organic carbon (e.g., acetate, pyruvate) to test for heterotrophic growth. The absence of growth in the inorganic medium but growth in the organic-amended medium indicates obligate heterotrophy.
 - **Advanced Cultivation:** For fastidious organisms, consider diffusion chambers or co-culture with other bacteria that may provide essential growth factors without providing a carbon source [60].

The following workflow diagram illustrates the integration of these key methodological approaches:



The Scientist's Toolkit: Essential Research Reagents and Materials

Successful differentiation of trophic strategies relies on specific reagents and instruments. Key materials for these experiments are listed below.

Table 2: Essential Reagents and Materials for Autotrophy Research


Item	Function/Application	Example Use Case
¹³C-Labeled Sodium Bicarbonate	Tracer for inorganic carbon assimilation in SIP experiments.	Quantifying CO ₂ fixation rates in Marinisomatota enrichments.
Inorganic Mineral Medium	Provides essential nutrients without organic carbon for cultivation assays.	Formulating a medium to test for autotrophic growth of a novel isolate.
Sterivex (0.22 μm) Cartridge Filters	For concentrating microbial biomass from large water volumes for DNA/SIP.	Collecting biomass from oligotrophic ocean waters for metagenomics [61].
DNA/RNA Isolation Kits (PowerSoil, etc.)	Nucleic acid extraction from environmental samples and cultures.	Preparing template for 16S rRNA amplicon or metagenomic sequencing.
Flow Cytometer	For precise cell counting and sorting to monitor growth or assess purity.	Tracking population growth in autotrophic cultivation experiments [59].
NanoSIMS	Provides high-resolution imaging and quantification of isotopic labels at the single-cell level.	Confirming ¹³ C-bicarbonate incorporation into individual Marinisomatota cells.
Metagenomic Assembly & Annotation Software (e.g., anvi'o, GTDB-Tk)	Reconstruction of MAGs and functional profiling.	Identifying carbon fixation pathway genes in Marinisomatota genomes [1] [4].

Case Study: Application to Marinisomatota

The global distribution and ecological success of Marinisomatota have long been enigmatic. Applying the above framework has been pivotal in redefining their functional diversity.

- **Genomic Evidence:** Analysis of 1,588 *Marinisomatota* genomes revealed three distinct metabolic strategies: **MS0** (possessing photoautotrophic potential via proteorhodopsin and CAM-like carbon fixation genes), **MS1** (heterotrophic with strong glycolytic pathways), and **MS2** (heterotrophic without glycolysis) [1]. This genomic diversity immediately signals that the phylum cannot be classified under a single trophic mode.
- **Ecological Interpretation:** The presence of light-harvesting and carbon fixation genes in the MS0 group suggests a **mixotrophic lifestyle**. This is hypothesized as an adaptation to nutrient-limited oceans, where the ability to harness light energy can supplement heterotrophic carbon acquisition, providing a competitive advantage in the deep chlorophyll maximum where light is dim but organic carbon is scarce [1]. True autotrophy (obligate dependence on this pathway) remains unconfirmed.
- **Research Imperative:** This case highlights the necessity of moving beyond genomic prediction to experimental validation. Future work must employ SIP and targeted cultivation of MS0 *Marinisomatota* to determine if they can indeed function as true autotrophs or if their autotrophic genes serve a different physiological purpose.

The following diagram summarizes the metabolic strategies identified within the *Marinisomatota* phylum and the key genes used for their classification:



Differentiating true autotrophy from heterotrophy, especially in versatile phyla like *Marinisomatota*, requires a rigorous, multi-pronged investigative framework. Reliance on any single line of evidence is insufficient. Genomic predictions must be validated by direct measurement of metabolic activity via SIP and cultivation. The application of this integrated approach is reshaping our understanding of the ocean's carbon cycle, revealing a world of microbial metabolic complexity where mixotrophy may be a key strategy for survival and success. For *Marinisomatota*, the confirmed presence of multiple metabolic strategies within the phylum underscores the importance of species- and strain-level analysis to accurately define their role in marine ecosystem function and biogeochemical processes [1] [62].

Strategies for Isolating and Maintaining Pure Cultures in the Lab

The phylum ***Marinisomatota*** (formerly known as *Marinimicrobia*, Marine Group A, and SAR406) represents a ubiquitous and abundant group of microorganisms in marine ecosystems, yet remains largely uncultured and physiologically enigmatic [2] [1]. Genomic analyses suggest they play significant roles in global biogeochemical cycles, with recent metagenomic studies revealing astonishing diversity—1,588 genomes representing 14 families, 31 genera, and 67 species, with relative abundances reaching up to 36.21% in low-latitude marine regions [1]. Despite this pervasive presence, a profound gap exists between their detected genomic diversity and our ability to cultivate them in laboratory settings, limiting our understanding of their functional roles in ocean ecosystems.

Closing this gap through advanced isolation and maintenance strategies is imperative for a comprehensive thesis on the functional diversity of *Marinisomatota*. While genomic data can predict metabolic potential, only through pure culture experimentation can researchers validate these predictions, elucidate physiological capabilities, and understand the complex interactions that define their ecological niches. This guide provides detailed methodologies for isolating and maintaining pure cultures of fastidious marine microorganisms, with specific application to *Marinisomatota*, enabling researchers to bridge the divide between genomic predictions and physiological reality.

Physiological Background and Ecological Context of *Marinisomatota*

Understanding the metabolic capabilities and environmental preferences of *Marinisomatota* is foundational to developing successful isolation strategies. Current knowledge, derived primarily from metagenome-assembled genomes (MAGs), indicates this group exhibits remarkable metabolic plasticity. Three distinct metabolic strategies have been identified: **MS0** (photoautotrophic potential), **MS1** (heterotrophic with pronounced glycolytic pathway), and **MS2** (heterotrophic without glycolysis) [1]. This metabolic diversity likely explains their success across diverse marine habitats, from the photic zone to the deep sea.

Certain Marinisomatota families, including S15-B10, TCS55, UBA1611, UBA2128, and UBA8226, possess genetic potential for light-dependent processes associated with Crassulacean acid metabolism [1]. This suggests some lineages may employ a mixotrophic strategy, combining heterotrophic and phototrophic capabilities—a crucial consideration when designing cultivation media and conditions. Their distribution patterns indicate adaptation to specific environmental gradients, with depth, nutrient availability, and light penetration serving as key selective factors [2].

The historical difficulty in cultivating Marinisomatota and similar marine lineages stems from our incomplete understanding of their specific nutritional requirements, symbiotic dependencies, and sensitivity to oxygen and pressure changes. Furthermore, many marine microorganisms exist in a state of metabolic dormancy or require specific signaling compounds for activation, making traditional cultivation approaches insufficient [63]. The strategies outlined below address these challenges through targeted, ecologically-informed approaches.

Sample Collection and Preservation Strategies

Proper collection and preservation of environmental samples is the critical first step in any isolation campaign, as it preserves community structure and viability for subsequent cultivation efforts.

Table 1: Comparison of Sample Preservation Methods for Microbial Diversity Studies

Preservation Method	Impact on Microbial Diversity	Impact on Community Structure	Recommended Storage Duration	Best Use Cases
Dry Ice (+ subsequent -80°C)	Minimal impact	Minimal impact	1 day on dry ice + 5 days at -80°C	Field collections, shipping
Liquid Nitrogen (+ subsequent -80°C)	Minimal impact	Minimal impact	1 day in LN2 + 5 days at -80°C	Critical samples, long-term viability
Freezing at -20°C	Moderate to high impact	Altered community structure	Not recommended beyond 3 days	Temporary holding only
Refrigeration at 4°C	Moderate to high impact	Altered community structure, especially rare taxa	Not recommended beyond 3 days	Short-term transport

For intestinal microbiota of marine fishes, which can serve as a model for host-associated Marinisomatota, dry ice and liquid nitrogen methods with subsequent transfer to -80°C storage demonstrated the least impact on microbial diversity and community structure [64]. Refrigeration at 4°C and freezing at -20°C resulted in suboptimal reproducibility and significantly altered community structure, particularly affecting rare microbial taxa [64]. These findings are applicable to sediment and water column sampling for Marinisomatota, as maintaining the integrity of rare community members is essential for isolating representatives of this typically low-abundance phylum.

Sample Collection Protocol for Marine Sediments

- **Sample Using Cores:** Collect sediment samples using push corers or box corers to maintain stratigraphic integrity. For hydrothermal sediments as in Guaymas Basin, samples are typically collected from multiple horizons (e.g., 0–2 cm, 3–4 cm, and 5–10 cm below seafloor) [5].
- **Anoxic Preservation:** Immediately transfer samples into wide-mouth glass bottles with minimal headspace to preserve anaerobic conditions crucial for many Marinisomatota lineages.
- **Temperature Control:** Maintain samples at *in situ* temperature during transport using insulated containers. For deep-sea samples, this may require cooling rather than warming.
- **Rapid Processing:** Begin isolation procedures within 24 hours of sample collection whenever possible, even when using optimal preservation methods.

Isolation Techniques and Cultivation Strategies

Enrichment Strategies

Strategic enrichment cultures can significantly increase the relative abundance of target organisms, making subsequent isolation feasible. For previously uncultured groups like Gaopeijiales (a marine lineage of Gemmatimonadota), aerobic enrichment successfully increased relative abundance from 0.37% in the original sample to 2.6% in the enrichment culture [63].

Table 2: Enrichment Culture Formulations for Targeting Different Metabolic Groups

Target Metabolism	Carbon Source	Electron Acceptor	Incubation Temperature	Key Additives	Potential Marinisomatota Group
Heterotrophic Polysaccharide Degraders	Mix of polysaccharides (chitin, cellulose, pectin, alginate)	Sulfate or oxygen	15-25°C (mesophilic)	Vitamin mix, trace elements	Families with multiple CAZymes [5]
Heterotrophic Protein Degraders	Bovine Serum Albumin (BSA)	Sulfate or oxygen	15-25°C (mesophilic)	Vitamin mix, trace elements	Putative peptide transporters
Mixotrophic Types	Bicarbonate + organic acids	Oxygen (microaerophilic)	15-25°C (mesophilic)	Vitamin mix, neutral density filters	MSo-type Marinisomatota [1]
Hadal Zone Adaptations	Aromatic compounds + lipids	Sulfate or nitrate	4-10°C (psychrophilic)	Antioxidants, pressure vessels	Deep-sea adapted lineages

For hydrothermal sediments, such as those from Guaymas Basin, successful enrichment cultures have been established at 60°C for 4 weeks under oxic, sulfate-reducing, and fermentative conditions [5]. However, for most Marinisomatota, moderate temperatures (15-25°C) are more appropriate, reflecting their typical pelagic and mesophilic habitats.

Direct Isolation Methods

Aerobic Enrichment and Dilution-to-Extinction Protocol:

- **Preparation of Inoculum:**
 - Create sediment slurries using a 1:10 ratio of sediment to anoxic artificial seawater medium [5].
 - For water column samples, concentrate cells via gentle filtration (0.22 µm filters) and resuspend in sterile medium.
- **Medium Formulation:**
 - Base saltwater medium should mimic natural seawater composition.
 - Include a carbon source cocktail reflecting Marinisomatota's predicted capabilities: combinations of polysaccharides (chitin, cellulose, pectin), organic acids (acetate, pyruvate), and bicarbonate [5] [1].
 - Add vitamin and trace element supplements, as marine oligotrophs often require specific growth factors.
- **Enrichment Incubation:**
 - Incubate under a range of oxygen conditions (oxic, microaerophilic, anoxic) to capture different metabolic types.
 - Use low nutrient concentrations (typical of marine environments) to avoid favoring fast-growing generalists.
- **Dilution-to-Extinction:**
 - Serially dilute enriched cultures in fresh medium (typically 10-fold serial dilutions).
 - Dispense into multi-well plates (96-well format is efficient).
 - Incubate for 4-12 weeks, monitoring for growth via turbidity or microscopic examination.
- **Purity Assessment:**
 - Use phase-contrast microscopy to check for morphological uniformity.
 - Employ 16S rRNA gene sequencing to confirm culture purity.
 - Use BOX-PCR profiling to genetically distinguish closely related strains [63].

High-Throughput Isolation Using Automated Systems

For processing large sample sets, automated systems can significantly improve isolation efficiency:

- **Robotic Liquid Handling:** Systems like the ROTOR robot (Singer Instrument Company) can perform high-throughput sample transfer, picking, and arraying in multi-well formats [65].
- **Arrayed Culture Libraries:** Cultures can be arrayed in 96-, 384-, or 1536-well format in multi-well plates containing liquid media or solid agar [65].
- **High-Throughput Imaging:** The Strain Library Imaging Protocol (SLIP) enables automated microscopy of large strain collections, with image acquisition completed within 4 minutes per 96-well plate [65].

Maintenance and Preservation of Pure Cultures

Optimal Growth Conditions for Marinisomatota

Based on related marine isolates and genomic predictions, Marinisomatota likely require specific conditions for long-term maintenance:

- **Salinity:** Most marine isolates require at least 1% (wt/vol) NaCl, with optimum growth at 3% (wt/vol) NaCl, similar to the novel Gaopeijiales strains which tolerated up to 7-8% NaCl [63].
- **Temperature:** Based on their distribution, most Marinisomatota likely prefer mesophilic temperatures (15-25°C), though some deep-sea lineages may require cooler temperatures (4-10°C).
- **Oxygen Conditions:** Marinisomatota likely include aerobic, microaerophilic, and anaerobic representatives. The recently isolated Gaopeijiales strains grew under aerobic and microaerophilic (8-9% O₂) conditions, but not under strict anaerobiosis [63].
- **Pressure:** For deep-sea isolates, consider using pressure reactors to maintain *in situ* pressure conditions, as this can significantly impact growth and metabolism.

Long-Term Preservation Methods

- **Cryopreservation at -80°C:**
 - Prepare cryoprotectant medium containing 10-15% glycerol or DMSO in growth medium.
 - Mix exponentially growing culture with cryoprotectant in equal volumes.
 - Use controlled-rate freezing when possible (cooling at approximately 1°C per minute to -40°C, then transfer to -80°C).
 - For long-term storage, maintain at -80°C or in liquid nitrogen vapor phase.
- **Lyophilization:**
 - Suspend cells in a lyoprotectant such as skim milk (10-20%) or sucrose (10-12%).
 - Flash-freeze in liquid nitrogen.
 - Lyophilize for 24-48 hours.
 - Seal under vacuum and store at 4°C or -20°C.

Characterization and Validation of Pure Cultures

Morphological and Phenotypic Characterization

- **Cell Imaging Techniques:**
 - **Phase Contrast Microscopy:** Useful for basic morphological assessment of live cells without staining [66].
 - **Ptychography:** A label-free, high-contrast imaging technique that provides quantitative phase information, particularly suitable for reporting cellular changes such as mitosis, apoptosis, and differentiation [66].
 - **Electron Microscopy:** For detailed ultrastructural analysis of cell morphology and surface features.
- **Phenotypic Microarrays:**
 - Test carbon source utilization profiles using BIOLOG plates or similar systems.
 - Determine temperature, pH, and salinity ranges for growth.
 - Test susceptibility to various antibiotics to establish genetic manipulation potential.

Genetic Validation and Monitoring

- **16S rRNA Gene Sequencing:** Confirm phylogenetic placement and culture purity.
- **BOX-PCR Profiling:** Generate strain-specific fingerprints to distinguish closely related isolates and monitor culture stability [63].
- **Whole Genome Sequencing:** Essential for confirming the identity of isolates and linking them to metagenomic data.

Essential Research Reagents and Equipment

Table 3: Research Reagent Solutions for Marine Microbial Cultivation


Reagent/Equipment	Function	Application Example	Specific Considerations
Anoxic Artificial Seawater Medium	Base medium for isolation	Creating sediment slurries and dilution series	Must be prepared anoxically with resazurin as redox indicator [5]

Reagent/Equipment	Function	Application Example	Specific Considerations
Polysaccharide Cocktail	Carbon source for heterotrophs	Enriching polysaccharide-degrading Marinisomatota	Include chitin, cellulose, pectin, alginate, chondroitin [5]
Vitamin and Trace Element Supplements	Essential cofactors	All cultivation attempts for oligotrophic marine bacteria	Critical for growth of fastidious marine microorganisms
96-Pin Replicators	High-throughput transfer	Inoculating large format agar plates	Enables processing of multiple samples simultaneously [65]
Large Agar Pads	Substrate for single-cell imaging	SLIP protocol for high-throughput microscopy	Same size as multi-well plates for efficient transfer [65]
TTL Signal Control System	Hardware communication	Automated microscopy acquisition	Reduces stage movement time by >80% [65]

Successfully isolating and maintaining pure cultures of Marinisomatota requires an integrated approach that combines ecologically-informed cultivation strategies with modern genomic tools. By using environmental genomic data to predict metabolic capabilities and environmental preferences, researchers can design targeted isolation strategies that specifically enrich for these elusive microorganisms. Once isolated, maintaining these cultures requires careful attention to their specific physiological requirements, particularly regarding salinity, oxygen tension, and nutrient concentrations.

The ability to cultivate Marinisomatota in pure culture will dramatically advance our understanding of their functional roles in marine ecosystems, allowing researchers to move beyond genomic predictions to physiological validation. This technical guide provides the foundational methodologies necessary to achieve this goal, contributing essential tools for a comprehensive thesis on the functional diversity of Marinisomatota in ocean ecosystems.

Experimental Workflow Diagram



Marinisomatota Isolation Workflow

Interpreting Ecological Niche Based on Genomic Potential vs. Expression Data

The ecological niche of a microorganism represents the total set of biological and environmental conditions where it can maintain stable population sizes. Traditional microbial ecology relied on taxonomic classifications from 16S rRNA gene surveys, which provided limited functional insights. The advent of omics technologies has revolutionized this paradigm by enabling researchers to move beyond mere presence/absence data to understand *functional potential* (genomic capacity encoded in DNA) and *functional expression* (active processes revealed via RNA and proteins).

This distinction is particularly crucial for understanding the ecology of ubiquitous marine bacterial phyla like **Marinisomatota** (formerly *Marinimicrobia*, *Marine Group A*, *SAR406*), which thrive across diverse oceanic zones from sunlit surface waters to the deep aphotic zone [2]. For such widely distributed taxa, genomic potential often reveals a broad metabolic capability, while expression data uncovers the specific strategies employed in response to local environmental conditions. Framing niche interpretation through this dual-lens approach provides unprecedented

resolution into the mechanisms driving microbial community assembly and ecosystem function, ultimately illuminating the functional diversity of Marinisomatota in global ocean ecosystems.

Methodological Framework: From Samples to Ecological Inference

Core Experimental Workflows

A robust analysis integrating genomic potential and expression data involves a multi-stage process, from sample collection to ecological modeling. The workflow below outlines the primary pathways for niche interpretation.




Diagram 1: Integrated workflow for niche interpretation from genomic and transcriptomic data.

The Scientist's Toolkit: Essential Research Reagents and Materials

Successful implementation of the workflow depends on specific laboratory and computational tools. The following table catalogs key reagents, software, and their functions critical for marine microbialomics studies.

Table 1: Essential Research Reagents and Computational Tools

Category/Item	Specific Examples	Function/Application
DNA/RNA Extraction Kit	DNeasy PowerSoil Kit (Qiagen) [67]	Standardized isolation of high-quality genomic DNA and total RNA from diverse marine samples (water, biofilm).
Library Prep Kits	Illumina DNA Prep, Illumina RNA Prep	Preparation of sequencing libraries for next-generation platforms like Illumina NovaSeq.
Sequencing Platform	Illumina NovaSeq [67]	High-throughput generation of metagenomic and metatranscriptomic short-read data (e.g., 150 bp paired-end).

Category/Item	Specific Examples	Function/Application
Assembly Software	MEGAHIT [67]	<i>De novo</i> assembly of quality-filtered sequencing reads into longer contiguous sequences (contigs).
Binning Tool	MaxBin2 [67]	Clustering of contigs into Metagenome-Assembled Genomes (MAGs) based on sequence composition and abundance.
Quality Assessment	CheckM [67]	Evaluation of MAG completeness and contamination using conserved single-copy marker genes.
Taxonomic Classifier	GTDB-Tk [67]	Accurate taxonomic assignment of MAGs against the Genome Taxonomy Database (GTDB).
Functional Annotator	Prokka [67]	Rapid annotation of coding sequences (CDS) within MAGs, identifying protein-coding genes, rRNAs, and tRNAs.
Abundance Profiler	CoverM [67]	Calculation of MAG abundance (read coverage) across different samples.

Data Integration and Interpretation: Contrasting Potential with Activity

Conceptual Framework for Niche Interpretation

The core challenge lies in synthesizing genomic and transcriptomic data into a coherent ecological narrative. The following conceptual diagram illustrates the relationship between a population's total genomic potential and its expressed niche in a given environment.




Diagram 2: The realized niche is formed when environmental filters shape gene expression from the total genomic potential.

Case Study: Metabolic Strategies in Widespread Marinisomatota

A 2025 study on Marinisomatota exemplifies this approach. Metagenomic analysis of 1,588 genomes revealed a broad genomic potential for diverse metabolic strategies, from photoautotrophy to heterotrophy [2]. However, metatranscriptomic data was crucial for identifying three distinct, active metabolic strategies (MS0, MS1, MS2) and understanding their distribution across different oceanic provinces. This integration showed how populations with similar genomic potential functionally diverge to occupy specific niches.

Table 2: Quantitative Comparison of Marinisomatota Metabolic Strategies [2]

Metabolic Strategy	Defining Genomic Potential	Key Expressed Functions (RNA)	Relative Abundance Range	Inferred Ecological Niche
MS0	Genes for light-dependent processes (Crassulacean acid metabolism, Moo169)	Expression of proteorhodopsin and CO2 fixation genes	0.18% to 36.21%	Mixotrophic; dominant in the translucent ocean zone
MS1	Full glycolytic pathway	High expression of glycolytic enzymes	Not Specified	Heterotrophic; specialized in carbon processing in nutrient-rich zones

Metabolic Strategy	Defining Genomic Potential	Key Expressed Functions (RNA)	Relative Abundance Range	Inferred Ecological Niche
MS2	Incomplete glycolytic pathway	Expression of alternative carbon catabolism pathways	Not Specified	Heterotrophic; adapted to nutrient-poor or aphotic zones

The data in Table 2 demonstrates a clear niche partitioning based on expressed function. While many *Marinisomatota* genomes hold the genetic blueprint for light utilization (genomic potential), only the MS0 group significantly expresses these genes in the photic zone, thereby realizing a distinct mixotrophic niche [2].

Advanced Modeling: Integrating Genomics with Environmental Gradients

Cutting-edge research further incorporates these data into models predicting responses to environmental change. The **genomic offset** metric quantifies the genetic mismatch a population may face under future climate scenarios by analyzing genotype-environment associations [68]. This can be coupled with niche suitability models based on current distribution. For example, cold-tolerant bird populations showed higher genomic offset but less niche suitability decline under future climates than warm-tolerant ones, revealing complex vulnerability patterns [68]. This genome-niche framework is directly applicable to marine microbes, allowing researchers to predict which *Marinisomatota* populations are most vulnerable to ocean warming or deoxygenation.

Detailed Experimental Protocols

Metagenomic Assembly and Binning for Genomic Potential

This protocol is adapted from methodologies used in recent studies of engineered and natural microbial systems [67].

- **Quality Control and Sequencing:** Generate 150-bp paired-end reads on an Illumina NovaSeq platform. Quality-trim raw reads using tools like Trimmomatic or Fastp to remove adapters and low-quality sequences.
- **Co-assembly:** Assemble the high-quality reads from all samples *co-assembled* or *individual-assembled* using MEGAHIT [67]. Retain contigs longer than 1000 bp for downstream analysis.
- **Bin Metagenome-Assembled Genomes (MAGs):** Bin contigs from each sample into MAGs using a tool like MaxBin2 [67]. This leverages sequence composition and sample abundance.
- **Assess MAG Quality:** Use CheckM to estimate the completeness and contamination of each MAG [67]. Retain only high-quality (completeness >90%, contamination <5%) and medium-quality (completeness >50%, contamination <5%) MAGs as per current community standards [67].
- **Dereplication:** Cluster MAGs from across all samples using dRep to obtain a non-redundant set of representative MAGs (rMAGs). A common threshold is 99% average nucleotide identity for secondary clustering [67].

Metatranscriptomic Analysis for Gene Expression

- **RNA Extraction and Library Preparation:** Extract total RNA from biomass filters. Remove ribosomal RNA (rRNA) from the total RNA sample using a depletion kit. Synthesize cDNA and prepare sequencing libraries for Illumina sequencing.
- **Read Mapping and Quantification:** Map the quality-controlled metatranscriptomic reads to the catalog of genes predicted from the MAGs (the "functional annotation" from Table 1). Use a tool like Salmon or HTSeq to quantify the number of reads mapping to each gene.
- **Normalization and Expression Profiling:** Normalize read counts to account for gene length and total library size (e.g., calculating TPM - Transcripts Per Million). This creates a gene expression profile for each sample, indicating which metabolic pathways are actively being transcribed.

Genomic Offset Calculation for Climate Vulnerability

This protocol, adapted from genomic studies on macro-organisms, provides a framework for forecasting microbial vulnerability [68].

- **Genotype-Environment Association:** Use a machine-learning method like gradientForest to identify SNPs in the rMAGs that are significantly associated with current spatial gradients in environmental variables (e.g., temperature, nutrient concentrations) [68].
- **Build a Prediction Model:** The algorithm learns a function that transforms the multidimensional environmental space into a multidimensional genetic space (allele frequencies).
- **Project Future Genomic Mismatch:** Use the trained model to predict the expected genetic composition under future climate scenarios (e.g., RCP 4.5 or 8.5 for 2050/2070). The **genomic offset** is calculated as the Euclidean distance between the current and future predicted genetic compositions for a given location, representing the amount of genetic adaptation required for population persistence [68].

Validating Functional Roles: How Marinisomatota Compare to Other Key Marine Microbes

Marine mixotrophy, the combination of autotrophic and heterotrophic metabolic strategies within a single organism, represents a critical adaptation for survival in nutrient-limited oceanic environments. This whitepaper examines the functional diversity of Marinisomatota (formerly known as Marinimicrobia, Marine Group A, and SAR406) in comparison with other mixotrophic marine bacteria. Through analysis of recent global-scale metagenomic studies and experimental data, we reveal the unique metabolic plasticity and ecological distribution of Marinisomatota, identifying three distinct metabolic modes (MSo, MS1, MS2) that demonstrate evolutionary adaptations to oceanic nutrient gradients. The implications of these findings for understanding microbial ecosystem functioning and potential biotechnological applications are discussed within the framework of ocean ecosystem research.

The functional diversity of marine microorganisms underpins global biogeochemical cycles and ecosystem stability. Within this framework, **Marinisomatota** have emerged as a ubiquitous and abundant group in global oceans, traditionally characterized as heterotrophic microorganisms but recently revealed to possess remarkable metabolic versatility [2]. These organisms inhabit a wide range of marine environments, from surface waters to the deep ocean, with relative abundances ranging from 0.18% to 36.21% across different oceanic provinces [2].

Mixotrophy represents a strategic advantage in the marine environment, allowing organisms to adapt to fluctuating nutrient conditions by combining carbon fixation through autotrophic processes with organic carbon assimilation through heterotrophic nutrition. While multiple bacterial lineages have developed mixotrophic capabilities, **Marinisomatota** exhibit particularly sophisticated adaptations that enable their persistence across diverse marine habitats, from the sunlit translucent zone to the dark aphotic layers of the ocean [2]. Understanding the contrasting lifestyles between Marinisomatota and other mixotrophic marine bacteria provides critical insights into the microbial mechanisms driving carbon and energy flow in ocean ecosystems.

Marinisomatota: Ecological Distribution and Metabolic Diversity

Taxonomic Diversity and Genomic Features

Recent large-scale metagenomic studies have dramatically expanded our understanding of Marinisomatota diversity. A 2025 analysis reconstructed **1,588 Marinisomatota genomes** from global ocean datasets, representing substantial diversity including one class, two orders, 14 families, 31 genera, and 67 species [2]. This taxonomic breadth underscores the evolutionary success of this group and highlights their significant genetic potential for diverse metabolic functions.

The genomic features of Marinisomatota reveal adaptations to their marine habitats. Among the 14 families identified, five (**S15-B10**, **TCS55**, **UBA1611**, **UBA2128**, and **UBA8226**) exhibit genetic potential for light-dependent processes associated with Crassulacean acid metabolism (MOO169) [2]. This metabolic capability represents a key differentiator from other marine mixotrophic bacteria and enables Marinisomatota to occupy unique ecological niches.

Distinct Metabolic Strategies

Research has identified three distinct metabolic strategies within Marinisomatota populations, demonstrating a spectrum of trophic adaptations:

Table 1: Metabolic Strategies in Marinisomatota

Metabolic Strategy	Trophic Mode	Key Characteristics	Ecological Distribution
MSo	Photoautotrophic potential	Capable of light-dependent carbon fixation	Primarily in translucent zone
MS1	Heterotrophic with enhanced glycolytic capacity	Pronounced glycolytic pathway; versatile energy generation	Transition zones between translucent and aphotic layers
MS2	Heterotrophic without glycolysis	Relies on alternative energy-yielding pathways	Predominantly in aphotic zones

The **MSo strategy** represents Marinisomatota with photoautotrophic potential, capable of harnessing light for carbon dioxide fixation and organic compound synthesis [2]. These organisms predominantly inhabit the translucent zone of the ocean where light penetration supports their energy needs. The **MS1 strategy** describes heterotrophic Marinisomatota with pronounced glycolytic capacity, enabling efficient processing of organic carbon sources while potentially maintaining limited autotrophic capabilities [2]. Finally, the **MS2 strategy** comprises heterotrophic organisms without glycolysis, relying instead on alternative metabolic pathways for energy generation in nutrient-limited environments [2].

The emergence of these distinct metabolic strategies likely represents an evolutionary response to nutrient limitations within oceanic environments, allowing Marinisomatota to exploit varied ecological niches across depth gradients and water masses [2].

Comparative Analysis: Marinisomatota vs. Other Mixotrophic Marine Bacteria

Metabolic Pathway Specialization

When compared to other mixotrophic marine bacteria, Marinisomatota exhibit distinct metabolic specializations that define their ecological roles:

Table 2: Metabolic Comparison of Marinisomatota with Other Mixotrophic Marine Bacteria

Characteristic	Marinisomatota	Other Mixotrophic Marine Bacteria
Primary Autotrophic Mechanism	Crassulacean acid metabolism (M00169) in specific families [2]	Diverse mechanisms including proteorhodopsin-based systems, anoxygenic photosynthesis
Carbon Fixation Pathways	Light-dependent CO2 fixation in MSo type [2]	Multiple pathways including Calvin cycle, reverse TCA cycle
Heterotrophic Capabilities	Ranges from glycolytic (MS1) to non-glycolytic (MS2) strategies [2]	Typically specialized for specific organic substrate groups
Environmental Distribution	Low-latitude marine regions, relative abundance 0.18-36.21% [2]	Varies by taxonomic group; often more restricted distribution
Genomic Features	1,588 genomes representing substantial diversity [2]	Variable genome sizes and coding density

Unlike many other mixotrophic bacteria that employ proteorhodopsin-based light energy capture or anoxygenic photosynthesis, Marinisomatota utilize **Crassulacean acid metabolism (CAM)**-like pathways in specific families, representing a unique adaptation within marine bacterial communities [2]. This metabolic distinction potentially provides competitive advantages in specific oceanic environments where traditional photosynthetic mechanisms are less efficient.

Ecological Niches and Biogeographic Patterns

Marinisomatota demonstrate distinct biogeographic patterns compared to other mixotrophic bacteria. They are **predominantly found in low-latitude marine regions** with specific adaptations to different depth zones [2]. Their distribution contrasts with other mixotrophic groups that may show more restricted geographic ranges or depth preferences.

Evidence from studies of isolated marine environments further highlights the unique position of Marinisomatota. Beneath the Ross Ice Shelf, a dark, oligotrophic environment, Marinisomatota were identified among the dominant microbial community members, alongside Proteobacteria, SAR324, Crenarchaeota, Chloroflexota, and Planctomycetota [25]. This presence in an extreme environment underscores their metabolic versatility and ability to thrive in habitats where traditional photosynthetic mixotrophs cannot persist.

Methodological Approaches: Experimental Protocols for Studying Marine Mixotrophy

Genomic Reconstruction and Metabolic Inference

The investigation of Marinisomatota metabolic strategies relies on sophisticated genomic approaches:




Figure 1: Workflow for Genomic Reconstruction of Metabolic Strategies

Sample Collection and Processing:

- Environmental samples collected across depth gradients (surface to bathypelagic) and geographic regions
- Filtration through sequential size fractions (0.2-0.8 μm for free-living; 0.8-20 μm for particle-attached) to capture different ecological niches [69]
- DNA extraction using optimized protocols for microbial community DNA preservation

Metagenomic Sequencing and Assembly:

- High-throughput sequencing using Illumina or PacBio platforms
- Assembly of sequencing reads into contigs using metaSPAdes or similar assemblers
- **Bin extraction** and refinement to obtain metagenome-assembled genomes (MAGs) [70]
- Quality assessment based on completeness (<82.33% average), contamination (<1.79% potential), and strain heterogeneity [70]

Metabolic Inference:

- Taxonomic classification against Genome Taxonomy Database (GTDB)
- Functional annotation using KEGG, Pfam, and custom databases
- Identification of key metabolic markers: Crassulacean acid metabolism genes, glycolytic pathway components, and electron transport chain genes [2]
- **Metatranscriptomic analysis** to validate gene expression and metabolic activity [2]

Computational Host Prediction for Viral Interactions

Understanding virus-host interactions provides insights into microbial ecology and metabolic constraints:




Figure 2: Host Prediction Methodology for Viral Ecology

This approach has revealed specific virus-host associations, with **39 viral sequences linked to Marinisomatota hosts** in bathypelagic ecosystems [69]. These interactions influence microbial community structure and metabolic functioning through the transfer of auxiliary metabolic genes (AMGs).

Table 3: Key Research Reagents and Computational Tools for Studying Marine Mixotrophy

Resource Category	Specific Tools/Reagents	Application/Function
Sequencing Technologies	Illumina NovaSeq, PacBio Sequel, Oxford Nanopore	High-throughput DNA sequencing for metagenomic and metatranscriptomic analysis
Bioinformatics Tools	GTDB-Tk, CheckV, VIBRANT, PHIST	Taxonomic classification, viral genome quality assessment, host prediction [69]
Reference Databases	GTDB, KEGG, Pfam, UniRef	Functional annotation and metabolic pathway inference [2]
Cultivation Media	Modified low-nutrient media, diffusion-based chambers	Isolation of previously uncultured marine bacteria [56]
Analytical Tools	VIBRANT, CheckV, PHIST	Viral identification, quality assessment, and host prediction [69]

Research Implications and Future Directions

The functional diversity of Marinisomatota has significant implications for understanding ocean ecosystem dynamics. Their metabolic versatility contributes to **carbon cycling across depth gradients**, potentially influencing carbon export efficiency and nutrient remineralization in the deep ocean [2]. The distinct metabolic strategies (MSO, MS1, MS2) represent adaptations to specific ecological niches, allowing Marinisomatota to persist across a range of marine environments.

Future research should focus on several critical areas:

- **Cultivation of representative strains** to validate metabolic predictions through physiological experiments
- **Integration of multi-omics data** (genomics, transcriptomics, proteomics, metabolomics) to build comprehensive models of metabolic networks
- **Investigation of viral influences** on Marinisomatota metabolism and evolution through analysis of auxiliary metabolic genes
- **Expansion of biogeographic sampling** to include underrepresented ocean regions and depth zones

The contrasting lifestyles of Marinisomatota compared to other mixotrophic marine bacteria highlight the complexity of microbial interactions in ocean ecosystems and underscore the need for continued research into the functional diversity of marine microorganisms.

Functional Redundancy and Uniqueness in the Marine Microbial Web

The concept of functional redundancy has long served as a foundational principle in marine microbial ecology, positing that the ocean's tremendous microbial diversity masks a fundamental metabolic simplicity wherein different taxa perform identical biochemical roles [71] [72]. This paradigm has significantly influenced how scientists model biogeochemical cycles, project ecosystem responses to environmental change, and interpret the ecological significance of microbial diversity. The "paradox of the plankton"—how an apparently limited range of resources can support an unexpectedly large number of species—has often been resolved by invoking this principle of redundancy [72]. However, emerging genomic evidence

now challenges this entrenched view, suggesting that the marine microbial web exhibits remarkable functional uniqueness, where taxonomic shifts produce consequential changes in ecosystem function [71]. This paradigm shift carries profound implications for predicting how ocean ecosystems will respond to anthropogenic pressures and climate change.

Within this reconceptualized framework, the **Marinisomatota** phylum (formerly known as Marinimicrobia, Marine Group A, and SAR406) emerges as a critical case study [2]. These ubiquitous and abundant microorganisms demonstrate how phylogenetic groups can exhibit substantial metabolic versatility, blurring the traditional boundaries between autotrophic and heterotrophic lifestyles and defying simple categorization. The functional diversity within this single phylum illustrates why a nuanced understanding of microbial ecology requires moving beyond broad assumptions of redundancy to appreciate the specialized adaptations and unique metabolic contributions of different taxa [8] [2]. This whitepaper synthesizes recent advances in our understanding of functional redundancy and uniqueness in marine microbial systems, with particular emphasis on Marinisomatota as a model for reevaluating ecosystem structure and function in ocean environments.

The Functional Redundancy Debate: From Theoretical Assumption to Empirical Testing

Defining the Spectrum of Redundancy

The debate over functional redundancy necessitates precise terminology. **Strict functional redundancy** describes the scenario where different microbial taxa share an identical set of metabolic functions and can readily replace one another without altering ecosystem processes [72]. In contrast, **partial functional redundancy** occurs when organisms share some specific functions but differ in others or in their ecological preferences (e.g., temperature optima, substrate affinities) [72]. Much of the historical assumption of redundancy in marine systems implicitly referred to strict redundancy, which provided a convenient simplification for modeling microbial contributions to biogeochemical cycles.

Evidence Challenging the Redundancy Paradigm

A landmark study directly tested the functional redundancy hypothesis by examining microbial communities across temporal and spatial scales in the northwestern Mediterranean Sea. The research revealed that shifts in taxonomic composition consistently altered the functional attributes of the communities, demonstrating that marine microbial diversity reflects a "tremendous diversity of microbial metabolism" rather than interchangeable parts [71] [72]. This finding fundamentally contradicts the notion of strict functional redundancy.

Crucially, this study employed a **read-based metagenomic approach** (using Commet software) that analyzed all sequencing reads rather than relying solely on annotated genes, thus capturing the full functional potential of the communities without bias toward known sequences [72]. When the researchers compared the functional profiles of different communities, they found strong coupling between taxonomic and functional composition, indicating that taxonomically distinct communities maintain distinct metabolic capabilities. This approach avoided the oversimplification inherent in methods that depend exclusively on reference databases, which are skewed toward cultured organisms and known functions [72].

Table 1: Key Evidence Challenging Strict Functional Redundancy in Marine Microbial Systems

Evidence Type	Findings	Implications
Metagenomic Monitoring	Strong link between community composition and functional attributes across temporal and spatial scales [71] [72]	Taxonomic shifts alter ecosystem functioning, refuting strict redundancy
Protein Diversity	Diverse microbial communities harbor high diversity of potential proteins [72]	Marine microbial diversity reflects extensive metabolic innovation
Analytical Approach	Working with all metagenomic reads (not just annotated genes) revealed similarity between taxonomic and functional composition [72]	Previous studies relying on known genes may have overestimated redundancy

Marinisomatota: A Case Study in Metabolic Versatility

Ecological Distribution and Genomic Diversity

Marinisomatota are ubiquitous and abundant in marine environments, with relative abundances ranging from 0.18% to 36.21% across low-latitude marine regions [2]. They inhabit diverse oceanic zones, from sunlit surface waters to the dark ocean interior, with certain members demonstrating capacity to harness light for carbon dioxide fixation while transitioning between photic and aphotic layers [2]. This ecological distribution across light and nutrient gradients has likely driven the evolutionary diversification observed within the phylum.

Recent genomic investigations have reconstructed **1,588 Marinisomatota genomes** from global ocean datasets, revealing extensive phylogenetic diversity encompassing one class, two orders, 14 families, 31 genera, and 67 species [2]. Among these, families such as S15-B10, TCS55, UBA1611, UBA2128, and UBA8226 exhibit genetic potential for light-dependent processes associated with Crassulacean acid metabolism (M00169), challenging the traditional characterization of Marinisomatota as exclusively heterotrophic [2].

Distinct Metabolic Strategies and Niche Specialization

Detailed analysis of Marinisomatota genomes has identified three distinct metabolic modes, demonstrating how functional diversity enables niche specialization within a single phylum [2]:

- **MS0 (Photoautotrophic Potential):** These lineages possess genetic machinery for light capture and carbon fixation, potentially enabling them to function as photoautotrophs under certain conditions.
- **MS1 (Heterotrophic with Enhanced Glycolytic Capacity):** These heterotrophic specialists exhibit enhanced glycolytic pathways for carbon processing, likely specializing in the breakdown of organic matter.
- **MS2 (Heterotrophic without Glycolysis):** These heterotrophs utilize alternative pathways for carbon metabolism, representing a distinct metabolic strategy from MS1 taxa.

The emergence of these specialized metabolic strategies likely represents an evolutionary response to nutrient limitation in oceanic ecosystems [2]. The coexistence of these functionally distinct types within the same phylum demonstrates that what might appear as redundancy at broad phylogenetic levels actually conceals significant functional specialization at finer taxonomic scales.


Table 2: Metabolic Strategies Identified in Marinisomatota [2]

Metabolic Strategy	Key Characteristics	Ecological Advantages	Relative Distribution
MS0	Photoautotrophic potential; capacity for light-dependent carbon fixation	Energy generation in light-limited environments; mixotrophic lifestyle	Predominant in regions with variable light conditions
MS1	Heterotrophic with enhanced glycolytic capacity	Efficient organic carbon processing in nutrient-rich microzones	Widespread across nutrient gradients
MS2	Heterotrophic without glycolysis; utilizes alternative pathways	Metabolic flexibility under energy limitation	Found in deeper water layers and oligotrophic regions

Methodologies for Investigating Functional Diversity

Metagenomic and Metatranscriptomic Approaches

Comprehensive investigation of Marinisomatota metabolic strategies requires integrated genomic and transcriptomic methodologies [2]. The following workflow outlines the key experimental and analytical steps:



Sample Collection and Processing: Marine samples are collected across spatial, depth, and temporal gradients to capture environmental variability. Biomass is typically collected via filtration (e.g., 0.22- μm pore-size Sterivex cartridges) and stored at -80°C until nucleic acid extraction [72].

DNA Extraction and Sequencing: Nucleic acids are extracted using commercial kits (e.g., AllPrep DNA/RNA kit, Qiagen) with modifications for marine samples, potentially including lysozyme and proteinase K treatment for enhanced cell lysis [72] [73]. For metagenomic analysis, libraries are prepared using kits such as Nextera XT DNA Sample Preparation Kit and sequenced on platforms such as Illumina HiSeq (2 $\times$ 100 bp paired-end) [72].

Bioinformatic Processing: Quality-controlled reads are assembled into contigs using tools such as IDBA-UD or MEGAHIT [74] [72]. For Marinisomatota-specific investigations, genome reconstruction from metagenomic data involves binning procedures to recover population genomes. Gene prediction is performed on contigs using tools such as MetaGeneAnnotator or Prodigal, followed by functional annotation against databases such as eggNOG, KEGG, and specialized metabolic pathway databases [74] [72].

Metabolic Classification: Metabolic strategies are inferred through the presence and completeness of key metabolic pathways, including carbon fixation genes, photosynthetic machinery, and heterotrophic carbon processing pathways [2]. Transcriptomic data can further reveal actively expressed pathways under different environmental conditions.

Large-scale investigation of marine microbial functional diversity benefits from integrated computational platforms. The **MASH-Ocean platform** (Microbiome Atlas/Sino-Hydrosphere for Ocean Ecosystem) represents one such resource, comprising 2,147 metagenomic samples with analysis modules for diversity, function, biogeography, and interaction networks [74]. This platform enables researchers to visualize global distribution patterns of specific microbial taxa and functional genes, facilitating investigation of functional diversity across oceanic gradients.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Essential Research Reagents and Materials for Marine Microbial Functional Diversity Studies

Item	Function/Application	Examples/Specifications
Sterivex Filter Cartridges	Biomass collection from seawater	0.22- μm pore-size GV Sterivex (Millipore) for microbial cell collection [72]

Item	Function/Application	Examples/Specifications
Nucleic Acid Extraction Kits	DNA/RNA co-extraction from microbial biomass	AllPrep DNA/RNA Kit (Qiagen) with lysozyme/proteinase K pretreatment [72]
Library Preparation Kits	Metagenomic library construction for sequencing	Nextera XT DNA Sample Preparation Kit (Illumina) [72]
Sequence Assembly Software	Reconstruction of contigs from sequencing reads	MEGAHIT v1.2.9 [74], IDBA-UD [72]
Gene Prediction Tools	Identification of protein-coding sequences	MetaGeneAnnotator [72], Prodigal v2.6.3 [74]
Functional Annotation Databases	Metabolic pathway assignment and classification	eggNOG [74], KEGG [74], GTDB-Tk v2.3.2 for taxonomy [74]
Metabolic Pathway Analysis	Inference of metabolic strategies from genomic data	Presence/absence of key genes (e.g., CBB cycle, glycolysis, CAs) [2]

Implications for Ocean Ecosystem Research and Biogeochemical Modeling

The recognition of limited functional redundancy in marine microbial communities carries profound implications for understanding and predicting ocean ecosystem responses to environmental change. Rather than functioning as buffered systems where microbial activities remain stable despite taxonomic shifts, marine ecosystems may be more susceptible to biodiversity loss than previously assumed [71]. The functional uniqueness of microbial taxa means that changes in community composition could directly alter biogeochemical cycling rates and pathways, with potential consequences for global climate processes [8] [71].

For researchers investigating the **functional diversity of Marinisomatota in ocean ecosystems**, these findings highlight the importance of strain-level metabolic differences and their ecological significance. The mixotrophic capabilities observed in certain Marinisomatota lineages [2] represent precisely the type of metabolic specialization that would be overlooked under a functional redundancy paradigm but which may critically influence carbon cycling in the ocean's interior. Future research should prioritize linking these metabolic capabilities to in situ activities through integrated metatranscriptomic and metaproteomic approaches, and exploring how Marinisomatota interactions with other microbial taxa shape broader ecosystem processes.

The marine microbial web represents a complex system of functionally distinctive taxa whose metabolic capabilities are tightly linked to their phylogenetic identity. The case of Marinisomatota illustrates how presumed functional redundancy often reflects insufficient resolution of metabolic diversity rather than true ecological equivalence. Moving forward, marine microbial ecology must embrace this complexity, developing models and approaches that account for functional uniqueness and its ecosystem consequences. Such refined understanding will be essential for predicting how ocean ecosystems will respond to ongoing global change and for harnessing marine microbial capabilities in biotechnological applications.

Crassulacean Acid Metabolism (CAM) represents a pivotal evolutionary innovation in plant biology, enhancing water-use efficiency and photosynthetic performance in arid environments. Recent genomic investigations have elucidated the complex genetic underpinnings of CAM, revealing a suite of core genes responsible for its characteristic diurnal acid metabolism. This whitepaper examines the functional diversity of CAM genes, with particular emphasis on their role in marine microbial systems including the *Marinisomatota* phylum. We synthesize current understanding of CAM pathway genetics, present quantitative analyses of gene expression patterns, and detail experimental methodologies for investigating CAM in non-model aquatic organisms. The findings demonstrate that CAM photosynthesis employs conserved genetic components across terrestrial and aquatic species, yet exhibits significant evolutionary plasticity in its regulatory mechanisms and enzyme recruitment strategies.

Crassulacean Acid Metabolism (CAM) is a carbon-concentrating mechanism that evolved independently in numerous plant lineages as an adaptation to environmental stressors [75]. The pathway temporally separates carbon acquisition from fixation: stomata open at night for CO2 uptake and fixation into organic acids (primarily malate), which are stored in vacuoles. During the day, stomata close to conserve water while the stored malate is decarboxylated, releasing CO2 for photosynthesis via the Calvin cycle [75]. While historically associated with terrestrial succulent plants in arid environments, CAM also exists in aquatic species where it solves problems of CO2 limitation rather than water conservation [76] [75].

In marine ecosystems, CAM-like metabolism may contribute to the functional diversity of microbial communities, including the *Marinisomatota* phylum. These bacteria inhabit environments with fluctuating nutrient availability where diurnal carbon concentration strategies could provide selective advantages. Understanding the genetic basis of CAM provides insights into metabolic innovation and the evolution of carbon concentration mechanisms across diverse biological systems.

Core CAM Genes and Their Functions

The CAM pathway relies on coordinated expression of genes encoding enzymes that facilitate nocturnal carboxylation, diurnal decarboxylation, and transporter proteins that facilitate metabolite movement across cellular compartments.

Key Enzymes in the CAM Pathway

Table 1: Core Enzymes in the CAM Photosynthetic Pathway

Enzyme	Abbreviation	Primary Function	Phase	Cellular Location
Phosphoenolpyruvate carboxylase	PEPC	Nocturnal CO ₂ fixation: carboxylates PEP to form oxaloacetate	Night	Cytosol
NAD-dependent malate dehydrogenase	NAD-MDH	Reduces oxaloacetate to malate using NADH	Night	Cytosol
Vacuolar ATPase	V-ATPase	Acidifies vacuole to facilitate malate storage	Night	Tonoplast
Vacuolar pyrophosphatase	V-PPase	Secondary proton pump for malate transport	Night	Tonoplast
NADP-dependent malic enzyme	NADP-ME	Diurnal decarboxylation of malate to pyruvate and CO ₂	Day	Chloroplast
Phosphoenolpyruvate carboxykinase	PEPCK	Alternative decarboxylation pathway enzyme	Day	Cytosol
Pyruvate phosphate dikinase	PPDK	Regenerates PEP from pyruvate	Day	Chloroplast

The primary nocturnal carboxylation is catalyzed by **phosphoenolpyruvate carboxylase (PEPC)**, which fixes HCO₃⁻ onto phosphoenolpyruvate (PEP) to form oxaloacetate [75]. This represents a critical entry point for carbon into the CAM cycle. In the aquatic lycophyte *Isoetes taiwanensis*, genomic analysis revealed recruitment of both plant-type and bacterial-type PEPCs, suggesting evolutionary flexibility in this key CAM component [76]. The resulting oxaloacetate is reduced to malate by **NAD-dependent malate dehydrogenase (NAD-MDH)**, and malate is transported into vacuoles for storage as malic acid [75].

Diurnal decarboxylation is mediated by **NADP-dependent malic enzyme (NADP-ME)** or, in some species, **phosphoenolpyruvate carboxykinase (PEPCK)**, which release CO₂ from malate for fixation by Rubisco in the Calvin cycle [75]. The resulting pyruvate is converted back to PEP via **pyruvate phosphate dikinase (PPDK)**, completing the cycle.

Transporter and Regulatory Genes

Beyond the core enzymes, CAM functionality depends on transporter genes, particularly those governing malate movement across the tonoplast. The **vacuolar ATPase (V-ATPase)** and **vacuolar pyrophosphatase (V-PPase)** establish the proton gradient that drives malate uptake into vacuoles at night and its release during the day [75].

Circadian clock genes form the regulatory backbone that synchronizes CAM gene expression with diurnal cycles. In *Isoetes*, core circadian regulators show distinct expression patterns compared to angiosperm CAM species, suggesting divergent evolutionary paths to achieving temporal separation of carboxylation and decarboxylation phases [76].

Genomic Features of CAM in Aquatic Systems

Distinctive Features of Aquatic CAM Genetics

Comparative genomics between terrestrial and aquatic CAM plants reveals both conserved elements and significant divergences in genetic architecture:

- **PEPC Gene Diversity:** Aquatic *Isoetes* species utilize a 'bacterial-type' PEPC alongside the 'plant-type' PEPC exclusively used in terrestrial CAM and C₄ plants [76]. This represents a fundamental difference in the genetic toolkit for carbon concentration.
- **Circadian Regulation Divergence:** While terrestrial CAM plants show conserved circadian control of CAM pathway genes, *Isoetes* exhibits substantially different regulatory patterns, suggesting alternative evolutionary trajectories for coordinating CAM rhythms [76].
- **Genome Organization:** The *Isoetes taiwanensis* genome shows even distribution of genes and repetitive elements, contrasting with the gene-rich regions typically found near chromosome ends in angiosperms [76]. This structural difference may influence regulation of metabolic pathways like CAM.

Evolutionary Origins and Horizontal Gene Transfer

The presence of bacterial-type PEPC in *Isoetes* raises questions about potential horizontal gene transfer events in the evolutionary history of aquatic CAM [76]. Marine ecosystems, with their high microbial density and frequent gene exchange, provide ideal conditions for such transfers. The *Marinisomatota* phylum, known for genomic plasticity and integrative elements, may possess similar acquired metabolic genes that enhance carbon concentration capabilities [77].

Table 2: Genomic Comparison of CAM Features Across Species

Genomic Feature	Terrestrial CAM Plants	Aquatic CAM Plants (Isoetes)	Marinisomatota
PEPC Type	Plant-type exclusively	Plant-type and bacterial-type	Unknown (potential bacterial-type)
Circadian Control	Highly conserved	Significantly diverged	Likely environmental sensing
Genome Organization	Gene-rich chromosomal regions	Even gene/repeat distribution	High plasticity with genomic islands
Key Decarboxylase	NADP-ME or PEPCK	NADP-ME predominant	Potential alternative mechanisms
Whole Genome Duplications	Multiple events in some lineages	Single ancient WGD (ISTEβ)	Not reported

Experimental Protocols for CAM Gene Investigation

Genomic Sequencing and Assembly

Protocol 1: High-Quality Genome Assembly for CAM Plants

- **DNA Extraction:** Use fresh leaf tissue from CAM-performing species. For aquatic plants, include root and leaf tissues to capture full genetic repertoire.
- **Multi-platform Sequencing:** Combine Illumina short-reads (for accuracy), Nanopore long-reads (for scaffolding), and Bionano optical mapping (for chromosomal structure) [76].
- **Genome Assembly:** Employ hybrid assembly approaches integrating all data types. For *Isoetes taiwanensis*, this yielded 204 scaffolds with N50 = 17.40 Mb, covering 90.13% of the predicted genome [76].
- **Gene Annotation:** Combine *ab initio* prediction, protein homology, and transcriptome evidence. Use BUSCO scores to assess completeness (target >90%) [76].

Diurnal Expression Analysis

Protocol 2: Time-Course Transcriptomics for CAM Gene Expression

- **Sample Collection:** Harvest tissue at 4-hour intervals over 24-hour cycles, with biological replicates (n≥3) for each time point [76].
- **RNA Extraction:** Use standardized methods with DNase treatment to remove genomic DNA contamination.
- **Library Preparation and Sequencing:** Prepare stranded RNA-seq libraries and sequence on Illumina platforms to minimum depth of 30 million reads per sample.
- **Bioinformatic Analysis:**
 - Map reads to reference genome using splice-aware aligners (e.g., STAR)
 - Quantify transcript abundance (e.g., with StringTie)
 - Identify differentially expressed genes across time points
 - Conduct phase analysis for circadian cycling of CAM genes

Functional Characterization of CAM Enzymes

Protocol 3: Enzyme Activity Assays for Key CAM Components

- **Protein Extraction:** Prepare crude extracts from tissue harvested at peak day/night phases.
- **PEPC Activity Assay:**
 - Reaction buffer: 50 mM HEPES-KOH (pH 8.0), 5 mM MgCl2, 5 mM NaHCO3, 0.2 mM NADH, 10 U malate dehydrogenase, 2 mM PEP
 - Monitor NADH oxidation at 340 nm for 5 minutes
 - Calculate activity as μmol NADH oxidized min⁻¹ mg⁻¹ protein
- **NADP-ME Activity Assay:**
 - Reaction buffer: 50 mM HEPES-KOH (pH 7.3), 5 mM MgCl2, 0.5 mM NADP⁺, 10 mM malate
 - Monitor NADP⁺ reduction at 340 nm for 5 minutes
- **Malate Quantification:** Use enzymatic test kits or HPLC to measure tissue malate content throughout diurnal cycle.

Research Reagent Solutions for CAM Studies

Table 3: Essential Research Reagents for CAM Investigation

Reagent/Category	Specific Examples	Research Application	Function in CAM Studies
Sequencing Kits	Illumina NovaSeq, Nanopore Ligation Sequencing	Whole genome sequencing	Genome assembly and variant discovery
RNA Library Prep	Illumina TruSeq Stranded mRNA	Transcriptome analysis	Diurnal gene expression profiling
Enzyme Assay Kits	PEPC Activity Assay Kit (Sigma MAK310), Malate Assay Kit (Sigma MAK067)	Biochemical characterization	Quantifying enzyme activities and metabolite flux
Antibodies	Anti-PEPC, Anti-NADP-ME (Agrisera)	Protein localization and quantification	Immunohistochemistry and western blotting
Metabolite Standards	Malic acid, Phosphoenolpyruvate (Sigma)	Metabolite profiling	HPLC and LC-MS quantification references
Bioinformatics Tools	BUSCO, OrthoFinder, ggplot2	Data analysis and visualization	Genome assessment, orthology identification, figure generation

Visualizing CAM Pathways and Experimental Workflows




Figure 1: CAM Biochemical Pathway Showing Nocturnal and Diurnal Phases




Figure 2: Experimental Workflow for Comprehensive CAM Gene Analysis

The genetic architecture of Crassulacean Acid Metabolism reveals both conserved core components and remarkable evolutionary flexibility across terrestrial and aquatic systems. Key CAM genes encoding PEPC, decarboxylases, and transporters show distinct evolutionary patterns in aquatic species like *Isoetes*, including recruitment of bacterial-type enzymes and divergent circadian regulation [76]. These findings highlight the potential for discovering novel carbon concentration mechanisms in marine microbial systems, including the *Marinisomatota* phylum.

Future research should prioritize functional characterization of CAM-like genes in marine bacteria, investigation of horizontal gene transfer events in carbon concentration pathways, and exploration of CAM gene regulatory networks across diverse aquatic organisms. The experimental frameworks and reagent resources outlined here provide foundational methodologies for advancing this research frontier. Understanding the genetic basis of CAM photosynthesis in marine systems not only elucidates fundamental biological mechanisms but also informs biotechnological applications in carbon capture and crop engineering for water-limited environments.

The phylum **Marinisomatota** (formerly recognized as Marinimicrobia, Marine Group A, and SAR406) represents a ubiquitous and abundant group of microorganisms in global ocean ecosystems [2]. Traditionally characterized as heterotrophic, emerging research reveals a complex functional diversity within this phylum, encompassing roles in both carbon fixation and the remineralization of organic matter [2]. Understanding the metabolic strategies that govern the trophic behaviors of *Marinisomatota* is critical for a holistic view of the ocean's biological carbon pump. This whitepaper synthesizes current research to elucidate the dual ecological impact of *Marinisomatota*, framing their activities within the broader context of functional diversity in marine microbial communities. The findings herein are based on metagenomic, metatranscriptomic, and metaproteomic analyses, providing an in-depth technical guide for researchers and scientists [2] [78].

Metabolic Strategies and Functional Diversity of *Marinisomatota*

Comprehensive analysis of 1,588 metagenomically-assembled *Marinisomatota* genomes from global open oceans has delineated a significant breadth of functional diversity within the phylum, which is classified into one class, two orders, 14 families, 31 genera, and 67 species [2]. These organisms are predominantly found in low-latitude marine regions, with relative abundances ranging from 0.18% to 36.21% [2].

Three distinct metabolic strategies (MS) have been identified, which represent adaptations to nutrient limitations in the ocean and underscore a potential for mixotrophy [2]:

- **MSo (Photoautotrophic Potential):** Characterized by the genetic capacity for light-dependent processes associated with Crassulacean acid metabolism (M00169). This strategy allows for carbon dioxide fixation and the synthesis of organic compounds, enabling *Marinisomatota* to thrive in the translucent zone and transition layers [2].
- **MS1 (Heterotrophic with Glycolysis):** A heterotrophic strategy with a pronounced glycolytic pathway for the breakdown of organic carbon [2].
- **MS2 (Heterotrophic without Glycolysis):** A heterotrophic strategy that operates without a strong glycolytic pathway, suggesting alternative mechanisms for carbon acquisition [2].

The emergence of these strategies highlights a critical interplay between life history traits and metabolic pathways in the evolution of novel nutritional groups [2].

Table 1: Metabolic Strategies and Distribution of *Marinisomatota*

Metabolic Strategy	Trophic Mode	Key Genetic or Metabolic Feature	Ecological Niche
MSo	Photoautotrophic Potential	Crassulacean acid metabolism (Moo169)	Translucent zone, Aphotic transition
MS1	Heterotrophic	Pronounced glycolytic pathway	Various marine depths
MS2	Heterotrophic	Lacks strong glycolysis	Various marine depths

Marinisomatota in the Context of Microbial Remineralization

The remineralization of particulate organic carbon (POC) is a critical process in the ocean's twilight zone (base of the euphotic zone to 1,000 m depth), determining the efficiency of carbon sequestration and global climate regulation [78]. Prokaryotes are responsible for 70-92% of POC remineralization [78]. While Marinisomatota is a key player, other bacterial groups also contribute significantly to this process, illustrating the functional diversity of microbial communities in the dark ocean.

Metaproteomic studies of POC in the Northwest Pacific Ocean have identified specific active bacterial groups and their metabolic pathways [78]. These studies reveal that particle-attached bacteria possess specialized extracellular hydrolytic enzymes that release carbon trapped in POC, and the expression levels and properties of these enzymes control the microbial breakdown rate [78].

Table 2: Key Microbial Players in POC Remineralization in the Twilight Zone


Microbial Order	Class	Key Hydrolases Produced	Primary Function in POC Degradation
Alteromonadales	Gammaproteobacteria	Proteases, Hydrolases	Degrades proteinaceous components and polysaccharides
Rhodobacterales	Alphaproteobacteria	Proteases, Hydrolases	Degrades proteinaceous components and polysaccharides
Enterobacterales	Gammaproteobacteria	Proteases, Hydrolases	Degrades proteinaceous components and polysaccharides; dominates in cold waters

The community composition and functional processes are strongly influenced by environmental factors, with temperature being a primary driver. For instance, Enterobacterales can replace Alteromonadales as the predominant remineralizing group under low-temperature conditions [78]. Furthermore, niche complementarity and species substitution among these bacterial groups ensure the efficient remineralization of POC across diverse marine environments [78].

Methodologies for Investigating Microbial Carbon Cycling

Metagenomic and Metatranscriptomic Workflow for Marinisomatota

The analysis of Marinisomatota's functional diversity relies on advanced genomic techniques. The following workflow outlines the process from sample collection to metabolic inference, which can be applied to other microbial groups as well.



Metaproteomic Analysis of POC-Associated Microbes

To directly assess the active microbial groups and their *in situ* metabolic activities in POC remineralization, a metaproteomic approach is employed. This method provides direct evidence of expressed proteins and functional pathways [78].

Detailed Experimental Protocol:

- **Large-Volume *In Situ* Sampling:** POC samples (size range: 0.7 to 200 μm) are collected from targeted depths (e.g., DCM, 100 m, 200 m, 500 m) using a large-volume *in situ* water transfer system to enrich sufficient biomass for analysis [78].
- **Protein Extraction and Digestion:** Proteins are extracted from the POC samples. A three-step search strategy against a comprehensive database like the Ocean Microbial Reference Gene Catalog (OM-RGC) is used to achieve high qualitative and quantitative protein coverage. This typically involves filter-aided sample preparation and digestion with trypsin [78].
- **Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS):** The digested peptides are separated by liquid chromatography and analyzed by tandem mass spectrometry to obtain peptide mass and fragmentation data [78].
- **Data Analysis and Protein Identification:** MS/MS spectra are searched against the protein database to identify peptides and proteins. The study cited identified an average of 7,177 proteins per sample, resulting in 24,967 nonredundant proteins. 97.28% of these were assigned a taxonomic origin [78].
- **Functional and Taxonomic Assignment:** Identified proteins are classified using databases like KEGG to reconstruct metabolic pathways and assigned to taxonomic groups to identify active key microbial players [78].

Research Reagent Solutions

The following reagents and materials are essential for executing the protocols described above.

Table 3: Essential Research Reagents and Materials

Reagent/Material	Function in Protocol
Ocean Microbial Reference Gene Catalog (OM-RGC)	A curated database used for protein and gene identification; provides a standardized reference for annotating metagenomic and metaproteomic data from marine environments [78].
GF/F Filters (0.7 µm)	Glass fiber filters used for the initial concentration and size-fractionation of particulate organic matter and microbial biomass from large volumes of seawater [78].
Trypsin	A protease enzyme used to digest extracted proteins into peptides for subsequent analysis by mass spectrometry [78].
KEGG Database	Bioinformatics database used for the functional annotation of identified genes and proteins, enabling reconstruction of metabolic pathways such as carbon fixation and glycolysis [2] [78].

The functional diversity of Marinisomatota has a profound and integrated impact on the ocean's carbon cycle. Through the metabolic strategies outlined, they contribute to both the input (via photoautotrophy) and output (via heterotrophy) of organic carbon in the water column. The mixotrophic potential of certain Marinisomatota lineages allows them to adapt to the fluctuating energy conditions of the oceanic water column, particularly in the transition zone between light and dark [2].

This functional versatility positions Marinisomatota as a key phylum within the broader context of marine microbial ecology. Their activities are complemented by other specialized heterotrophic bacteria like Alteromonadales and Rhodobacterales, which are primary agents of POC remineralization [78]. The collective metabolic output of these diverse groups, regulated by environmental drivers like temperature, ultimately determines the efficiency of the biological carbon pump. Understanding the functional diversity of Marinisomatota and related microbes is, therefore, essential for accurately modeling global carbon fluxes and predicting the ocean's response to environmental change.

The functional diversity of microbial life is a cornerstone of ocean ecosystem function. Within this complex community, the phyla **Proteobacteria** and **Bacteroidota** (formerly Bacteroidetes) have been extensively studied and are often treated as model systems for understanding microbial ecology and metabolism in marine environments [79] [80]. These model phyla provide a foundational benchmark against which the biology of less-characterized groups can be measured. This guide frames the ecological and metabolic strategies of the ubiquitous yet enigmatic phylum **Marinisomatota** (formerly Marinimicrobia) within the context of the well-established functional profiles of these model phyla [2].

Marinisomatota are abundant and globally distributed in marine environments, particularly in the deep ocean, but their metabolic capabilities and ecological roles are not as fully defined as those of Proteobacteria and Bacteroidota [2] [4]. By employing a comparative genomics approach, benchmarking against model phyla allows researchers to generate testable hypotheses about the potential functions of Marinisomatota in nutrient cycling, energy acquisition, and ecosystem productivity. This whitepaper provides a technical framework for such comparative analyses, detailing essential experimental protocols, data interpretation strategies, and reagent solutions to advance research in this field.

Ecological Dominance and Functional Profiles of Model Phyla

Proteobacteria: The Metabolic Generalists

As one of the most abundant and diverse bacterial phyla in the ocean, Proteobacteria play a wide array of roles in carbon, nitrogen, and sulfur cycling. Their ecological success is linked to tremendous metabolic versatility, allowing them to thrive across diverse marine habitats, from nutrient-rich coastal areas to oligotrophic open oceans [81].

Bacteroidota: Specialists in High-Molecular-Weight Organic Matter Degradation

Bacteroidota are critically important for the marine carbon cycle due to their specialization in degrading particulate organic matter, such as proteins and polysaccharides. Genomic analyses reveal specific adaptations for this lifestyle, including an extensive arsenal of hydrolytic enzymes [80]. **Table 1** summarizes the key genomic features that underpin their ecological strategies.

Table 1: Comparative Genomic and Ecological Features of Model Marine Phyla

Feature	Proteobacteria	Bacteroidota	Marinisomatota (Context)
Primary Ecological Role	Metabolic generalists; involved in multiple biogeochemical cycles	Specialists in degradation of high-molecular-weight organic matter (proteins, polysaccharides)	Traditionally considered heterotrophic, with recent evidence for photoautotrophic potential [2]
Characteristic Metabolic Traits	Diverse aerobic heterotrophy, chemolithotrophy, photosynthesis	High numbers of peptidases, glycoside hydrolases (GHs), glycosyl transferases	Three distinct metabolic strategies identified: photoautotrophic (MS0), heterotrophic with glycolysis (MS1), heterotrophic without glycolysis (MS2) [2]

Feature	Proteobacteria	Bacteroidota	Marinisomatota (Context)
Genomic Adaptations	Versatile core metabolism, diverse substrate utilization pathways	Genes for gliding motility, adhesion proteins, TonB-dependent transporters	Presence of genes for Crassulacean acid metabolism (CAM) in some families (e.g., S15-B10, TCS55) [2]
Habitat Preference	Free-living in water column, particle-associated	Particle-associated, especially during algal blooms	Low-latitude marine regions, relative abundance 0.18% to 36.21% [2]

Benchmarking Marinisomatota Against Model Phyla

Metabolic Strategy Classification in Marinisomatota

The functional diversity of Marinisomatota can be decoded by benchmarking their genomic signatures against the established profiles of Proteobacteria and Bacteroidota. A recent analysis of 1,588 Marinisomatota genomes revealed three distinct metabolic strategies, demonstrating a functional plasticity that overlaps with and diverges from the model phyla [2].


The **MSo** strategy shows photoautotrophic potential, a trait uncommon in the heterotrophic Bacteroidota but present in some Proteobacteria (e.g., *Rhodobacterales*). The **MS1** and **MS2** strategies are heterotrophic, akin to the primary lifestyles of both Proteobacteria and Bacteroidota, but with a key differentiation: MS1 utilizes a pronounced glycolytic pathway, while MS2 does not. This suggests niche partitioning in carbon source utilization, potentially reducing direct competition with the highly efficient heterotrophic pathways of Bacteroidota [80] [2].

Table 2: Metabolic Strategies and Functional Potential in Marinisomatota

Metabolic Strategy	Defining Characteristics	Putative Ecological Niche	Benchmarking Against Model Phyla
MSo	Photoautotrophic potential; genes for Crassulacean acid metabolism (CAM)	Inorganic carbon fixation in the translucent zone	Functional overlap with photoautotrophic Proteobacteria; distinct from primary Bacteroidota strategy
MS1	Heterotrophic with a pronounced glycolytic pathway	Utilization of labile organic carbon in nutrient-rich microzones	Parallels the generalist heterotrophy of many Proteobacteria
MS2	Heterotrophic without glycolysis	Utilization of alternative, possibly more recalcitrant, carbon sources in nutrient-poor conditions	Diverges from the classic glycolytic model, suggesting a unique niche distinct from both model phyla

A Conceptual Workflow for Functional Benchmarking

The following diagram illustrates a structured workflow for benchmarking a less-studied phylum (like Marinisomatota) against model marine phyla, integrating genomic, environmental, and experimental data.



Essential Experimental Methodologies

Genome-Resolved Metagenomics for Cataloging Diversity

Principle: This culture-independent method allows for the reconstruction of Metagenome-Assembled Genomes (MAGs) directly from environmental samples (e.g., seawater, sediment), enabling the study of uncultured microorganisms like many Marinisomatota [4].

Detailed Protocol:

- **Sample Collection and Sequencing:** Filter a large volume of seawater (e.g., 2 L) through a 0.22 μm polycarbonate membrane to capture microbial cells [81]. Flash-freeze the filters in liquid nitrogen and store at -80°C . Extract total genomic DNA using a commercial kit (e.g., PowerWater DNA Isolation Kit). Perform whole-genome shotgun sequencing on an Illumina or PacBio platform.
- **Bioinformatic Processing and MAG Reconstruction:**
 - **Quality Control:** Use tools like Trimmomatic to remove adapters and low-quality bases from raw sequencing reads [82].
 - **Assembly:** Assemble quality-filtered reads into longer contigs using a metaSPAdes or MEGAHIT.
 - **Binning:** Group contigs into putative MAGs based on sequence composition (k-mer frequency, GC content) and abundance across samples using tools like MetaBAT2 or MaxBin2.
 - **Quality Assessment:** Check MAG quality using CheckM. Aim for medium to high-quality drafts (e.g., >50% completeness, <10% contamination) [4].
 - **Taxonomic Classification:** Classify MAGs against a reference database such as the Genome Taxonomy Database (GTDB) [82] [4].

16S rRNA Gene Amplicon Sequencing for Community Profiling

Principle: Sequencing a hypervariable region of the 16S rRNA gene provides a cost-effective method to profile microbial community composition and relative abundance, though with lower taxonomic resolution than metagenomics [82].

Detailed Protocol:

- **PCR Amplification:** Amplify the target region (e.g., V3-V4 with primers 341F/806R, or the near-full-length gene with ONT27F/ONT1492R) from the extracted environmental DNA [82] [81].
- **Library Preparation and Sequencing:** This can be done via:
 - **Illumina MiSeq:** For short-read, high-accuracy sequencing of the V3-V4 region [81].
 - **Oxford Nanopore GridION:** For long-read, full-length gene sequencing, which can offer improved taxonomic resolution [82].
- **Bioinformatic Analysis:**
 - **Demultiplexing:** Assign sequences to samples based on barcodes.
 - **Denoising & ASV/OTU Picking:** Use DADA2 (for Amplicon Sequence Variants - ASVs) in QIIME2 or Mothur (for Operational Taxonomic Units - OTUs at 97% identity) to cluster sequences into taxonomic units [82] [81].
 - **Taxonomy Assignment:** Assign taxonomy to ASVs/OTUs using reference databases like SILVA, GreenGenes2, or NCBI [82].

Culture-Dependent Isolation and Characterization

Principle: Cultivation provides pure bacterial strains for detailed physiological and genetic studies, enabling the functional validation of genomic predictions and the discovery of novel species [79].

Detailed Protocol (as applied to marine turtles, adaptable for water samples):

- **Sample Homogenization:** Aseptically homogenize the sample (e.g., fecal matter, marine snow particles) in sterile seawater [79].
- **Culture Conditions:** Plate serial dilutions of the homogenate onto marine-specific agar media (e.g., 2216E medium, supplemented with sodium acetate or jellyfish extract). Incubate plates under both **aerobic** and **anaerobic** conditions at a relevant temperature (e.g., 28°C) to capture a wider diversity of organisms [79].
- **Strain Purification and Identification:** Pick morphologically distinct colonies and re-streak repeatedly on fresh media to obtain pure cultures. Extract genomic DNA from pure colonies.
- **16S rRNA Gene Sequencing:** Amplify and sequence the nearly full-length 16S rRNA gene using primers 27F and 1492R. Compare the resulting sequences against public databases (EzTaxon, NCBI BLAST) for identification and to check for novelty [79].


The Scientist's Toolkit: Key Research Reagents & Materials

Table 3: Essential Reagents and Kits for Marine Microbiome Research

Item	Specific Example	Function/Application
DNA Extraction Kit	PowerWater DNA Isolation Kit (MP BIO)	Designed to efficiently extract microbial DNA from water filters, overcoming inhibitors common in marine environments [81].
PCR Enzymes	KAPA 3G Kit (Illumina); LongAmp Taq 2X Master Mix (Nanopore)	Polymerases optimized for amplicon generation for specific sequencing platforms (short-read vs. long-read) [82].
Sequencing Primers	341F/806R (V3-V4); ONT27F/ONT1492R (full-length 16S)	Target specific regions of the 16S rRNA gene for amplification prior to sequencing [82].
Culture Media	2216E Marine Agar, supplemented with sodium acetate or jellyfish extract	A standard nutrient-rich medium for cultivating heterotrophic marine bacteria under aerobic or anaerobic conditions [79].
Bioinformatic Databases	SILVA, GTDB, NCBI RefSeq, Pfam	Used for taxonomic classification of sequences and functional annotation of genes and genomes [80] [82] [4].

Visualization of Metabolic Strategies and Niche Partitioning

The metabolic strategies of Marinisomatota can be conceptualized as adaptations to nutrient gradients in the ocean. The following diagram illustrates how the different strategies (MS0, MS1, MS2) may occupy distinct niches relative to the model phyla.



Conclusion

The functional diversity of Marinisomatota, particularly their mixotrophic capabilities and distinct metabolic strategies (MS0, MS1, MS2), underscores their critical and previously underestimated role in ocean biogeochemical cycles. Their adaptation to nutrient limitation through metabolic flexibility represents a key evolutionary innovation for survival in the oligotrophic ocean. The application of advanced genomic and cultivation techniques has been instrumental in moving this phylum from a 'microbial dark matter' candidate to a group with defined ecological functions. For biomedical and clinical research, the vast, untapped genetic repertoire of Marinisomatota, as part of the broader marine microbial diversity, presents a promising frontier for bioprospecting. Future research should focus on isolating novel strains, experimentally validating their metabolic pathways, and screening their genomes for biosynthetic gene clusters that could yield new antimicrobial peptides, enzymes, or other therapeutic compounds, thereby bridging the gap between marine microbial ecology and drug discovery pipelines.

References

1. Ecological diversity and metabolic strategies of widespread ... [https://link.springer.com/article/10.1007/s42995-025-00293-x]
2. Ecological diversity and metabolic strategies of widespread ... [https://pubmed.ncbi.nlm.nih.gov/40919477/]
3. Expanding the World of Marine Bacterial and Archaeal Clades [https://pmc.ncbi.nlm.nih.gov/articles/PMC4705458/]
4. Global marine microbial diversity and its potential in ... [https://www.nature.com/articles/s41586-024-07891-2]
5. Degradation of biological macromolecules supports ... [https://www.nature.com/articles/s41396-021-01026-5]
6. New globally distributed bacterial phyla within the FCB ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9727166/]
7. Microbial ecosystems and ecological driving forces in the ... [https://www.sciencedirect.com/science/article/pii/S009286742401479X]
8. Advancing ecological and environmental biology research [https://pmc.ncbi.nlm.nih.gov/articles/PMC12413372/]
9. Long-term patterns of an interconnected core marine microbiota [https://environmentalmicrobiome.biomedcentral.com/articles/10.1186/s40793-022-00417-1]
10. Quantitatively Partitioning Microbial Genomic Traits among ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC6714891/]
11. Depth-dependent microbial metagenomes sampled in the ... [https://www.nature.com/articles/s41597-024-02939-4]
12. Ultrahigh resolution MS1/MS2-based reconstruction of ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9667311/]
13. Emergent Relationships Between the Functional Diversity ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC11897942/]
14. Mass-spectrometry based metabolomics - PubMed Central - NIH [https://pmc.ncbi.nlm.nih.gov/articles/PMC12105183/]
15. A mass spectrum-oriented computational method for ion ... [https://www.nature.com/articles/s41467-023-37539-0]
16. Knowledge and data-driven two-layer networking for ... [https://www.nature.com/articles/s41467-025-63536-6]
17. Mixotroph ecology: More than the sum of its parts - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC6442617/]
18. Marine mixotrophy increases trophic transfer efficiency, mean ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC4801304/]
19. Marine mixotrophs exploit multiple resource pools to ... [https://www.us-ocb.org/marine-mixotrophs-exploit-multiple-resource-pools-to-balance-supply-and-demand/]
20. 揭秘海洋中广泛分布的Marinisomatota：生态多样性与代谢 ... [https://www.ebiotrade.com/newsf/2025-4/20250422045245682.htm]
21. Nutrient and carbon limitation drive broad-scale patterns of ... [https://www.us-ocb.org/nutrient-c-limitation-patterns-mixotrophy/]
22. 虫黄藻介导了珊瑚的营养可塑性 [https://www.tio.org.cn/OWUP/html_mobile/kycg/20240115/3512.html]
23. 青岛能源所揭示海洋聚球藻与异养菌群长期共存下互利共生 ... [https://qibebt.cas.cn/news/kyjz/202210/t20221001_6519773.html]
24. 科研进展 | 铁磷限制调控单细胞固氮生物对海洋酸化的响应 [https://mel.xmu.edu.cn/info/1012/33948.htm]
25. Phylogenetically and functionally diverse microorganisms ... [https://www.nature.com/articles/s41467-021-27769-5]
26. Exploring omics strategies for drug discovery from ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12394500/]
27. Metagenomics Unveils Viral Biodiversity [https://virologyresearchservices.com/2025/10/28/metagenomics-unveils-viral-biodiversity/]
28. Potential applications and future prospects of ... [https://www.sciencedirect.com/science/article/abs/pii/S0378111925005098]
29. The microbiome of the Black Sea water column analyzed by ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC8067304/]
30. Skin metatranscriptomics reveals a landscape of variation ... [https://www.nature.com/articles/s41587-025-02797-4]
31. Metatranscriptomic and functional metagenomic analysis of ... [https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2013.00340/full]
32. Metagenomic analysis reveals Northwest Pacific Ocean as ... [https://www.sciencedirect.com/science/article/abs/pii/S0269749125013119]
33. New globally distributed bacterial phyla within the FCB ... [https://www.nature.com/articles/s41467-022-34388-1]
34. Metagenomics-assembled genomes reveal microbial ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12174138/]
35. Phylogenetically and functionally diverse microorganisms ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC8748734/]
36. Emergent Relationships Between the Functional Diversity ... [https://pubmed.ncbi.nlm.nih.gov/40071437/]
37. Microbial dark matter spearheading the biogeochemical ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12343361/]
38. Ecological diversity and metabolic strategies of widespread ... [https://ui.adsabs.harvard.edu/abs/2025MLST....7..523X/abstract]
39. Seasonal variability in community structure and metabolism of ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12507015/]
40. Clear niche partitioning of nitrite-oxidizing bacteria from the ... [https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-025-02192-w]
41. Exploring marine-derived bacterial compounds targeting ... [https://www.nature.com/articles/s41598-025-96411-x]

- [42. Bioprospecting and marine 'omics': surfing the deep blue ...](https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2024.1362697/full) [https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2024.1362697/full]
- [43. Novel Bioactive Compounds From Marine Sources as a ...](https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2022.832957/full) [https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2022.832957/full]
- [44. In silico bioprospecting of receptors associated with the ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11162988/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC11162988/]
- [45. In silico bioprospecting of receptors for Oligoventin](https://www.sciencedirect.com/science/article/abs/pii/S0927776524007318) [https://www.sciencedirect.com/science/article/abs/pii/S0927776524007318]
- [46. MAGdb: a comprehensive high quality MAGs repository for ...](https://genomebiology.biomedcentral.com/articles/10.1186/s13059-025-03711-6) [https://genomebiology.biomedcentral.com/articles/10.1186/s13059-025-03711-6]
- [47. Microbial dark matter spearheading the biogeochemical ...](https://www.sciencedirect.com/science/article/pii/S266651742500094X) [https://www.sciencedirect.com/science/article/pii/S266651742500094X]
- [48. Microbial dark matter spearheading the biogeochemical ...](https://pubmed.ncbi.nlm.nih.gov/40808764/) [https://pubmed.ncbi.nlm.nih.gov/40808764/]
- [49. Mock Plant Communities and a Large Mammal Case Study ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC12414724/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC12414724/]
- [50. Assessing the efficiency of COI primers in eDNA ...](https://www.sciencedirect.com/science/article/abs/pii/S014113625005070) [https://www.sciencedirect.com/science/article/abs/pii/S014113625005070]
- [51. Multi-marker comparative analysis of 18S, ITS1, and ITS2 ...](https://www.frontiersin.org/articles/10.3389/fbinf.2025.1690766/full) [https://www.frontiersin.org/articles/10.3389/fbinf.2025.1690766/full]
- [52. METABOLIC: high-throughput profiling of microbial genomes ...](https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-021-01213-8) [https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-021-01213-8]
- [53. In-silico structural and functional annotation of hypothetical ...](https://www.sciencedirect.com/science/article/abs/pii/S0010482525012880) [https://www.sciencedirect.com/science/article/abs/pii/S0010482525012880]
- [54. Semantic design of functional de novo genes from a ...](https://www.nature.com/articles/s41586-025-09749-7) [https://www.nature.com/articles/s41586-025-09749-7]
- [55. Genomic potential for mercury biotransformation in marine ...](https://www.nature.com/articles/s41467-025-63808-1) [https://www.nature.com/articles/s41467-025-63808-1]
- [56. Advancing ecological and environmental biology research](https://link.springer.com/article/10.1007/s42995-025-00319-4) [https://link.springer.com/article/10.1007/s42995-025-00319-4]
- [57. High-Performance Genome Annotation for a Safer and ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11946116/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC11946116/]
- [58. and heterotrophic processes in Mn-rich marine sediments](https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2025.1639181/full) [https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2025.1639181/full]
- [59. Flow cytometric analysis of autotrophic picoplankton](https://www.sciencedirect.com/science/article/pii/S058095170130051X) [https://www.sciencedirect.com/science/article/pii/S058095170130051X]
- [60. Cultivating marine bacteria under laboratory conditions](https://pmc.ncbi.nlm.nih.gov/articles/PMC9428589/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC9428589/]
- [61. Inter-comparison of marine microbiome sampling protocols](https://www.nature.com/articles/s43705-023-00278-w) [https://www.nature.com/articles/s43705-023-00278-w]
- [62. Environmental preferences and interspecific interactions of ...](https://www.sciencedirect.com/science/article/pii/S0929139325006857) [https://www.sciencedirect.com/science/article/pii/S0929139325006857]
- [63. Description of the first marine-isolated member of the under ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11651109/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC11651109/]
- [64. Effects of Different Preservation Methods on the Structure ...](https://pubmed.ncbi.nlm.nih.gov/39804371/) [https://pubmed.ncbi.nlm.nih.gov/39804371/]
- [65. Strain Library Imaging Protocol: high-throughput, ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC5831406/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC5831406/]
- [66. Ptychography – a label free, high-contrast imaging ...](https://www.nature.com/articles/srep02369) [https://www.nature.com/articles/srep02369]
- [67. Niche differentiation in microbial communities with stable ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC10987969/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC10987969/]
- [68. The combination of genomic offset and niche modelling ...](https://www.nature.com/articles/s41467-022-32546-z) [https://www.nature.com/articles/s41467-022-32546-z]
- [69. Water mass age structures the auxiliary metabolic gene ...](https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-023-01547-5) [https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-023-01547-5]
- [70. Global marine microbial diversity and its potential in ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11390488/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC11390488/]
- [71. A Strong Link Between Marine Microbial Community ...](https://pubmed.ncbi.nlm.nih.gov/29925880/) [https://pubmed.ncbi.nlm.nih.gov/29925880/]
- [72. A strong link between marine microbial ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC6155072/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC6155072/]
- [73. Frontiers | Real-Time PCR Quantification and Diversity of... Analysis](https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2011.00253/full) [https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2011.00253/full]
- [74. MASH-Ocean 1.0: Interactive platform for investigating microbial ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11183159/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC11183159/]
- [75. Crassulacean acid metabolism](https://en.wikipedia.org/wiki/Crassulacean_acid_metabolism) [https://en.wikipedia.org/wiki/Crassulacean_acid_metabolism]
- [76. Underwater CAM photosynthesis elucidated by Isoetes ...](https://www.nature.com/articles/s41467-021-26644-7) [https://www.nature.com/articles/s41467-021-26644-7]
- [77. Novel integrative elements and genomic plasticity in ocean ...](https://www.sciencedirect.com/science/article/pii/S0092867422015197) [https://www.sciencedirect.com/science/article/pii/S0092867422015197]
- [78. Illuminating Key Microbial Players and Metabolic Processes ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC8478463/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC8478463/]
- [79. Analysis of the gut-specific microbiome from critically ...](https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2025.1671894/full) [https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2025.1671894/full]
- [80. Ecology of marine Bacteroidetes: a comparative genomics ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC3635232/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC3635232/]
- [81. Bacterial taxonomic and functional profiles from Bohai Sea ...](https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1139950/full) [https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1139950/full]
- [82. Benchmarking 16S rRNA Gene-Based Approaches to ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC12367389/) [https://pmc.ncbi.nlm.nih.gov/articles/PMC12367389/]