Navigating Nature's Pharmacy: Charting the Natural Product Chemical Space for Modern Drug Discovery

Generated at: 2026-04-03
Source URL: https://www.natprodchem.com/posts/navigating-natures-pharmacy-charting-the-natural-product-chemical-space-for-modern-drug-discovery

Navigating Nature's Pharmacy: Charting the Natural Product Chemical Space for Modern Drug Discovery

Abstract

This article provides a comprehensive overview for researchers and drug development professionals on the strategic exploration of natural product (NP) chemical space to uncover novel therapeutic leads. It covers the foundational concept of the biologically relevant chemical space (BioReCS), where NPs occupy unique and underexplored regions compared to synthetic libraries. The piece details cutting-edge methodological approaches, including AI-driven screening, genomics, and high-throughput assays, and addresses significant challenges such as supply, characterization, and regulatory hurdles. By validating the success of NPs through approved drugs and comparative analyses, the article underscores NPs' irreplaceable role in addressing unmet medical needs, particularly in antimicrobial and anticancer therapy, and outlines a future roadmap for integration with innovative technologies.

Defining the Biologically Relevant Chemical Space: Why Natural Products Are Unique

The concept of **chemical space (CS)**, also referred to as the "chemical universe," is a foundational concept in modern drug discovery and many other chemical disciplines [1]. While often used intuitively, chemical space is formally defined as a **multidimensional space** where molecular properties—both structural and functional—define coordinates and relationships between compounds [1]. Within this vast universe, the **Biologically Relevant Chemical Space (BioReCS)** comprises the subset of molecules with biological activity, spanning both beneficial compounds (therapeutics) and detrimental ones (toxins) [1].

The exploration of chemical space is particularly crucial for drug discovery, as the theoretical number of possible small organic molecules below 500 Da is estimated to exceed 10^60 structures [2]. This immense size makes comprehensive experimental screening impossible, necessitating intelligent navigation strategies to identify promising regions for bioactive molecule discovery, especially those inspired by or derived from **natural products** [3].

Defining the Biologically Relevant Chemical Space (BioReCS)

Conceptual Framework of BioReCS

BioReCS encompasses all molecules that interact with biological systems, creating a complex landscape of chemical subspaces (ChemSpas) distinguished by shared structural or functional features [1]. This space includes not only drug-like molecules but also **agrochemicals, flavor and odor chemicals, food components, and natural products** [1]. A critical aspect of BioReCS is that it includes compounds with both desirable and undesirable biological effects, including **promiscuous binders, poly-active molecules, and toxic compounds** [1].

Systematic study of BioReCS requires **molecular descriptors** that define the dimensionality of the space, with the choice of descriptors depending on project goals, compound classes, and dataset characteristics [1]. The rise of machine learning has further driven the development of novel molecular representations that can efficiently navigate these complex spaces [1].

Key Databases for BioReCS Exploration

Chemical compound databases serve as essential resources for exploring BioReCS. The table below summarizes major public databases covering different regions of the biologically relevant chemical space.

Table 1: Representative Public Compound Databases Covering Different Regions of BioReCS

Database Name	Primary Focus	Key Applications
ChEMBL [1]	Bioactive small molecules, primarily organic compounds	Major source for poly-active and promiscuous structures; drug discovery
PubChem [1]	Bioactive small molecules with extensive annotations	Biological activity analysis; chemical biology research

Database Name	Primary Focus	Key Applications
InertDB [1]	Curated and AI-generated inactive compounds	Defining non-biologically relevant chemical space; negative data for machine learning
Dark Chemical Matter [1]	Compounds repeatedly inactive in HTS assays	Understanding chemical features associated with lack of bioactivity

Mapped and Underexplored Regions of BioReCS

The exploration of BioReCS has been uneven, with certain regions receiving extensive attention while others remain largely uncharted:

- **Heavily Explored Regions**: The chemical space of **drug-like small organic molecules** and **natural products** has been extensively characterized [1]. Related areas such as **small peptides** and other **beyond Rule of 5 (bRo5)** entities are also reasonably well-mapped [1].
- **Underexplored Regions**: Several chemically and biologically important classes remain underrepresented, including **metal-containing molecules** (often filtered out by standard cheminformatics tools), **large natural products, macrocycles, protein-protein interaction (PPI) modulators, PROTACs, and mid-sized peptides** [1]. Many of these fall into the bRo5 category and present unique modeling challenges [1].
- **Dark Regions**: BioReCS also includes "gray-to-dark" areas containing compounds with undesirable biological effects, such as **toxic chemicals** [1]. These regions have received less attention but are vital for understanding what separates harmful from beneficial compounds.

Chemical Space Exploration Strategies for Drug Discovery

Navigating Chemical Space with Computational Tools

The vastness of chemical space necessitates sophisticated computational approaches for efficient navigation. Several algorithmic strategies have been developed to handle trillion-sized compound collections:

Table 2: Key Algorithmic Approaches for Chemical Space Exploration

Algorithm	Search Principle	Key Applications
FTrees [4]	Fuzzy pharmacophore similarity	Identifying close analogs with similar pharmacophore properties
SpaceLight [4]	Molecular fingerprint similarity (ECFP/CSFP)	High-throughput similarity screening using Tanimoto metrics
SpaceMACS [4]	Maximum common substructure (MCS)	Scaffold-based similarity searching and analysis

These algorithms enable researchers to identify close neighbors of known bioactive compounds within massive virtual chemical spaces. For example, screening FDA-approved drugs against the eXplore chemical space (containing 2.8 trillion virtual molecules) demonstrated that these methods can retrieve high-similarity analogs for a significant percentage of known drugs, providing starting points for drug optimization campaigns [4].

Natural Product-Informed Exploration

Natural products represent a privileged region of BioReCS, having evolved through biological selection processes to interact with macromolecular targets [3]. Strategies for **natural product-informed exploration** of chemical space include:

- **Structural simplification**: Creating simplified analogs of complex natural products while retaining core bioactive elements [3].
- **Biology-inspired synthesis**: Using natural product biosynthetic pathways as inspiration for synthetic library design [3].
- **Chemical space mapping**: Positioning natural products within broader chemical space to identify underrepresented structural regions [3].

These approaches have enabled the discovery of novel bioactive molecules that might not have been identified through traditional screening methods, providing access to distinctive regions of BioReCS [3].

Experimental and Computational Methodologies

Universal Descriptors for Cross-Chemical Space Analysis

The structural diversity across BioReCS presents challenges for consistent chemical space analysis using traditional descriptors optimized for specific compound classes [1]. Ongoing efforts aim to develop **universal molecular descriptors** that can accommodate diverse chemical types:

- **Molecular quantum numbers**: Provide a unified framework for molecular representation [1].
- **MAP4 fingerprint**: Designed to accommodate entities ranging from small molecules to biomolecules and metabolomic data [1].

- **Neural network embeddings**: Derived from chemical language models, these show promise in encoding chemically meaningful representations [1].

Workflow for Chemical Space Exploration in Drug Discovery

The following diagram illustrates a generalized workflow for exploring chemical space in drug discovery, particularly emphasizing natural product-inspired approaches:



Figure 1: Workflow for Natural Product-Informed Drug Discovery

Addressing pH-Dependent Chemical Space

A critical consideration in BioReCS exploration is the **pH-dependent nature** of many bioactive compounds [1]. Most chemoinformatics analyses assume neutral charge states, yet approximately 80% of contemporary drugs are ionizable under physiological conditions [1]. This ionization significantly impacts solubility, permeability, absorption, distribution, toxicity, and target binding, necessitating methods that account for charged species in chemical space analysis [1].

Visualization and Analysis of Chemical Space

Dimensionality Reduction for Chemical Space Mapping

The high-dimensional nature of chemical space requires **dimensionality reduction techniques** for visualization and interpretation [1]. Common approaches include:

- **Principal Component Analysis (PCA)**: Projects chemical space into lower dimensions based on variance maximization [2].
- **Self-Organizing Maps (SOMs)**: Neural network-based approach for producing low-dimensional representations [2].
- **t-Distributed Stochastic Neighbor Embedding (t-SNE)**: Particularly effective for visualizing high-dimensional data in two or three dimensions [5].
- **UMAP (Uniform Manifold Approximation and Projection)**: Preserves more of the global structure compared to t-SNE [5].

These visualization approaches enable researchers to identify clusters of compounds with similar properties, locate sparsely populated regions of chemical space that may represent opportunities for novel discovery, and understand the relationship between natural products and synthetic compounds [5] [2].

Table 3: Essential Research Tools for Chemical Space Exploration

Tool/Category	Specific Examples	Function in BioReCS Exploration
Chemical Databases	ChEMBL, PubChem, ZINC, GDB [1] [2]	Source of annotated chemical structures and bioactivity data

Tool/Category	Specific Examples	Function in BioReCS Exploration
Similarity Search Tools	FTrees, SpaceLight, SpaceMACS [4]	Identify analogs and nearby compounds in chemical space
Molecular Descriptors	ECFP, MAP4, Molecular Quantum Numbers [1]	Numeric representations encoding chemical structure
Visualization Platforms	Chemical cartography tools, SOM implementations [5]	2D/3D projection of high-dimensional chemical space
Virtual Screening	Docking, Pharmacophore screening [6]	Computational prioritization of compounds for testing

Applications in Natural Product-Based Drug Discovery

Integrating Multi-Omics Data with Chemical Space Analysis

Modern drug discovery increasingly leverages **network-based multi-omics integration** to understand complex biological systems and their interaction with chemical space [7]. These approaches combine various molecular data types (genomics, transcriptomics, proteomics) with biological networks (protein-protein interaction, drug-target interaction) to better predict drug responses, identify novel targets, and facilitate drug repurposing [7].

For natural product research, this means positioning natural compounds within broader biological context networks, connecting their chemical structures to target networks, metabolic pathways, and phenotypic effects [7]. Method categories include:

- **Network propagation/diffusion models**
- **Similarity-based integration approaches**
- **Graph neural networks**
- **Network inference models** [7]

Case Study: Natural Product-Inspired Discovery

Research has demonstrated that **natural product-informed exploration** of chemical space enables the discovery of distinctive and novel bioactive small molecules [3]. These approaches help focus molecular discovery on biologically relevant regions of chemical space, increasing the likelihood of identifying useful chemical probes and therapeutic candidates [3].

The relationship between natural products, chemical space exploration, and drug discovery can be visualized as follows:



Figure 2: Natural Product-Informed BioReCS Exploration

Future Directions and Challenges

The exploration of BioReCS faces several important challenges and opportunities:

- **Descriptor Development**: There remains a pressing need for systematic molecular fingerprints that can handle biomaterials, inorganic molecules, and other underexplored compound classes [1].
- **Standardization Tools**: Initiatives like the proposed **ChemSpace Tool** for non-targeted analysis aim to standardize the reporting of chemical space coverage and improve method comparability [8] [9].
- **Integration of Dark Matter**: More comprehensive inclusion of negative data (inactive compounds) and poorly characterized regions will refine the boundaries of BioReCS [1].
- **Temporal and Spatial Dynamics**: Future methods may incorporate the dynamic nature of biological systems and their interaction with chemical space [7].

As these challenges are addressed, the systematic exploration of biologically relevant chemical space, particularly regions inspired by natural products, will continue to drive innovation in drug discovery and chemical biology.

Natural products (NPs) from plants, animals, and microorganisms have served as a cornerstone of pharmacotherapy throughout human history, providing a rich source of structurally diverse and biologically active compounds for treating human diseases [10] [11]. These secondary metabolites represent an invaluable chemical resource, with over half of approved small-molecule drugs originating directly or indirectly from natural product scaffolds [12] [10]. The structural complexity and evolutionary optimization of natural products for biological interaction make them exceptionally suited for drug discovery, particularly for challenging targets such as protein-protein interactions [1] [12].

Within the framework of exploring natural product chemical space for drug discovery research, this review examines the biologically relevant chemical space (BioReCS) of natural products, which encompasses molecules with both beneficial and detrimental biological activities [1]. Current databases document over 1.1 million natural products that display high structural diversity and complexity, frequently featuring glycosylation and halogenation patterns that distinguish them from synthetic compounds [12]. Despite a declining discovery rate of novel structures, natural products continue to offer unique scaffolds that occupy broader chemical spaces than synthetic compounds, positioning them as an indispensable resource for addressing current therapeutic challenges [12] [10].

Historical Foundations and Contemporary Significance

Historical Context and Industrial Perspectives

The relationship between natural products and human medicine dates back to ancient healing traditions, with well-documented use in Ayurvedic medicine, Traditional Chinese Medicine (TCM), Japanese Kampo, and European phytotherapy [10]. These traditional systems provided the initial framework for exploring nature's pharmacopeia, with many modern drugs tracing their origins to ethnobotanical and ethnopharmacological knowledge [11].

The pharmaceutical industry's engagement with natural products has experienced significant fluctuations over recent decades. The 1990s witnessed a "Green Rush" in natural product research, driven by advancements in high-throughput screening (HTS) and isolation technologies that enabled systematic exploration of biodiversity [11]. This period saw substantial investment in bioprospecting initiatives targeting terrestrial and marine organisms for novel drug leads. However, in the early 2000s, most major pharmaceutical companies terminated or significantly reduced their HTS and natural product discovery programs in favor of combinatorial chemistry and rational drug design approaches [11].

Contemporary analysis reveals that the relatively low productivity of purely synthetic approaches has quietly repositioned pharmacognosy back into the drug discovery mainstream [11]. Current estimates indicate that approximately 50% of FDA-approved medications between 1981–2006 were natural products or synthetic derivatives inspired by natural products, highlighting their enduring impact despite fluctuating industrial interest [13]. This reemergence recognizes that natural products offer structural complexity and biological relevance that remains challenging to replicate through purely synthetic approaches [12] [11].

Quantitative Impact of Natural Products in Modern Therapeutics

Table 1: Therapeutic Areas Significantly Influenced by Natural Product-Derived Drugs

Therapeutic Area	Representative Drugs	Natural Source	Clinical Significance
Oncology	Paclitaxel, Docetaxel, Trabectedin	Pacific Yew Tree, European Yew, Marine Tunicate	Taxanes represent cornerstone therapies for various cancers; marine-derived agents offer novel mechanisms
Infectious Diseases	Penicillins, Tetracyclines, Erythromycin	Fungi, Soil Bacteria	Foundation of anti-infective therapies with diverse mechanisms against pathogens
Immunosuppression	Cyclosporine, Fingolimod	Soil Fungus, Fungus *Isaria sinclairii*	Revolutionized organ transplantation; advanced multiple sclerosis treatment
Neurological Disorders	Galantamine, Huperzine A	Daffodil bulbs, Chinese Herb *Huperzia serrata*	Acetylcholinesterase inhibition for Alzheimer's management

Table 2: Structural and Property Comparisons Between Natural Products and Synthetic Compounds

Property	Natural Products	Synthetic Compounds	Biological Implications
Structural Complexity	High (multiple chiral centers, intricate ring systems)	Moderate to Low	Enhanced target selectivity and novel binding modes
Molecular Weight	Broader distribution, including bRo5 space	Typically focused on lower MW	Access to challenging target classes like PPIs
Oxygen Atoms	Higher count	Lower count	Improved hydrogen bonding capacity

Property	Natural Products	Synthetic Compounds	Biological Implications
Stereochemical Complexity	High	Variable to Low	Biological specificity and metabolic stability
Chemical Space Coverage	Broader, underexplored regions	Narrower, focused on drug-like space	Access to novel bioactive scaffolds

Analysis of natural product chemical space reveals distinct structural characteristics that contribute to their biological success. Natural products frequently exhibit higher stereochemical complexity, greater abundance of oxygen atoms, and more varied ring systems compared to synthetic compounds [12]. These properties enable natural products to interact with complex biological targets through unique binding modes often inaccessible to synthetic libraries [1] [12]. Marine natural products, for instance, demonstrate particularly novel scaffolds with potent bioactivities, exemplified by the development of trabectedin from a marine tunicate [11].

Exploring Natural Product Chemical Space

Chemoinformatic Characterization of Natural Product Diversity

Systematic exploration of natural product chemical space requires robust chemoinformatic approaches to characterize structural diversity, bioactivity patterns, and source-related characteristics. Natural products exhibit distinct chemical features based on their biological origins, with marine-derived compounds generally displaying higher molecular weight and hydrophobicity compared to terrestrial counterparts [12]. NPs from extreme environments such as deep-sea ecosystems and extremophiles frequently reveal novel scaffolds with unique bioactivities, highlighting the value of biodiversity exploration in drug discovery [12].

The concept of the biologically relevant chemical space (BioReCS) provides a framework for understanding natural products' privileged status in therapeutic development. BioReCS encompasses all molecules with biological activity—both beneficial and detrimental—spanning drug discovery, agrochemistry, sensory chemistry, and toxicological domains [1]. Within this framework, natural products occupy regions characterized by high structural diversity and complexity, often distinct from synthetic compounds [1] [12].

Key chemoinformatic analyses have revealed that natural products contain a higher prevalence of unique ring systems with different atom compositions and connectivity compared to synthetic molecules [12]. This structural novelty translates to diverse biological interactions and mechanisms of action. Furthermore, natural products frequently undergo specific biochemical modifications such as glycosylation and halogenation that enhance their biological activities and target affinity [12].

Underexplored Regions of Natural Product Chemical Space

Despite extensive research, significant regions of natural product chemical space remain underexplored, presenting opportunities for future discovery. Several compound classes are notably underrepresented in current databases and drug discovery efforts:

- **Metal-containing molecules**: Often excluded during standard data curation due to optimization of chemoinformatics tools for small organic compounds [1]
- **Macrocycles** (compounds containing rings of ≥12 atoms): Complex structures with potential for modulating challenging target classes [1]
- **Protein-protein interaction (PPI) modulators**: Larger molecular frameworks capable of disrupting complex protein interfaces [1]
- **Beyond Rule of 5 (bRo5) compounds**: Natural products frequently violate traditional drug-like filters while maintaining bioavailability [1] [12]

These structurally complex natural products often fall into the beyond Rule of 5 (bRo5) category, presenting challenges for synthesis and optimization but offering unique opportunities for addressing difficult therapeutic targets [1]. Recent studies have begun systematically characterizing these underrepresented regions, including peptides, agrochemicals, metallodrugs, macrocycles, and PPI modulators [1].

Methodological Framework for Natural Product-Based Drug Discovery

Experimental Workflows and Isolation Strategies

The systematic investigation of natural products for drug discovery follows established experimental workflows that integrate traditional knowledge with modern analytical techniques. The process typically begins with source selection guided by ethnobotanical knowledge, ecological considerations, or biodiversity surveys, followed by careful specimen collection and authentication [10] [11].

Table 3: Key Methodologies in Natural Product Isolation and Characterization

Method Category	Specific Techniques	Applications in NP Drug Discovery
Extraction & Fractionation	Bioassay-guided fractionation, Solvent-solvent partitioning, Liquid-liquid chromatography	Selective enrichment of bioactive compounds from complex mixtures
Compound Isolation	High-performance liquid chromatography (HPLC), Countercurrent chromatography, Flash chromatography	Purification of individual natural products from crude extracts
Structure Elucidation	NMR spectroscopy (1D/2D), Mass spectrometry (MS), X-ray crystallography	Determination of molecular structure and stereochemistry
Bioactivity Screening	High-throughput screening (HTS), Phenotypic assays, Target-based assays	Identification of biologically active natural products

Bioassay-guided fractionation represents a cornerstone approach, wherein biological activity tracking directs the isolation of active constituents from complex natural extracts [13]. This method ensures that purification efforts focus on compounds with relevant biological effects, increasing the efficiency of lead identification. Advances in analytical technologies, particularly NMR and mass spectrometry, have dramatically accelerated the structure elucidation process, enabling determination of complex structures with minimal material [13] [11].

The following workflow diagram illustrates the integrated experimental and computational approach for natural product-based drug discovery:



Computational and Artificial Intelligence Approaches

The integration of computational methods has transformed natural product research, enabling more efficient exploration of chemical space and prediction of bioactivity. Computer-aided drug design (CADD) approaches, particularly artificial intelligence (AI) and machine learning (ML), have demonstrated significant utility in navigating the complex chemical space of natural products [13].

AI-driven approaches include:

- **Virtual screening** of natural product libraries against protein targets
- **De novo design** of natural product-inspired compounds
- **ADMET prediction** for early assessment of drug-like properties

- **Structural classification** and **dereplication** to identify novel scaffolds

Machine learning algorithms, including support vector machines (SVMs), neural networks, and decision trees, enable pattern recognition in complex structure-activity relationship data [13]. Deep learning approaches, particularly convolutional neural networks (CNNs) and recurrent neural networks (RNNs), facilitate analysis of molecular structures and prediction of bioactive conformations [13]. Natural language processing (NLP) techniques further enhance these approaches by extracting relevant information from scientific literature, patents, and natural product databases [13].

The following diagram illustrates the integration of AI technologies in natural product drug discovery:



Lead Identification and Optimization Strategies

From Hit to Lead: Experimental Protocols

The transition from initial bioactive natural product hits to viable lead compounds requires systematic approaches to evaluate and optimize chemical structures. Lead identification begins with validating biological activity through dose-response experiments and specificity assessments, followed by comprehensive characterization of absorption, distribution, metabolism, excretion, and toxicity (ADMET) properties [14].

High-throughput screening (HTS) and ultra-high-throughput screening (UHTS) methodologies enable efficient evaluation of extensive natural product libraries, with capacity reaching up to 100,000 assays per day using automated robotic systems [14]. These approaches offer significant advantages over traditional screening methods, including enhanced automation, reduced sample volumes, improved sensitivity, and cost savings in reagents and culture media [14].

Hit validation involves rigorous assessment of:

- **Potency** (IC_{50}, EC_{50} values)
- **Selectivity** against related targets
- **Cytotoxicity** and general cellular toxicity
- **Chemical stability** under assay conditions
- **Solubility** and **aggregation potential**

Confirmed hits progress to lead optimization, where medicinal chemistry strategies enhance desirable properties while mitigating limitations. The lead optimization phase involves synthesis and characterization of analog structures, evaluation using biochemical assays (e.g., Irwin's test for neurobehavioral assessment, Ames test for genotoxicity), and detailed analysis of drug-induced metabolism through metabolic profiling [14].

Structure-Activity Relationship (SAR) and Analog Design

Structure-activity relationship (SAR) studies form the foundation of natural product optimization, systematically exploring how structural modifications influence biological activity and drug-like properties. SAR analysis identifies critical pharmacophoric elements—the specific molecular features essential for biological activity—and guides strategic modifications to enhance efficacy and reduce toxicity [15].

Key strategies in natural product analog design include:

- **Direct chemical manipulation**: Adding, removing, or swapping functional groups; making isosteric replacements; adjusting ring systems [14]
- **Scaffold simplification**: Reducing structural complexity while retaining essential pharmacophoric elements [15]
- **Bioisosteric replacement**: Substituting functional groups with alternatives that maintain biological properties but improve pharmacokinetics [15]
- **Fragment-based design**: Deconstructing complex natural products into simpler fragments that retain key features [15]

The iterative process of analog design and optimization follows a cyclical approach of design, synthesis, testing, and refinement. This process continues until compounds achieve the optimal balance of potency, selectivity, and drug-like properties required for preclinical development [15].

Table 4: Key Research Reagent Solutions for Natural Product Drug Discovery

Reagent/Category	Specific Examples	Function in NP Research
Analytical Standards	Certified reference materials, Deuterated solvents, Quantitative NMR standards	Compound identification, quantification, method validation
Bioassay Kits	Enzyme inhibition assays, Cell viability assays, Receptor binding assays	Biological activity assessment, mechanism elucidation
Chromatography Materials	HPLC columns, Solid-phase extraction cartridges, Countercurrent chromatography solvents	Compound separation, purification, and enrichment
Molecular Biology Reagents	Protein expression systems, Enzyme substrates, Reporter gene assays	Target identification and validation, mechanism studies
Computational Tools	Molecular docking software, QSAR programs, Cheminformatics platforms	Virtual screening, property prediction, SAR analysis

Emerging Trends and Future Perspectives

Technological Innovations and Paradigm Shifts

The field of natural product drug discovery is experiencing significant transformation through the integration of emerging technologies and interdisciplinary approaches. Several key trends are shaping the future of this field:

Artificial Intelligence and Cheminformatics: AI-driven approaches are revolutionizing natural product research through enhanced pattern recognition in complex chemical and biological data [13]. Chemical language models and neural network embeddings generate chemically meaningful representations that can reconstruct molecular structures or predict properties, accelerating the identification of promising bioactive molecules [1]. The development of universal molecular descriptors, such as molecular quantum numbers and the MAP4 fingerprint, enables more consistent analysis of natural product chemical space across diverse compound classes [1].

Integration of Multi-Omics Technologies: Genomic, transcriptomic, and metabolomic approaches provide unprecedented insights into biosynthetic pathways and ecological functions of natural products [10]. These technologies facilitate the identification of gene clusters responsible for natural product biosynthesis, enabling heterologous expression and engineering of novel analogs [12].

Exploration of Underexplored Biodiversity: Research continues to focus on extreme environments (deep-sea, deserts, polar regions) and symbiotic relationships (endophytic fungi, microbial symbionts) as sources of novel natural products with unique scaffolds and bioactivities [12]. These ecosystems offer chemical diversity distinct from traditional sources, with marine natural products particularly promising for anticancer and antiviral applications [13].

Addressing Current Challenges and Limitations

Despite promising advances, natural product drug discovery faces several persistent challenges that require innovative solutions:

Supply and Sustainability Issues: Many natural products occur in minute quantities in their source organisms, creating supply challenges for development and large-scale production [13]. Sustainable sourcing strategies, including cultivation, partial synthesis, and biotechnology approaches, are essential for addressing ecological concerns and ensuring consistent supply [11].

Technical Complexities in Characterization: The structural complexity of natural products presents challenges for synthesis, structural elucidation, and optimization [13]. Advances in synthetic methodologies, analytical technologies, and computational prediction are gradually overcoming these barriers, making complex natural products more accessible for drug discovery [15].

Data Integration and Quality: The lack of standardized data quality and reporting in natural product research hampers data mining and reproducibility [10]. Initiatives to improve data curation, implement standardized protocols, and develop integrated databases are critical for advancing the field [1] [12].

The historical legacy of natural products as a pillar of pharmacotherapy continues to evolve through the integration of traditional knowledge with contemporary scientific approaches. As technological innovations provide new tools for exploring natural product chemical space, the unique structural features and biological relevance of natural products ensure their continued importance in addressing current and future therapeutic challenges. By leveraging advances in AI, omics technologies, and synthetic biology, researchers can unlock the full potential of nature's chemical diversity for the development of next-generation therapeutics.

Natural products (NPs) and their derivatives have historically been a cornerstone of pharmacotherapy, accounting for over 60% of all small-molecule drugs approved between 1981 and 2014 [16] [17]. Despite this proven utility, synthetic compounds (SCs) dominate most commercial screening libraries, constrained by decade-old conventions like Lipinski's Rule of Five and synthetic accessibility [16]. This preference persists even as challenging biological targets—such as protein-protein interactions, nucleic acid complexes, and antibacterial modalities—often remain recalcitrant to libraries of drug-like molecules [18].

The fundamental advantage of NPs lies in their evolutionary origin. As products of natural selection, they have co-evolved to interact with biological macromolecules, encoding inherent biological relevance and an ability to explore a broader swath of biologically relevant chemical space [19] [20]. Consequently, NPs exhibit structural features—such as increased molecular complexity, higher fractions of sp^3-hybridized carbons, and greater stereochemical density—that are often underrepresented in synthetic libraries [18] [16]. This manuscript demonstrates how Principal Component Analysis (PCA) serves as a powerful computational tool to visualize and quantify this superior diversity, providing a compelling rationale for reintegrating NPs into modern drug discovery pipelines.

Theoretical Foundations of Chemical Space Visualization

Defining Chemical Space and the Chemical Multiverse

In chemoinformatics, **chemical space** is defined as a multi-dimensional descriptor space where each molecule is represented by a numerical vector encoding aspects of its structure or physicochemical properties [21]. The concept of a **chemical multiverse** acknowledges that the chemical space of a single dataset is not unique; it is a "group of multiple chemical spaces, each defined by a given set of descriptors" [21]. The visual representation of this space, therefore, depends critically on the chosen descriptors and dimensionality reduction techniques.

Principal Component Analysis as a Dimensionality Reduction Tool

PCA is a mathematical method for dimensionality reduction that transforms a multidimensional dataset into a new set of orthogonal axes called principal components (PCs) [22]. These components are linear combinations of the original descriptors, with the first PC (PC1) capturing the maximum variance in the data, the second PC (PC2) capturing the next highest variance, and so on [22]. By projecting high-dimensional data onto a two- or three-dimensional plot, PCA allows for intuitive visualization of similarities, differences, and patterns within compound collections with minimal loss of information [22]. When applied to collections of NPs and SCs, PCA vividly reveals the distinct regions these classes occupy and their relative diversity.

Comparative Analysis of Natural Products and Synthetic Compounds

Time-Dependent Evolution of Structural Properties

A comprehensive, time-dependent chemoinformatic analysis comparing NPs from the Dictionary of Natural Products with SCs from 12 databases reveals distinct evolutionary trajectories. NPs discovered over time have become **larger, more complex, and more hydrophobic** [19]. Specifically, descriptors of molecular size—including molecular weight, molecular volume, and the number of heavy atoms—show a consistent upward trend in NPs, a phenomenon attributed to advances in separation and purification technologies [19].

Conversely, the physicochemical properties of SCs have been constrained within a narrower range, largely governed by drug-like rules and synthetic accessibility [19]. **Table 1** summarizes key differentiating properties based on analyses of hundreds of thousands of compounds [19] [16].

Table 1: Key Physicochemical and Structural Differences Between Natural Products and Synthetic Compounds

Property	Natural Products (NPs)	Synthetic Compounds (SCs)
Molecular Size	Generally larger; size increasing over time [19]	Smaller; constrained by drug-like rules [19]
Fraction of sp^3 Carbons (Fsp3)	Higher, indicating more 3D character [16]	Lower, indicating more flat, aromatic structures [18]
Stereochemical Complexity	Higher number of stereocenters [19] [16]	Fewer stereocenters [18]
Ring Systems	More rings, larger fused rings, more non-aromatic rings [19]	More aromatic rings (e.g., benzene derivatives) [19]
Oxygen & Nitrogen Content	More oxygen atoms [19]	More nitrogen atoms [19]
Biological Relevance	High, due to evolutionary selection [19] [20]	Broader synthetic pathways but declining relevance [19]

Visualizing Diversity with PCA and TMAP

A PCA analysis utilizing 16 two-dimensional structural descriptors on a combined ~390,000 NPs and SCs clearly demonstrates the greater structural variability of NPs [16]. The NPs occupy a broader, more dispersed region in the PCA plot, particularly evident in properties like the fraction of sp^3 carbon atoms (Fsp3), a key metric of molecular complexity [16].

Another powerful visualization tool is the Tree MAP (TMAP), a two-dimensional tree-based clustering algorithm built for large-scale data. When clusters are generated using molecular fingerprints (MHFP), NPs occupy vast structural areas that are largely unexplored by synthetic molecules [16]. The TMAP visualization further corroborates that NPs are structurally more complex, not only in Fsp3 but also in features like the number of spiroatoms [16].

Experimental Protocol for Chemical Space Analysis

This section provides a detailed methodology for reproducing the chemical space comparisons described in this review.

Data Collection and Curation

1. Source Natural Product Databases:

- **UNPD (Universal Natural Products Database):** Available as CSV format [16].
- **TCM Database@Taiwan:** Available as MOL2 file [16].
- **NP Atlas:** Available as CSV for download [16].
- **FooDB:** A public database containing food chemicals and their flavors [21].

2. Source Synthetic Compound Database:

- **ZINC Database:** A popular resource for commercially available compounds. Use the "ZINC in-stock" subset for readily available synthetic molecules [16].

3. Data Cleaning and Standardization:

- **Remove undesired molecules:** Filter out compounds with a molecular weight (MW) < 150 Da or > 1000 Da to focus on a drug-like range that considers cell permeability [16].
- **Standardize structures:** Use cheminformatics toolkits like RDKit or the MolVS library to canonicalize SMILES strings, neutralize charges, and generate canonical tautomers [21].
- **Handle multi-component molecules:** Split salts and other multi-component structures, retaining only the largest fragment [16] [21].
- **Remove terminal sugars:** For a more accurate assessment of the bioactive aglycon, utilize a deglycosylation tool to remove terminal sugar moieties from NPs [16].
- **Filter elements:** Remove molecules containing elements outside a defined set (e.g., H, B, C, N, O, F, Si, P, S, Cl, Se, Br, I) [16] [21].

Descriptor Calculation

Calculate the following 16 two-dimensional molecular descriptors for each standardized compound. This can be accomplished using software such as ChemAxon's Instant JChem or the RDKit library in Python [22] [16] [21].

Table 2: Essential Molecular Descriptors for PCA of Chemical Space

Descriptor	Description	Interpretation in NP/SC Context
MW	Molecular Weight	NPs are generally larger [19].
LogP	Partition coefficient (octanol/water)	Measures lipophilicity [22].
TPSA	Topological Polar Surface Area	Related to polarity and hydrogen bonding [22].
a_acc	Number of hydrogen bond acceptors	NPs often have more oxygen atoms [19].
a_don	Number of hydrogen bond donors	NPs often have more donors [22].
a_heavy	Number of heavy atoms	Indicator of molecular size [19].
b_rotR	Fraction of rotatable bonds	Related to molecular flexibility [22].
a_nN	Number of nitrogen atoms	SCs are often richer in nitrogen [19].
a_nO	Number of oxygen atoms	NPs are often richer in oxygen [19].
FCharge	Sum of formal charges	Influences solubility and interactions.
a_aro	Number of aromatic atoms	SCs typically have more aromatic character [19].
chiral	Number of chiral centers	NPs have higher stereochemical complexity [16].
rings	Number of rings	NPs tend to have more ring systems [19].
stereo	Number of stereocenters	Key indicator of NP complexity [22] [16].
fsp3	Fraction of sp^3 hybridized carbons	Critical measure of 3D complexity; higher in NPs [18] [16].
a_spiro	Number of spiro atoms	Indicator of complex ring fusions; higher in NPs [16].

Performing Principal Component Analysis

- **Data Matrix Preparation:** Compile all calculated descriptors into a matrix where rows represent compounds and columns represent the 16 descriptors. Standardize the data (e.g., z-score normalization) so that each descriptor has a mean of 0 and a standard deviation of 1 to prevent variables with larger scales from dominating the analysis.
- **PCA Execution:** Perform PCA on the standardized matrix using statistical software or programming environments like R or Python (with libraries such as scikit-learn). The analysis will output the principal components and the proportion of variance explained by each.
- **Visualization:** Generate 2D or 3D scatter plots using the first two or three principal components. Color-code the data points by origin (NP vs. SC) and by key properties like Fsp3 to visually decode the structural patterns.

The following workflow diagram summarizes the experimental protocol for chemical space analysis:



The Scientist's Toolkit: Essential Research Reagents and Software

Table 3: Key Software and Resources for Chemical Space Analysis

Tool/Resource	Type	Function in Analysis
RDKit	Open-source Cheminformatics Library	Data standardization, descriptor calculation, and fingerprint generation [16] [21].
Instant JChem	Commercial Cheminformatics Platform	Management of chemical data, batch calculation of physicochemical parameters [22].
R / Python (scikit-learn)	Programming Environments	Performing Principal Component Analysis and statistical computations [22].
VCC Lab ALOGPS	Web Service	Calculating additional properties like logP and aqueous solubility [22].
MolVS	Open-source Library	Standardizing molecular structures (tautomers, charges, fragments) [21].
FooDB	Public Database	Source of natural product structures, particularly food-related chemicals [21].
ZINC Database	Public Database	Source of commercially available synthetic compound structures [16].

Implications for Drug Discovery and Library Design

The clear visualization of NP diversity has direct, practical implications for drug discovery. The finding that NPs explore vast, biologically relevant regions of chemical space that SCs do not reach provides a strong rationale for designing new libraries that capture these underrepresented features [18] [20]. Several strategies have emerged to bridge this gap:

- **Biology-Oriented Synthesis (BIOS):** Uses NP scaffolds as starting points for library design, keeping compounds close to validated chemical space [20].
- **Pseudo-Natural Products (PNPs):** Combines NP fragments in novel arrangements not found in nature, creating new scaffolds that retain biological relevance while exploring uncharted territory [19] [20].

- **Diversity-Oriented Synthesis (DOS):** Aims to generate high structural diversity, often incorporating NP-like features such as high Fsp3 and stereochemical complexity [20].

PCA can guide these efforts by quantifying how well a new library penetrates NP-like regions of chemical space. As demonstrated in one study, analyzing the component loadings can identify which structural parameters (e.g., number of oxygen atoms, stereochemical density, Fsp3) most influence the separation between NPs and SCs. Chemists can then target these specific parameters through synthetic modification to "shift" their compounds towards the NP region of the PCA plot [22]. This data-driven approach enables a more rational and effective exploration of nature's vast chemical repertoire for drug discovery.

Principal Component Analysis provides an unambiguous visual and quantitative demonstration of the superior structural diversity inherent in natural products compared to synthetic chemical libraries. The broader distribution of NPs in chemical space, characterized by greater molecular complexity, stereochemical richness, and distinct physicochemical properties, underscores their immense and irreplaceable value for drug discovery. By leveraging PCA as a guide for library design and analysis, researchers can move beyond the constraints of traditional drug-like chemical space, harnessing the evolutionary wisdom encoded in natural products to develop novel therapeutics for the most challenging biological targets.

Heavily Explored vs. Underexplored Regions in the NP Chemical Universe

Natural products (NPs) represent a vast and structurally diverse resource for drug discovery, comprising over 173,000 known structures that have evolved to interact with biological systems [23]. The concept of "chemical space" refers to a multidimensional universe where molecular properties define coordinates and relationships between compounds, with the biologically relevant chemical space (BioReCS) encompassing molecules with demonstrated biological activity [1]. Within this framework, natural products occupy a strategic position, as they largely adhere to the Rule of Five while simultaneously exploring regions of chemical space not covered by synthetic compounds and available screening collections [24]. This renders them a valuable, unique, and necessary component of screening libraries used in drug discovery. Analyses of 10,495 natural products and 5,757 trade drugs reveal that natural products possess 1,748 different ring systems compared to 807 different ring systems found in trade drugs, demonstrating their superior structural diversity [23]. Despite this proven potential, significant portions of the natural product chemical universe remain underexplored, creating opportunities for discovering novel bioactive compounds with resistance-breaking properties and new mechanisms of action, particularly in challenging therapeutic areas like antimicrobial resistance [25].

Charting the NP Chemical Universe: Classification Approaches

Structural and Biosynthetic Classification Frameworks

The systematic organization of natural products enables effective navigation of their chemical space. Several classification approaches have been developed, with **structural classification of natural products (SCONP)** emerging as a powerful organizing principle [26]. SCONP arranges the scaffolds of natural products in a tree-like fashion, providing both an analysis- and hypothesis-generating tool for the design of natural product-derived compound collections [26]. This approach facilitates the identification of biologically relevant subfractions of chemical space and has been successfully applied in the development of novel inhibitor classes, such as selective and potent inhibitors of 11β-hydroxysteroid dehydrogenase type 1 with cellular activity [26].

Alternative classification systems group natural products according to recurring structural features. For instance, **flavonoid compounds** are oxygenated derivatives of a specific aromatic ring structure, while alkaloids containing an indole ring are classified as **indole alkaloids** [27]. These structural classifications complement biosynthetic organization systems, which categorize compounds based on their metabolic pathways of origin within producing organisms [27]. Each classification approach offers distinct advantages for drug discovery, with structural systems enabling scaffold-based diversity analysis and biosynthetic systems facilitating genomics-guided discovery.

Computational Tools for NP Chemical Space Navigation

Computational methods have become indispensable for mapping and navigating natural product chemical space. **ChemGPS-NP** and **Scaffold Hunter** represent two widely used tools that enable researchers to explore biologically relevant NP chemical space in a focused and targeted fashion [24]. These cheminformatics platforms help bridge the gap between computational methods and compound library synthesis, integrating cheminformatics and chemical space analyses with synthetic chemistry and biochemistry to successfully identify novel small molecule modulators of protein function [24].

The analytical power of these tools stems from their ability to process multidimensional molecular descriptors that define the dimensionality of chemical space [1]. Recent advances include the development of more universal molecular descriptors, such as **MAP4 fingerprints** and neural network embeddings from chemical language models, which can accommodate entities ranging from small molecules to biomolecules [1]. These tools are particularly valuable for identifying "holes" in existing screening data sets—regions of chemical space that can and should be explored by chemistry and biology to discover new bioactive compounds [24].

Quantitative Comparison: Explored vs. Underexplored NP Regions

Table 1: Structural and Property-Based Comparison of Natural Products and Trade Drugs

Characteristic	Natural Products	Trade Drugs	Data Source
Average Molecular Weight	356	360	[23]
Average log P value	2.9	2.5	[23]
Number of Ring Systems	1,748	807	[23]
Rule-of-5 Violations	Similar percentage	Similar percentage	[23]
Hydrogen Bond Donors	Fewer per molecule	More per molecule	[23]
Bridgehead Atoms	Much higher number	Lower number	[23]
Chiral Centers	Many more per molecule	Fewer per molecule	[23]

Table 2: Heavily Explored vs. Underexplored Regions of NP Chemical Space

Aspect	Heavily Explored Regions	Underexplored Regions	Research Implications
Structural Classes	Flavonoids, indole alkaloids, opium alkaloids, common scaffold systems	Macrocycles, RiPPs (ribosomally synthesized and post-translationally modified peptides), metallodrugs	New structural motifs with potentially novel mechanisms of action [27] [28] [1]
Source Organisms	Soil-derived actinomycetes, terrestrial plants	Microbes from extreme environments, marine symbionts, cyanobacteria, hot sulfur springs	Unique biosynthetic pathways and enzymatic transformations [25]
Chemical Space Properties	Drug-like properties, Rule-of-5 compliance, well-characterized pharmacology	Beyond Rule of 5 (bRo5) compounds, protein-protein interaction inhibitors, PROTACs	Challenges in synthesis and optimization, but potential for targeting difficult therapeutic areas [28] [1]
Discovery Approaches	Bioactivity-guided fractionation, traditional natural product chemistry	Genome mining, metabolomics, bioengineering, synthetic biology	Access to cryptic biosynthetic gene clusters and previously inaccessible chemical diversity [25] [23]

The data reveal that while natural products share many drug-like properties with trade drugs, they explore significantly more structural diversity, particularly in complex ring systems and stereochemistry [23]. This structural complexity contributes to their biological relevance but also presents challenges for synthesis and modification. The underexplored regions of NP chemical space are characterized by structural classes that fall outside traditional drug-like property space, source organisms from extreme or difficult-to-access environments, and novel biosynthetic pathways [25] [28] [1].

Heavily Explored Regions of NP Chemical Space

Traditional Natural Product Classes and Scaffolds

Certain classes of natural products have been extensively investigated due to their historical therapeutic success and relative accessibility. **Flavonoids** and **alkaloids** represent two such heavily explored families, with well-established biosynthetic pathways, known pharmacological activities, and extensive structure-activity relationship data [27]. These compounds typically exhibit favorable drug-like properties, with molecular weights and log P values falling within ranges comparable to approved drugs [23]. The structural classification of natural products (SCONP) has further illuminated that certain molecular scaffolds recur frequently among known natural products, creating regions of chemical space that have been systematically explored for drug discovery [26].

The heavy exploration of these regions is evidenced by the fact that more than 100 marketed macrocycle drugs are almost exclusively derived from natural products, yet this structural class remains poorly explored within targeted drug discovery efforts [28]. Similarly, natural products have contributed significantly to approved drugs across multiple therapeutic areas: 78% of antibacterial drugs, 75% of platelet aggregation inhibitors, 61% of anticancer drugs, 48% of anti-hypotensive drugs, 47.6% of antiulcer drugs, and 32.5% of anti-inflammatory drugs have a natural origin [23]. This extensive exploration has generated robust structure-activity relationship data for these compound classes but has also led to diminishing returns in discovering truly novel chemotypes from traditional sources.

Limitations and Repetition in Explored Regions

The heavy focus on specific natural product classes and source organisms has resulted in significant redundancy in discovery efforts. Recent analyses indicate that although the total number of characterized natural products has increased over the last decades, only a small percentage of recently discovered compounds possess previously unknown chemical structures [25]. This repetition stems from several factors: the repeated isolation of

known compounds from related species, the focus on easily cultivable microorganisms from similar ecological niches, and the application of standardized extraction and isolation procedures that selectively capture certain chemical classes while missing others.

This redundancy presents a substantial challenge for drug discovery, particularly in areas like antibiotic development where structurally new chemicals are urgently required for resistance-breaking properties [25]. The known natural product chemical space likely represents only "the tip of the iceberg," with significant biosynthetic potential remaining concealed in underexplored organisms, environments, and biosynthetic pathways [25]. Overcoming this limitation requires deliberate exploration of untapped regions of NP chemical space through innovative approaches and technologies.

Underexplored Regions of NP Chemical Space

Structural Classes with Discovery Potential

Several structural classes of natural products remain underexplored despite their significant potential for drug discovery. **Macrocycles**, defined as compounds containing rings of 12 or more atoms, represent a particularly promising yet underexploited structural class [28]. These compounds provide diverse functionality and stereochemical complexity in a conformationally pre-organized ring structure, which can result in high affinity and selectivity for protein targets while preserving sufficient bioavailability to reach intracellular locations [28]. Macrocycles have demonstrated repeated success when addressing targets that have proved highly challenging for standard small-molecule drug discovery, especially in modulating macromolecular processes such as protein-protein interactions [28].

Other underexplored structural classes include **ribosomally synthesized and post-translationally modified peptides (RiPPs)**, which exhibit remarkable structural diversity and bioactivities [25]. Recent research has identified ribosomally derived lipopeptides containing distinct fatty acyl moieties as a promising area for exploration [25]. Additionally, **metal-containing natural products** represent a structurally and functionally important class that is commonly excluded from standard chemoinformatics analyses due to modeling challenges [1]. The difficulty of modeling these regions of BioReCS should not justify their exclusion from systematic exploration, as they may offer unique therapeutic opportunities.

Underexplored Source Organisms and Environments

The biosynthetic potential of certain microbial groups and extreme environments remains largely untapped. **Cyanobacteria** and microbes that colonize extreme habitats represent talented but neglected natural product producers [25]. These organisms often possess unique biosynthetic pathways evolved to produce specialized metabolites under challenging environmental conditions, resulting in chemical structures not found in organisms from conventional sources.

Recent advances in metagenomics have revealed that the wealth of publicly available (meta)genomes conceals significant biosynthetic potential that has yet to be elucidated [25]. One comprehensive study of the global ocean microbiome uncovered extensive biosynthetic diversity, with thousands of new biosynthetic gene clusters identified in marine microorganisms [25]. The isolation of natural products from habitats and organisms previously thought to lack natural product biosynthesis potential (e.g., hot sulfur springs) further supports the hypothesis that known natural product chemical space represents only a fraction of what exists in nature [25].

Dark Chemical Matter and Inactive Compounds

A particularly intriguing underexplored region of biologically relevant chemical space consists of so-called **"dark chemical matter"** – compounds that have repeatedly failed to show activity in high-throughput screening assays [1]. These molecules represent the non-biologically relevant portions of chemical space and provide crucial boundary conditions for understanding bioactivity. Recent efforts have led to the development of InertDB, a curated collection of 3,205 experimentally confirmed inactive compounds supplemented with 64,368 putative inactive molecules generated using deep generative artificial intelligence models [1]. Understanding why these compounds lack activity can provide equally valuable insights for drug discovery as studying successful bioactive molecules.

Experimental Protocols for Exploring Underexplored NP Space

Genomics-Guided Discovery Workflow

The integration of genomic information with natural product chemistry has emerged as a powerful approach for targeted exploration of underexplored regions of NP chemical space. The following protocol outlines a genomics-guided discovery workflow:

Table 3: Research Reagent Solutions for Genomics-Guided NP Discovery

Research Reagent	Function/Application	Experimental Role
Metagenomic DNA Libraries	Source of biosynthetic gene clusters from unculturable microorganisms	Provides access to genetic potential of microbial communities without cultivation [25]

Research Reagent	Function/Application	Experimental Role
Heterologous Expression Systems	Host organisms for expressing foreign biosynthetic gene clusters	Enables production of natural products from unculturable or genetically intractable organisms [25]
Bioinformatics Tools (e.g., antiSMASH)	Identification and analysis of biosynthetic gene clusters in genomic data	Guides target selection and predicts structural features of encoded natural products [25]
Mass Spectrometry Platforms	Detection and structural characterization of novel natural products	Links biosynthetic gene clusters to their metabolic products through metabolomics [23]

Protocol Steps:

- **Sample Collection and DNA Extraction**: Collect environmental samples from underexplored niches (e.g., extreme environments, marine sediments). Extract high-molecular-weight DNA suitable for metagenomic library construction [25].
- **Sequencing and Bioinformatic Analysis**: Perform whole-metagenome sequencing using Illumina or PacBio platforms. Analyze sequence data using specialized bioinformatics tools (e.g., antiSMASH, PRISM) to identify biosynthetic gene clusters with novel architectures [25].
- **Heterologous Expression**: Clone promising biosynthetic gene clusters into suitable expression hosts (e.g., *Streptomyces coelicolor*, *E. coli*). Optimize expression conditions to activate silent gene clusters [25].
- **Metabolite Analysis and Isolation**: Compare metabolic profiles of expression hosts containing target gene clusters against control strains using LC-HRMS. Isplicate novel compounds using bioactivity-guided or mass-guided fractionation [25] [23].
- **Structural Elucidation and Bioactivity Testing**: Determine structures of novel compounds using NMR, MS/MS, and other spectroscopic techniques. Evaluate bioactivity against target disease models, with particular attention to resistance-breaking antimicrobial activity [25].



Bioengineering and Synthetic Biology Approaches

Bioengineering provides powerful methods to access underexplored regions of natural product chemical space through targeted modification of biosynthetic pathways:

Protocol Steps:

- **Pathway Refactoring**: Redesign complete biosynthetic gene clusters using synthetic biology principles to optimize expression and enable genetic manipulation [25].
- **Combinatorial Biosynthesis**: Exchange domains in modular biosynthetic enzymes (e.g., polyketide synthases, nonribosomal peptide synthetases) to create novel hybrid pathways [25].
- **Precursor-Directed Biosynthesis**: Supplement producing organisms with non-natural substrate analogs to shunt biosynthesis toward novel derivatives [25].
- **Enzyme Engineering**: Apply directed evolution or structure-based design to modify substrate specificity of key biosynthetic enzymes, enabling production of "non-natural" natural products [25].
- **Pathway Activation**: Employ genetic techniques (promoter engineering, regulatory gene overexpression) to activate silent/cryptic biosynthetic gene clusters [25].

Visualization of NP Chemical Space Exploration Strategies

The systematic exploration of underexplored regions of natural product chemical space requires an integrated approach combining multiple scientific disciplines and methodologies. The following diagram illustrates the workflow for discovering novel natural products from underexplored sources, highlighting the interdisciplinary nature of modern natural product research:



Table 4: Essential Research Tools and Resources for NP Chemical Space Exploration

Tool/Resource Category	Specific Examples	Application in NP Research
Chemical Space Navigation	ChemGPS-NP, Scaffold Hunter	Guide exploration of biologically relevant NP chemical space in a focused and targeted fashion [24]
Bioinformatics Platforms	antiSMASH, PRISM, MIBiG	Identify and analyze biosynthetic gene clusters in genomic and metagenomic data [25]
Analytical Technologies	LC-HRMS, MS Imaging, NMR	Detect, characterize, and visualize natural products in complex biological matrices [23]
Genomic Resources	Metagenomic libraries, Heterologous expression systems	Access biosynthetic potential of unculturable microorganisms and engineer biosynthetic pathways [25]
Specialized Compound Libraries	Macrocyclic libraries, RiPP libraries, Dark chemical matter collections	Focus screening efforts on underexplored regions of chemical space [28] [1]

The systematic exploration of natural product chemical space represents a crucial frontier in drug discovery, particularly as resistance to existing therapies grows and challenging targets require innovative chemical solutions. While heavily explored regions of NP chemical space have provided numerous therapeutic agents, they face diminishing returns in yielding truly novel chemotypes. In contrast, underexplored regions—including macrocycles, RiPPs, metabolites from extreme environments, and cryptic biosynthetic pathways—offer significant opportunities for discovering compounds with resistance-breaking properties and novel mechanisms of action. The integrated application of genomics, bioinformatics, synthetic biology, and advanced analytical technologies provides powerful methods to navigate and populate these underexplored regions of natural product chemical space. As computational tools continue to evolve and our understanding of biosynthetic pathways expands, targeted exploration of these underexplored regions will play an increasingly important role in addressing unmet medical needs through natural product-inspired drug discovery.

The concept of the biologically relevant chemical space (BioReCS) provides a foundational framework for modern drug discovery. BioReCS encompasses all molecules with biological activity—both beneficial and detrimental—spanning diverse application areas including drug discovery, agrochemistry, and natural product research [1]. This chemical universe is vast, with estimates suggesting the existence of up to 10^60 drug-like compounds, creating a fundamental challenge for researchers seeking to identify novel therapeutic agents [29]. Within this expansive universe, natural products (NPs) occupy a particularly privileged region, characterized by unique structural complexity and high relevance to human biology. Analyses reveal that over half of approved small-molecule drugs originate directly or indirectly from natural products, highlighting their enduring importance [12].

The structural and physicochemical properties of natural products differ significantly from typical synthetic compounds. Natural products often feature greater stereochemical complexity, higher sp^3-hybridized carbon counts, more oxygen atoms, and intricate ring systems that confer sophisticated three-dimensional architectures [30]. These characteristics enable natural products to interact with challenging biological targets, including protein-protein interactions, which have proven difficult to modulate with conventional synthetic compounds [30]. Despite the known structural diversity of natural products, current databases document approximately 1.1 million natural products, with only about 10% readily obtainable for experimental testing, creating a significant accessibility challenge [12]. This gap between known structures and readily testable compounds represents a critical bottleneck in natural product-based drug discovery.

Table 1: Key Characteristics of Natural Product Chemical Space

Property	Natural Products	Synthetic/Drug-like Compounds
Structural Complexity	High (multiple stereocenters, complex ring systems)	Variable, often lower
sp^3 Hybridized Carbons	Higher fraction (Fsp^3)	Lower fraction
Oxygen Content	High	Variable
Number of Aromatic Rings	Generally lower	Generally higher
Relevance to Drug Discovery	>50% of approved drugs NP-derived	Foundation of combinatorial libraries
Readily Accessible Compounds	~10% of known structures	High percentage

Charting Unexplored Territories in Natural Product Chemical Space

Analytical Tools for Navigation

Systematic exploration of natural product chemical space requires specialized computational tools that can map its complex topography. Platforms such as ChemGPS-NP and Scaffold Hunter enable researchers to navigate biologically relevant regions in a focused manner, identifying both densely populated and sparsely explored areas [24]. These cheminformatic tools employ dimensionality reduction techniques to project high-dimensional

chemical descriptor data into visualizable and interpretable formats, allowing researchers to identify structural patterns and anomalies across large compound collections [1].

Recent advances in molecular representation have been critical for effective chemical space analysis. While traditional descriptors were optimized for small organic molecules, newer approaches like the MAP4 fingerprint and neural network embeddings from chemical language models offer more universal representations that can accommodate diverse molecular classes ranging from small molecules to peptides and even metallodrugs [1]. These improved descriptors facilitate more meaningful comparisons across different regions of chemical space and enable the identification of truly novel scaffolds with potential bioactivity.

Underexplored Regions and Dark Chemical Matter

Analysis of natural product chemical space reveals several underexplored regions with high potential for drug discovery. Certain structural classes remain underrepresented in current screening collections, including metal-containing molecules, large and complex natural products, macrocycles, protein-protein interaction (PPI) modulators, PROTACs, and mid-sized peptides [1]. Many of these compounds fall into the "beyond Rule of 5" (bRo5) category, presenting both challenges and opportunities for drug development [1].

Marine natural products represent another distinctive region of chemical space, characterized by larger molecular weights and greater hydrophobicity compared to their terrestrial counterparts [12]. Particularly interesting are natural products derived from deep-sea and extremophile organisms, which often display novel scaffolds and notable bioactivities honed by adaptation to unique environmental conditions [12]. The continued discovery of such structurally distinct compounds highlights the value of exploring diverse biological sources.

Beyond these structural classes, the concept of "dark chemical matter"—compounds that consistently show no activity in high-throughput screens—provides valuable negative data that helps define the boundaries of BioReCS [1]. Similarly, databases of curated inactive compounds, such as InertDB, which includes both experimentally determined and AI-generated putative inactive molecules, contribute to our understanding of the structural features that separate bioactive from non-bioactive chemical space [1].

Table 2: Key Public Databases for Exploring Natural Product Chemical Space

Database	Scope	Key Features
COCONUT (Collection of Open Natural Products)	Comprehensive NP collection	>400,000 fully characterized natural products [31]
ChEMBL	Bioactive drug-like molecules	Extensive biological activity annotations [1]
PubChem	Chemical substances and bioactivities	Large repository with screening data [1]
Super Natural II	Natural products	Includes predicted bioactivity and pathways [12]
NAPRORE-CR	Costa Rican natural products	Geographically focused NP database [12]
PeruNPDB	Peruvian natural products	Regional focus for drug screening [12]

Strategic Approaches to Populating Chemical Space through Synthesis

Natural Product-Inspired Synthesis Strategies

Several sophisticated synthesis strategies have emerged to systematically populate promising regions of natural product chemical space with novel compounds. These approaches leverage the structural information encoded in natural products while introducing significant diversity to explore surrounding chemical space.

Biology-Oriented Synthesis (BIOS) proceeds from the premise that natural products are "privileged structures" with inherent biological relevance. This strategy employs natural products as starting points for designing focused libraries that retain core structural elements of the original bioactive compound while introducing strategic modifications [32] [30]. For example, Waldmann's development of an oxepane-based library inspired by bioactive natural products like heliannuol B and zoapatanol led to the discovery of novel Wnt signaling modulators that interact with the previously undrugged target Vangl1 [32]. BIOS libraries typically contain fewer compounds than traditional combinatorial libraries but demonstrate higher hit rates due to their foundation in evolutionarily validated scaffolds.

Diversity-Oriented Synthesis (DOS) aims to generate broad structural diversity through branching reaction pathways that produce compounds with varied skeletons and stereochemistries from common intermediates [32] [30]. Unlike target-oriented synthesis, DOS employs forward synthetic analysis to create structurally complex and diverse libraries that populate expansive regions of chemical space. A prominent example includes Schreiber's work generating a library of 2,070 macrolactone-based small molecules, which led to the discovery of robotnikinin—a potent inhibitor of the Hedgehog signaling pathway with potential applications in cancer treatment [30]. DOS libraries are particularly valuable for phenotypic screening campaigns where the biological targets may not be fully characterized.

Pharmacophore-Directed Retrosynthesis (PDR) represents a more recent strategy that integrates synthetic planning with the identification of key structural features essential for bioactivity [32]. This approach begins with a retrospective analysis of structure-activity relationships to identify critical pharmacophoric elements, then designs synthetic routes that maximize opportunities to generate analogs exploring variations in these key features. PDR aims to balance the efficiency of total synthesis with the systematic investigation of structure-activity relationships throughout the synthetic process.



Diagram 1: NP-Inspired Synthesis Strategies

Advanced Methodologies for Late-Stage Molecular Diversification

Beyond comprehensive library synthesis strategies, recent methodological advances enable precise molecular editing that can dramatically expand accessible chemical space from advanced intermediates. These approaches are particularly valuable for lead optimization phases where subtle structural modifications can significantly improve drug properties.

Skeletal Editing techniques allow direct modification of molecular frameworks through atom insertion, deletion, or exchange. A groundbreaking example is the development of sulfenylcarbene-mediated carbon atom insertion into N-heterocycles, which enables the transformation of existing drug scaffolds into new candidates by adding just one carbon atom at room temperature under metal-free conditions [33]. This method achieves yields up to 98% and is compatible with DNA-encoded library technology, making it particularly valuable for late-stage diversification of lead compounds [33]. The ability to perform such precise molecular surgery represents a paradigm shift in medicinal chemistry, potentially reducing drug development costs by enabling efficient renovation of existing molecular structures rather than requiring de novo synthesis.

Ring Distortion of Natural Products capitalizes on the complex ring systems found in many natural products by subjecting them to reaction conditions that dramatically rearrange their core structures. This approach can generate diverse, natural product-like compounds that would be challenging to access through conventional synthesis. The resulting libraries maintain the three-dimensional complexity and fraction of sp^3-hybridized carbons characteristic of bioactive natural products while exploring unprecedented structural space around the original scaffold.

Hybrid Natural Products combine structural elements from two or more biologically active natural products to create novel compounds with potentially enhanced or dual activities. This strategy mimics nature's own evolutionary approach, as exemplified by the potent anticancer natural product vincristine, which represents a hybrid of the simpler alkaloids vindoline and catharanthine [30]. Synthetic hybridization enables the exploration of chemical space between known bioactive regions, potentially yielding compounds with novel mechanisms of action.

Table 3: Experimental Approaches for Chemical Space Exploration

Methodology	Key Reagents/Techniques	Applications	Experimental Considerations
Skeletal Editing	Sulfenylcarbene reagents, metal-free conditions	Late-stage functionalization, lead optimization	Bench-stable reagents, room temperature operation, compatibility with DNA-encoded libraries [33]
DOS	Branching pathways, multicomponent reactions, complexity-generating transformations	Library generation for phenotypic screening	Aim for ≤5 steps, incorporate stereochemical diversity, use pluripotent intermediates [30]

Methodology	Key Reagents/Techniques	Applications	Experimental Considerations
BIOS	Natural product-inspired building blocks, target-oriented synthesis	Focused library for specific target classes	Prioritize privileged NP scaffolds, employ divergent synthesis from common intermediates [32]
Ring Distortion	Lewis acids, oxidants, rearranging conditions	Generating complexity and diversity from NP starting materials	Use stable, readily available NPs, employ multiple reaction conditions on single substrate [30]

Computational and Experimental Workflows for Targeted Exploration

Integrated Screening Protocols

The effective exploration of natural product chemical space requires tight integration between computational prediction and experimental validation. Advanced screening protocols leverage the complementary strengths of both approaches to efficiently navigate vast molecular libraries toward promising bioactive compounds.

Active Learning with Alchemical Free Energy Calculations represents a powerful workflow that combines machine learning with physics-based binding affinity predictions. In this approach, an initial set of compounds is evaluated using computationally intensive alchemical free energy calculations, which provide highly accurate binding affinity estimates [29]. These data then train machine learning models that can rapidly predict affinities for much larger compound libraries. The most promising compounds from these predictions (selected through strategies like greedy selection or mixed uncertainty sampling) are subsequently validated through additional free energy calculations, creating an iterative refinement cycle that efficiently converges toward high-affinity binders [29]. This approach dramatically reduces the computational resources required to screen large libraries while maintaining high prediction accuracy.

High-Throughput Synthetic Platform technologies have emerged to accelerate the production and analysis of compound libraries. For instance, Blair and colleagues developed an approach that simplified the analysis of thousands of simultaneous reactions by focusing on molecular fragments, reducing analysis time from two months to a single day while generating 5,000 new chemicals through 20,000 reactions [34]. Such platforms address the critical bottleneck in chemical space exploration by enabling rapid synthesis and characterization of diverse compound collections, making large-scale investigation of underexplored chemical regions practically feasible.



Diagram 2: Integrated Screening Workflow

AI-Enhanced Natural Product Discovery

Artificial intelligence has revolutionized natural product discovery by enabling the generation of novel natural product-like structures that expand beyond known chemical space. Deep generative models, particularly recurrent neural networks (RNNs) with long short-term memory (LSTM) units trained on natural product SMILES representations, can produce vast libraries of novel yet biologically relevant structures [31]. One such effort generated 67 million natural product-like molecules—a 165-fold expansion over the approximately 400,000 known natural products—while maintaining distributions of natural product-likeness scores similar to authentic natural products [31].

These AI-generated compound libraries significantly expand the accessible chemical space for drug discovery, populating regions with structural novelty while maintaining the favorable physicochemical properties associated with natural products. The generated compounds exhibit expanded physiochemical and structural space compared to known natural products, as visualized through t-SNE projections of molecular descriptors [31]. This approach effectively inverts the traditional discovery process by first generating promising virtual structures that can then be prioritized for synthesis and testing, potentially uncovering entirely new classes of bioactive compounds.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 4: Essential Research Reagents and Materials for Chemical Space Exploration

Reagent/Material	Function	Application Examples
Sulfenylcarbene Reagents	Carbon atom insertion into N-heterocycles	Late-stage skeletal editing of drug candidates [33]
DNA-Encoded Library (DEL) Components	Facilitating high-throughput screening of billions of compounds	Building diverse libraries for protein binding screens [33]
Pluripotent Building Blocks	Branching point substrates for DOS	Generating diverse scaffolds from common intermediates [30]
Solid-Supported Phosphonates	Facilitating parallel synthesis and purification	DOS library generation with simplified workup [30]
Natural Product Fragment Libraries	Starting points for BIOS and hybrid molecules	Exploring NP-inspired regions of chemical space [12]
Bench-Stable Carbene Precursors	Safe, metal-free carbene generation	Sustainable skeletal editing compatible with DEL [33]
Chromatin/Nucleosome Assembly Kits	Creating biologically relevant screening substrates	Targeting epigenetic mechanisms in drug discovery [34]

The systematic exploration of natural product chemical space represents a paradigm shift in drug discovery, moving from serendipitous finding to rational design. By integrating advanced cheminformatic analysis with innovative synthetic methodologies, researchers can now navigate the vast landscape of possible drug-like molecules with unprecedented precision and efficiency. The strategies outlined—from biology-oriented and diversity-oriented synthesis to skeletal editing and AI-generated molecular design—provide a comprehensive toolkit for populating underexplored yet biologically relevant regions of chemical space.

Future advances will likely focus on improving the integration between computational prediction and experimental validation, further accelerating the discovery cycle. Additionally, as synthetic methodologies continue to evolve, particularly those enabling late-stage functionalization and skeletal editing, the ability to fine-tune molecular properties while maintaining core bioactivity will become increasingly sophisticated. The ongoing development of open-access natural product databases and analysis tools will further democratize access to chemical space exploration, potentially unlocking novel therapeutic opportunities for challenging disease targets. Through the continued bridging of chemical space analysis and novel synthesis, drug discovery can more effectively harness the rich structural diversity evolved in nature while expanding into entirely new regions of chemical space with designed synthetic compounds.

Modern Arsenal for Exploration: From AI and Omics to High-Throughput Screening

Leveraging AI and Machine Learning for Target Identification and Property Prediction

The exploration of natural product (NP) chemical space represents a frontier of untapped therapeutic potential, historically plagued by insurmountable complexity and scale. The process of identifying bioactive compounds and predicting their properties from millions of candidate structures has been a protracted, resource-intensive endeavor. The integration of **Artificial Intelligence (AI)** and **Machine Learning (ML)** is now fundamentally reshaping this landscape, transforming NP-based drug discovery from a slow, empirical process into a predictive, data-driven science [35] [13]. These technologies are enabling researchers to navigate the vast, intricate chemical space of NPs—a space estimated to contain over 1060 drug-like molecules—with unprecedented speed and precision [36] [37]. This whitepaper provides an in-depth technical guide to the core AI/ML methodologies revolutionizing target identification and property prediction within the context of natural product drug discovery, detailing experimental protocols, benchmarking data, and essential computational tools for the modern research scientist.

The Natural Product Cheminformatics Landscape

Natural products are chemical compounds or substances produced by living organisms, including plants, animals, and microorganisms [13]. They have served as a rich source of biologically active compounds, with approximately 50% of FDA-approved medications between 1981 and 2006 being NPs or their synthetic derivatives [13]. However, the discovery of drugs derived from NPs presents numerous challenges, including the limited availability of bioactive molecules, the complexity of molecular structures, low yields of promising compounds, and the labor-intensive process of isolation and structural elucidation [13].

The accelerating growth of make-on-demand and virtual chemical libraries provides unprecedented opportunities but also creates a fundamental bottleneck. While these libraries now contain >70 billion readily available molecules, the number of possible drug-like molecules is estimated to be

more than 1060, exceeding the size of chemical libraries evaluated in early drug discovery by many orders of magnitude [37]. This disparity highlights the critical need for more efficient virtual screening approaches capable of evaluating these vast chemical libraries [37].

Table 1: Key Challenges in Natural Product Drug Discovery and AI-Driven Solutions

Challenge	Traditional Approach	AI/ML Solution	Impact
Dereplication	Manual literature review & experimental comparison	AI-powered database mining & pattern recognition [38]	Reduces redundant discovery efforts
Target Identification	Bioassay-guided fractionation	Predictive bioactivity modeling & reverse docking [13] [38]	Accelerates hypothesis generation
Property Prediction	Empirical structure-activity relationship (SAR) studies	Quantitative Structure-Activity/Property Relationship (QSA/PR) models [39] [38]	Enables *in silico* ADMET profiling
Chemical Space Exploration	Limited library screening	Deep generative models & latent space navigation [40] [41]	Expands access to novel scaffolds

AI's role in this domain is rapidly expanding. Analysis of the CAS Content Collection, the largest human-curated collection of published scientific information, found over 600,000 scientific publications related to natural product research since 2010, with a notable increase in AI applications [38]. The most common AI application in natural products is in anti-tumor agents, followed by antiviral and antibacterial agents [38].

AI-Driven Target Identification and Binding Affinity Prediction

Machine Learning-Guided Docking Screens

Structure-based virtual screening of ultralarge libraries has identified ligands of important therapeutic targets, but evaluating massive libraries requires substantial computational resources [37]. A breakthrough strategy combining machine learning and molecular docking enables rapid virtual screening of databases containing billions of compounds, reducing the computational cost by more than 1,000-fold [37].

The core protocol involves training a classification algorithm to identify top-scoring compounds based on molecular docking of a subset (e.g., 1 million compounds) to the target protein. The conformal prediction framework then makes selections from the multi-billion-scale library, drastically reducing the number of compounds requiring explicit docking scoring [37]. In application to a library of 3.5 billion compounds, this protocol successfully identified ligands of G protein-coupled receptors (GPCRs), one of the most important families of drug targets [37].

Table 2: Performance Benchmark of ML-Guided Docking Workflow [37]

Metric	Standard Docking	ML-Guided Docking	Improvement
Library Size	3.5 billion compounds	3.5 billion compounds	-
Compounds Docked	3.5 billion	~25-30 million	>100-fold reduction
Computational Cost	~493 trillion complexes predicted	Not specified	>1,000-fold reduction
Sensitivity	Baseline (100%)	87-88%	Maintains high recall
Experimental Hit Rate	Target-dependent	Successfully identified GPCR ligands	Confirmed utility

Experimental Protocol: ML-Accelerated Virtual Screening Pipeline

Objective: To identify potential ligands for a target protein from an ultralarge chemical library while reducing computational requirements by >100-fold.

Input Requirements:

- Target protein structure (experimental or predicted)
- Ultralarge chemical library (e.g., Enamine REAL Space, ZINC15)
- Docking software (e.g., AutoDock, Schrödinger)
- ML libraries (e.g., CatBoost, PyTorch)

Methodology:

- **Initial Docking Screen**: Prepare and dock a representative subset of 1 million compounds from the larger library against the prepared protein structure.

- **Model Training**: Train a classifier (CatBoost with Morgan2 fingerprints recommended) to distinguish between active and inactive compounds based on the docking scores from the initial screen. The energy threshold for the active class is typically set based on the top-scoring 1% of the initial screen.
- **Conformal Prediction**: Apply the Mondrian conformal prediction framework to the entire multi-billion compound library using the trained model. This step assigns normalized P values to each compound, indicating their likelihood of being active.
- **Compound Selection**: Based on a chosen significance level (ε), the framework divides compounds into virtual active, virtual inactive, or uncertain categories. The virtual active set is selected for explicit docking.
- **Experimental Validation**: The top-ranked compounds from the final docking screen are procured and validated using binding assays (e.g., CETSA for cellular target engagement) and functional assays [37] [42].



Advanced Molecular Representation and Property Prediction

Molecular Descriptors for QSAR Modeling

The predictive power of AI in property prediction hinges on effective molecular representations. Quantitative Structure-Activity Relationship (QSAR) modeling correlates numerical molecular descriptors with biological activity or physicochemical properties [39]. These descriptors are classified by dimensions:

- **1D Descriptors**: Encode global properties like molecular weight and atom counts [39].
- **2D Descriptors**: Include topological indices encoding connectivity patterns [39].
- **3D Descriptors**: Capture molecular shape, volume, and electrostatic potential maps [39].
- **4D Descriptors**: Account for conformational flexibility using ensembles of molecular structures [39].

Modern AI approaches have expanded these to include learned representations. Deep learning techniques generate "deep descriptors" derived from molecular graphs or SMILES strings without manual engineering, capturing abstract and hierarchical molecular features [39].

Table 3: Molecular Representations in AI-Driven Drug Discovery [39] [43]

Representation	Description	AI Applications	Advantages	Limitations
SMILES Strings	1D linear notation of chemical structure	RNN, LSTM, Transformers (ChemBERTa) [43]	Simple, compact, widely supported	Non-unique, sensitive to syntax errors

Representation	Description	AI Applications	Advantages	Limitations
Molecular Fingerprints	Bit vectors indicating substructure presence (ECFP4, Morgan)	Classical ML, Deep Learning [37] [43]	Fixed-length, suitable for similarity search	Lack 3D stereochemical detail
Molecular Graphs	Atom-bond networks with nodes and edges	Graph Neural Networks (GNN, GCN, GAT) [39] [43]	Preserves atomic connectivity and topology	Computationally expensive
3D Representations	Atomic coordinates and spatial relationships	SchNet, DimeNet, GeoMol [43]	Captures stereochemistry and shape	Requires conformer generation

Experimental Protocol: Building a Robust QSAR Model

Objective: To develop a predictive QSAR model for ADMET (Absorption, Distribution, Metabolism, Excretion, and Toxicity) properties or bioactivity of natural products.

Data Curation:

- Collect experimental bioactivity/ADMET data from public databases (ChEMBL, PubChem, DrugCentral) or in-house assays.
- Apply strict curational filters for endpoint accuracy and chemical structure standardization.
- Divide dataset into training (80%), calibration (optional), and test (20%) sets, ensuring structural diversity.

Feature Engineering & Model Training:

- Compute molecular descriptors (e.g., using RDKit, DRAGON) or generate learned representations (graph embeddings, SMILES-based transformers).
- Apply dimensionality reduction (PCA, RFE) or feature selection (LASSO, mutual information) to eliminate redundant variables.
- Train multiple ML algorithms (Random Forest, SVM, Gradient Boosting, Deep Neural Networks) using cross-validation to optimize hyperparameters.
- For deep learning approaches, use appropriate architectures: Graph Neural Networks for molecular graphs, RNN/Transformers for SMILES, or hybrid models.

Model Validation & Interpretation:

- Evaluate performance using both internal (Q^2, ROC-AUC) and external validation metrics on the held-out test set.
- Apply interpretability methods (SHAP, LIME) to identify which structural features contribute most to predictions, generating testable hypotheses for medicinal chemistry optimization [39].

Deep Generative Models for Exploring Natural Product Space

Expanding Confined Chemical Space

The therapeutic potential of natural products is often confined to specific regions of chemical space. Deep generative models provide an alternative approach to explore wider drug-like chemical spaces [40] [41]. These models can generate novel molecular structures with desired properties, capturing the chemical space of known drugs while expanding into unexplored territories [40].

Conditional generative models, such as the Conditional Randomized Transformer with molecular fingerprints as a condition, can perform guided exploration in drug-like chemical space [40]. The combination of quantitative estimation of drug-likeness (QED) and quantitative estimate of protein-protein interaction targeting drug-likeness (QEPPI) can cover a larger drug-like space than either metric alone [40].

Experimental Protocol: De Novo Molecular Design with Generative AI

Objective: To generate novel, synthetically accessible natural product-inspired compounds with optimized properties.

Model Selection & Training:

- **Architecture Choices**: Conditional Variational Autoencoders (CVAE), Generative Adversarial Networks (GANs), or Transformer-based models (e.g., ChemBERTa) [40] [43].
- **Training Data**: Curate a high-quality dataset of known natural products and bioactives (e.g., from COCONUT, NPASS, or proprietary libraries).
- **Conditioning**: Define desired properties for conditioning (e.g., high QED, specific target prediction, low toxicity).

Generation & Optimization:

- Sample from the latent space of the trained model to generate novel molecular structures.
- Use transfer learning to fine-tune generative models on specific natural product sub-families (e.g., alkaloids, terpenoids).
- Apply reinforcement learning or Bayesian optimization to steer generation toward multi-parameter optimization (potency, solubility, metabolic stability).

Validation & Synthesis Planning:

- Filter generated molecules using synthetic accessibility (SA) scores and retrosynthesis tools (e.g., ASKCOS, AiZynthFinder) [41].
- Select top candidates for in silico validation (docking, ADMET prediction) before proceeding to synthesis and biological testing [40].



The Scientist's Toolkit: Essential Research Reagents & Databases

Successful implementation of AI-driven NP discovery requires access to specialized computational tools, databases, and experimental reagents. The following table catalogs essential resources referenced in contemporary literature.

Table 4: Essential Research Reagents & Computational Tools for AI-Driven NP Discovery

Resource Name	Type	Function/Application	Reference
Enamine REAL Space	Chemical Library	>70 billion make-on-demand compounds for virtual screening	[37]
CAS Content Collection	Database	Human-curated collection of published scientific information on NPs	[38]
CETSA (Cellular Thermal Shift Assay)	Experimental Method	Validates direct target engagement in intact cells and tissues	[42]
CatBoost	ML Algorithm	Gradient boosting classifier optimal for molecular fingerprint data	[37]
RDKit	Cheminformatics	Open-source toolkit for descriptor calculation & cheminformatics	[37] [39]
NRPSpredictor2	Web Server	Predicts substrate specificity of NP biosynthetic enzymes using ML	[38]
AutoDock, SwissADME	Software Platform	Molecular docking and ADMET property prediction	[42]

Resource Name	Type	Function/Application	Reference
QSARINS	Software	Development and validation of classical QSAR models	[39]
NuBBE Database	Database	Specialized natural product database from Brazilian biodiversity	[38]

Future Perspectives and Concluding Remarks

The integration of AI and ML into natural product research marks a paradigm shift from serendipitous discovery to rational, predictive exploration. The emerging "lab-in-a-loop" concept, where AI algorithms are continuously refined using real-world experimental data, promises a future of autonomous, adaptive, and exponentially accelerating drug discovery [43]. This closed-loop, self-improving ecosystem represents the next frontier, transforming drug development from a linear, human-driven process into a cyclical, AI-driven process with human oversight [43].

However, challenges remain in the widespread adoption of these technologies. Data quality and standardization continue to be significant hurdles, particularly for natural products with complex stereochemistry and limited available data [38]. Model interpretability, regulatory acceptance, and the need for large-scale experimental validation of AI-generated designs are additional areas requiring continued focus [39] [41]. As these challenges are addressed, AI-driven exploration of natural product space will undoubtedly unlock novel therapeutic avenues, harnessing the best of what nature has to offer to address human disease.

High-Throughput Screening (HTS) represents an automated, robust approach to rapidly testing large collections of molecules for bioactivity, holding particular promise in antibacterial drug discovery where over 50% of marketed antibiotics originate from natural products (NPs) [44]. The screening of natural product libraries (NPLs) presents unique challenges and opportunities due to the complex chemical nature of NP extracts, which contain a plethora of molecules at varying concentrations with potential for antagonistic or synergistic biological activities [44]. Within this paradigm, two primary screening philosophies have emerged: cellular target-based HTS (CT-HTS) and molecular target-based HTS (MT-HTS). The selection between these approaches carries significant implications for hit identification, validation, and subsequent development within the broader context of exploring natural product chemical space for drug discovery research. This technical guide examines both methodologies, their experimental protocols, and their application in modern drug discovery pipelines.

Core HTS Approaches: Cellular vs. Molecular Target Screening

Cellular Target-Based HTS (CT-HTS)

Definition and Principle: CT-HTS, also referred to as whole cell-based or phenotypic screening, utilizes intact living cells to identify compounds that produce a desired phenotypic response, such as bacterial cell death or inhibition of viral replication, without prior knowledge of the specific molecular target [44] [45].

Key Characteristics:

- **System Complexity:** Screens compounds against the entire cellular system, including all potential molecular targets, membranes, and regulatory networks.
- **Target Identification:** Identifies intrinsically active compounds but requires secondary screening for target deconvolution and elimination of non-specific cytotoxic compounds [44].
- **Physiological Relevance:** Maintains biological context including cellular permeability, efflux mechanisms, and metabolic activation, potentially leading to higher clinical translation success [46].
- **Hit Validation:** Confirmed hits demonstrate biological activity in a physiologically relevant environment but may have unknown mechanisms of action.

Molecular Target-Based HTS (MT-HTS)

Definition and Principle: MT-HTS employs isolated molecular targets – typically purified proteins, enzymes, or nucleic acids – to identify compounds that interact with these specific biomolecules through binding or functional modulation [44] [45].

Key Characteristics:

- **Reductionist Approach:** Focuses on specific, well-defined molecular interactions, typically using purified target proteins in controlled in vitro conditions.
- **Mechanistic Clarity:** Provides immediate information about mechanism of action at the molecular level.
- **Technical Considerations:** May fail to identify compounds that require cellular activation or those whose activity depends on complex cellular contexts; may produce hits with poor cellular permeability or those susceptible to efflux [44].
- **Interference Challenges:** Requires secondary screening to eliminate pan assay interference molecules (PAINS) that produce false positives through non-specific binding or assay interference [44].

Table 1: Comparative Analysis of Cellular vs. Molecular Target HTS Approaches

Parameter	Cellular Target HTS (CT-HTS)	Molecular Target HTS (MT-HTS)
Screening System	Whole living cells (bacterial, fungal, mammalian)	Purified proteins, enzymes, or nucleic acids
Target Knowledge	Not required; phenotypic outcome driven	Required prior to screening
Physiological Context	Full physiological context maintained	Minimal to no physiological context
Hit Rate for NPs	~0.3% (with polyketides) [44]	Variable; typically lower than CT-HTS
Primary Advantage	Identifies compounds with cellular activity	Reveals specific molecular mechanisms
Primary Challenge	Target deconvolution required	May not translate to cellular activity
Secondary Screening	Eliminate non-specific cytotoxics	Eliminate PAINS and promiscuous binders

Advanced HTS Strategy: Mechanism-Informed Phenotypic Screening

A hybrid approach has emerged that combines advantages of both CT-HTS and MT-HTS through mechanism-informed phenotypic screening, most commonly implemented as reporter gene assays [44]. These assays utilize cells engineered with reporter constructs that produce measurable signals (e.g., luminescence, fluorescence) when specific pathways of interest are modulated. For example, the ATAD5-luciferase HTS assay identifies genotoxic compounds by exploiting the stabilization of ATAD5 protein following DNA damage [46]. This approach maintains physiological context while providing information about the signaling pathways with which hits interact, effectively bridging the gap between purely phenotypic and purely target-based screening [44].

Other innovative reporter systems include:

- **Imaging-based HTS:** Identifies antibacterial agents based on film formation ability or using reporters of antibacterial activity (e.g., adenylate kinase release upon cell lysis) [44].
- **Fluorescence anisotropy:** Screens compounds interacting with cell membrane lipids, enabling identification of agents targeting lipid II and interacting proteins (PBP1b, FtsW, and MurJ) [44].
- **Virulence-targeting HTS:** Screens for inhibitors of quorum-sensing and virulence factors without directly killing bacteria, potentially reducing selective pressure for resistance development [44].



Diagram 1: HTS Approaches for NP Library Screening. This workflow illustrates the three main strategies for screening natural product libraries, with their respective advantages and limitations.

Experimental Protocols and Methodologies

CT-HTS Protocol: Bacterial Growth Inhibition Screening

Objective: Identify natural product extracts that inhibit growth of pathogenic bacterial strains.

Materials and Reagents:

- Bacterial strains (e.g., ESKAPE pathogens: *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Pseudomonas aeruginosa*, *Enterobacter* species) [44]
- 384-well or 1536-well microtiter plates
- Natural product library (extracts or purified compounds in DMSO)
- Liquid handling robotics
- Multichannel pipettes or dispensers
- Culture media (appropriate for bacterial strains)

- Microplate spectrophotometer or fluorimeter

Procedure:

- **Assay Plate Preparation:**
  - Transfer 20-50 nL of natural product extracts from stock plates to assay plates using acoustic dispensing or pin tools [47].
  - Use 384-well or 1536-well formats with final DMSO concentration ≤1% to maintain cell viability.

- **Cell Seeding and Compound Exposure:**

  - Prepare bacterial inoculum in mid-log phase (OD600 ≈ 0.5) in appropriate culture media.
  - Dilute bacterial culture to approximately 5×10^5 CFU/mL in fresh media.
  - Dispense 50 µL bacterial suspension into each well of assay plate using liquid dispenser.
  - Include controls: media-only (background), DMSO-only (negative control), reference antibiotic (positive control).

- **Incubation and Signal Detection:**

  - Incubate plates at 37°C with appropriate humidity for 16-24 hours.
  - Measure bacterial growth by optical density (OD600) or using resazurin-based viability dyes.
  - For fluorescence-based readouts: Add resazurin (0.02 mg/mL final concentration), incubate 2-4 hours, measure fluorescence (Ex560/Em590).

- **Data Analysis:**

  - Calculate percent inhibition relative to controls: % Inhibition = [(Negative Control - Test Sample)/(Negative Control - Background)] × 100
  - Apply quality control metrics (Z'-factor > 0.5 indicates robust assay) [47]
  - Select hits showing >70% inhibition for confirmation studies.

MT-HTS Protocol: Enzyme Inhibition Screening

Objective: Identify natural product extracts that inhibit specific bacterial enzyme targets (e.g., DNA gyrase, topoisomerase, transpeptidases).

Materials and Reagents:

- Purified enzyme target
- Enzyme substrate and cofactors
- Detection reagents (fluorogenic or chromogenic)
- 384-well low-volume microplates
- Assay buffer optimized for enzyme activity
- Natural product library (pre-clarified if crude extracts)

Procedure:

- **Assay Optimization:**
  - Determine enzyme KM for substrate and optimal enzyme concentration for linear reaction kinetics.
  - Establish signal-to-background ratio (>3:1) and Z'-factor (>0.5) in miniaturized format.

- **Screening Reaction Setup:**

  - Transfer 100 nL natural product extracts to assay plates.
  - Add 5 µL enzyme solution in assay buffer to all wells.
  - Pre-incubate enzyme with compounds for 15-30 minutes at room temperature.

- **Reaction Initiation and Detection:**

  - Initiate reaction by adding 5 µL substrate solution.
  - Incubate for appropriate reaction time (determined during optimization).
  - Stop reaction if necessary or measure continuous kinetic readings.
  - Detect product formation using appropriate method:
    - Fluorescence: Measure fluorescence intensity with appropriate filters
    - Absorbance: Measure absorbance change at specific wavelength
    - Luminescence: Add detection reagent and measure luminescent signal

- **Data Analysis:**

- Calculate percent enzyme inhibition: % Inhibition = [1 - (Test Signal - Background)/(Negative Control - Background)] × 100
- Apply hit selection criteria (typically >50% inhibition at screening concentration)
- Confirm dose-response for primary hits in secondary screening

Table 2: Key Research Reagent Solutions for NP HTS Campaigns

Reagent Category	Specific Examples	Function in HTS	Application Notes
Detection Systems	Resazurin, ATP-lite, GFP reporters	Cell viability and metabolic activity assessment	Resazurin preferred for bacterial screens due to linear range [46]
Reporters	Luciferase, β-lactamase, fluorescent proteins	Pathway-specific reporter gene assays	ATAD5-luciferase for genotoxicity screening [46]
Cellular Systems	ESKAPE pathogens, DT40 cell lines, primary cells	Physiological context for screening	DNA-repair-deficient DT40 lines for genotoxin screening [46]
Molecular Targets	Purified enzymes, protein-protein interactions	Target-specific screening	Fluorescence anisotropy for lipid II binding proteins [44]
Automation Tools	Liquid handlers, plate stackers, detectors	Enable high-throughput processing	Robotic systems can test >100,000 compounds daily [47]

HTS Workflow and Quality Control

The successful implementation of HTS for natural product libraries requires careful attention to workflow design and quality control measures throughout the process.



Diagram 2: HTS Workflow with Quality Control Gates. The screening process incorporates quality control checkpoints at each stage to ensure identification of high-quality hits from natural product libraries.

Critical Quality Control Parameters

Assay Robustness Metrics:

- **Z'-factor:** Comprehensive assay quality metric accounting for dynamic range and data variation; Z' > 0.5 indicates excellent assay suitable for HTS [47].

- **Signal-to-Background Ratio:** Minimum 3:1 ratio required for reliable hit identification.
- **Coefficient of Variation (CV):** <10% for intra-plate and inter-plate reproducibility.
- **Strictly Standardized Mean Difference (SSMD):** Recently proposed for improved assessment of data quality in HTS assays, particularly for screens with replicates [47].

Hit Selection Criteria:

- **For primary screens without replicates:** z-score method or SSMD, which capture data variability based on the assumption that every compound has the same variability as a negative reference [47].
- **For confirmatory screens with replicates:** t-statistic or SSMD that directly estimates variability for each compound without relying on strong assumptions [47].
- **Robust methods:** z-*score, SSMD*, B-score, and quantile-based methods address outlier sensitivity common in HTS experiments [47].

The strategic selection between cellular and molecular target approaches for HTS of natural product libraries represents a fundamental decision point in drug discovery. CT-HTS offers the advantage of physiological relevance and identification of cellularly active compounds, while MT-HTS provides mechanistic clarity and target engagement information. The emerging paradigm of mechanism-informed phenotypic screening bridges these approaches, offering both physiological context and pathway-specific information. As natural products continue to play a crucial role in addressing the antibiotic resistance crisis and other therapeutic challenges, the intelligent application of these HTS methodologies – coupled with robust quality control and hit validation protocols – will maximize the potential of exploring natural product chemical space for drug discovery research. Future directions will likely see increased integration of artificial intelligence, advanced bioinformatics, and innovative screening technologies to further enhance the efficiency and success of NP-based drug discovery campaigns.

The declining efficiency of purely target-based drug discovery has catalyzed a resurgence in phenotypic screening. However, the limited translatability of simple phenotypic observations has necessitated an evolution in strategy. This whitepaper details the paradigm of **Mechanism-Informed Phenotypic Drug Discovery (MIPDD)**, a hybrid approach that integrates the physiological relevance of phenotypic observation with molecular-level mechanistic insight. Framed within the critical context of exploring **Natural Product (NP) chemical space**, we demonstrate how MIPDD leverages the unique physicochemical properties of NPs to identify novel therapeutic leads. This technical guide provides a comprehensive overview of the conceptual foundation, experimental methodologies, and practical implementation of MIPDD, with a specific focus on its application in antiviral and anticancer drug discovery.

Modern drug discovery has been dominated by target-based approaches, but their high attrition rates have prompted a critical re-evaluation. Analysis of cancer drug origins reveals that while the majority of approved small-molecule drugs originated from target-based discovery, very few were discovered entirely by 'classical' phenotypic screening [48]. This highlights a fundamental challenge: traditional phenotypic screens, often reliant on nonspecific readouts like cytotoxicity, are insufficient for identifying drugs with novel, therapeutically translatable mechanisms [48].

This realization has spurred the development of **Mechanism-Informed Phenotypic Drug Discovery (MIPDD)**, defined as the use of phenotypic assays designed around specific molecular pathways and targets, employing disease-relevant cellular models [48]. MIPDD aims to determine the causal relationships between target inhibition and phenotypic effects, opening new avenues for understanding cancer biology and discovering drugs with optimal molecular mechanisms of action [48].

Concurrently, the exploration of **biologically relevant chemical space (BioReCS)** has identified natural products as occupants of unique regions not represented by synthetic medicinal chemistry compounds [49] [50] [24]. Their structural rigidity, lower aromaticity, and high degree of stereochemistry make them exceptional starting points for MIPDD campaigns, providing a strategic advantage in identifying first-in-class therapeutics [49].

Table 1: Comparison of Drug Discovery Approaches

Feature	Classical Phenotypic Screening	Target-Based Screening	Mechanism-Informed Phenotypic Screening (MIPDD)
Primary Focus	Observable phenotypic change without prior target knowledge [51]	Modulation of a predefined molecular target [51]	Phenotypic change informed by underlying molecular mechanisms [48]
Discovery Bias	Unbiased, allows novel target identification [51]	Hypothesis-driven, limited to known pathways [51]	Hypothesis-guided, informed by disease biology
Mechanism of Action	Often unknown initially, requires deconvolution [51]	Defined from the outset [51]	Informed by pathway context, facilitating deconvolution
Chemical Library Strategy	Diverse libraries, emphasis on natural products [52] [49]	Focused libraries for specific target classes	Biased diversity, leveraging NP chemical space for specific phenotypes [52]

The Conceptual Framework of MIPDD

Core Principles and Definitions

Mechanism-informed phenotypic screening represents a neoclassic strategy that merges the best attributes of phenotypic and target-based approaches. Its core principle is the use of **mechanistically defined cellular models for therapeutically translatable cancer phenotypes** [48]. This involves:

- **Pathway-Focused Phenotypes:** Moving beyond general cytotoxicity to assess phenotypes tied to specific molecular pathways, such as synthetic lethality in genetically defined cancers or specific virulence processes in antivirals [48] [53].
- **Model Relevance:** Employing physiologically relevant models, including cocultures, 3D organoids, and patient-derived cells, which better recapitulate the disease context [51].
- **Chemical Bias:** Leveraging chemical libraries, particularly those rich in natural products, that are pre-enriched for bioactivity and the ability to modulate complex phenotypes [52] [49].

The Role of Natural Product Chemical Space

Natural products are a critical component for populating MIPDD screening libraries. Computational analyses using tools like ChemGPS-NP have demonstrated that NPs and synthetic bioactive compounds "differ notably in coverage of chemical space" [49] [50]. Key characteristics of NP chemical space include:

- **Structural Rigidity and Complexity:** NPs are generally more structurally rigid and contain fewer aromatic rings than synthetic medicinal chemistry compounds, favoring high-affinity target binding [49].
- **Lead-Like Properties:** A significant proportion of NPs are "lead-like" and occupy regions of chemical space sparsely populated by synthetic compounds, indicating high potential for novel target discovery [49] [24].
- **Synergistic Potential:** NPs consist of multiple constituents that can act on a variety of targets, potentially culminating in additive or synergistic therapeutic effects, making them ideal for combination therapy discovery [52].

Table 2: Key Characteristics of Natural Products in Chemical Space

Property	Finding	Implication for MIPDD
Rule of Five Compliance	~60% of unique NPs have no R05 violations; NP-derived drugs are equally split between R05 compliant and violators [49]	NPs are a valuable source for both conventional oral drugs and "beyond Rule of 5" therapeutics
Chemical Space Coverage	NPs cover regions not represented by synthetic medicinal chemistry databases (e.g., WOMBAT) [49] [50]	Provides access to novel scaffolds and mechanisms not found in standard synthetic libraries
Structural Features	NPs are less flexible and contain fewer aromatic rings than synthetic medicinal chemistry compounds [49]	Favors binding to challenging target classes like protein-protein interactions

Experimental Methodologies and Protocols

A Phenomics Workflow for Antiviral Discovery

The following workflow, adapted from a study on human coronavirus 229E, exemplifies a modern MIPDD approach for identifying host-targeting antivirals [54].



Protocol Details:

- **Biological Model and Infection:** Use MRC-5 human lung fibroblasts seeded in multiwell plates. After overnight attachment, inoculate cells with CoV-229E at a suitable MOI for 1 hour to maximize viral entry. Remove virus-containing media before compound addition to model post-infection treatment [54].
- **Compound Treatment:** Add a diverse compound library, enriched with natural products, and incubate for 48 hours. This duration allows for full development of both viral infection and compound-induced phenotypic changes.

- **Modified Cell Painting Staining:** Fix cells and perform a multiplexed fluorescent staining. This protocol modifies the standard Cell Painting by replacing the live-cell mitochondrial stain (MitoTracker) with an antibody against the viral nucleoprotein (NP). This allows simultaneous detection of infection status and detailed morphological profiling [54].
- **High-Content Imaging and Analysis:** Image plates using a high-content microscope. An automated image analysis pipeline (e.g., in CellProfiler) is used to segment individual cells and classify them as infected or non-infected based on the NP antibody signal. Subsequently, extract 1,441 morphological features from the five Cell Painting channels for each cell [54].
- **Data Analysis and Hit Identification:** Use multivariate statistical methods like Principal Component Analysis (PCA) and Partial Least-Squares Discriminant Analysis (PLS-DA). Effective antiviral compounds are identified by their ability to shift the morphological profile of infected cells back towards the non-infected state in the multivariate model [54].

Mechanism-Informed Screening in Oncology

In cancer drug discovery, MIPDD moves beyond simple proliferation assays. Key strategies include [48]:

- **Synthetic Lethal Screens:** Using engineered human tumor cells to identify compounds that selectively kill cancer cells bearing specific mutations (e.g., RAS, p53) while sparing wild-type cells.
- **Phenotypic Screens for Pathway Modulation:** Employing reporter cell lines or specific biomarkers to track the modulation of a particular pathway of interest, even if the exact target is unknown. This represents a middle ground between purely phenotypic and fully target-based screening.
- **Delayed-Death Phenotyping:** In antimalarial discovery, this approach identifies compounds with a "delayed death" phenotype, where parasite inhibition occurs in the second lifecycle generation, indicating a potential mechanism involving the apicoplast organelle [53].

The Scientist's Toolkit: Essential Research Reagents and Solutions

Successful implementation of MIPDD relies on a suite of specialized reagents and tools. The following table details key components for establishing these assays.

Table 3: Essential Research Reagents for MIPDD Assays

Reagent / Solution	Function / Purpose	Application Example
Cell Painting Dye Set	Multiplexed fluorescent staining for capturing a wide spectrum of morphological features [54]	General phenomic profiling; antiviral phenomics [54]
Anti-Viral Nucleoprotein Antibody	Specific detection of virus-infected cells at a single-cell level within a phenotypic assay [54]	Classifying infection status in host-cell morphological profiling [54]
3D Organoid / Spheroid Cultures	Physiologically relevant models that better mimic tissue architecture and disease context [51]	Oncology screening using more predictive in vitro models [48] [51]
iPSC-Derived Cell Models	Patient-specific disease modeling and drug screening for complex diseases [51]	Neurological disease modeling, personalized medicine applications
ChemGPS-NP / Scaffold Hunter	Computational tools for navigating and analyzing natural product chemical space [24]	Guiding the selection of NP-enriched screening libraries [49] [24]
High-Content Imaging System	Automated microscopy and image analysis for quantitative multiparametric data extraction	Essential for all image-based phenotypic screening workflows [51] [54]

Integration with Virulence-Targeting Screens

The MIPDD approach is highly applicable to antimicrobial discovery, particularly in targeting virulence mechanisms. This shifts the focus from essential pathogen viability to disarming its ability to cause disease, a strategy that may impose less selective pressure for resistance.

In antimalarial research, this has translated to developing robust phenotypic screens against diverse lifecycle stages beyond the symptomatic asexual blood stage, such as exoerythrocytic stages and transmission-blocking gametocytes [53]. The core logic of this approach is mapped below:



This strategy has been successfully implemented to discover compounds like the spiroindolone KAE609, which targets P-type cation-transporter ATPase4 (PfATP4) and demonstrates rapid parasiticidal activity [53].

Mechanism-Informed Phenotypic Drug Discovery represents a powerful and necessary evolution in the search for novel therapeutics. By integrating the physiological relevance of phenotypic observation with growing molecular understanding of disease pathways, MIPDD increases the probability of identifying high-quality leads with novel mechanisms of action. The strategic integration of this approach with the systematic exploration of natural product chemical space creates a synergistic partnership. NPs provide a source of unique, pre-validated scaffolds that populate biologically relevant but otherwise sparsely occupied regions of chemical space, while MIPDD offers a sophisticated framework to effectively probe and decode their complex biological activities. As technological advances in high-content imaging, automated image analysis, and biologically complex model systems continue, the implementation and success of MIPDD are poised to expand, firmly establishing its role in the future of drug discovery.

The relentless pursuit of new therapeutic compounds has driven researchers to delve into the vast chemical space of natural products, which have served as a cornerstone for drug development for decades. Plant natural products, or specialized metabolites, play a vital role in this endeavor, with many clinically important drugs such as the anticancer agents topotecan (derived from camptothecin) and etoposide (derived from podophyllotoxin) originating from plant sources [55]. Historically, the discovery of these compounds relied on bioactivity-guided fractionation approaches, which are increasingly hampered by the high rate of compound re-discovery [56]. The natural products discovery field has therefore begun a decisive shift away from these traditional methods toward strategies that capitalize on large-scale -omics technologies [56].

This transformation is powered by the integration of genomics and metabolomics, which provides a comprehensive framework for elucidating biosynthetic pathways. Genomics reveals the blueprint of an organism's biosynthetic potential, while metabolomics captures the chemical expression of that potential under specific conditions [56]. The convergence of these disciplines generates vast datasets, and the application of advanced computational tools, machine learning, and data analytics has become crucial for processing and interpreting this information to uncover intricate regulatory networks and identify key components of biosynthetic pathways [55]. This in-depth technical guide explores how the power of genomics and metabolomics is being harnessed to unlock biosynthetic pathways, framing these advancements within the critical context of exploring natural product chemical space for modern drug discovery research.

Genomic Foundations: Decoding Biosynthetic Blueprints

Genomics provides the foundational blueprint for biosynthetic pathway discovery by enabling the identification and annotation of Biosynthetic Gene Clusters (BGCs)—genomic loci that co-localize genes encoding the enzymes responsible for producing a specialized metabolite. The first step in genomic exploration involves obtaining high-quality sequence data. While Illumina next-generation sequencing (NGS) offers high fidelity and low cost, its short reads can result in fragmented assemblies. Advanced single-molecule sequencing technologies like Pacific Biosciences (PacBio) and Oxford Nanopore generate longer reads, which are invaluable for assembling complete BGCs, despite their higher per-base error rates [56].

Once a genome is sequenced and assembled, the critical task of BGC identification begins. This is accomplished using sophisticated bioinformatic algorithms that scan genomic data for signature biosynthetic genes. Several tools have been developed for this purpose, each with distinct strengths and applications [56].

Table 1: Key Computational Tools for Biosynthetic Gene Cluster Identification

Tool Name	Primary Application	Methodology Overview
antiSMASH [56]	Bacteria, Fungi, Plants	Uses a library of profile Hidden Markov Models (pHMMs) to detect >50 classes of BGCs; widely considered a industry standard.
plantiSMASH [57]	Plants	A specialized derivative of antiSMASH using modified rules tailored to plant genomes.
PRISM [56]	Bacteria & Fungi	Employs pHMMs and machine learning to predict BGCs and the chemical structures of their products.
SMURF [56]	Fungi	pHMM-based tool designed for fungal genome mining.
CO-OCCUR [56]	Phylogenetically diverse fungi	Identifies accessory biosynthetic genes through frequency and co-occurrence analysis around core genes, complementing other tools.

These tools function by identifying core biosynthetic genes, such as those for polyketide synthases (PKSs) and non-ribosomal peptide synthetases (NRPSs), and then scanning the surrounding genomic region for additional genes encoding tailoring enzymes, transporters, and regulators [56]. The output is a map of an organism's biosynthetic potential, which often reveals a surprising abundance of uncharacterized BGCs, even in well-studied organisms [56]. This highlights the vastness of unexplored natural product chemical space and provides a genetic starting point for discovery.

Experimental Protocol: Genome Mining for BGC Identification

1. DNA Extraction and Sequencing:

- Isolate high-molecular-weight genomic DNA from the target organism using a standardized protocol (e.g., CTAB method for plants).
- Assess DNA quality and integrity using spectrophotometry (e.g., Nanodrop) and gel electrophoresis.
- Proceed with whole-genome sequencing using a platform that suits project goals. A hybrid approach using Illumina for high accuracy and PacBio/Oxford Nanopore for long-read scaffolding is often optimal for complete BGC assembly [56].

2. Genome Assembly and Annotation:

- Assemble raw sequencing reads into contigs using assemblers like SPAdes or Canu.
- Scaffold contigs into chromosomes where possible using linkage information.
- Annotate the assembled genome using pipelines such as the NCBI Eukaryotic Genome Annotation Pipeline or BRAKER, which predict gene models and assign putative functions.

3. BGC Identification and Analysis:

- Input the annotated genome file (in GenBank or FASTA format) into a BGC prediction tool such as antiSMASH.
- Analyze the output to identify the types and genomic locations of BGCs.
- Manually inspect the predicted BGCs, paying close attention to the domain architecture of core biosynthetic enzymes (e.g., PKS and NRPS domains) to hypothesize about the potential chemical scaffold of the metabolite product.



Diagram 1: Genomic workflow for BGC identification.

Metabolomic Technologies: Capturing Chemical Phenotypes

Metabolomics delivers the complementary chemical phenotype by providing a snapshot of the entire set of metabolites in a biological system at a given time. In the context of natural product research, it focuses on the secondary metabolites actually produced by the organism, offering a direct readout of biosynthetic pathway activity [56]. The workflow is typically divided into pre-analytical, analytical, and post-analytical stages, with careful standardization at each phase being critical for robust and reproducible results [58].

The pre-analytical phase involves sample collection, handling, and storage. Factors such as collection tubes, centrifugation steps, freeze-thaw cycles, and storage conditions must be standardized using Standardized Operating Procedures (SOPs) to minimize variability and ensure data accurately reflects endogenous metabolite levels [58]. For MS-based metabolomics, sample preparation involves extracting metabolites from proteins and other matrix components, a process that should be automated where possible to reduce human error [58].

The analytical heart of modern metabolomics is mass spectrometry (MS), often coupled with separation techniques like liquid chromatography (LC) or gas chromatography (GC). MS is favored for its high sensitivity and specificity, allowing for the measurement of thousands of metabolites in small sample volumes [58]. Two primary analytical approaches are employed:

- **Targeted Metabolomics:** This approach focuses on the precise quantification of a predefined set of metabolites. A prominent example is the use of the Biocrates AbsoluteIDQ p180 Kit, which enables the quantification of a panel of 188 metabolites, including amino acids, biogenic amines, acylcarnitines, and lipids [59]. This method is highly reproducible and ideal for biomarker validation.
- **Untargeted Metabolomics:** This hypothesis-generating approach aims to comprehensively profile as many metabolites as possible in a sample without bias. It is particularly powerful for discovering novel metabolites and generating new hypotheses about their roles in disease or drug mechanisms [58].

Table 2: Core Metabolomics Instrumentation and Reagents

Category / Item	Function / Description	Application in Pathway Elucidation
LC-MS / GC-MS System	Separates (chromatography) and detects (mass spectrometry) complex metabolite mixtures.	Workhorse platform for profiling metabolite extracts; enables detection of thousands of features.
Biocrates AbsoluteIDQ p180 Kit [59]	Standardized kit for targeted quantification of 188 plasma metabolites.	Provides highly reproducible quantitative data for defined metabolite classes; used for biomarker studies.

Category / Item	Function / Description	Application in Pathway Elucidation
High-Resolution MS (Orbitrap, FTICR-MS) [57]	Mass spectrometers with very high mass accuracy and resolution.	Critical for determining precise molecular formulae of unknown metabolites from untargeted data.
Nuclear Magnetic Resonance (NMR) [58]	Non-destructive, quantitative analytical technique.	Useful for structural elucidation and quantifying abundant metabolites; complements MS data.
Ion Mobility Spectrometry [58]	Separates ions based on their size, shape, and charge.	Adds an additional separation dimension, helping to resolve structurally similar isomers.

Experimental Protocol: Untargeted Metabolomics for Pathway Discovery

1. Sample Preparation and Extraction:

- Flash-freeze biological tissue (e.g., plant root) in liquid nitrogen and homogenize to a fine powder.
- Weigh a precise amount of powdered tissue and add a cold extraction solvent (e.g., methanol:water:chloroform in a 2.5:1:1 ratio) to comprehensively extract metabolites of diverse polarities.
- Vortex vigorously, sonicate in a cold water bath, and centrifuge to pellet insoluble debris.
- Transfer the supernatant (containing metabolites) to a new vial and dry under a gentle stream of nitrogen gas.
- Reconstitute the dried extract in a solvent compatible with the LC-MS system (e.g., 100 µL of 80% methanol).

2. LC-MS Data Acquisition:

- Inject the reconstituted sample onto a reversed-phase UHPLC column (e.g., C18) coupled to a high-resolution mass spectrometer (e.g., Orbitrap).
- Use a gradient elution method (e.g., water to acetonitrile, both with 0.1% formic acid) to separate metabolites.
- Acquire data in both positive and negative ionization modes to maximize metabolite coverage. Data-Dependent Acquisition (DDA) mode can be used to fragment the most abundant ions and collect MS/MS spectra for structural annotation.

3. Data Pre-processing and Metabolite Annotation:

- Process raw LC-MS data using software (e.g., XCMS, MS-DIAL) for peak picking, alignment, and integration to create a feature table (containing mass-to-charge ratio, retention time, and intensity for each detected ion).
- Annotate metabolite features by querying their accurate mass and MS/MS spectra against public databases such as LOTUS, a comprehensive resource of natural products [57].
- Statistical analysis (e.g., PCA, OPLS-DA) is then performed to identify metabolites that differ significantly between sample groups (e.g., induced vs. control tissues).



Diagram 2: Metabolomics workflow for chemical phenotyping.

Integrated Multi-Omics: From Correlation to Causation in Pathway Reconstruction

The true power of -omics approaches is realized when genomics and metabolomics datasets are integrated, moving beyond correlation to predict causal relationships within biosynthetic pathways. This integration allows researchers to simultaneously identify expressed secondary metabolites and link them to their biosynthetic machinery [56]. One of the primary strategies for integration is **co-expression analysis**, which identifies genes and metabolites that show correlated abundance patterns across different samples (e.g., different tissues, treatments, or time points) [57]. A gene whose expression profile closely mirrors the accumulation of a specific metabolite is a strong candidate for encoding an enzyme involved in that metabolite's biosynthesis.

Cutting-edge computational tools are now automating this integration process. A leading example is **MEANtools**, a systematic and unsupervised computational workflow that predicts candidate metabolic pathways *de novo* [57]. MEANtools integrates mass features from metabolomics data and transcripts from transcriptomics data. It uses a mutual rank-based correlation method to identify highly correlated metabolite-transcript pairs and then leverages known biochemical reaction rules from databases like RetroRules to assess whether correlated transcripts encode enzymes that can catalyze reactions between correlated metabolites [57]. This allows the pipeline to construct putative biosynthetic pathways from the integrated data, generating testable hypotheses.

Another powerful integration method is the **Genome-Wide Association Study (GWAS) of metabolites**, known as **mQTL (metabolite Quantitative Trait Loci) mapping**. This approach identifies genomic regions associated with natural variation in metabolite levels [59]. In a study on pigs, mQTL mapping successfully identified 97 genomic loci associated with the levels of 126 metabolites, directly linking genetic variants to metabolic phenotypes and uncovering genes involved in specific metabolic pathways [59].

Experimental Protocol: Integrated Multi-Omics with MEANtools

1. Paired Sample Collection and Multi-Omics Data Generation:

- Collect a series of samples from the same organism under different conditions (e.g., different tissues, time points post-elicitation, different light regimes). This provides the variation necessary for correlation analysis.
- For each sample in the series, simultaneously generate:
  - **Transcriptomic Data:** RNA sequencing (RNA-seq) to obtain gene expression counts.
  - **Metabolomic Data:** Untargeted LC-MS to obtain a list of mass features (mass-to-charge ratio and intensity).

2. Data Integration and Pathway Prediction with MEANtools:

- Format the input data as required by MEANtools: a metabolomic feature table and a transcriptomic count matrix.
- MEANtools will then [57]:
  - **Correlate:** Calculate the mutual rank correlation between every mass feature and every transcript across the sample set.
  - **Annotate:** Query the LOTUS database to find potential structural matches for the mass features.
  - **Connect:** Access the RetroRules database to retrieve known enzymatic reaction rules and associated Enzyme Commission (EC) numbers.
  - **Predict:** Identify trios where a transcript (encoding an enzyme with an EC number) is highly correlated with two mass features whose chemical structures are logically connected by that enzyme's reaction rule.
- The output is a set of predicted metabolic pathways, where nodes are mass features and edges are enzymatic reactions supported by correlated transcripts.

3. Hypothesis Testing and Validation:

- Prioritize the predicted pathways based on the strength of correlation and biochemical plausibility.
- Clone the candidate genes and heterologously express them in a system like *E. coli* or yeast.
- Test the enzymatic activity of the purified recombinant proteins against the predicted substrate metabolites.
- Use CRISPR-Cas9 or RNAi to knock out the candidate gene in the host organism and confirm the loss of the metabolite product.



Diagram 3: Multi-omics integration for pathway prediction.

Applications in Drug Discovery: From Target Identification to Precision Medicine

The application of integrated genomics and metabolomics is revolutionizing the drug discovery and development pipeline, offering powerful approaches from initial target identification to the realization of precision medicine.

In the **target identification** phase, metabolomics can reveal specific metabolic pathways that are altered in disease states. For instance, an unbiased discovery metabolomics approach can characterize the molecular heterogeneity of complex diseases like type 2 diabetes mellitus (T2DM), identifying distinct patient subtypes with different underlying metabolic disturbances [58]. This can pinpoint specific enzymes or metabolic regulators as novel therapeutic targets. Genomics complements this by identifying genetic variants associated with both disease risk and metabolite levels (mQTLs), providing orthogonal evidence for a target's validity and highlighting potential mechanisms of action [59].

In **natural product-based drug discovery**, the integration of -omics technologies directly addresses the challenge of efficiently linking bioactive compounds to their BGCs. By combining metabolomic profiling with genomic mining, researchers can prioritize uncharacterized BGCs that are active under specific conditions and associated with the production of novel chemical scaffolds [55] [56]. This strategy efficiently guides the isolation and characterization of new lead compounds from the vast "dark matter" of uncharacterized natural product space.

Finally, metabolomics plays a crucial role in **biomarker discovery** for patient stratification and treatment response prediction—a cornerstone of precision medicine. The identification of genetically influenced metabolites (GIMs) provides a powerful class of biomarkers. As demonstrated in pig models, these stable molecular phenotypes are highly heritable and can be used to dissect complex traits [59]. In humans, metabolomic signatures can define individual "metabotypes," enabling the stratification of patient populations to predict drug response and optimize therapeutic outcomes [58]. This ensures that the right natural product-derived drug or other therapy is delivered to the right patient.

Building and Curating High-Quality Natural Product Libraries for Screening

Natural Products (NPs) represent an indispensable source of chemical diversity for drug discovery, providing greater structural variety than standard synthetic chemistry and unique opportunities for identifying novel low molecular weight lead compounds [60]. A detailed analysis of FDA-approved drugs between 1981 and 2019 reveals that natural products, their direct derivatives, or synthetic drugs incorporating pharmacophoric groups of active secondary metabolites constitute approximately 56.1% of all approved drugs, with particularly significant contributions to anticancer (69.6%), antibacterial (58%), and antiviral (37.6%) therapies [60]. This remarkable success stems from the evolutionary optimization of NPs for biological interaction, often resulting in complex three-dimensional structures rich in sp^3-hybridized carbon atoms and stereocenters that cover chemical space regions largely inaccessible to purely synthetic compounds [61]. The screening of natural product libraries therefore offers significant advantages for finding novel therapeutic agents, but realizing this potential requires meticulous attention to library construction, curation, and management. This technical guide outlines comprehensive methodologies for building and maintaining high-quality NP libraries specifically framed within the context of systematically exploring natural product chemical space for drug discovery research.

Composition and Sourcing of Natural Product Libraries

Library Components and Structural Considerations

Modern natural product libraries encompass diverse physical forms and structural types, each with distinct advantages for screening campaigns. These libraries typically include crude extracts from plants, marine organisms, and microorganisms; prefractionated extracts that reduce complexity while preserving synergistic interactions; and pure natural product compounds [62] [60]. A particularly promising approach involves the deconstruction of NPs into fragments and their recombination into unprecedented pseudo-natural product frameworks, which retain NP-inspired features while extending into novel structural and functional space [63]. When designing a library, careful consideration must be given to the balance between complexity and screening compatibility. Crude extracts offer the fullest representation of natural chemical diversity but may present challenges in dereplication and identification of active constituents, while pure compounds provide immediate structural information but require significant upfront investment in isolation and characterization.

Table 1: Composition of Representative Natural Product Libraries

Library/Source	Type	Scale/Size	Key Characteristics	References
COCONUT 2.0	Database (Virtual)	695,133 distinct structures	Comprehensive collection of open natural products; extensive chemical space coverage	[61]
LANaPDB	Database (Virtual)	13,578 compounds	Focus on Latin American biodiversity; non-duplicate natural products	[61]
MEDINA	Physical Library	>200,000 extracts	Microbial-derived natural products from diverse global environments	[62]
NCI Natural Products Repository	Physical Library	230,000+ crude extracts; 400+ purified compounds	One of world's most comprehensive collections; includes traditional Chinese medicine extracts	[62]
University of Michigan Natural Products Discovery Core	Physical Library	45,000+ natural product extracts (NPEs)	Metadata-enabled with chemical and genetic profiles; HTS-formatted	[62]
NatureBank, Griffith University	Physical Library	18,000+ extracts; 90,000+ fractions; 100+ pure compounds	Australian biodiversity focus; lead-like enhanced libraries	[62]

Fragment Libraries and Chemical Space Analysis

Fragment-based approaches have emerged as powerful tools for efficiently exploring NP chemical space. Recent research has generated comprehensive fragment libraries from large NP databases, with 2,583,127 fragments derived from COCONUT and 74,193 fragments from LANaPDB [61]. These fragments, typically obtained using methods like the Retrosynthetic Combinatorial Analysis Procedure (RECAP), adhere to the "rule of three" (RO3) for fragment-based drug design: molecular weight ≤300 Da, rotatable bonds ≤3, topological polar surface area ≤60 $Å^2$, LogP ≤3, hydrogen-bond acceptors ≤3, and hydrogen-bond donors ≤3 [61]. Analysis reveals that only 1.5% of COCONUT fragments and 2.5% of LANaPDB fragments fulfill all RO3 criteria, highlighting both the structural complexity of natural products and the need for careful curation to optimize fragment libraries for screening [61]. When compared to synthetic fragment libraries, NP-derived fragments occupy distinct regions of chemical space, often exhibiting greater stereochemical complexity and scaffold diversity that can provide unique starting points for drug discovery programs focused on challenging therapeutic targets.

Table 2: Performance Metrics of Natural Product Fragment Libraries

Library	Initial Fragments	Fragments After Standardization	Fragments Fulfilling RO3	Percentage Fulfilling RO3
COCONUT	2,583,127	2,583,127	38,747	1.5%
LANaPDB	74,193	74,193	1,832	2.5%
CRAFT	1,214	1,202	176	14.6%
Enamine	12,505	12,496	8,386	67.1%
ChemDiv	74,721	72,356	16,723	23.1%

Regulatory Framework and Ethical Sourcing

The access and use of biological resources for natural product library development must comply with international and national regulations governing genetic resources and associated traditional knowledge. The United Nations 1992 Convention on Biological Diversity (CBD) and its supplementary Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits (ABS) establish the legal framework requiring mutually agreed terms between source countries and users [60]. These agreements typically include provisions for prior informed consent, benefit-sharing arrangements, and respect for the rights of indigenous communities and traditional knowledge holders. In Brazil, for example, Law 13.123/15 and the National System for the Management of Genetic Resources and Associated Traditional Knowledge (SisGen) regulate research and development involving Brazilian biodiversity, requiring registration of activities and establishing that foreign researchers must collaborate with Brazilian institutions [60]. Similar frameworks exist in other biodiverse countries, creating a complex regulatory landscape that necessitates careful legal assessment during library planning. Negotiating appropriate access and benefit-sharing agreements can be time-consuming but represents an essential ethical and legal foundation for sustainable natural product research that respects national sovereignty and contributes to conservation efforts.

Library Construction Methodologies

Sample Preparation and Extraction Protocols

The construction of high-quality natural product libraries begins with meticulous sample preparation, which directly influences phytochemical composition and screening outcomes. For plant-based libraries, critical considerations include proper taxonomic identification by qualified botanists with voucher specimen deposition in recognized herbaria, optimal collection timing that accounts for seasonal and diurnal variation in metabolite production, and appropriate preservation methods such as freeze-drying or controlled drying to prevent degradation [60]. Extraction strategies should be designed to maximize chemical diversity while maintaining compatibility with screening platforms, typically employing a sequential approach with solvents of increasing polarity (e.g., hexane, dichloromethane, ethyl acetate, methanol, water) [60]. For microorganism-derived libraries, specialized isolation media and cultivation conditions are essential to access the full biosynthetic potential, as standard laboratory conditions may not activate silent gene clusters responsible for producing many bioactive metabolites. Advanced techniques such as co-cultivation, OSMAC (one strain many compounds) approaches, and genomic mining can significantly enhance chemical diversity from microbial sources.

Library Formatting and Quality Control

Standardized formatting and rigorous quality control are essential for generating reproducible screening data from natural product libraries. Most modern libraries are formatted in 96-well or 384-well plates compatible with high-throughput screening robotics, with typical concentrations of 1-10 mg/mL for extracts and 1-10 mM for pure compounds in dimethyl sulfoxide (DMSO) [62]. Quality control measures should include chemical profiling using HPLC-UV/PDA/ELSD and/or LC-MS to verify composition and stability, determination of dry weight for extract normalization, and assessment of potential interferants such as tannins, pigments, or non-specific binding compounds that may produce false positives in certain assay formats [60]. For pure compound libraries, purity assessment (typically ≥95% by HPLC) and structural confirmation (via NMR and HRMS) are essential, along with curation of associated metadata including natural source, isolation method, physicochemical properties, and known biological activities [62]. The ChromaDex Natural Compound Libraries exemplify this approach, offering extensively characterized fractions that preserve cross-fraction synergy while providing detailed compositional data [62].

Screening Methodologies for Natural Product Libraries

Affinity Selection Mass Spectrometry (AS-MS) Workflow

Affinity selection mass spectrometry has emerged as a powerful label-free biophysical method for identifying ligands from complex natural product libraries against various biological targets, including soluble proteins, membrane proteins, nucleic acids, and nucleic acid-protein complexes [64]. AS-MS interrogates non-covalent target-ligand complexes in a non-functional assay, simultaneously identifying multiple ligands with different mechanisms of action, including orthosteric and allosteric binders [64]. The methodology involves four critical stages: (1) static incubation of the target with the natural product library, typically with the target in molar excess to avoid competition effects; (2) separation of target-ligand complexes from unbound components; (3) dissociation of ligands from the complexes; and (4) identification of ligands by mass spectrometry [64]. This

approach offers significant advantages over traditional bioactivity-guided fractionation by reducing false positives and avoiding activity loss through multiple fractionation steps.



Solution-Based AS-MS: Ultrafiltration Methodology

Ultrafiltration represents a particularly effective solution-based AS-MS technique that separates target-ligand complexes from unbound molecules based on size exclusion through specialized membranes with controlled porosity [64]. In a typical implementation, the target protein is incubated with the natural product library at low micromolar concentrations optimal for detecting high-affinity ligands with specific binding interactions. Following equilibrium establishment, ultrafiltration membranes with molecular weight cutoffs between 500-500,000 Da retain the larger ligand-protein complexes while allowing unbound molecules to pass through [64]. Ligands are subsequently dissociated using denaturing conditions such as methanol or acetonitrile with volatile organic acids (e.g., formic acid), maintaining compatibility with subsequent LC-MS analysis. This approach has been successfully applied to identify bioactive natural products, such as the discovery of botulin, lanosterol, and quercetin as 5-lipoxygenase ligands from Inonotus obliquus extracts [64]. The methodology offers advantages in maintaining native protein conformation during screening and applicability to diverse target classes, though careful optimization of filtration conditions is necessary to prevent membrane fouling or non-specific binding.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Key Research Reagent Solutions for Natural Product Library Screening

Reagent/Resource	Function/Application	Implementation Example	Considerations	
Ultrafiltration Membranes	Separation of target-ligand complexes from unbound molecules	Molecular weight cutoffs 500-500,000 Da for protein-ligand complex retention	Pore size uniformity, minimal non-specific binding, chemical compatibility	[64]
Immobilization Supports	Ligand fishing with immobilized targets	Magnetic microbeads (MagMASS), chromatographic resins	Retention of target activity after immobilization, ligand accessibility	[64]

Reagent/Resource	Function/Application	Implementation Example	Considerations	
Mass Spectrometry Platforms	Detection and identification of bound ligands	LC-MS systems with high resolution and mass accuracy	Sensitivity, dynamic range, compatibility with dissociation solvents	[64]
Bioaffinity Chromatography Systems	Zonal or frontal chromatography for ligand disclosure	Solid-supported proteins for "functional chromatography"	Retention time precision, breakthrough curve analysis	[64]
Natural Product Libraries	Sources of diverse 3D molecular features for screening	Commercial sources (e.g., ChromaDex, MicroSource) or custom collections	Chemical diversity, annotation quality, regulatory compliance	[62] [60]
Fragment Libraries	Fragment-based drug design starting points	COCONUT, LANaPDB, CRAFT, commercial vendors	Rule of three compliance, synthetic accessibility, structural diversity	[61]

Building and curating high-quality natural product libraries for drug discovery represents both a significant technical challenge and a substantial opportunity to access unique chemical space with proven therapeutic relevance. Success in this endeavor requires integrated expertise spanning taxonomy, natural product chemistry, analytical methodology, screening technology, and regulatory compliance. The future of natural product-based drug discovery will increasingly leverage computational approaches for virtual screening, chemical space analysis, and bioactivity prediction, while advanced screening technologies like AS-MS enhance the efficiency of ligand identification from complex mixtures. Furthermore, innovative strategies such as pseudo-natural product design, which recombines biosynthetically unrelated NP fragments into novel scaffolds, promise to extend accessible chemical space beyond naturally occurring structures while retaining desirable NP-like properties [63]. By applying the systematic approaches outlined in this guide—from ethical sourcing and standardized library construction to advanced screening methodologies—research institutions and pharmaceutical companies can fully leverage the remarkable structural and functional diversity of natural products to address unmet medical needs through novel therapeutic agents.

The exploration of **natural products (NPs)** represents a cornerstone of drug discovery, with a significant portion of modern small-molecule drugs originating from or being inspired by natural compounds [65]. However, the traditional drug discovery pipeline is notoriously protracted, often exceeding **12 years** and costing more than **$1.8 billion USD** on average [66]. The vastness of biologically relevant chemical space, coupled with challenges in sourcing and characterizing NPs, necessitates more efficient discovery approaches [49].

In silico technologies have emerged as powerful tools to address these challenges. **Computer-Aided Drug Discovery (CADD)** leverages computational power to streamline hit identification and optimization, dramatically reducing the time and cost associated with early-stage discovery [67] [66]. This technical guide details the core methodologies of **virtual screening (VS)** and **molecular dynamics (MD)** simulations, framing them within the strategic imperative to efficiently navigate the unique and biologically relevant chemical space occupied by natural products [68] [49].

Virtual Screening for Navigating Natural Product Chemical Space

Virtual screening is a computational technique for identifying potential hit compounds from vast digital libraries. Its application is particularly valuable for exploring NPs, which are often structurally complex and sparsely represented in conventional screening collections [49].

The Rationale for Natural Products in Drug Discovery

Natural products are pre-validated by nature, possessing evolutionary optimization for interaction with biological macromolecules. Analyses reveal that NPs occupy **distinct regions of chemical space** compared to synthetic medicinal chemistry compounds. They are typically more structurally rigid and possess a lower degree of aromaticity, offering access to novel scaffolds that can circumvent pre-existing intellectual property and overcome the limitations of conventional chemical libraries [49]. It is estimated that about **two-thirds** of modern small-molecule drugs are related to natural compounds [65].

Key Virtual Screening Methodologies

Two primary VS approaches are employed, often in tandem, for efficient hit identification.

- **Ligand-Based Virtual Screening (LBVS):** This method is used when the 3D structure of the target protein is unknown but a set of active ligands is available. It relies on comparing molecular descriptors or fingerprints to identify new compounds with similar properties [69]. Advanced LBVS platforms, such as the BIOPTIC B1 system, utilize transformer-based models trained on massive molecular datasets (e.g., **160 million molecules**) to create potency-aware embeddings. These enable ultra-high-throughput screening of multi-billion compound libraries

in mere minutes, demonstrating success in prospective campaigns for targets like **LRRK2** for Parkinson's disease, yielding novel sub-micromolar binders (Kd = **110 nM**) [69].

- **Structure-Based Virtual Screening (SBVS):** This approach requires a 3D structure of the biological target, typically derived from X-ray crystallography, NMR, or homology modeling. The primary technique used in SBVS is **molecular docking**, which predicts the preferred orientation and binding affinity (scoring) of a small molecule within a target's binding site [67] [65]. Docking simulations help understand molecular-level interactions and are crucial for identifying hit leads from natural product libraries [67].

Table 1: Key Databases for Virtual Screening of Natural Products

Database Name	Description	Content Focus	Utility in VS
ZINC15 [70]	Curated database of commercially available compounds.	100+ million compounds in ready-to-dock, 3D formats.	Primary source for purchasable screening compounds.
ChEMBL [70] [71]	Manually curated database of bioactive molecules.	Drug-like molecules with bioactivity data.	Ligand-based screening and model training.
PubChem [70]	NCBI repository of chemical molecules and bioactivities.	Massive collection of compounds and bioassay results.	Similarity searching (2D/3D) and bioactivity data.
Dictionary of Natural Products (DNP) [70] [49]	Comprehensive and fully-edited NP database.	Over **170,000** compounds of natural origin.	Definitive source for natural product structures and data.
TCM Database [70]	Database on Traditional Chinese Medicine.	**~170,000** compounds with 2D/3D structural files.	Virtual screening of natural product libraries.

Integrated Virtual Screening Workflow

The following diagram illustrates a synergistic VS workflow that integrates both ligand-based and structure-based methods to efficiently mine natural product chemical space.



Molecular Dynamics for Hit Validation and Optimization

While molecular docking provides a static snapshot of binding, **Molecular Dynamics (MD)** simulations model the dynamic behavior of molecules over time, offering critical insights into stability, conformational changes, and binding mechanisms that are essential for hit-to-lead optimization [65].

Fundamentals of MD Simulations

MD simulations numerically integrate Newton's second law of motion for all atoms in a system, typically involving a protein-ligand complex solvated in water. This allows researchers to observe the time-dependent evolution of the molecular system [72]. Key parameters must be carefully set to

ensure simulation stability and accuracy:

- **Time Step:** The interval for calculating atomic movements. A value of **1-2 femtoseconds (fs)** is suitable for systems with light atoms (e.g., hydrogen), while **5 fs** may be adequate for metallic systems [72].
- **Ensemble:** The thermodynamic conditions of the simulation.
  - **NVE:** Constant Number of particles, Volume, and Energy (microcanonical). Preserves total energy [72].
  - **NVT:** Constant Number of particles, Volume, and Temperature (canonical). Uses thermostats (e.g., Langevin, Nosé-Hoover) to maintain temperature [72].
  - **NpT:** Constant Number of particles, Pressure, and Temperature (isothermal-isobaric). Uses barostats to control pressure [72].

MD Simulation Protocol for a Protein-Ligand Complex

A typical MD workflow for validating a natural product hit bound to its target protein involves the following steps [65] [72]:

- **System Preparation:** Obtain the 3D structure of the protein-ligand complex from docking studies. Add missing hydrogen atoms and assign protonation states at physiological pH.
- **Solvation and Ion Addition:** Place the complex in a simulation box (e.g., a cubic or rhombic dodecahedron box) filled with water molecules (e.g., TIP3P model). Add ions (e.g., Na^+, Cl^-) to neutralize the system's charge and mimic physiological ion concentration (~0.15 M).
- **Energy Minimization:** Perform a steepest descent or conjugate gradient minimization to relieve any steric clashes or structural strains introduced during the setup, converging until the maximum force is below a specified threshold (e.g., 1000 kJ/mol/nm).
- **System Equilibration:**
  - *NVT Ensemble:* Heat the system from 0 K to the target temperature (e.g., 310 K) over **100 ps**, using a thermostat (e.g., Langevin with a friction coefficient of 1 ps^{-1}).
  - *NpT Ensemble:* Adjust the box dimensions to achieve the correct density at the target temperature and pressure (1 bar) over **100 ps**, using a barostat (e.g., Berendsen).
- **Production Run:** Conduct an extended MD simulation in the NpT ensemble for a duration sufficient to capture the relevant biological processes (typically **50-500 nanoseconds** or more). The trajectory—containing atomic coordinates saved at regular intervals (e.g., every 10 ps)—is used for analysis.
- **Trajectory Analysis:** Analyze the saved trajectory to calculate:
  - **Root Mean Square Deviation (RMSD)** of the protein and ligand to assess structural stability.
  - **Root Mean Square Fluctuation (RMSF)** to identify flexible regions.
  - **Ligand-protein hydrogen bonding** occupancy and lifetime.
  - **Binding free energy** using methods like MM/PBSA or MM/GBSA.

Table 2: Critical MD Parameters and Recommended Settings for Stability

Parameter	Description	Recommended Settings	Rationale
Time Step [72]	Interval for numerical integration.	**1-2 fs** (systems with H-bonds); **5 fs** (metallic systems).	Prevents instability; ensures energy conservation.
Force Field [73]	Mathematical functions for atomic interactions.	**COMPASS II**, CHARMM, AMBER, OPLS.	Determines accuracy of interatomic forces.
Temperature Control [72]	Thermostat for NVT/NpT ensembles.	**Langevin**, Nosé-Hoover, Bussi.	Correctly samples the canonical ensemble.
Pressure Control [72]	Barostat for NpT ensemble.	Berendsen, Parrinello-Rahman.	Maintains correct system density.
Simulation Length [65]	Duration of production run.	**50 - 500 ns** (varies by system).	Captures relevant biological dynamics and stability.
Periodic Boundary Conditions (PBC) [72]	Mimics an infinite system.	Cubic or rhombic dodecahedron box.	Eliminates edge effects; uses a finite number of atoms.

MD Simulation Workflow

The following diagram outlines the sequential steps involved in setting up and running an MD simulation for a protein-ligand complex.



The Scientist's Toolkit: Essential Research Reagents and Software

Successful implementation of in silico drug discovery relies on a suite of specialized software tools and databases.

Table 3: Essential In Silico Research Tools for Natural Product Drug Discovery

Tool Category	Representative Examples	Key Function	Application in NP Discovery
Molecular Docking Software [67] [65]	AutoDock Vina, GOLD, Glide	Predicts binding pose and affinity of ligands.	Structure-based screening of NP libraries against therapeutic targets.
MD Simulation Engines [72] [73]	ASE (Atomic Simulation Environment), GROMACS, AMBER, NAMD	Performs molecular dynamics simulations.	Validating binding stability and mechanism of NP hits.
Cheminformatics & ML Libraries [71]	RDKit, Open Babel, scikit-learn	Handles chemical data and builds ML models.	Processing NP structures; building LBVS models with ECFP fingerprints.
Homology Modeling Tools [65]	MODELLER, SWISS-MODEL	Predicts 3D protein structures from amino acid sequences.	Generating target models for SBVS when experimental structures are unavailable.
Bioactivity Databases [70] [71]	ChEMBL, BindingDB, PubChem BioAssay	Provides experimental bioactivity data.	Training and validating machine learning models for target prediction.

The integration of **virtual screening** and **molecular dynamics** provides a powerful, synergistic framework for accelerating the discovery of bioactive hits from the vast and structurally diverse universe of natural products. By leveraging these in silico tools, researchers can efficiently navigate biologically relevant chemical space, prioritize the most promising NP-derived leads with novel scaffolds, and gain deep mechanistic insights into their interactions with therapeutic targets. This computational approach de-risks and informs subsequent experimental validation, paving the way for the development of new drugs inspired by nature's intricate chemistry. As these methodologies continue to advance, particularly with the integration of machine learning and increasing computational power, their role in unlocking the full potential of natural products for drug discovery will only become more profound.

Navigating the Pipeline: Overcoming Technical and Regulatory Hurdles

Technical Barriers in Screening, Isolation, and Characterization of Complex Mixtures

The exploration of **natural products** represents a cornerstone in drug discovery, offering access to a vast and structurally diverse **chemical space** that is largely untapped by synthetic compound libraries [49]. Natural product libraries are a source of diverse **3D molecular features** furnishing an array of biological functions and are resourceful in furnishing scaffolds for drug discovery research [64]. However, the very complexity that makes

these mixtures so valuable also presents significant technical challenges throughout the discovery pipeline. The process of prospecting active molecules from these complex mixtures is classically performed by **bio-guided isolation**, but this is intensive work that can be hampered by false positive results and loss of activity through multiple fractionation steps and repetitive bioassays [64]. This whitepaper examines the principal technical barriers in screening, isolation, and characterization of complex natural product mixtures within the context of exploring **biologically relevant chemical space (BioReCS)** for drug discovery research, and details emerging technological solutions overcoming these limitations.

The Challenge of Navigating Biologically Relevant Chemical Space

Defining Biologically Relevant Chemical Space for Natural Products

The concept of **chemical space (CS)** or **chemical universe** refers to the theoretical multidimensional space encompassing all possible chemical compounds, where molecular properties define coordinates and relationships between compounds [1]. Within this vast universe, the **biologically relevant chemical space (BioReCS)** comprises molecules with biological activity—both beneficial and detrimental [1]. Natural products exhibit distinct and privileged occupancy within BioReCS, populating regions that often lack representation in synthetic medicinal chemistry databases [49].

Comprehensive analyses reveal that natural products possess distinctive physicochemical properties that differentiate them from synthetic compounds and drugs, while largely adhering to the Rule of Five, which renders them a valuable and necessary component of screening libraries for drug discovery [24]. Studies using the chemical space navigation tool **ChemGPS-NP** have demonstrated that natural products cover unique regions of chemical space not adequately explored by conventional medicinal chemistry compounds, indicating these regions represent promising yet underexplored territory for drug discovery [49].

Table 1: Chemical Space Comparison: Natural Products vs. Medicinal Chemistry Compounds

Property	Natural Products	Medicinal Chemistry Compounds	Analysis Method
Structural Rigidity	Generally more structurally rigid [49]	Generally more flexible [49]	ChemGPS-NP Principal Component 4 (PC4) [49]
Aromaticity	Lower degree of aromaticity [49]	Higher degree of aromaticity [49]	ChemGPS-NP Principal Component 2 (PC2) [49]
Size & Lipophilicity	Similar distribution [49]	Similar distribution [49]	ChemGPS-NP Principal Components 1 & 3 (PC1, PC3) [49]
Scaffold Diversity	High scaffold diversity, unique topologies [50]	Limited scaffold diversity, focused around historical targets [50]	Scaffold topology analysis [50]

Underexplored Regions in BioReCS

Certain regions of BioReCS remain underexplored due to significant technical challenges in their investigation. These include:

- **Macrocycles and bRo5 Compounds**: Large and complex natural products, macrocycles (compounds containing rings of ≥12 atoms), and other mid-sized peptides often fall into the **beyond Rule of 5 (bRo5)** category, presenting challenges for conventional screening and isolation protocols [1].
- **Metal-Containing Molecules**: These are frequently excluded from standard chemoinformatic analyses because most modeling tools are optimized for small organic compounds [1].
- **Compounds with Undesirable Effects**: BioReCS includes "dark regions" containing compounds with toxic or other detrimental biological effects, which have naturally received less research focus [1].

Technical Barriers in Screening Complex Mixtures

The Library Quality and Preparation Hurdle

The success of any screening campaign is fundamentally dependent on the quality of the chemical library. Constructing high-quality natural product libraries—whether from microbial, plant, marine, or other sources—is a costly and technically challenging endeavor [74]. These libraries can be composed of **crude extracts**, **semi-pure fractions**, or **single purified natural products**, each design carrying distinctive advantages and disadvantages [74]. Crude extract libraries have lower resource requirements for sample preparation but demand significant effort for the subsequent identification of bioactive constituents. Pre-fractionated libraries balance preparation effort with a shortened active principle identification timeline, while purified natural product libraries require substantial upfront resources but simplify the hit detection process to that of synthetic single-component libraries [74].

The Dereplication Imperative

A critical step in natural product screening is **dereplication**—the process of rapidly identifying known compounds present in a mixture to avoid redundant rediscovery [74]. This process is essential for prioritizing novel leads and allocating resources efficiently. The use of **mass spectrometry** and **HPLC-mass spectrometry** together with spectral databases serves as a powerful tool in the chemometric profiling of bio-sources for natural product production [74]. High-throughput, high-sensitivity flow NMR is also emerging as a valuable tool in this area [74].

Advanced Screening Technologies

To overcome the limitations of traditional bioassay-guided fractionation, advanced screening technologies have been developed that directly probe ligand-target interactions.

Affinity Selection Mass Spectrometry (AS-MS)

Affinity selection mass spectrometry (AS-MS) is a consolidate **high-throughput screening (HTS)** technique that interrogates non-covalent target-ligand complexes as a non-functional assay [64]. It is a label-free biophysical method that discloses binders solely by mass spectrometry data, providing conditions for chemical annotation of the identified ligands [64]. A key advantage is its ability to identify several ligands exhibiting multiple mechanisms of action against the same target, including orthosteric and allosteric ligands [64].

The AS-MS workflow involves four major stages [64]:

- **Static Incubation**: The biological target is incubated with the natural product library.
- **Separation**: Removal of non-binding mixture components.
- **Dissociation**: Ligands are released from the target-ligand complex.
- **Identification**: Dissociated ligands are analyzed by LC-MS.



Figure 1: AS-MS Workflow for Natural Product Screening

AS-MS can be implemented in various formats, primarily categorized into solution-based and immobilized target approaches [64]. Each method presents distinct advantages and limitations, which must be considered when designing a screening campaign.

Table 2: AS-MS Methodologies: Comparative Analysis

Method Type	Specific Techniques	Key Features	Considerations
Solution-Based	Size exclusion chromatography (SEC), Ultrafiltration, Vacuum filtration, Gel filtration [64]	Target remains in native state; Suitable for soluble proteins [64]	Potential for ligand loss with rapid off-rates [64]
Immobilized Target	Affinity capture MS (AC-MS), Magnetic microbeads (MagMass), Ligand-fishing [64]	Target can be recycled; Controlled washing conditions [64]	Potential for target denaturation during immobilization [64]

Ultrafiltration-based AS-MS has been successfully applied to explore 5-lipoxygenase (5-LOX) ligands in *Inonotus obliquus*, leading to the identification of botulin, lanosterol, and quercetin as promising molecules [64].

NMR-Based Screening

NMR spectroscopy offers unique opportunities for screening complex mixtures due to its unbiased nature and rich structural information content [75]. Unlike mass spectrometry, NMR is less biased toward specific compound classes, providing relatively uniform detection across diverse metabolites [75]. Techniques such as **SAR by NMR** and **STD-NMR** have been effectively utilized to screen molecular libraries directly in mixtures, without the need for prior separation [74] [75].

Technical Barriers in Isolation and Characterization

The Instability and Low Abundance Challenge

A significant barrier in natural product research is the isolation of compounds that are chemically unstable or present in minute quantities. Traditionally, achieving high purity through chromatographic fractionation was deemed essential for successful NMR characterization, but this approach excluded metabolites not robust enough to survive chromatography [75]. Furthermore, activity-guided isolation may overlook biologically important compounds that act synergistically rather than individually [75].

Advanced Characterization Technologies

NMR Spectroscopic Analysis of Mixtures

NMR spectroscopy has undergone a paradigm shift, evolving from a technique relegated primarily to pure compounds to a powerful tool for characterizing complex metabolite mixtures [75]. This approach is particularly valuable for identifying otherwise inaccessible small molecules, such as compounds prone to chemical decomposition that cannot be isolated [75].

Pioneering work analyzing unfractionated biofluids has demonstrated the power of this approach. For example:

- Analysis of fresh, unfractionated poison gland secretion from *Myrmicaria* ants led to the identification of myrmicarin 430A, a new family of heptacyclic alkaloids that rapidly decomposes upon exposure to air [75].
- Using reduced-volume cryogenic probes, researchers performed "single insect NMR" on defensive secretions from individual walking sticks, revealing novel monoterpene dialdehydes and demonstrating chemical diversity at the level of individual animals [75].
- Screening of unfractionated spider venom samples using 2D NMR spectroscopy led to the discovery of sulfated nucleosides as major venom components in several spider species, components that had been missed by earlier chromatographic approaches [75].

Hyphenated Analytical Techniques

The various available **hyphenated techniques** (e.g., GC-MS, LC-PDA, LC-MS, LC-FTIR, LC-NMR, LC-NMR-MS, CE-MS) have made possible the pre-isolation analyses of crude extracts or fractions from different natural sources [76]. These integrated systems enable:

- On-line detection of natural products
- Dereplication of known compounds
- Chemotaxonomic studies and chemical fingerprinting
- Quality control of herbal products
- Metabolomic studies [76]

Ultra-high-performance liquid chromatography coupled with quadrupole-Orbitrap high-resolution mass spectrometry has been successfully applied for comprehensive chemical constituent analysis of complex natural products like *Ranunculus sceleratus* L. [77].

The Scientist's Toolkit: Essential Research Reagent Solutions

Success in navigating the technical barriers of natural product research requires specialized reagents, materials, and instrumentation. The following toolkit details essential resources for effective screening, isolation, and characterization.

Table 3: Research Reagent Solutions for Natural Product Research

Tool/Reagent	Function/Application	Technical Specifications
Ultrafiltration Membranes	Separation of target-ligand complexes from unbound molecules in AS-MS [64]	Molecular weight cutoffs 500-500,000 Da; Compatible with centrifugal force, vacuum, or pressure [64]

Tool/Reagent	Function/Application	Technical Specifications
Immobilized Target Platforms	Ligand fishing using immobilized biological targets on solid supports [64]	Magnetic microbeads (MagMass); Functionalized chromatography resins [64]
Cryogenic NMR Probes	Enhanced sensitivity for NMR analysis of limited samples or low-abundance compounds [75]	1-mm HTS cryogenic probes providing 25-fold greater sensitivity than conventional probes [75]
Hyphenated System Columns	Chromatographic separation prior to mass spectrometric or NMR detection [76] [77]	UHPLC columns compatible with high-resolution MS systems (e.g., Quadrupole-Orbitrap) [77]
Chemical Dereplication Databases	Rapid identification of known compounds to avoid redundant rediscovery [74]	Spectral databases for MS and NMR; Dictionary of Natural Products (DNP) [74] [49]

Integrated Workflow for Modern Natural Product Research

To maximize efficiency in exploring natural product chemical space, an integrated workflow incorporating advanced technologies at each stage is essential. The following diagram illustrates a modern approach that combines chemical and biological profiling for functional annotation of complex natural product mixtures.



Figure 2: Integrated Workflow for Natural Products Research

This workflow emphasizes the integration of **chemical profiling** and **biological screening** data early in the process, enabling informed prioritization of leads before committing to resource-intensive isolation efforts. Advances in **chemoinformatics** tools and **molecular networking (MN)** allow researchers to relate the presence or absence of specific metabolites to observations of biological phenotypes in profiling assays [78]. This integrated systems biology approach provides a broad perspective on the biological roles of all metabolites in complex samples, ultimately accelerating the identification of novel therapeutic candidates from nature's chemical treasure trove.

The technical barriers in screening, isolation, and characterization of complex natural product mixtures remain significant, yet technological advances are rapidly overcoming these historical limitations. Methods such as **affinity selection mass spectrometry** and **NMR-based mixture screening** are transforming the screening landscape, while **hyphenated analytical techniques** and **advanced chemoinformatic tools** are accelerating the isolation and characterization process. By adopting integrated workflows that combine chemical and biological profiling, researchers can more effectively navigate the **biologically relevant chemical space** occupied by natural products, bridging the gap between computational methods and experimental validation. As these technologies continue to evolve, natural products will maintain their essential role in drug discovery, providing novel chemical scaffolds with unique properties that continue to elude conventional synthetic approaches.

Natural products (NPs) and their derivatives have historically been a prolific source of therapeutic agents, accounting for a significant proportion of FDA-approved small-molecule drugs [79] [12] [80]. These compounds, derived from plants, microorganisms, and marine organisms, exhibit remarkable structural diversity and complexity that often surpasses synthetic chemical libraries [12]. However, a critical challenge persists in natural product-based drug discovery: the **resupply problem**. Transitioning a natural compound from a "screening hit" through a "drug lead" to a "marketed drug" creates exponentially increasing demands for compound amount, which frequently cannot be met by re-isolation from original biological sources due to limited availability, environmental concerns, and unsustainable harvesting practices [79].

This whitepaper examines how sustainable sourcing and synthetic biology approaches are solving the natural product resupply problem within the broader context of exploring natural product chemical space for drug discovery. With over 1.1 million natural products currently documented in databases—only approximately 10% of which are readily purchasable—the scientific community faces significant challenges in accessing these compounds for comprehensive research and development [12]. We explore integrated strategies that combine advanced biotechnology, bioinformatics, and engineering principles to create reliable, scalable, and environmentally responsible resupply pipelines, thereby enabling the continued utilization of nature's chemical richness in pharmaceutical development.

Understanding the Chemical Space and Sourcing Challenges of Natural Products

Natural products occupy a broader chemical space compared to synthetic compounds, characterized by higher structural complexity, increased stereochemical diversity, and distinct physicochemical properties [12] [80]. Analyses of natural product databases reveal that these compounds typically feature more chiral centers, higher oxygen content, greater molecular rigidity, and aliphatic ring systems that contrast with the predominance of aromatic rings in synthetic libraries [80]. This structural diversity directly contributes to their biological relevance and success as drug leads, but also complicates their chemical synthesis and resupply.

Quantitative Analysis of Natural Product Properties

Table 1: Comparative Analysis of Natural Product Properties Versus Synthetic Compounds

Property	Pure Natural Products (PNP)	Natural Products & Derivatives (SNP)	Synthetic Compounds
Molecular Weight	393.9	409.2	Typically <500
ClogP	2.3	3.7	<5
H-bond Donors	2.7	1.4	≤5
H-bond Acceptors	6.6	6.4	≤10
Ring Count	3.6	3.5	Variable
Rotatable Bonds	5.2	6.1	Variable
Chiral Atoms	5.5	1.4	Fewer
Lipinski Violations ≥2	18%	10%	<10%

Source: Adapted from Life Chemicals Natural Product-like Compound Library analysis [80]

The resupply challenge is compounded by several factors: limited natural availability, with many bioactive compounds present in minute quantities in their source organisms; environmental sustainability concerns regarding large-scale harvesting of sensitive species; and the structural complexity of natural products that often makes traditional chemical synthesis economically unviable [79] [12]. Furthermore, certain natural products originate from organisms that are difficult to cultivate or from extreme environments such as deep-sea ecosystems, presenting additional practical challenges for sustainable sourcing [12].

Synthetic Biology Solutions for Natural Product Resupply

Synthetic biology applies engineering principles to biological systems, creating engineered biological platforms that can address the resupply challenge through multiple approaches. This field has reoriented natural product drug discovery by enabling the development of microbial biofactories and engineered biosynthetic pathways that can produce complex natural products sustainably and at scale [81].

Metabolic Engineering and Pathway Refactoring

The foundational approach in synthetic biology involves identifying and transferring entire biosynthetic gene clusters from native producers to heterologous hosts such as *E. coli* or *S. cerevisiae*. This strategy was pioneered with the discovery that giant biosynthetic units, such as the 28-protein module that synthesizes erythromycin in *Actinomycetes*, could be isolated and implemented in host organisms [81]. Success in this area requires extensive pathway engineering, including codon optimization, promoter engineering, and balancing enzyme expression levels to avoid burdening the host metabolism.

The artemisinin bioproduction project represents a landmark achievement in this domain. Through sophisticated metabolic engineering, researchers developed yeast strains capable of producing artemisinic acid, a precursor to the antimalarial drug artemisinin, which traditionally was extracted from the sweet wormwood plant (*Artemisia annua*) with significant supply chain limitations [81]. This project demonstrated the viability of synthetic biology for producing complex plant-derived natural products in microbial systems, establishing a blueprint for numerous subsequent efforts.

Genome Mining for Novel Enzyme Discovery

Advances in genome sequencing and bioinformatics have enabled **genome mining** approaches that identify cryptic biosynthetic gene clusters in microbial genomes, revealing enzymes capable of performing novel chemical transformations [82]. This strategy has been particularly valuable for discovering enzymes that catalyze **stereodivergent transformations**, providing access to diverse stereoisomers of natural product scaffolds that might be difficult to obtain through chemical synthesis [82].

Table 2: Key Bioinformatic Tools for Natural Product Biosynthetic Gene Cluster Analysis

Tool Name	Primary Function	Application in Resupply Solutions
antiSMASH	Identification & analysis of biosynthetic gene clusters	Prediction of natural product pathways from genomic data
SMURF	Similar function to antiSMASH	Genome mining for secondary metabolite pathways
Natural Product-Likeness Scorer	Computational assessment of natural product similarity	Prioritization of compounds for library development
GDB-17	Enumeration of possible organic molecules	Virtual exploration of synthesizable chemical space
SANCDB	Curated database of natural compounds & analogs	Resource for natural product discovery & optimization

Source: Compiled from multiple references [36] [12] [81]

Genome mining has uncovered enzymes with noncanonical activities that exhibit unusual stereoselectivities, significantly expanding the toolbox available for biocatalytic production of natural products [82]. These enzymes can process diverse substrate scopes, enabling the generation of products with distinct stereochemical markers that are crucial for pharmaceutical efficacy. The discovery of such enzymes through genome mining provides new biocatalytic tools that can be integrated into synthetic biology platforms for natural product synthesis.

Engineering Genetic Circuits and Synthetic Cellular Models

Synthetic biology employs engineered genetic circuits to create cellular factories with precisely controlled behaviors. These circuits typically comprise three elements: an inducer (small molecule, ligand, or light), a genetically encoded circuit that processes the input signal, and an output (such as a reporter gene or target natural product) [81]. Such systems can be designed for dynamic regulation of metabolic fluxes, improving titers of desired compounds by avoiding the accumulation of intermediate metabolites that might be toxic to the host cell.

Synthetic cellular models can also function as screening platforms for both target-based and phenotypic-based drug discovery approaches [81]. These systems can be designed to incorporate human drug targets or disease-relevant pathways, enabling direct screening of natural product libraries while simultaneously developing production strains for hit compounds. This integration of discovery and production represents a powerful paradigm for accelerating natural product-based drug development.

Experimental Protocols for Synthetic Biology Approaches

Protocol: Genome Mining for Biosynthetic Gene Clusters

This protocol outlines the key steps for identifying novel natural product biosynthetic pathways through genome mining:

- **Genome Sequencing and Assembly**: Sequence target organism genomes using Illumina, PacBio, or Oxford Nanopore technologies to obtain high-quality draft or complete genomes. For complex environmental samples, perform metagenomic sequencing.
- **Bioinformatic Analysis**: Utilize specialized tools such as antiSMASH (Antibiotics & Secondary Metabolite Analysis Shell) or SMURF (Secondary Metabolite Unknown Regions Finder) to identify biosynthetic gene clusters (BGCs) encoding natural product pathways [81]. These tools detect characteristic signature sequences of polyketide synthases (PKS), non-ribosomal peptide synthetases (NRPS), and other biosynthetic systems.
- **Comparative Genomics**: Perform phylogenomic analysis to identify unique or divergent BGCs by comparing against databases of known gene clusters. Prioritize clusters with novel architectures or in silent/silenced genomic regions.

- **Heterologous Expression**: Clone prioritized BGCs into suitable expression vectors (e.g., BAC, cosmic, or artificial chromosome vectors) and introduce into heterologous hosts such as *Streptomyces coelicolor*, *E. coli*, or *S. cerevisiae* [81]. Optimize expression through promoter engineering and ribosome binding site modification.
- **Metabolite Analysis**: Characterize compounds produced by recombinant strains using LC-MS/MS and NMR spectroscopy. Compare spectral data against natural product databases to identify novel compounds.
- **Pathway Engineering**: Refactor the BGC for improved production titers by optimizing codon usage, removing regulatory elements, and balancing expression of individual genes.

Protocol: Metabolic Engineering for Natural Product Production

This protocol provides a framework for engineering microbial hosts for natural product production:

- **Host Selection**: Choose an appropriate microbial host (*E. coli*, *S. cerevisiae*, *B. subtilis*) based on the target natural product's biosynthetic requirements, including precursor availability, cofactor requirements, and potential toxicity.
- **Pathway Design**: Design biosynthetic pathways using bioinformatic tools, breaking down the target molecule into biosynthetic steps and identifying or engineering enzymes for each transformation.
- **Genetic Construct Assembly**: Assemble genetic constructs using modern DNA assembly methods (Golden Gate, Gibson Assembly, CRISPR/Cas9). Include appropriate regulatory elements (promoters, RBS, terminators) for balanced expression.
- **Host Transformation and Screening**: Introduce constructs into the host organism and screen for production using analytical methods (HPLC, LC-MS). Employ high-throughput screening methods when applicable.
- **Strain Optimization**: Implement iterative cycles of the Design-Build-Test-Learn paradigm to improve production titers. Strategies include:
    - Modifying precursor supply pathways
    - Engineering cofactor regeneration systems
    - Removing competing pathways
    - Implementing dynamic regulatory controls
- **Bioprocess Development**: Scale up production from laboratory flasks to bioreactors, optimizing process parameters (pH, temperature, aeration, feeding strategies) for maximum yield and productivity.

Visualization of Synthetic Biology Workflows

The following diagrams illustrate key synthetic biology workflows for solving the natural product resupply problem.

Biosynthetic Pathway Engineering Workflow



Diagram Title: Biosynthetic Pathway Engineering Workflow

Genetic Circuit Design for Natural Product Synthesis



Diagram Title: Genetic Circuit Design Framework

The Scientist's Toolkit: Research Reagent Solutions

Implementation of synthetic biology approaches for natural product resupply requires specialized research reagents and tools. The following table details key resources available to scientists working in this field.

Table 3: Essential Research Reagents and Tools for Natural Product Synthetic Biology

Tool/Reagent Category	Specific Examples	Function & Application
Natural Product Libraries	Selleck Natural Product Library (3,673 compounds) [83]; Life Chemicals Natural Product-like Libraries (15,000+ compounds) [80]	High-throughput screening for bioactivity; source of lead compounds
Specialized Databases	Super Natural II; Coconut 2.0; Dictionary of Natural Products; SANCDB; NAPRORE-CR [12]	Cheminformatic analysis; chemical space exploration; target prediction
Genome Mining Tools	antiSMASH; SMURF; BAGEL; PRISM [81]	Identification of biosynthetic gene clusters; pathway prediction
Heterologous Host Systems	*E. coli* BAP1; *S. cerevisiae*; *Streptomyces coelicolor*; *Bacillus subtilis* [81]	Production chassis for heterologous expression of biosynthetic pathways
Genetic Engineering Tools	CRISPR-Cas9; Gibson Assembly; Golden Gate Assembly; SEVA plasmids [81]	Genetic manipulation; pathway construction; genome editing
Bioinformatic Resources	Natural Product-Likeness Calculator; Synthetic Accessibility Score (SAS) tools [36] [80]	Assessment of compound natural product similarity; synthetic feasibility evaluation
Metabolic Modeling Software	COBRApy; OptFlux; GEMs for host organisms [81]	Prediction of metabolic fluxes; identification of engineering targets

Synthetic biology provides a powerful and expanding toolkit for addressing the longstanding resupply problem in natural product-based drug discovery. By leveraging advances in metabolic engineering, genome mining, and genetic circuit design, researchers can create sustainable production platforms for complex natural products that would otherwise be inaccessible for development as therapeutic agents. These approaches are complemented by sophisticated cheminformatic analyses that help prioritize the most promising natural product scaffolds for development [12].

The future of natural product resupply will likely involve increased integration of artificial intelligence and machine learning approaches to predict biosynthetic pathways, optimize enzyme function, and design engineered production strains [36] [84]. Additionally, the exploration of previously untapped natural sources, including organisms from extreme environments and microbial dark matter, will continue to expand the accessible natural product chemical space [12]. As these technologies mature, synthetic biology platforms will become increasingly central to natural product-based drug discovery, enabling reliable, scalable, and sustainable access to nature's chemical diversity for the development of next-generation therapeutics.

The Nagoya Protocol and Access and Benefit-Sharing (ABS) Regulations

The Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising from their Utilization (ABS) is an international agreement that establishes a transparent legal framework for the effective implementation of one of the three objectives of the Convention on Biological Diversity (CBD): the fair and equitable sharing of benefits arising from the utilization of genetic resources [85]. For researchers exploring natural products for drug discovery, this protocol has profound implications. Natural products (NPs) and their derivatives represent a significant source of therapeutic agents, accounting for approximately 56.1% of all drugs approved by the FDA between 1981 and 2019 [60]. The protocol recognizes that each country has sovereign rights over the genetic resources within its jurisdiction and aims to ensure that benefits arising from their use are shared fairly and equitably [85].

In practical terms, "genetic resources" within this context include any material of plant, animal, microbial, or other origin containing functional units of heredity that possesses actual or potential value, including their derivatives [85]. For drug discovery professionals, this encompasses the biological materials typically used in natural product research, from which novel bioactive compounds are often isolated. The protocol also covers "traditional knowledge" associated with genetic resources, defined as the knowledge, innovations, and practices of indigenous and local communities relevant for the utilization of genetic resources [85]. This dual coverage creates both obligations and opportunities for researchers seeking to explore the vast chemical space of natural products for therapeutic development.

Core Principles and Regulatory Framework

Fundamental Concepts and Definitions

The Nagoya Protocol operates on several foundational principles that researchers must understand:

- **Prior Informed Consent (PIC)**: The permission given by the provider country, in this case the Competent National Authority, to the user prior to accessing genetic resources, based on all necessary information about the intended use and terms of access [85] [86].
- **Mutually Agreed Terms (MAT)**: The negotiated agreement between the provider and user that establishes the fair and equitable sharing of benefits arising from the utilization of the genetic resources [85].
- **Internationally Recognized Certificate of Compliance (IRCC)**: The document that serves as evidence that genetic resources have been accessed in accordance with PIC and that MAT have been established [87] [85].
- **Utilization of Genetic Resources**: The protocol defines utilization as "conducting research and development on the genetic and/or biochemical composition of genetic resources, including through the application of biotechnology" [85]. This definition directly encompasses drug discovery activities involving natural products.

Global Implementation and Variations

The Nagoya Protocol entered into force on 12 October 2014 and has been implemented through various national legislations [85]. While the core principles remain consistent, specific requirements can vary significantly between countries. As of 2020, the CBD and Nagoya Protocol on ABS had been ratified, accorded to, approved, or accepted by 196 and 123 countries, respectively [88]. More recent developments include India's Biological Diversity (Access and Benefit Sharing) Regulation 2025, which has expanded its scope to include Digital Sequence Information (DSI) [86].

Brazil represents another important case study, having established its legal framework through Law 13.123/15 and the National System for the Management of Genetic Resources and Associated Traditional Knowledge (SisGen) [60]. A notable feature of Brazilian legislation is that foreign researchers can access native biodiversity only if associated with public or private Brazilian scientific and technological research institutions, which must take responsibility for registering the activity [60].

Table 1: Key Implementation Differences in Select Jurisdictions

Jurisdiction	Governing Legislation	Unique Requirements	Special Considerations
European Union/UK	Regulation (EU) No 511/2014; The Nagoya Protocol (Compliance) Regulations 2015	Due diligence declarations at research funding and commercialization stages	User compliance monitoring by Office for Product Safety and Standards [85]
India	Biological Diversity Act, 2002; ABS Regulation 2025	Benefit sharing slabs based on annual turnover; includes DSI	Prior intimation to National Biodiversity Authority for resource access [86]
Brazil	Law 13.123/15; SisGen registry	Mandatory association with Brazilian institutions for foreign researchers	Registry replaces previous authorization system, reducing bureaucratization [60]

Compliance Workflow for Researchers

Due Diligence Determination Protocol

Researchers must follow a systematic approach to determine their obligations under the Nagoya Protocol. The following workflow outlines the key decision points in the compliance process:



Access Pathways and Compliance Requirements

Depending on how genetic resources are obtained, researchers must follow different compliance pathways. The Nagoya Protocol distinguishes between direct access (obtaining resources directly from the country of origin) and indirect access (obtaining resources from a third party such as a collaborator or registered collection) [85].

For **direct access**, researchers must:

- Using the ABS Clearing-House, determine whether the access measures include requirements to obtain PIC and MAT for the genetic resource [85].
- If required, apply for PIC by submitting the required information to the identified entry points and stakeholders of the provider country [85].
- Negotiate MAT with the Competent National Authority, which will issue a national permit or its equivalent [85].
- Obtain an Internationally Recognized Certificate of Compliance (IRCC) generated through the ABS Clearing-House [87].

For **indirect access**, researchers must:

- Inquire about the proper method to obtain the genetic resource from the intermediary [85].
- Confirm whether PIC and MAT were established by the intermediary when the resources were originally accessed [85].
- Obtain PIC and MAT documentation from the intermediary, typically in the form of an IRCC or equivalent information [85].
- Verify that the transfer and intended utilization are covered by the existing PIC and MAT conditions [85].

Documentation and Record-Keeping Requirements

Compliance with the Nagoya Protocol requires meticulous record-keeping. Users of genetic resources accessed under the protocol are required to seek, keep, and transfer to subsequent users specific information for a period of 20 years following the end of the period of use [85]. Required records include:

- The date and place the genetic resources and associated traditional knowledge were acquired
- A description of the items acquired, using unique identifiers where available
- The source from which the items were obtained
- Whether the items are subject to rights and obligations regarding access and benefit sharing

- Any decision made regarding the access, as well as the mutually agreed terms of access

Table 2: Essential Documentation for Nagoya Protocol Compliance

Document Type	Purpose	When Required	Retention Period
Internationally Recognized Certificate of Compliance (IRCC)	Evidence that genetic resources were accessed in accordance with PIC and MAT	For all utilization of genetic resources covered by the protocol	20 years after end of use [85]
Due Diligence Declaration	Declaration of compliance with ABS requirements to regulatory authorities	At research funding receipt and commercialization stages [85]	20 years after end of use
Mutually Agreed Terms (MAT)	Contract specifying benefit-sharing arrangements	Before accessing genetic resources for utilization	20 years after end of use
Prior Informed Consent (PIC)	Permission from provider country	Before accessing genetic resources	20 years after end of use
Transfer Documentation	Records of any transfer of genetic resources to third parties	When providing genetic resources to other researchers	20 years after end of use [85]

Benefit-Sharing Mechanisms and Calculations

Forms of Benefit Sharing

The Nagoya Protocol requires fair and equitable sharing of benefits arising from the utilization of genetic resources. Benefit-sharing can take various forms, categorized as monetary and non-monetary benefits [85] [86].

Monetary benefits may include:

- Access fees/fair and equitable sharing of royalties per each product
- Up-front and milestone payments
- Payment of relevant salaries and preferential terms for researchers
- Contributions to research funding
- Joint ownership of relevant intellectual property rights

Non-monetary benefits may include:

- Sharing of research and development results
- Collaboration in scientific research and development programs
- Participation in product development
- Admission to and/or contribution to the development of research facilities
- Training and exchange of expertise and knowledge

Calculation Frameworks

Recent regulatory developments have introduced more precise frameworks for calculating benefit-sharing obligations. India's 2025 ABS Regulations, for example, delineate slabs based on the annual turnover of the person/industry utilizing the resources [86]. The regulations also specify that for biological resources having high conservation or economic value (such as red sanders or agarwood), the benefit sharing shall not be less than 5% of the proceeds of the auction or sale amount or the purchase price and could be more than 20% in case of commercial use [86].

For intellectual property commercialization, if a person commercializes a product based on IPR developed using biological resources, they must share a monetary benefit up to 1% of the annual gross ex-factory sale price (excluding taxes), depending on the sector and case specifics [86].

Integration with Natural Product Drug Discovery Workflows

Modified Natural Product Library Development Protocol

The drug discovery workflow for natural products must be adapted to incorporate ABS compliance at each stage. The following diagram illustrates a Nagoya-compliant natural product drug discovery pipeline:



Practical Implementation in Library Creation

The National Cancer Institute's (NCI) Program for Natural Product Discovery provides an exemplary model of implementing ABS principles in large-scale natural product library development. The NCI produces a library of 1,000,000 partially purified natural product fractions for distribution to the research community [88]. The program adheres to collection agreements based on the NCI Letter of Collection (LOC), which stipulates equitable benefit sharing from commercial products derived from discoveries, irrespective of whether a formal agreement has been signed by each participating source country [88].

For researchers creating smaller-scale libraries, essential practices include:

- **Proper annotation**: Accurately tag (e.g., barcoded labels) and document each collection with the collecting institution, collector(s), taxonomy and taxonomist(s), location coordinates, date and time, and any relevant field notes [88].
- **Voucher specimens**: Collect and preserve voucher specimens to encourage efforts to keep the categorization and naming of samples current with changes in taxonomy [88].
- **Metadata collection**: Establish a database for sample tracking, possible recollection of sourced material, and conservation understanding [88].

Cheminformatics and ABS Compliance

Modern cheminformatics approaches can support ABS compliance while facilitating natural product discovery. With over 1.1 million natural products documented in current databases, chemoinformatics analysis reveals that natural products occupy broader chemical spaces than synthetic compounds [12]. However, the limited availability of NPs (only ~10% purchasable) and redundancy in known scaffolds pose major challenges in NP research [12].

Specialized natural product databases provide valuable resources for research that can complement ABS compliance:

- **LOTUS, COCONUT, SuperNatural-II, NPASS**: Contain comprehensive information on naturally existing compound structures, molecular physicochemical properties, and molecular descriptors [89].
- **SistematX, PeruNPDB, NAPRORE-CR**: Regional databases that may incorporate ABS considerations for specific geographical sources [12].

These databases enable similarity analysis and virtual screening approaches that can help researchers prioritize genetic resources for access, potentially reducing the need for extensive physical sampling while still exploring chemical diversity.

Research Reagent Solutions and Experimental Tools

Table 3: Essential Research Tools for Nagoya-Compliant Natural Product Research

Tool/Category	Specific Examples	Function in NP Drug Discovery	ABS Compliance Relevance
ABS Compliance Platforms	ABS Clearing-House [87]	Platform for exchanging information on access and benefit-sharing	Provides IRCCs, information on country requirements, and regulatory updates
Natural Product Databases	COCONUT, SuperNatural II, NPASS, LOTUS [89]	Chemical information on natural product structures and properties	Enables preliminary screening and reduces unnecessary physical access to resources
Cheminformatics Tools	ChemSAR [90], DeepAutoQSAR [91]	SAR modeling and molecular property prediction	Facilitates efficient use of accessed materials through in silico approaches
Molecular Representation	RDKit, CDK, OpenBabel, PaDEL [90]	Molecular descriptor calculation and fingerprint generation	Enables comprehensive analysis of chemical space from limited samples
Sample Management Systems	NCI's Natural Product Repository [88]	Physical storage and distribution of natural product fractions	Maintains chain of custody and transfer documentation
Regulatory Tracking	SisGen (Brazil), NBA (India) portals [60] [86]	Country-specific compliance documentation	Manages national regulatory requirements for specific jurisdictions

The Nagoya Protocol and ABS regulations represent a critical framework that researchers must integrate into their natural product drug discovery workflows. While presenting additional complexity, these regulations enable ethical and sustainable exploration of natural product chemical space. Future developments in this area will likely include:

- **Increased focus on Digital Sequence Information (DSI)**: Recent regulations, such as India's 2025 ABS Rules, now include DSI within their scope, creating new considerations for researchers working with genomic data [86].
- **Multilateral benefit-sharing mechanisms**: Development of global systems for fair sharing across borders, in line with the Nagoya Protocol's objectives [86].
- **Enhanced computational approaches**: Leveraging artificial intelligence for target prediction and virtual screening to maximize the value of accessed genetic resources while minimizing unnecessary collection [92] [12].
- **Integration with emerging screening technologies**: Combining ABS-compliant natural product libraries with high-throughput screening and bioassay methods to identify novel bioactive compounds [88].

For researchers, success in this evolving landscape requires understanding both the scientific and regulatory dimensions of natural product discovery. By implementing robust compliance workflows from the outset and leveraging modern cheminformatics tools, drug discovery professionals can continue to explore the rich chemical space of natural products while ensuring fair and equitable benefit sharing with provider countries and communities.

The exploration of natural products for drug discovery is akin to searching for a needle in a haystack, with the added challenge that most of the easily discoverable "needles" have already been found. **Rediscovery**, the repeated identification of known compounds, represents a significant bottleneck in natural product-based drug discovery, consuming valuable resources and time. The underlying issue stems from the immense scale of **chemical space**—the theoretical space containing all possible organic molecules. Research indicates that while known chemical databases contain millions of compounds, this represents only a tiny fraction (<0.1%) of the possible small molecule structures that could be synthesized and tested [93]. This vast unexplored territory offers tremendous opportunity but also presents formidable challenges in navigation. The concept of the **chemotype**—a chemically distinct entity defined by its specific composition of secondary metabolites—provides a crucial framework for addressing this challenge [94]. By focusing on novel chemotype discovery rather than simply new compound isolation, researchers can implement strategic approaches to systematically explore uncharted regions of chemical space and minimize redundant rediscovery of known chemical scaffolds.

Defining Chemotypes and Their Role in Chemical Diversity

Chemotype Fundamentals

A **chemotype** is defined as a chemically distinct entity within a species that demonstrates consistent differences in its secondary metabolite profile, largely under genetic control [94]. Importantly, chemotypes may be morphologically indistinguishable, with their chemical differences arising from minor genetic or epigenetic variations that nonetheless produce significant changes in chemical phenotype. In practical terms, chemotypes are often classified based on the most abundant secondary metabolite produced by an individual organism. For instance, *Thymus vulgaris* (thyme)

demonstrates seven distinct chemotypes characterized by whether thymol, carvacrol, linalool, geraniol, sabinene hydrate, α-terpineol, or eucalyptol dominates its essential oil composition [94]. This classification system, while useful, has limitations, as it may oversimplify complex chemical profiles where multiple compounds contribute significantly to biological activity.

Chemotypes in Diversity Assessment

The systematic analysis of chemotypes provides a powerful alternative to traditional molecular descriptor-based approaches for assessing chemical diversity. **Chemotype-based diversity analysis** offers several distinct advantages for addressing rediscovery. By focusing on molecular scaffolds or core structures, chemotype analysis enables researchers to quantify and maximize the structural diversity of compound libraries, ensuring that screening collections encompass the broadest possible range of chemical skeletons [95]. Studies have demonstrated that diversity selection algorithms based on chemotype analysis can outperform traditional methods using molecular fingerprints, retrieving a larger share of the chemotypes contained in a library when selecting subsets of compounds [95]. This approach is particularly valuable for designing general-purpose screening libraries against novel targets with limited prior structural information, as it maximizes the probability of identifying novel bioactive scaffolds with minimal compound throughput.

Table 1: Comparison of Diversity Assessment Methods

Method	Basis	Advantages	Limitations
Chemotype Analysis	Molecular scaffolds/core structures	Intuitive interpretation; maximizes structural diversity; efficient library design	May oversimplify complex structures
Molecular Fingerprints	Binary representation of structural features	Comprehensive structural representation; well-established algorithms	Less intuitive; may miss scaffold diversity
Molecular Quantum Numbers (MQN)	42 integer descriptors counting atom/bond types, polarity, topology	Simple, universal chemical space classification; easily identifiable features	Newer approach with limited track record

Computational Strategies for Novel Chemotype Discovery

Active Learning and Alchemical Free Energy Calculations

Active learning represents a powerful machine learning strategy for efficiently navigating chemical space by iteratively selecting the most informative compounds for experimental evaluation. When combined with first-principles based **alchemical free energy calculations**, this approach enables targeted exploration of regions containing high-affinity binders while explicitly evaluating only a small subset of a large chemical library [96]. The protocol typically involves an iterative cycle where, at each iteration, a carefully chosen fraction of compounds undergoes computational evaluation, and the resulting affinity data trains machine learning models to improve predictions for subsequent rounds. This strategy has been successfully applied to identify high-affinity phosphodiesterase 2 (PDE2) inhibitors, robustly identifying a large fraction of true positives while dramatically reducing the computational resources required compared to exhaustive screening [96].

Hybrid Dynamic Pharmacophore Models

Dynamic hybrid pharmacophore models (DHPM) represent an innovative approach that addresses the limitations of conventional pharmacophore models by incorporating protein flexibility and multiple binding sites. Unlike conventional pharmacophore models generated from single binding sites, DHPMs capture the combined interaction features of different binding pockets, enabling identification of novel chemotypes that simultaneously engage multiple regions of a target [97]. The development of DHPMs typically involves molecular dynamics simulations of target structures with ligands bound to adjacent sites (e.g., cofactor binding site and substrate binding site), trajectory clustering to identify stable interaction features, and generation of pharmacophore hypotheses that represent the combined binding characteristics. This approach has demonstrated success in identifying structurally diverse compounds with improved binding strength and drug-like properties compared to those identified through conventional methods [97].



Chemical Space Enumeration and Mapping

The comprehensive **enumeration of chemical space** from first principles provides a foundational approach for systematic exploration of novel chemotypes. Initiatives such as the Chemical Universe Generated Databases (GDB) have demonstrated that almost all small molecules (>99.9%) have never been synthesized and remain available for exploration [93]. The classification and representation of this chemical space using systems such as **Molecular Quantum Numbers (MQN)**—42 integer value descriptors that count elementary molecular features including atom and bond types, polar groups, and topological characteristics—enable intuitive navigation and identification of underrepresented regions [93]. By mapping known natural products and synthetic compounds within this framework, researchers can identify "white spaces" in chemical space that represent opportunities for novel chemotype discovery, strategically targeting these regions through synthesis or focused natural product sourcing.

Table 2: Computational Strategies for Novel Chemotype Discovery

Strategy	Key Methodology	Application in Novel Chemotype Discovery	Technical Requirements
Active Learning with Free Energy Calculations	Iterative ML model training with alchemical free energy calculations	Efficient navigation toward high-affinity chemotypes in large libraries	MD simulation expertise; ML infrastructure
Hybrid Dynamic Pharmacophore Models	MD simulations of multi-site binding; hybrid pharmacophore generation	Identification of chemotypes binding multiple target sites simultaneously	MD simulation capability; pharmacophore modeling tools
Chemical Space Enumeration & Mapping	Systematic enumeration using MQN descriptors; chemical space visualization	Targeted exploration of underrepresented chemical regions	Large-scale computing; cheminformatics expertise

Experimental Approaches for Isolating Novel Chemotypes

Advanced Extraction Technologies

The initial extraction process critically influences the range of chemotypes accessible from biological source material. While traditional methods like maceration, percolation, and Soxhlet extraction remain valuable, **contemporary extraction techniques** offer improved efficiency, reduced extraction times, and decreased solvent consumption [98]. Key advanced methods include:

- **Ultrasound-assisted extraction**: Utilizes ultrasonic energy to enhance cell wall disruption and improve solvent penetration, typically reducing extraction time and temperature requirements.

- **Microwave-assisted extraction**: Employs microwave energy to rapidly heat the sample, accelerating desorption of compounds from the matrix while using less solvent than conventional methods.
- **Pressurized solvent extraction**: Uses solvents at elevated temperatures and pressures to maintain them in liquid state above their normal boiling points, significantly improving extraction efficiency and speed.

These methods must be carefully optimized to prevent degradation of labile natural products, as temperature selection is crucial for maintaining compound stability [98]. The choice of extraction method should be guided by the chemical properties of the target compounds and the nature of the biological matrix.

Green Solvent Systems for Expanded Chemotype Access

The selection of extraction solvents plays a pivotal role in determining the quality, quantity, and selectivity of isolated compounds, thereby directly influencing the range of accessible chemotypes. Traditional organic solvents present significant disadvantages including volatility, toxicity, and environmental concerns [98]. The development of **green solvent systems** represents a crucial advancement for accessing novel chemotypes while addressing sustainability and safety concerns:

- **Natural deep eutectic solvents (NADES)**: These solvent systems typically consist of natural primary metabolites (e.g., choline chloride combined with sugars, organic acids, or alcohols) that form eutectic mixtures with superior extraction properties for various natural product classes.
- **Ionic liquids**: These designer solvents offer tunable properties through selection of appropriate cation-anion combinations, enabling selective extraction of specific chemotype classes.
- **Supercritical and subcritical fluids**: Supercritical CO_2 offers tunable solvating power by varying temperature and pressure, while subcritical water exhibits altered polarity and improved extraction efficiency for more polar compounds.

The principle of "like dissolves like" remains fundamental to solvent selection, with solvents having polarity values near the target solute's polarity generally performing better [98]. However, innovative solvent systems can overcome these traditional limitations, enabling access to previously challenging chemotypes.

Dereplication Strategies Early in the Isolation Workflow

Dereplication—the process of rapidly identifying known compounds in complex mixtures—represents a critical strategy for minimizing rediscovery early in the isolation pipeline. Modern dereplication approaches typically combine sophisticated analytical techniques with database searching:

- **LC-MS/MS and LC-HRMS**: Liquid chromatography coupled with tandem mass spectrometry or high-resolution mass spectrometry provides structural information that can be matched against natural product databases.
- **NMR-based dereplication**: Advanced NMR techniques, including DOSY, LC-NMR, and microcryoprobe NMR, enable structure elucidation directly in complex mixtures with minimal purification.
- **Database integration**: Automated searching against comprehensive natural product databases (e.g., AntiBase, Dictionary of Natural Products) facilitates rapid identification of known compounds.

Implementation of robust dereplication protocols at the earliest possible stage of extraction and fractionation enables researchers to prioritize fractions containing potentially novel chemotypes, effectively allocating resources to the most promising leads.

Integrated Workflow: From Biological Material to Novel Chemotypes

A comprehensive, integrated workflow is essential for systematic isolation of novel chemotypes while minimizing rediscovery. The following diagram illustrates a strategic approach that combines computational and experimental methods:



This integrated workflow emphasizes the critical importance of **interdisciplinary collaboration** between biologists, chemists, and computational scientists, which has been shown to significantly advance natural product research [98]. For example, partnerships between chemical engineers and biologists have clarified relationships between extraction methods and the biological activity of natural compounds like C-phycocyanin [98]. Similarly, collaboration between natural product chemists and computational chemists enables the effective application of chemical space mapping and virtual screening to guide experimental isolation efforts.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Essential Research Reagents and Materials for Novel Chemotype Isolation

Reagent/Material	Function/Application	Considerations for Novel Chemotype Discovery
Natural Deep Eutectic Solvents	Green extraction medium with tunable properties	Enhanced extraction of specific chemotype classes; reduced environmental impact
Ionic Liquids	Designer solvents for selective extraction	Customizable for target compound polarity; improved selectivity
Supercritical CO_2	Non-polar extraction medium	Tunable solvating power; minimal solvent residues; temperature-sensitive compounds
Hybrid Silica Materials	Chromatographic stationary phases	Enhanced separation of complex natural product mixtures
Chiral Stationary Phases	Enantioseparation of natural products	Resolution of stereoisomers; access to enantiomerically pure chemotypes
Molecularly Imprinted Polymers	Selective solid-phase extraction	Target-specific isolation; reduced matrix interference
LC-MS/MS Systems	Dereplication and structure elucidation	Rapid identification of known compounds; prioritization of novel leads
Microcryoprobe NMR	Structure elucidation of limited samples	Enhanced sensitivity for rare or minor novel chemotypes

The systematic isolation of novel chemotypes from natural sources requires a multifaceted strategy that integrates computational guidance with experimental innovation. By leveraging approaches such as active learning protocols, dynamic hybrid pharmacophore models, and comprehensive chemical space mapping, researchers can strategically navigate the vast terrain of unexplored chemistry [96] [93] [97]. Simultaneously, advances in extraction technologies, green solvent systems, and early-stage dereplication provide the experimental tools necessary to access and identify novel

chemical entities efficiently [98]. The continuing development of interdisciplinary collaborations promises to further enhance our ability to explore natural product chemical space, addressing the persistent challenge of rediscovery while unlocking new sources of valuable bioactive compounds for drug discovery and other applications. As these methodologies evolve and integrate, they create a powerful framework for systematic exploration of nature's chemical diversity, ensuring that natural product research continues to contribute novel chemotypes to the drug discovery pipeline.

Optimizing ADMET Properties through Early In Silico Prediction

The high failure rate of drug candidates in late development stages poses a significant challenge for the pharmaceutical sector, with poor pharmacokinetics and toxicity accounting for numerous setbacks. Early integration of *in silico* Absorption, Distribution, Metabolism, Excretion, and Toxicity (ADMET) prediction now serves as a crucial strategy to mitigate this risk. This approach is particularly vital within the context of exploring **natural product chemical space**, a proven source of structurally diverse compounds with privileged biological activities. By employing computational ADMET tools at the initial stages of drug discovery, researchers can effectively filter out molecules with unfavorable pharmacokinetic profiles, prioritize promising candidates from complex natural product extracts, and guide the optimization of lead compounds. This whitepaper provides an in-depth technical guide to the methodologies, applications, and limitations of early *in silico* ADMET prediction, with a specific focus on its role in unlocking the therapeutic potential of natural products for drug discovery research [99] [24] [100].

Natural products represent a heterogeneous group of compounds with diverse molecular properties that often occupy regions of **chemical space** not explored by synthetic compound libraries [24]. These molecules largely adhere to the rule-of-five, rendering them a valuable and necessary component of screening libraries [24]. However, a significant hurdle in identifying secondary metabolites from medicinal microbes is the presence of "sleeping gene clusters" – silent biosynthetic pathways that remain unactivated under standard laboratory conditions [99]. Techniques such as abiotic stress, including exposure to heavy metals, can activate these clusters to elicit novel secondary metabolites with enhanced pharmacological profiles [99].

The high rate of medication failures underscores the importance of ADMET evaluation early in the drug design process [100]. Selecting appropriate experimental data for ADMET prediction and applying it effectively in the context of physiological characteristics remains challenging [100]. *In silico* ADMET models, built on verified experimental datasets using key classifying factors and molecular descriptors, provide a powerful solution to these challenges [100]. When framed within the exploration of natural product chemical space, early ADMET prediction becomes an indispensable tool for identifying and optimizing novel therapeutic agents from biological sources.

Computational Methodologies for ADMET Prediction

Computer-aided drug design (CADD) methods are broadly categorized into structure-based drug design (SBDD) and ligand-based drug design (LBDD) approaches [101]. These computational tools interpret and guide experiments to expedite the antibiotic drug design process, and by extension, the discovery of drugs from natural products [101].

Structure-Based Drug Design (SBDD)

SBDD methods analyze macromolecular target 3-dimensional structural information, typically of proteins or RNA, to identify key sites and interactions important for biological function [101]. This information guides the design of drugs that can compete with essential interactions involving the target [101]. Key SBDD methodologies include:

- **Molecular Dynamics (MD) Simulations**: These simulations use force fields to estimate the energy and forces associated with drug-protein complexes [101]. Commonly used MD codes include CHARMM, AMBER, NAMD, GROMACS, and OpenMM [101].
- **Virtual Screening (VS)**: This technique screens large *in silico* compound databases to identify potential binders for a query target [101]. Docking software such as DOCK, AutoDock, and AutoDock Vina are commonly used for this purpose [101].
- **Binding Site Identification**: When no information on the binding site of a target is available, programs like SILCS (Site Identification by Ligand Competitive Saturation), FINDSITE, and ConCavity can identify putative binding sites by considering geometrical match and binding energy [101].

Ligand-Based Drug Design (LBDD)

LBDD methods focus on known ligands for a target to establish a relationship between their physiochemical properties and biological activities, known as a **structure-activity relationship** (SAR) [101]. This information guides optimization of known drugs or design of new drugs with improved activity [101]. Key software packages covering both SBDD and LBDD capabilities include Discovery Studio, OpenEye, Schrödinger, and MOE [101].

Benchmarking ADMET Predictors

Recent advancements in ADMET prediction include comprehensive benchmarking of predictors, particularly those leveraging foundation models [102]. Evaluation protocols now employ sophisticated data splitting strategies to test model generalization:

- **Random Split**: Tests a model's general interpolation ability through standard random data partitioning [102].

- **Scaffold Split**: Separates molecules based on core chemical structure to test generalization to new chemical scaffolds [102].
- **Perimeter Split**: Creates scenarios where test sets are intentionally dissimilar from training data to test extrapolation capabilities [102].

Technical paradigms for ADMET predictors include **end-to-end deep learning models** (e.g., GNNs and Transformers) that automatically learn feature representations, and **feature-based classical models** that rely on expert-engineered molecular descriptors [102]. Roughness Index variants (MODI, SARI, ROGI) help analyze model performance and dataset difficulty [102].

Experimental Protocols for Integrated Discovery

This section details a representative methodology for discovering natural products with optimized ADMET properties, combining metal stress elicitation with computational validation.

Based on research from Streptomyces sp. SH-1312 [99]

Objective: To elicit secondary metabolite production through heavy metal stress and evaluate their pharmacological activities.

Materials:

- **Bacterial Strain**: Actinobacteria *Streptomyces* sp. SH-1312 [99]
- **Culture Medium**: Gause's medium [99]
- **Elicitors**: Cobalt (Co^{2+}), Zinc (Zn^{2+}), and mixed metals (Co^{2+} + Zn^{2+}) at concentrations of 0.5 mM to 4 mM [99]
- **Extraction Solvent**: Ethyl acetate (EtOAc) [99]

Methodology:

- Inoculate strain SH-1312 in Gause's medium containing metal ions at varying concentrations [99].
- Incubate in a rotatory shaker at 180 rpm for 10 days at 28°C [99].
- Separate mycelium and culture broth using a gauze filter [99].
- Extract culture broth with EtOAc (2 × 200 mL) [99].
- Analyze extracts using HPLC to identify stress-induced metabolites [99].
- Isplicate active compounds and determine structures using NMR spectroscopy [99].
- Evaluate antioxidant and cytotoxic activities of purified compounds [99].
- Perform ADMET predictions and molecular docking studies [99].

Workflow Visualization

The following diagram illustrates the integrated experimental-computational workflow for natural product discovery with early ADMET optimization:



Diagram 1: Integrated discovery workflow for natural products with ADMET optimization.

ADMET Prediction Protocol

Objective: To predict ADMET properties of candidate molecules using computational tools.

Materials:

- **Software**: ADMET prediction tools (e.g., those benchmarked in [102])
- **Datasets**: Curated ADMET datasets with experimental verification [100]
- **Descriptors**: Molecular descriptors and fingerprints for quantitative structure-activity relationship (QSAR) modeling [100]

Methodology:

- **Data Curation and Preprocessing**:
  - Collect and standardize molecular structures [102].
  - Generate relevant molecular descriptors [102].
  - Apply strategic data splitting (random, scaffold, perimeter) for model validation [102].
- **Model Selection and Training**:
  - Choose appropriate algorithm based on data size and complexity [102].
  - Options include deep learning models (GNNs, Transformers) or classical machine learning (Random Forest, XGBoost) [102].
  - Train models using curated ADMET datasets [102].
- **Prediction and Validation**:
  - Input candidate molecule structures into trained models [100].
  - Generate predictions for key ADMET endpoints [100].
  - Validate predictions with experimental data where available [100].

Case Study: Anhydromevalonolactone (MVL) from Streptomyces sp. SH-1312

The metal stress approach successfully elicited production of anhydromevalonolactone (MVL) from *Streptomyces* sp. SH-1312, a metabolite absent in normal culture conditions [99]. This case demonstrates the power of integrated methodology for discovering compounds with favorable ADMET profiles.

Pharmacological Profile of MVL

Table 1: Antioxidant and Cytotoxic Activities of Anhydromevalonolactone (MVL)

Assay Type	Specific Assay	IC_{50} Value (μg/mL)	Standard Used	Standard IC_{50} (μg/mL)
Antioxidant	DPPH Scavenging	19.65 ± 5.7	Ascorbic Acid	6.52 ± 4.92
	NO Inhibition	15.49 ± 4.8	Ascorbic Acid	8.44 ± 4.17
	OH• Inhibition	19.65 ± 5.22	Gallic Acid	6.26 ± 6.39
	Iron Chelation	19.38 ± 7.11	EDTA	10.20 ± 6.54
Cytotoxic (PC3 Cell Lines)	24-hour exposure	35.81 ± 4.2	-	-
	48-hour exposure	23.29 ± 3.8	-	-
	72-hour exposure	16.25 ± 6.5	-	-

MVL exhibited remarkable antioxidant activities across multiple assays and demonstrated time-dependent cytotoxic activity against PC3 cancer cell lines [99]. Further mechanistic studies revealed that MVL exerts pharmacological efficacy by **upregulation of P53 and BAX** while **downregulation of BCL-2** expression, indicating induction of apoptotic pathways [99].

ADMET and Molecular Docking Results

Table 2: ADMET Profile and Molecular Docking of MVL

Property Category	Specific Property	Result for MVL
Toxicity	Hepatotoxicity	Safer profile
	Cytochrome Inhibition	Safer profile
	Cardiotoxicity	Non-cardiotoxic
Molecular Docking	Target Protein	Binding Energy

Property Category	Specific Property	Result for MVL
	P53	Good binding energy in active region
	BAX	Good binding energy in active region

During ADMET predictions, MVL displayed a favorable safety profile with no significant hepatotoxicity, cytochrome inhibition, or cardiotoxicity concerns [99]. Molecular docking studies confirmed that MVL binds in the active region of target proteins P53 and BAX [99]. The research triumphantly announced a prodigious effect of heavy metals on actinobacteria with fringe benefits as a key tool for MVL production with a strong pharmacological and pharmacokinetic profile [99].

Essential Research Reagents and Computational Tools

Successful implementation of early ADMET optimization requires specific research reagents and computational resources. The following table details key components for establishing this workflow.

Table 3: Research Reagent Solutions for ADMET-Optimized Natural Product Discovery

Category	Item	Function/Application
Biological Materials	Actinobacteria strains (e.g., *Streptomyces* sp.)	Source of diverse secondary metabolites with pharmaceutical potential [99]
	Gause's Medium	Specialized culture medium for actinobacteria cultivation [99]
Elicitors	Heavy metal ions (Co^{2+}, Zn^{2+})	Abiotic stress agents to activate silent gene clusters and enhance metabolite production [99]
Analytical Tools	HPLC with PDA/UV detector	Metabolic profiling, purification, and quantification of elicited compounds [99]
	NMR spectroscopy	Structural elucidation of novel natural products [99]
Computational Resources	ADMET prediction tools (e.g., AutoGluon, TabPFNv2)	Automated training and prediction of pharmacokinetic and toxicity properties [102]
	Molecular docking software (e.g., AutoDock Vina)	Prediction of ligand-target interactions and binding affinities [101]
	Chemical databases (e.g., ZINC)	Source of compound structures for virtual screening and model training [101]
Specialized Software	MD simulation packages (e.g., CHARMM, AMBER)	Simulation of molecular dynamics and protein-ligand interactions [101]
	CADD platforms (e.g., Schrödinger, MOE)	Integrated suites for computer-aided drug design [101]

Signaling Pathway Visualization

The following diagram illustrates the apoptotic mechanism identified for MVL, demonstrating how computational predictions align with experimental findings:



Diagram 2: MVL mechanism of action through apoptotic pathway regulation.

The integration of early *in silico* ADMET prediction into natural product drug discovery represents a paradigm shift in pharmaceutical development. By leveraging computational tools to evaluate pharmacokinetic and toxicity properties at initial stages, researchers can efficiently navigate the complex chemical space of natural products, prioritize candidates with the highest therapeutic potential, and reduce late-stage attrition rates. The case study of MVL production through metal stress elicitation demonstrates the power of this integrated approach, where compounds with favorable ADMET profiles can be identified and optimized before extensive laboratory investment. As ADMET prediction models continue to advance through benchmarked frameworks and sophisticated machine learning algorithms, their role in exploring biologically relevant natural product chemical space will become increasingly indispensable for discovering novel therapeutic agents with optimized pharmacological properties.

Standardization and Quality Control in Natural Product Library Construction

The exploration of **natural product chemical space** offers a powerful strategy for discovering novel therapeutic agents. Natural products (NPs) are a heterogeneous group of compounds with diverse molecular properties that often occupy regions of chemical space not explored by standard synthetic compounds while largely adhering to drug-like principles [24]. This chemical diversity makes them a **valuable, unique, and necessary component** of screening libraries for drug discovery. However, the full potential of this diversity can only be realized through rigorous **standardization and quality control** during library construction. Without these controls, the inherent complexity of natural product sources leads to irreproducible results, misidentified activities, and ultimately, failed discovery efforts.

This technical guide outlines the critical standardized processes and quality control measures required to construct biologically relevant natural product libraries that effectively populate and navigate chemical space for drug discovery research. By implementing these protocols, researchers can transform raw biodiversity into **systematically organized, well-characterized libraries** capable of supporting high-throughput screening (HTS) campaigns and generating reliable, reproducible data for downstream development.

Regulatory and Ethical Considerations in Sample Acquisition

Compliance with International Frameworks

The construction of natural product libraries begins with ethical and regulated sample acquisition. The access and use of biological resources must be mutually agreed upon between the researcher and the country of origin, which maintains sovereign rights over these resources [60]. The **Convention on Biological Diversity (CBD)** established at the 1992 United Nations Conference provides the foundational framework based on three pillars: conservation of biological diversity, sustainable use of its components, and fair and equitable sharing of benefits derived from genetic resources [60].

The **Nagoya Protocol** on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Deriving from Their Utilization supplements the CBD and provides a legal framework for benefit sharing [103]. As of 2020, the CBD and Nagoya Protocol have been ratified or accepted by 196 and 123 countries respectively [103]. Researchers must secure all necessary permits—including collection, shipping, and export permits—before initiating fieldwork, as this process can be time-consuming but is essential for legal and ethical compliance.

National Implementation and Benefit Sharing

Countries rich in biodiversity have implemented specific legal frameworks to regulate access to their genetic resources. For example, Brazil established Law 13.123/15 and the **National System for the Management of Genetic Resources and Associated Traditional Knowledge**

(SisGen) to facilitate compliance with CBD principles [60]. A significant aspect of this framework requires foreign researchers to collaborate with Brazilian scientific institutions, which must assume responsibility for registering the activity [60]. These regulations emphasize **fair and equitable benefit-sharing** arrangements that must be negotiated with source countries, potentially including technology transfer, royalty agreements, or capacity building initiatives.

Standardized Protocols for Sample Collection and Documentation

Field Collection and Voucher Specimens

Proper specimen collection and documentation are fundamental to creating traceable and reproducible natural product libraries. The collection process must include the creation of **voucher specimens** that are accurately tagged (e.g., with barcoded labels) and documented with essential metadata [103]. This documentation should include:

- Collecting institution and collector(s) identities
- Taxonomy and taxonomist(s) identification
- Precise location coordinates (GPS)
- Date and time of collection
- Relevant ecological field notes and habitat characteristics

These voucher specimens must be deposited in recognized herbariums or collections to maintain taxonomic verification and enable future recollection efforts. Comprehensive metadata collection establishes the foundation for **sample tracking databases** that support both scientific reproducibility and compliance with regulatory requirements.

Sample Processing and Storage

Following collection, biological samples require careful processing to preserve chemical integrity. While specific protocols vary by organism type (plant, marine, microbial), general principles include:

- Rapid processing after collection to prevent degradation
- Standardized drying methods (e.g., lyophilization for marine organisms, air-drying for plants)
- Uniform particle size reduction through milling or grinding
- Controlled storage conditions (typically -20°C or below) with limited light exposure
- Documentation of processing steps and parameters in sample metadata

Standardization across these steps minimizes chemical variation between batches and ensures consistent quality throughout the library's lifetime.

Extraction and Fractionation Methodologies

Standardized Extraction Protocols

Extraction protocols must balance comprehensive metabolite recovery with reproducibility and compatibility with downstream screening platforms. Recent methodological advances have improved extraction efficiency while streamlining workflow:

Table 1: Standardized Extraction Methods for Natural Product Libraries

Method	Principle	Advantages	Applications
Pressurized Liquid Extraction	Uses high pressure and temperature	Reduced solvent usage, faster processing	Ideal for solid plant materials
Ultrasound-Assisted Extraction	Applies ultrasonic energy	Enhanced extraction efficiency, minimal thermal degradation	Suitable for thermolabile compounds
Microwave-Assisted Extraction	Uses microwave energy	Rapid, selective heating, reduced extraction time	Effective for polar compounds
Supercritical Fluid Extraction	Employ supercritical CO_2	Solvent-free, tunable selectivity	Valuable for lipophilic compounds

The US National Cancer Institute's Natural Product Repository, one of the world's largest collections, generates between 15,000 and 20,000 extracts annually through high-throughput processing methods, demonstrating the scalability of standardized approaches [103].

Prefractionation Strategies for Enhanced Screening

Prefractionation significantly improves screening outcomes by reducing complexity and concentrating minor metabolites. Various chromatographic techniques are employed:

- **Solid Phase Extraction (SPE)**: Provides rapid fractionation with reusable cartridges
- **Counter-Current Chromatography (CCC)**: Offers high recovery rates without solid support
- **High Performance Liquid Chromatography (HPLC)**: Delivers high-resolution separation
- **Supercritical Fluid Chromatography (SFC)**: Uses environmentally friendly solvents

Prefractionated libraries demonstrate **improved screening performance** through higher confidence in hit rates, enhanced biological activity from concentrated minor metabolites, sequestration of nuisance compounds, and streamlined downstream processes [103]. The NCI's Cancer Moonshot program exemplifies large-scale implementation, producing a library of approximately 1,000,000 partially purified natural product fractions in 384-well plates for distribution to the research community [103].

Quality Control and Characterization Measures

Analytical Quality Control Standards

Robust quality control requires comprehensive analytical characterization to ensure batch-to-batch consistency and compound integrity. The following table outlines essential analytical methods:

Table 2: Analytical Quality Control Methods for Natural Product Libraries

Method	Quality Control Parameters	Acceptance Criteria
LC-MS/MS	Metabolic profiling, identity confirmation	Retention time stability, characteristic mass fragments
NMR Spectroscopy	Structural confirmation, purity assessment	Signal-to-noise ratio, absence of contaminant peaks
HPLC-UV/ELSD	Chromatographic fingerprint, purity	Peak area consistency, resolution of critical pairs
Standardized Bioassay	Biological activity baseline	Activity within historical control ranges

Implementing these analytical controls enables the detection of degradation, contamination, or other inconsistencies that could compromise screening results.

Chemical Space Mapping and Diversity Assessment

Systematic assessment of chemical space coverage ensures library diversity and drug-likeness. Computational analysis demonstrates that natural products cover a much larger volume of chemical diversity space than combinatorial compounds [60]. Principal Component Analysis (PCA) using descriptors including AlogP, molecular weight, hydrogen bond donors/acceptors, rotatable bonds, and ring systems reveals that **natural products occupy distinct regions** of chemical space often complementary to synthetic compounds [104]. Research indicates that approximately 52% of natural products comply with Lipinski's "Rule of Five" for drug-likeness, while 71.8% meet at least three of the four criteria [104], confirming their relevance to drug discovery.

Experimental Protocols for Library Construction and Screening

Standardized Workflow for Library Construction

The following diagram illustrates the complete standardized workflow for constructing natural product libraries:



Detailed Extraction and Fractionation Protocol

Protocol: Standardized Solid-Phase Extraction for Natural Product Prefractionation

Background: This protocol describes a standardized solid-phase extraction method for fractionating natural product extracts into distinct chemical fractions based on polarity, reducing complexity for biological screening [103].

Materials and Reagents:

- C18 reversed-phase SPE cartridges (e.g., 10g/60mL capacity)
- HPLC-grade methanol
- HPLC-grade water
- HPLC-grade acetonitrile
- Ethyl acetate (HPLC-grade)
- Hexane (HPLC-grade)
- Natural product extract (100-500mg dissolved in appropriate solvent)
- Vacuum manifold system
- Collection tubes (pre-labeled)

Equipment:

- Analytical balance (0.1mg sensitivity)

- pH meter
- Centrifuge (capable of 5000 × g)
- Nitrogen evaporator
- Volumetric pipettes and flasks

Procedure:

- **Conditioning**: Pass 2 column volumes of methanol through the SPE cartridge, followed by 2 column volumes of water. Maintain column hydration throughout.
- **Sample Loading**: Dissolve the natural product extract in minimal volume of water-methanol (9:1). Load onto column carefully without drying bed.
- **Fraction Elution** (collect each fraction separately):
  - Fraction 1 (non-polar): Elute with 2 column volumes of hexane
  - Fraction 2 (low-mid polarity): Elute with 2 column volumes of hexane:ethyl acetate (1:1)
  - Fraction 3 (mid polarity): Elute with 2 column volumes of ethyl acetate
  - Fraction 4 (mid-high polarity): Elute with 2 column volumes of ethyl acetate:methanol (1:1)
  - Fraction 5 (polar): Elute with 2 column volumes of methanol
  - Fraction 6 (aqueous): Elute with 2 column volumes of water:methanol (1:9)
- **Concentration**: Evaporate fractions to dryness under nitrogen gas at 40°C.
- **Reconstitution**: Weigh each fraction and reconstitute in DMSO at 10mg/mL for screening.

Critical Notes:

- Maintain consistent flow rates (1-2mL/min) throughout the procedure
- Include blank controls processed identically without natural product extract
- Document exact weights and recovery percentages for each fraction
- Store fractions at -20°C in sealed plates with desiccant

Validation: Validate the fractionation scheme using standard compounds with known polarity profiles. Assess fraction quality by TLC or LC-MS to confirm distinct chemical profiles between fractions.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Essential Research Reagents for Natural Product Library Construction

Category	Specific Items	Function & Importance
Extraction Solvents	Methanol, ethanol, ethyl acetate, hexane, water	Comprehensive extraction of diverse metabolite classes with varying polarity
Chromatography Media	C18 reversed-phase, silica gel, Sephadex LH-20, ion-exchange resins	Fractionation and purification based on chemical properties
Analytical Standards	Natural product standards, internal standards (e.g., umbelliferone)	Quality control, method validation, quantification
Stabilizing Agents	DMSO, glycerol, ascorbic acid, butylated hydroxytoluene (BHT)	Compound stabilization, prevention of degradation during storage
Storage Materials	384-well plates, amber glass vials, septa, desiccants	Long-term sample integrity maintenance, prevention of moisture/light damage
Bioassay Reagents	Cell culture media, assay buffers, enzyme substrates, detection reagents	Standardized biological screening across library specimens

Standardization and quality control transform the inherent chemical diversity of natural sources into reliable, screening-ready libraries that effectively explore natural product chemical space. By implementing the comprehensive framework outlined in this guide—encompassing ethical collection, standardized processing, rigorous quality control, and systematic documentation—researchers can construct natural product libraries that deliver reproducible results and identify novel bioactive compounds with drug development potential. As computational tools like ChemGPS-NP and Scaffold Hunter continue to evolve [24], their integration with well-characterized physical libraries will further enhance our ability to navigate chemical space and address unmet medical needs through natural product-inspired drug discovery.

Proven Success and Competitive Advantage: NPs in the Clinic

Natural Products (NPs) and their derivatives have been a cornerstone of medicine for centuries, evolving from ancient herbal remedies to the discovery of transformative drugs like morphine and quinine [105]. The mid-20th century marked a 'golden age' for antibiotic discovery from natural sources, which subsequently expanded into other therapeutic areas [105]. Despite a shift in focus towards technological advances and synthetic compound libraries in the late 20th century, natural products remain an indispensable source of molecular innovation. This whitepaper provides a quantitative analysis of the significant contribution of NPs to the pharmaceutical landscape, particularly to FDA-approved drugs, framing this contribution within the essential context of exploring natural product chemical space for modern drug discovery research. The extensive structural diversity and complexity of NPs, which frequently exhibit unique glycosylation and halogenation patterns, render them invaluable for probing biologically relevant chemical space and identifying novel therapeutic agents [12].

Quantitative Analysis of NP-Derived Drug Approvals

Global Approval Trends from 2014 to 2024

A comprehensive review of drugs approved globally between January 2014 and the end of 2024 provides a clear metric for the contribution of natural products. Among all 579 drugs approved in this period, 56, or **9.7%**, were classified as NPs or NP-derived (NP-D) [105]. This total comprises 44 New Chemical Entities (NCEs), representing 7.6% of all approvals and 11.3% of all NCEs, and 12 NP-Antibody Drug Conjugates (NP-ADCs), accounting for 2.1% of all approvals and 6.3% of all New Biological Entities (NBEs) [105]. The annual number of new NP-D NCEs and NP-ADCs has fluctuated, averaging five approvals per year since 2014 [105]. This data underscores the consistent and vital role of NPs in filling pharmaceutical pipelines, even amidst a growing number of biological therapies.

Table 1: Global Drug Approvals and NP-Derived Contributions (2014-2024)

Category	Total Approvals	NP-Derived Approvals	Percentage of Total	Percentage of Subcategory
All Drugs	579	56	9.7%	-
New Chemical Entities (NCEs)	388	44	7.6%	11.3%
New Biological Entities (NBEs)	191	12 (NP-ADCs)	2.1%	6.3%

Recent Launches and Clinical Pipeline

The momentum for NP-derived drugs continues. Between January 2014 and June 2025, a total of 58 NP-related drugs were launched globally [105]. This figure includes 45 NP and NP-D new chemical entities and 13 NP-antibody drug conjugates, highlighting the successful integration of natural product warheads with advanced biologic platforms [105]. Looking forward, the clinical pipeline remains robust. As of the end of December 2024, **125 NP and NP-D compounds** were identified as undergoing clinical trials or in the registration phase [105]. Notably, thirty-three new pharmacophores not previously found in approved drugs are currently in development, signaling ongoing innovation in this field, although the discovery of truly novel pharmacophores has slowed, with only one discovered in the past 15 years [105].

Navigating the Chemical Space of Natural Products

Structural Characteristics and Diversity

The chemical space occupied by natural products is distinct from that of synthetic compounds. Current databases document over **1.1 million** unique natural products, which display high structural diversity and complexity [12]. NPs frequently incorporate complex ring systems, glycosylation, and halogenation, features that are often underrepresented in synthetic screening libraries [12]. Chemoinformatic analyses consistently show that NPs occupy a broader chemical space than synthetic compounds while largely adhering to the Rule-of-Five, which predicts favorable oral bioavailability [24]. This makes them a valuable, unique, and necessary component of screening libraries for drug discovery [24].

Chemical Space by Biological and Ecological Source

The structural features of NPs are heavily influenced by their source organisms and environments, creating unique subspaces within the broader NP chemical universe:

- **Marine vs. Terrestrial NPs**: Marine natural products are generally larger and more hydrophobic than their terrestrial counterparts [12].
- **Extreme Environments**: NPs derived from deep-sea and other extreme environments often exhibit novel scaffolds and potent bioactivities, making them a critical frontier for discovery [12].
- **Distinct Chemotypes**: Structural characteristics vary noticeably based on biological origin, leading to distinct chemotypes from plants, microbes, and marine organisms [12].

Table 2: Key Databases for Navigating Natural Product Chemical Space

Database Name	Scope and Specialization	Primary Application in Research
Super Natural II	A comprehensive database of natural products [12].	Virtual screening and chemoinformatic analysis.
Dictionary of Marine Natural Products	Focuses on compounds isolated from marine organisms [12].	Research on marine-derived chemical space.
Coconut	An open, curated database of natural products [12].	Comparative analysis and accessible data for sourcing.
Natural Products Repository of Costa Rica (NAPRORE-CR)	Geographically focused open-access database [12].	Exploring region-specific chemical diversity.
PeruNPDB	The Peruvian Natural Products Database for in silico screening [12].	Drug discovery from traditionally sourced compounds.

Experimental Workflows for NP-Based Drug Discovery

The journey from a natural source to a drug candidate involves a multidisciplinary workflow that integrates traditional techniques with modern technological approaches. The following diagram visualizes this integrated process.



Figure 1. Integrated Workflow for Natural Product-Based Drug Discovery

Detailed Methodologies for Key Experimental Protocols

Bioassay-Guided Isolation and Structure Elucidation

This foundational protocol is critical for identifying active compounds from complex natural extracts [105].

- **Preparation of Crude Extracts**: Source material (e.g., plant tissue, marine sponge, microbial culture) is lyophilized and ground. The powder is sequentially extracted with solvents of increasing polarity (e.g., hexane, dichloromethane, ethyl acetate, methanol/water) to fractionate compounds based on hydrophobicity.
- **High-Throughput Bio-screening**: Each crude extract fraction is screened against a panel of therapeutic targets (e.g., specific enzymes, cell-based phenotypic assays). Fractions showing significant activity (e.g., IC50 < 10 μg/mL) are prioritized for further investigation.
- **Iterative Fractionation and Purification**: Active fractions are subjected to chromatographic techniques, including:
  - **Normal-Phase and Reversed-Phase Flash Chromatography**: For rapid fractionation.
  - **High-Performance Liquid Chromatography (HPLC)**: For high-resolution purification, typically using C18 columns and water/acetonitrile or methanol gradients. After each purification step, fractions are re-assayed to track the active compound(s).
- **Structure Elucidation of Pure Active Compounds**:
  - **Mass Spectrometry (MS)**: High-resolution MS (HRMS) is used to determine the exact molecular mass and molecular formula.
  - **Nuclear Magnetic Resonance (NMR) Spectroscopy**: A suite of 1D (1H, 13C) and 2D (COSY, HSQC, HMBC) NMR experiments is performed to determine planar structure, relative stereochemistry, and atomic connectivity.
  - **X-ray Crystallography**: If suitable crystals can be obtained, this technique provides unambiguous confirmation of the absolute stereochemistry.

Chemoinformatic Analysis of NP Chemical Space

This computational protocol characterizes the structural landscape of NPs to guide discovery [12].

- **Data Curation and Standardization**: NP structures are compiled from public (e.g., Coconut, Super Natural II) and commercial (e.g., Dictionary of Natural Products) databases. Structures are standardized (e.g., neutralization, removal of duplicates, standardization of tautomers) using toolkits like RDKit to ensure data integrity.
- **Molecular Descriptor Calculation**: A set of molecular descriptors relevant to drug-likeness and chemical diversity is calculated for each compound. Key descriptors include:
  - **Physicochemical Properties**: Molecular weight, calculated LogP (cLogP), number of hydrogen bond donors/acceptors, topological polar surface area (TPSA).
  - **Complexity and Scaffold Descriptors**: Number of rotatable bonds, fraction of sp3 carbons (Fsp3), molecular frameworks, and ring systems.
- **Chemical Space Visualization and Mapping**: Dimensionality reduction techniques, such as Principal Component Analysis (PCA) or t-distributed Stochastic Neighbor Embedding (t-SNE), are applied to the descriptor data to project the high-dimensional chemical space into 2D or 3D maps. Tools like ChemGPS-NP can be used to navigate this space and compare the position of NPs relative to synthetic compounds and approved drugs [24].
- **Diversity and Cluster Analysis**: Molecular similarity metrics (e.g., Tanimoto coefficient based on molecular fingerprints) are used to cluster NPs. This analysis helps identify densely populated regions of chemical space (scaffold redundancy) and, more importantly, under-explored "dark regions" that may harbor novel bioactivity.

Successful navigation of NP chemical space and development of NP-derived drugs rely on a suite of specialized reagents, databases, and tools.

Table 3: Essential Research Toolkit for NP Drug Discovery

Tool / Reagent Category	Specific Examples	Function and Application
NP Sourcing & Databases	Dictionary of Marine Natural Products [12], Super Natural II [12], NAPRORE-CR [12]	Provides curated structural and source data for virtual screening and chemoinformatic analysis.
Chromatography Media	C18 reversed-phase silica gel, Sephadex LH-20	Purification and fractionation of complex natural extracts. C18 separates by hydrophobicity; Sephadex LH-20 separates by size and polarity in organic solvents.
Analytical Standards	Commercially available NP libraries (e.g., MicroSource Spectrum)	Used as benchmarks in HPLC-MS for dereplication (early identification of known compounds to avoid rediscovery).
Cheminformatic Software	RDKit, ChemGPS-NP [24], Scaffold Hunter [24]	Calculates molecular properties, visualizes chemical space, and analyzes scaffold diversity to prioritize novel structures.
Target Prediction Tools	AI-driven platforms (e.g., SuperPred, SEA)	Predicts potential protein targets for a NP based on its structural similarity to known ligands, generating testable hypotheses for MoA studies.

The quantitative data presented herein leaves no doubt: natural products continue to be a significant and indispensable source of new chemical entities and pharmacophores for the pharmaceutical industry. The approval of 45 NP-derived NCEs in just over a decade, coupled with a robust pipeline of 125 clinical-stage candidates, firmly establishes their enduring value [105]. However, the declining discovery rate of novel pharmacophores signals a need for a strategic shift [105]. The future of NP drug discovery lies in the sophisticated navigation of its vast chemical space. This requires a renewed emphasis on **bioassay-guided isolation** coupled with detailed **mode of action studies** to identify new drug leads [105], integrated with advanced **chemoinformatic approaches** to map diversity and target unexplored regions [12] [24]. Leveraging artificial intelligence for target prediction, exploring untapped species and extreme environments, and mitigating the challenges of compound redundancy and availability are the key strategies that will unlock the next generation of life-saving medicines from nature's chemical treasury [12].

Natural products (NPs) and their derivatives represent an invaluable resource in the anticancer drug discovery pipeline, accounting for over half of all approved anticancer medicines [106]. Their unparalleled chemical diversity provides a vast resource for discovering novel compounds with enhanced efficacy and safety profiles [107]. Drug repurposing—identifying new therapeutic applications for existing drugs—has emerged as a strategy to significantly shorten the traditional 13-15 year drug development pathway at a fraction of the cost [108]. This whitepaper explores prominent case studies of artemisinin, ivermectin, and other natural products undergoing investigation as anticancer agents, focusing on their mechanisms of action, experimental evidence, and research methodologies relevant to drug development professionals.

Artemisinin and Its Derivatives: From Antimalarial to Anticancer Agent

Chemical Features and Pharmacological Profile

Artemisinin is a sesquiterpene lactone isolated from *Artemisia annua* L. (qinghao) and is characterized by a crucial endoperoxide bridge essential for its biological activity [109] [106]. Due to limitations in artemisinin's solubility and bioavailability, several derivatives have been developed, including **dihydroartemisinin (DHA)**, **artesunate (ART)**, **artemether**, and **arteether** [109] [106]. Artesunate is rapidly hydrolyzed to the active metabolite DHA under physiological conditions [106]. The peroxide group is activated by heme or intracellular iron, leading to the generation of cytotoxic reactive oxygen species (ROS) and carbon-centered radicals, which mediate both antimalarial and anticancer effects [109] [108].

Key Anticancer Mechanisms and Evidence

Artemisinin derivatives exhibit multifaceted anticancer activity through several interconnected mechanisms:

- **Induction of Cell Death:** Artemisinin derivatives promote multiple forms of programmed cell death, including apoptosis, autophagy, and ferroptosis [110] [106]. In lung cancer cells, DHA induces ferroptosis and apoptosis, with IC50 values of 19.68 μM in PC9 cells and 7.08 μM in NCI-H1975 cells [106]. Artemisinin itself induces apoptosis in A549 and H1299 lung cancer cells with IC50 values of 28.8 μg/mL and 27.2 μg/mL, respectively [106].
- **Inhibition of Cell Proliferation and Cell Cycle Arrest:** These compounds suppress proliferation and arrest the cell cycle at various checkpoints. In hepatocellular carcinoma (HCC) cells, DHA induces G2/M phase arrest by reducing cyclin B and CDC25C levels while inducing p21 [109]. Conversely, in hepatic stellate cells, DHA causes S-phase accumulation [109].
- **Anti-metastatic and Anti-angiogenic Effects:** Artemisinin derivatives inhibit cancer cell invasion, metastasis, and angiogenesis, crucial processes for tumor growth and dissemination [108] [106].
- **Sensitization to Conventional Therapy:** They demonstrate synergistic effects when combined with other chemotherapeutic agents, potentially reversing multidrug resistance [108].

Table 1: Anticancer Activity of Artemisinin and Its Derivatives Across Cancer Types

Cancer Type	Compound	Experimental Model	Key Findings	IC50 / Effective Concentration
Breast Cancer	Dihydroartemisinin (DHA)	MCF-7 cells	Suppressed proliferation, induced autophagy and pyroptosis, targeted cancer stem cells [106].	129.1 μM (24 h) [106]
	Artemisinin	MCF-7 cells	Suppressed cell growth, induced ferroptosis [106].	396.6 μM (24 h) [106]
	Artesunate	MCF-7 cells	Induced apoptosis [106].	83.28 μM (24 h) [106]
Lung Cancer	Artemisinin	A549 cells	Inhibited proliferation and metastasis, induced apoptosis [106].	28.8 μg/mL [106]
	Dihydroartemisinin	PC9 cells	Induced ferroptosis and apoptosis, inactivated STAT3 [106].	19.68 μM (48 h) [106]
	Artemisinin derivative 4	H1299 cells	Antiproliferation effect, induced ferroptosis [106].	0.09 μM [106]

Cancer Type	Compound	Experimental Model	Key Findings	IC50 / Effective Concentration
Liver Cancer	Artemisinins	HepG2, PLC/PRF/5 cells	Induced G2/M cell cycle arrest [109].	Varies by compound and cell line [106]
	Dihydroartemisinin	Hepatic stellate cells	Induced S-phase cell cycle arrest [109].	Varies by cell line [106]

Detailed Experimental Protocol: Assessing Antiproliferative Effects and Cell Cycle Arrest

Objective: To evaluate the *in vitro* antiproliferative activity of an artemisinin derivative and determine its effect on the cell cycle.

Materials:

- **Test Compound:** Artesunate or Dihydroartemisinin (e.g., from Sigma-Aldrich) [110].
- **Cell Lines:** Cancer cell lines relevant to the research (e.g., MCF-7 breast cancer, A549 lung cancer).
- **Culture Medium:** DMEM or RPMI-1640, supplemented with 10% Fetal Bovine Serum (FBS) and 1% penicillin-streptomycin [111].
- **Assay Reagents:** MTT or Sulforhodamine B (SRB) dye for cell viability/proliferation [111].
- **Cell Cycle Analysis Reagents:** Propidium Iodide (PI) or commercial cell cycle staining kits (e.g., FxCycle PI/RNase Staining Solution) [111].
- **Equipment:** CO_2 incubator, microplate reader, flow cytometer.

Methodology:

- **Cell Seeding and Treatment:**
  - Harvest exponentially growing cells and seed them at a density of 2,000-5,000 cells/well in 96-well plates for viability assays or $1x10^5$ cells/well in 6-well plates for cell cycle analysis [111].
  - After 24 hours of incubation, treat the cells with a concentration gradient of the artemisinin derivative. Include a vehicle control and a positive control.
  - Incubate for 24, 48, and 72 hours.

- **Cell Viability/Proliferation Assay (SRB Assay):**

  - At each time point, terminate the experiment by fixing cells with cold 40% trichloroacetic acid (TCA) for 1 hour [111].
  - Wash plates 3-5 times with tap water to remove TCA and air-dry.
  - Stain the fixed cells with 0.4% (w/v) SRB solution for 30 minutes [111].
  - Remove unbound dye by washing with 1% acetic acid and air-dry.
  - Solubilize the protein-bound dye with 10 mM Tris base solution [111].
  - Measure the absorbance at 492 nm or 510 nm using a microplate reader [111].
  - Calculate the percentage of cell growth inhibition and the IC50 value using non-linear regression analysis.

- **Cell Cycle Analysis by Flow Cytometry:**

  - After 48 hours of treatment, harvest the cells (including the floating cells) by trypsinization [111].
  - Wash the cell pellet with ice-cold Phosphate Buffered Saline (PBS) and fix with 70% ethanol overnight at -20°C [111].
  - Centrifuge to remove ethanol, wash with PBS, and resuspend the cell pellet in 500 µL of PBS containing PI/RNase staining solution [111].
  - Incubate for 30 minutes at room temperature in the dark.
  - Analyze the DNA content using a flow cytometer (e.g., BD FACSCanto II). A minimum of 10,000 events per sample should be acquired [111].
  - Use software (e.g., BD FACSDiva, ModFit) to determine the percentage of cells in the G_0/G_1, S, and G_2/M phases of the cell cycle.

Diagram 1: Experimental workflow for cell viability and cycle analysis.

Ivermectin: Repurposing an Antiparasitic Agent for Oncology

Background and Chemical Profile

Ivermectin (IVM) is a macrolide antiparasitic drug derived from avermectin, composed of 80% 22,23-dihydroavermectin-B1a and 20% 22,23-dihydroavermectin-B1b [112]. Its discoverers won the Nobel Prize in Physiology or Medicine in 2015. Beyond its established role in treating river blindness and scabies, IVM has demonstrated potent anticancer effects in various *in vitro* and *in vivo* models [112].

Key Anticancer Mechanisms and Evidence

Ivermectin exerts its anticancer activity through a multi-target mechanism:

- **Inhibition of Proliferation and Induction of Cell Death:** IVM inhibits proliferation and promotes apoptosis, autophagy, and pyroptosis in cancer cells [112]. This is often linked to the inhibition of the PAK1 kinase and the regulation of multiple downstream signaling pathways [112].
- **Reversal of Multidrug Resistance (MDR):** A significant finding is IVM's ability to reverse P-glycoprotein (P-gp)-mediated multidrug resistance. It achieves this not by direct inhibition but by suppressing P-gp expression via the EGFR/ERK/Akt/NF-κB pathway [113].
- **Targeting Cancer Stem Cells (CSCs) and Tumor Microenvironment:** IVM can inhibit tumor stem cells and modulate the tumor microenvironment, for instance, by enhancing the release of immunogenic molecules like HMGB1 [112].
- **Synergistic Combination with Chemotherapy:** IVM shows optimal efficacy when combined with other chemotherapeutic drugs, sensitizing resistant cancer cells [112] [113].

Table 2: Documented Anticancer Effects of Ivermectin Across Cancer Models

Cancer Type	Experimental Model	Key Findings	IC50 / Effective Dose
Cholangiocarcinoma (CCA)	KKU214 (Gem-sensitive) & KKU214GemR (Gem-resistant) cells	Inhibited proliferation & colony formation; Gem-resistant cells were more sensitive [111].	KKU214: 11.41 μM (48 h); KKU214GemR: 4.05 μM (48 h) [111]
Colorectal Cancer	HCT-8/VCR (Vincristine-resistant) cells	Reversed chemoresistance *in vitro* and *in vivo*; reduced P-gp expression [113].	*In vivo:* 2 mg/kg/day + VCR [113]
Breast Cancer	MCF-7/ADR (Adriamycin-resistant) cells	Reversed chemoresistance *in vitro* and *in vivo* via EGFR/ERK/Akt/NF-κB pathway [113].	*In vivo:* 2 mg/kg/day + ADR [113]
Chronic Myeloid Leukemia	K562/ADR (Adriamycin-resistant) cells	Reversed chemoresistance in xenograft mouse model [113].	*In vivo:* 2 mg/kg/day + ADR [113]
Gastric Cancer	MKN1, SH-10-TC (YAP1-high) cells	Inhibited proliferation in a YAP1-dependent manner [112].	Sensitive in YAP1-high cells [112]

Detailed Experimental Protocol: Evaluating MDR ReversalIn Vivo

Objective: To investigate the ability of ivermectin to reverse multidrug resistance in a xenograft mouse model.

Materials:

- **Test Compounds:** Ivermectin, Chemotherapeutic drug (e.g., Vincristine, Adriamycin) [113].
- **Cells:** Drug-resistant cancer cell line (e.g., HCT-8/VCR colorectal cancer cells) [113].
- **Animals:** Immunodeficient mice (e.g., nude BALB/c mice for solid tumors, NOD/SCID mice for leukemia) [113].
- **Vehicle:** 0.9% NaCl with DMSO for suspension [113].
- **Equipment:** HPLC system, calipers, Western blot apparatus [113].

Methodology:

- **Model Establishment and Grouping:**
  - Subcutaneously inject 1x10^7 drug-resistant cells (e.g., HCT-8/VCR) into the flank of each mouse [113].
  - When tumor volumes reach approximately 100 mm^3, randomize mice into groups (n=6): Vehicle control, Chemotherapy alone (e.g., VCR 0.2 mg/kg/day), IVM alone (2 mg/kg/day), and Combination (IVM + Chemotherapy) [113].
  - Administer drugs via intraperitoneal injection daily.

- **Tumor Monitoring and Analysis:**
  - Measure tumor dimensions with calipers every three days. Calculate volume using the formula: $V = (\text{length} \times \text{width}^2) / 2$ [113].
  - After 27 days, euthanize the animals, excise and weigh the tumors [113].
  - Process tumor tissues for further analysis (IHC, IF, Western blot).

- **Mechanistic Analysis (P-gp Expression):**
  - Homogenize tumor tissues and lyse in RIPA buffer with protease inhibitors [113].
  - Separate proteins by SDS-PAGE and transfer to a PVDF membrane [113].
  - Probe the membrane with primary antibodies against P-gp, followed by HRP-conjugated secondary antibodies [113].
  - Detect signals using chemiluminescence. Re-probe the membrane with an antibody for a loading control (e.g., GAPDH, β-actin).

- **Drug Accumulation Analysis (HPLC):**

- To confirm increased chemotherapeutic drug retention in tumors, analyze tumor homogenates for drug concentration (e.g., Vincristine) using High-Performance Liquid Chromatography (HPLC) [113].

Diagram 2: Ivermectin mechanism for reversing multidrug resistance via the EGFR pathway.

Beyond Single Agents: The Broader Landscape of Marine and Natural Product Anticancer Agents

The exploration of natural product chemical space extends beyond terrestrial plants to marine organisms. Marine cyanobacteria, for instance, are prolific sources of potent anticancer agents [114]. Key developments include:

- **Tubulin-Targeting Agents:** The dolastatins, originally isolated from a mollusk that feeds on cyanobacteria, led to the development of antibody-drug conjugate (ADC) payloads. Six ADCs with cyanobacterial cytotoxins have been approved, highlighting the clinical impact of this resource [114] [92].
- **Novel Mechanisms of Action (MOA):** Recent discoveries include gatorbulin, which acts on a novel pharmacological site on tubulin. Other compounds like apratoxin and coibamide A modulate cotranslational translocation at the Sec61 complex in the endoplasmic reticulum [114].
- **Epigenetic and Proteasome Modulators:** Cyanobacterial compounds such as largazole and santacruzamate A target class I histone deacetylases (HDACs), while carmaphycins are proteasome inhibitors [114].

This expanding pipeline, which also includes SERCA inhibitors and mitochondrial cytotoxins, underscores the richness of marine natural products for discovering agents with novel mechanisms to overcome drug resistance [114].

The Scientist's Toolkit: Essential Reagents and Models

Table 3: Key Research Reagent Solutions for Investigating Natural Product Anticancer Agents

Reagent / Material	Function in Research	Specific Examples / Notes
Cell Line Panels	*In vitro* screening for cytotoxicity, mechanism studies, and resistance modeling.	MCF-7 (breast), A549 (lung), HCT-8 (colorectal), KKU214 (cholangiocarcinoma), and their drug-resistant variants (e.g., MCF-7/ADR, HCT-8/VCR) [113] [111].
Xenograft Mouse Models	*In vivo* evaluation of efficacy, toxicity, and drug resistance reversal.	Nude mice for solid tumors; NOD/SCID mice for leukemias [113].
Antibodies for Western Blot / IHC	Mechanistic analysis of signaling pathways and target expression.	Antibodies against P-gp, p-EGFR, p-Akt, p-ERK, NF-κB, PAK1, YAP1, and cleaved caspases [112] [113].
Flow Cytometry Reagents	Analysis of cell cycle, apoptosis, and surface markers.	Propidium Iodide (PI), Annexin V-FITC, commercial kits (e.g., FxCycle PI/RNase) [113] [111].
Cytotoxicity Assay Kits	High-throughput assessment of cell viability and proliferation.	MTT, Sulforhodamine B (SRB), and Cell Counting Kit-8 (CCK-8) [111].
HPLC Systems	Quantifying drug concentrations in biological samples (pharmacokinetics) and analyzing compound purity.	Used to measure chemotherapeutic drug accumulation in cells and tissues [113].

Artemisinin, ivermectin, and marine-derived cytotoxins exemplify the immense potential of natural products in anticancer drug discovery. The evidence supports their multi-target mechanisms, which include inducing various forms of cell death, overcoming multidrug resistance, and modulating the tumor microenvironment. Future efforts should focus on integrating advanced methodologies such as artificial intelligence, high-throughput screening, and chemical biology to explore novel NP targets and accelerate development [92]. The successful clinical translation of these repurposed drugs and novel natural products will depend on robust, well-designed clinical trials that validate preclinical findings, ultimately increasing the accessibility and affordability of cancer therapies globally [108].

The exploration of **chemical space** for novel drug leads represents a fundamental challenge in modern drug discovery. This whitepaper provides a comparative analysis of two primary approaches: screening libraries of **natural products (NPs)** and those comprising **synthetic compounds**. Natural products, chemical entities produced by living organisms, are pre-validated by nature and have been the historical source of a majority of novel drug classes and essential medicines [17] [49]. In contrast, synthetic libraries, constructed using methodologies like combinatorial chemistry and diversity-oriented synthesis (DOS), offer advantages in terms of scalability and modularity [115] [36]. Framed within the broader thesis of exploring natural product chemical space for drug discovery, this analysis examines the structural diversity, hit-rate performance, and practical applications of these distinct yet complementary strategies, providing drug development professionals with a data-driven guide for library selection and design.

Chemical Space and Structural Diversity

Fundamental Differences in Chemical Space Occupancy

Computational analyses reveal that natural products and synthetic compounds derived from medicinal chemistry efforts occupy notably different regions of biologically relevant chemical space.

Table 1: Chemical Property and Structural Feature Comparison

Characteristic	Natural Products (NPs)	Synthetic/Bioactive Medicinal Chemistry Compounds
General Rigidity	Structurally more rigid [49]	Generally more flexible [49]
Aromaticity	Lower degree of aromaticity [49]	Higher degree of aromaticity [49]
Structural Complexity	Higher structural complexity and more stereocenters [116] [36]	Typically less complex [36]
Adherence to Ro5	~60% have no Lipinski's Rule of 5 (Ro5) violations; many remain bioavailable despite violations [49]	Designed for Ro5 compliance to ensure oral bioavailability [36]
Coverage of Uniqueness	Populate unique, sparsely explored regions of chemical space [49] [24]	Often cluster in over-sampled regions of chemical space [49]

Tools like **ChemGPS-NP** have mapped these differences, showing that NPs cover regions that lack representation in typical medicinal chemistry libraries, such as the **World Drug Index (WOMBAT)** database [49]. This unique occupancy is attributed to evolutionary selection, which optimizes NPs for interactions with biological macromolecules, rendering them a valuable component for any screening library aimed at discovering novel bioactive compounds [49] [24].

Quantitative Assessment of Scaffold Diversity

Scaffold diversity is a critical metric for assessing the potential of a compound library to yield novel hits. Analyses using frameworks like the **Murcko framework** and **Scaffold Tree** hierarchies provide quantitative measures.

Table 2: Scaffold Diversity of Standardized Compound Libraries (41,071 compounds each)

Compound Library	Number of Unique Murcko Frameworks	Number of Unique Level 1 Scaffolds	Relative Structural Diversity
TCMCD (Natural Product Library)	4,289	5,134	Highest Complexity
Chembridge	5,268	6,441	High
Mucle	4,953	6,123	High
VitasM	4,866	5,978	High
ChemicalBlock	4,911	6,032	High
Enamine	4,522	5,654	Medium
Maybridge	3,987	4,956	Medium

The **Traditional Chinese Medicine Compound Database (TCMCD)**, a representative NP library, demonstrates the highest structural complexity among the libraries studied [116]. However, its scaffolds are more conserved, resulting in fewer unique frameworks and Level 1 scaffolds compared to some highly diverse synthetic libraries like **Chembridge** and **Mucle** [116]. This suggests that while NPs introduce high-value, complex scaffolds, synthetic libraries can offer a greater raw number of distinct core structures, highlighting a key trade-off.

Hit Rates and Success in Drug Discovery

The ultimate validation of a screening library lies in its ability to produce viable hits and successful drugs. The historical and contemporary data strongly favor natural products in this regard.

- **Novel Drug Classes**: Natural products and their structural analogues have historically made a major contribution to pharmacotherapy, especially for cancer and infectious diseases, and are the richest source of novel compound classes [17] [49].
- **Hit Identification Strategy**: **Property-based similarity calculations** can identify natural product neighbors of approved drugs. Several of the NPs revealed by this method were confirmed to exhibit the same activity as their drug neighbors, illustrating a viable strategy for lead

identification from a natural product starting point [49].

- **Performance in Screening**: Analyses confirm that bioactive collections and NP libraries come closest to populating the biologically relevant regions of chemical space [49]. This is because their structures have been optimized through evolution for biological interactions, which increases the probability of identifying a hit during screening compared to synthetic compounds, which may be designed with a greater emphasis on synthetic feasibility and drug-like rules rather than inherent bioactivity [17] [36].

Experimental Approaches and Workflows

Diversity-Oriented Synthesis (DOS) for Synthetic Libraries

The **Build/Couple/Pair (B/C/P)** strategy in DOS is a powerful method for generating skeletally diverse synthetic libraries with NP-like complexity [115].



Figure 1: The Build/Couple/Pair (B/C/P) workflow in Diversity-Oriented Synthesis (DOS) for generating skeletally diverse compound libraries, such as lactams, from commercially available building blocks [115].

A computational-aided DOS workflow can be implemented using platforms like **KNIME** to generate a library of lactams, a privileged scaffold in drug discovery [115]. The process involves:

- **Building Blocks**: Retrieving commercially available amine and carboxylic acid building blocks from vendors like **Enamine** [115].
- **Coupling**: Performing intermolecular coupling via amide bond formation to generate linear precursors.
- **Pairing**: Employing different intramolecular pairing reactions (e.g., cyclization) to generate skeletally diverse macrocyclic and non-macrocyclic lactams.
- **Analysis**: The resulting library is then evaluated for novelty, diversity, and drug-like properties compared to existing NPs and drugs from databases like **ChEMBL** [115].

Combinatorial Refactoring of Natural Product Biosynthesis

For natural products, a major challenge is the heterologous expression of their often large and complex biosynthetic gene clusters. Modern synthetic biology addresses this through **combinatorial DNA assembly and refactoring**.



Figure 2: Refactoring the C. jejuni Pgl pathway using combinatorial DNA assembly to optimize heterologous production in E. coli [117].

A seminal application of this approach is the refactoring of the *Campylobacter jejuni* N-glycosylation (*pgl*) pathway in *E. coli* [117]. The experimental protocol is as follows:

- **Deconstruction**: The 10 coding sequences (CDSs: *gne, pglA, pglC, pglD, pglE, pglF, pglH, pglI, pglJ, pglK*) responsible for heptasaccharide biosynthesis are selected and removed from their native regulatory context [117].
- **Combinatorial Assembly**: Using a method like **Start-Stop Assembly**, each CDS is assembled combinatorially with mixtures of synthetic promoters and ribosome-binding sites (RBSs) of varying strengths. This creates a vast library of expression constructs for the pathway [117].
- **Screening**: The library is transformed into an *E. coli* host, and a rapid screen is employed to identify clones that outperform the native, unmodified *pgl* cluster in terms of glycan and glycoconjugate production [117].
- **Validation**: The best-performing refactored loci are combined with plasmids expressing the oligosaccharyltransferase PglB and a target acceptor protein. Glycosylation efficiency and glycoconjugate yield are then quantified, demonstrating the success of the refactoring approach [117].

The Scientist's Toolkit: Essential Research Reagents

Table 3: Key Research Reagent Solutions for Glycoengineering and Library Screening

Reagent / Tool	Source / Example	Function / Application
Oligosaccharyltransferase PglB	*Campylobacter jejuni*	Key enzyme for in vivo N-linked protein glycosylation in bacterial glycoengineering [117] [118] [119].
PNGase F & PNGase A	New England Biolabs	Amidases for enzymatic deglycosylation; gold standard for validating N-glycosylation status of proteins from mammalian (F) and plant/insect (A) systems [120].
ChemGPS-NP	Public Web Resource	Chemical space navigation tool for comparing and visualizing the location of compounds in a property-based reference space [49] [24].
Build/Couple/Pair (B/C/P)	Nielsen and Schreiber, 2007	A systematic DOS strategy for generating skeletally diverse small molecule libraries from simple building blocks [115].
Combinatorial DNA Assembly (Start-Stop)	Taylor et al., 2019	Scarless modular DNA assembly system for constructing combinatorial libraries of multigene expression constructs [117].
KNIME Analytics Platform	Open Source	Platform for designing and executing computational workflows, e.g., for generating virtual libraries of lactams [115].

Reagent / Tool	Source / Example	Function / Application
Traditional Chinese Medicine Compound Database (TCMCD)	Academic Source	A curated database of NPs used for analysis of structural complexity and scaffold diversity in comparative studies [116].

The comparative analysis between natural product and synthetic libraries reveals a landscape of compelling synergies rather than simple superiority. **Natural products** provide access to evolutionarily pre-validated, complex chemotypes that occupy unique and biologically relevant regions of chemical space, leading to historically high success rates in delivering novel drugs. **Synthetic libraries**, particularly those designed using DOS principles, offer unparalleled capacity for exploring vast regions of chemical space and generating high counts of unique, often lead-like scaffolds in a controlled and scalable manner. The future of productive drug discovery lies in the strategic integration of both paradigms. This can be achieved by using computational tools like **ChemGPS-NP** to identify sparsely populated, biologically relevant areas of chemical space and then employing advanced synthetic biology to refactor NP pathways or sophisticated DOS to populate these regions with novel, synthetically tractable compounds. This integrated approach maximizes the chances of discovering novel, effective, and developable small-molecule therapeutics.

The Rising Role of NP-Derived Payloads in Antibody-Drug Conjugates (ADDCs)

Antibody-Drug Conjugates (ADCs) represent a transformative class of targeted cancer therapeutics that combine the specificity of monoclonal antibodies with the potent cytotoxicity of small-molecule payloads. The structural architecture of ADCs comprises three fundamental components: a monoclonal antibody for target recognition, a chemical linker ensuring stability, and a cytotoxic payload responsible for ultimate tumor cell eradication [121] [122]. Within this sophisticated framework, natural products (NPs) and their derivatives have emerged as indispensable payload sources, contributing significantly to the clinical success of ADC technology.

Natural products have served as historic cornerstones in oncology drug discovery, with over half of approved small-molecule drugs originating directly or indirectly from NP origins [12]. This dominance extends powerfully into the ADC landscape, where NP-derived payloads constitute the majority of currently approved conjugates. The inherent biological compatibility, structural complexity, and potent mechanisms of action exhibited by natural products make them ideal candidates for ADC payload development [92] [121]. These compounds often demonstrate exquisite targeting of fundamental cellular processes, including microtubule dynamics and DNA integrity, with potencies 100 to 1000-fold greater than conventional chemotherapeutics [121].

The exploration of NP chemical space continues to yield valuable insights for ADC development. Current databases document over 1.1 million natural products displaying remarkable structural diversity and complexity, frequently featuring glycosylation and halogenation patterns [12]. NPs occupy broader chemical spaces than synthetic compounds and exhibit distinct characteristics based on their origins, with marine-derived NPs often displaying larger molecular weights and greater hydrophobicity than their terrestrial counterparts [12]. This review examines the rising role of NP-derived payloads within ADC development, framed within the broader context of exploring natural product chemical space for drug discovery research.

Clinical Landscape of NP-Derived Payloads in Approved ADCs

Dominant Payload Classes and Their Origins

The ADC clinical landscape is predominantly populated by natural product-derived payloads, which can be broadly categorized into several mechanistic classes. **Table 1** summarizes the key characteristics of NP-derived payload classes used in approved ADCs.

Table 1: Natural Product-Derived Payload Classes in Approved ADCs

Payload Class	Representative Payloads	Natural Product Origin	Mechanism of Action	Approved ADC Examples
Tubulin Inhibitors	DM1, DM4 (maytansinoids)	*Maytenus serrata* (African shrub)	Inhibits microtubule assembly, disrupting cell division	Trastuzumab emtansine (Kadcyla), Mirvetuximab soravtansine (ELAHERE)
Tubulin Inhibitors	Monomethyl auristatin E (MMAE), Monomethyl auristatin F (MMAF)	Dolastatin 10 (marine peptide from *Dolabella auricularia*)	Inhibits tubulin polymerization, preventing mitosis	Brentuximab vedotin (Adcetris), Enfortumab vedotin (Padcev)
DNA-Damaging Agents	Calicheamicin	*Micromonospora echinospora* (bacterial source)	DNA double-strand breaks via enediyne core	Gemtuzumab ozogamicin (Mylotarg), Inotuzumab ozogamicin (Besponsa)
Topoisomerase I Inhibitors	Exatecan (DXd derivative), SN-38	Camptothecin (*Camptotheca acuminata* tree)	Stabilizes topoisomerase I-DNA cleavage complexes	Trastuzumab deruxtecan (Enhertu), Sacituzumab govitecan (Trodelvy)

Payload Class	Representative Payloads	Natural Product Origin	Mechanism of Action	Approved ADC Examples
DNA Alkylators	SG3199 (PBD dimer)	Pyrrolobenzodiazepines (streptomyces species)	DNA minor groove cross-linking	Loncastuximab tesirine (Zynlonta)

The market dominance of NP-derived payloads is evident in commercial ADC therapeutics. As of 2024, twelve ADCs have received FDA approval, with eight achieving this milestone in the last five years alone, signaling a maturation of the field [123]. Monomethyl auristatin E (MMAE) represents one of the most successful NP-derived payloads, capturing 41% of the current ADC payload market share [124]. The auristatins originate from the marine peptide dolastatin 10, isolated from the sea hare *Dolabella auricularia*, demonstrating how exploration of diverse ecological niches yields valuable therapeutic compounds [121].

ADC Payload Market Distribution

The commercial impact of NP-derived payloads extends across target indications and therapeutic areas. **Table 2** quantifies the market distribution of ADC payloads and their clinical applications based on 2024 market data.

Table 2: Market Distribution of ADC Payloads and Applications (2024)

Parameter	Category	Market Share (%)	Projected CAGR (%)	Key NP-Derived Payloads
Payload Type	Monomethyl Auristatin E (MMAE)	41	-	Marine-derived tubulin inhibitor
	Camptothecin derivatives	18	-	Plant-derived topoisomerase I inhibitors
	DM1/DM4 (Maytansinoids)	12	-	Plant-derived tubulin inhibitors
Target Indication	Breast Cancer	44.7	-	T-DM1, T-DXd
	Lung Cancer	-	31.6	T-DXd, Sacituzumab govitecan
Therapeutic Area	Solid Tumors	71	76 (2035 projection)	Various NP-derived payloads
	Hematological Cancers	29	24 (2035 projection)	Auristatins, Calicheamicin

The global ADC market, valued at $12.30 billion in 2024, is projected to reach $28.41 billion by 2035, representing a compound annual growth rate (CAGR) of 6.4% [124]. This growth is substantially fueled by the continued innovation in NP-derived payloads, particularly as applications expand beyond hematological malignancies to dominate solid tumor therapeutics, which currently account for 71% of the ADC market share [124].

Mechanisms of Action and Resistance Pathways

Molecular Mechanisms of NP-Derived Payloads

NP-derived payloads exert their cytotoxic effects through targeting essential cellular processes, with two primary mechanisms dominating the clinical landscape: microtubule disruption and DNA damage.

Microtubule inhibitors, including auristatins (MMAE, MMAF) and maytansinoids (DM1, DM4), bind to tubulin and prevent polymerization into microtubules, disrupting mitotic spindle formation and arresting cell division during mitosis [121]. These compounds demonstrate exceptional potency, with IC50 values in the picomolar to nanomolar range against susceptible tumor cells.

DNA-damaging agents encompass structurally diverse NP-derived compounds including calicheamicins, duocarmycins, and pyrrolobenzodiazepines (PBDs). Calicheamicin, an enediyne antibiotic, binds to the DNA minor groove and generates double-strand breaks via a radical-mediated mechanism [123]. PBD dimers, such as SG3199 in loncastuximab tesirine, form covalent cross-links between opposing strands of DNA, preventing strand separation and essential processes like transcription and replication [123].

The following diagram illustrates the sequential mechanism of action of ADCs from cellular binding to payload-mediated cytotoxicity:



Figure 1: ADC Mechanism of Action from Cellular Binding to Payload-Mediated Cytotoxicity

Resistance Mechanisms to NP-Derived Payloads

Despite their potency, the therapeutic efficacy of NP-derived payloads is often limited by the emergence of resistance. A primary mechanism involves drug efflux mediated by ATP-binding cassette (ABC) transporters, particularly P-glycoprotein (P-gp) [122]. These transmembrane proteins recognize and actively export payloads from tumor cells, reducing intracellular accumulation and diminishing cytotoxicity. Multiple NP-derived payloads, including MMAE, DM1, DM4, and calicheamicin, have been identified as P-gp substrates [122].

Additional resistance mechanisms include:

- **Antigen modulation**: Reduced target antigen expression or mutations in extracellular domains limit ADC binding and internalization [123] [122].
- **Altered intracellular trafficking**: Modifications in lysosomal processing or ADC recycling pathways prevent efficient payload liberation [122].
- **Enhanced DNA repair**: Upregulation of DNA repair pathways counteracts the effects of DNA-damaging payloads [123].

The following diagram illustrates the key resistance mechanisms that impair ADC efficacy:



Figure 2: Key Resistance Mechanisms Limiting ADC Efficacy

Technical Workflows for NP-Derived Payload Evaluation

Payload Potency Assessment

Evaluating the cytotoxicity of NP-derived payloads requires sophisticated bioanalytical approaches due to their exceptional potency. The standard workflow involves multiple complementary techniques:

In vitro cell viability assays form the cornerstone of payload potency assessment. These include:

- **Cell proliferation assays** (MTT, CellTiter-Glo, Alamar Blue) to determine IC50 values
- **Apoptosis analysis** via flow cytometry with Annexin V/propidium iodide staining
- **Cell cycle analysis** to identify mitotic arrest or DNA damage responses

Advanced mechanistic studies provide deeper insights into payload activity:

- **Immunofluorescence microscopy** to visualize microtubule disruption or DNA damage foci
- **Western blotting** to detect activation of DNA damage response pathways (γH2AX, p53)
- **Comet assays** to quantify DNA strand breaks for DNA-damaging payloads

The exceptional potency of NP-derived payloads necessitates highly sensitive detection methods. "Free payload concentrations are typically low, requiring more sensitive, innovative methodologies. Simple sample preparations such as protein precipitation often need to be replaced with more elaborate ones such as liquid-liquid extraction or solid phase extraction, in combination with the use of the most sensitive triple quad mass spectrometers" [125].

Bioanalytical Challenges in Payload Quantification

Accurate measurement of NP-derived payloads presents unique technical challenges that require specialized methodologies:

Chromatographic interference must be carefully managed. "Large molecule ADCs, with their much bigger contact surfaces, will have much more retention and generally will not interfere on a reversed-phase liquid chromatography (LC) system. This is regardless of whether or not the molecule is left intact during sample preparation" [125].

Payload stability considerations are critical for accurate quantification. "ADC concentrations can be orders of magnitude higher than payload concentrations. Due to the excess of ADC in study samples, even the smallest amount of ADC degradation will result in huge payload biases over time" [125].

Sample preparation optimization is essential for reliable results. "Incorporating a high organic flush phase in the mobile phase gradient is necessary to prevent column contamination or interference from slow moving large molecules. This flush phase will wash off the ADC and any other high retentive matrix constituents after each injection" [125].

Emerging Trends: Dual-Payload ADCs and Novel NP Scaffolds

Dual-Payload ADC Platforms

A revolutionary advancement in ADC technology involves the incorporation of two distinct NP-derived payloads within a single conjugate, designed to overcome resistance mechanisms and enhance antitumor efficacy. The field has recently "exploded" with at least 15 dual-payload ADCs disclosed in preclinical presentations as of 2025 [126].

Table 3 highlights promising dual-payload ADC candidates in development:

Table 3: Emerging Dual-Payload ADCs in Development

ADC Candidate	Company	Target	Payload Combination	Development Stage
KH815	Chengdu Kanghong	TROP2	Topo1 inhibitor + RNA pol 2 inhibitor	Phase 1 (first in human)
DXC018	Hangzhou Dac	HER2 x HER2	Topo1 inhibitor + antimetabolite inhibitor	Preclinical
Unnamed	Sutro	Undisclosed	Topo1 inhibitor + PARP inhibitor	Preclinical
JSKN021	Jiangsu Alphamab	EGFR x HER3	Topo1 inhibitor + MMAE	Preclinical
IMD2113	Affinity Biopharmaceutical	EGFR x TROP2	Undisclosed dual mechanism	Preclinical

The rationale for dual-payload strategies centers on overcoming resistance: "patients treated with an ADC can relapse not only through loss of the target antigen, but also by developing resistance to the payload that an ADC uses" [126]. However, significant challenges remain in optimizing linker technology specifically for dual-payload configurations and managing potential increased toxicity profiles [126].

Exploring Untapped NP Chemical Space

Future innovation in NP-derived payloads depends on accessing novel chemical scaffolds from underexplored biological sources. Several promising approaches include:

Marine and extremophile natural products: "Marine NPs are larger and more hydrophobic than terrestrial counterparts, while deep-sea and extremophile-derived NPs show novel scaffolds and bioactivities" [12]. These environments represent rich reservoirs for discovering payloads with unique mechanisms.

AI-enabled NP discovery: "Integrating advanced methodologies, such as artificial intelligence (AI), high-throughput screening, chemical biology, bioinformatics, gene regulation, the highly accurate non-labeling chemical proteomics approach to explore novel NPs targets" [92] will accelerate payload identification.

Addressing NP availability: "The limited availability of NPs (only ~10% purchasable) and the redundancy in known scaffolds pose major challenges in NP research" [12]. Future strategies highlight integrating multidimensional databases and exploring untapped species and extreme environments to uncover unique bioactive compounds [12].

Successful development of ADCs with NP-derived payloads requires specialized reagents and technical capabilities. The following table details essential research tools and their applications:

Table 4: Essential Research Reagents and Resources for ADC Payload Development

Reagent/Resource	Function/Application	Key Considerations
High-Potency Payload Standards (MMAE, DM1, Calicheamicin)	ADC assembly and analytical reference standards	Require specialized containment facilities; typically >95% purity [127]
Cleavable Linkers (Valine-Citrulline, GGFG)	Enable intracellular payload release	Account for ~70% of ADC market; provide plasma stability with tumor-specific cleavage [128]
Site-Specific Conjugation Technologies (Engineered cysteines, unnatural amino acids)	Generate homogeneous ADC products with defined DAR	Growing at >30% CAGR; reduce heterogeneity-related toxicity [128]
Triple Quadrupole Mass Spectrometers	Quantify free payload concentrations in plasma	Essential for detecting low payload levels; newest models offer 3-4x sensitivity improvements [125]
Specialized Chromatography (Reversed-phase with high aqueous mobile phases)	Separate payloads from ADC molecules	Requires high organic flush phases to prevent column contamination [125]
Cell-Based Potency Assays (CellTiter-Glo, MTT)	Determine payload and ADC cytotoxicity	Must include resistant cell lines to assess P-gp mediated efflux [127] [122]
Natural Product Databases (Super Natural II, Dictionary of Natural Products)	Source structural and bioactivity data for NP discovery	Contain >1.1 million compounds; only ~10% are readily purchasable [12]

Natural product-derived payloads continue to dominate the ADC landscape, driven by their unparalleled potency, diverse mechanisms of action, and proven clinical efficacy. The integration of NP chemical space exploration with advanced ADC engineering approaches—including site-specific conjugation, novel linker technologies, and emerging dual-payload strategies—promises to address current limitations in resistance and therapeutic index. Future directions will increasingly leverage AI-enabled NP discovery, exploration of underexplored biological sources, and innovative

bioanalytical methods to quantify payload dynamics with unprecedented precision. As the ADC field continues to mature, NP-derived payloads will remain indispensable components in the targeted therapy arsenal, offering renewed hope for addressing challenging malignancies through their exquisite targeting of fundamental biological processes.

Antimicrobial resistance (AMR) represents one of the most severe global health threats of the 21st century, directly challenging the efficacy of modern medicine. The unchecked use and abuse of traditional antibiotics have precipitated this crisis, leading to increased treatment failures and mortality rates [129]. In response, the **World Health Organization (WHO)** has prioritized the research and development of new antimicrobial agents. **Natural Products (NPs)**, with their vast and evolutionarily refined chemical diversity, have emerged as a beacon of hope. This whitepaper details how the systematic exploration of **natural product chemical space** provides powerful, innovative strategies to reconstruct the antibiotic pipeline and combat AMR [129] [30]. We outline the most promising NP-inspired approaches, provide detailed experimental methodologies, and visualize the critical pathways and workflows, offering a technical guide for researchers and drug development professionals.

The AMR Crisis and the WHO Priority Pathogens

The AMR crisis is exacerbated by the rapid development of resistance mechanisms in bacteria, including drug efflux pumps, modification of antibiotic targets, and enzymatic degradation of the drugs themselves [130]. The COVID-19 pandemic has further intensified this threat, leading to constrained antibiotic treatment options and surging resistance rates [131].

To focus global research efforts, the WHO has established a **Bacterial Priority Pathogens List**, categorized by urgency level [131]:

Critical Priority

- *Acinetobacter baumannii* (carbapenem-resistant)
- *Enterobacterales* (carbapenem-resistant and 3rd-generation cephalosporin-resistant)

High Priority

- *Pseudomonas aeruginosa* (carbapenem-resistant)
- *Staphylococcus aureus* (methicillin-resistant, MRSA)
- *Enterococcus faecium* (vancomycin-resistant)
- *Salmonella Typhi* (fluoroquinolone-resistant)
- *Neisseria gonorrhoeae* (cephalosporin and/or fluoroquinolone-resistant)

Medium Priority

- *Streptococcus pneumoniae* (macrolide-resistant)
- *Haemophilus influenzae* (ampicillin-resistant)
- *Group A/B Streptococci* (macrolide/penicillin-resistant)

This list serves as a crucial guide for targeting research on novel antimicrobial agents, with NPs showing significant activity against these resilient pathogens [131].

Exploring Natural Product Chemical Space: A Strategic Framework

The chemical space of NPs is inherently "biologically relevant," as these molecules have evolved through natural selection to interact with biological macromolecules [132]. This makes them ideal starting points for drug discovery. Several sophisticated strategies have been developed to systematically explore this space, moving beyond simply isolating novel compounds from nature.

Biology-Oriented Synthesis (BIOS)

Biology-Oriented Synthesis (BIOS) uses the structural scaffolds of known NPs as inspiration. The core principle is the systematic simplification of complex NP scaffolds into core structures that retain biological relevance but are synthetically more accessible [133] [30].

- **Workflow**: NP Scaffold → Computational Simplification (e.g., using SCONP algorithm or Scaffold Hunter) → Design of Synthetically Tractable, NP-inspired Scaffolds → Synthesis of Focused Libraries → Biological Screening.
- **Application Example**: Inspired by the NP sodwanone S, a library based on a bicyclic oxepane scaffold was synthesized. Screening identified "Wntepane," a novel small-molecule activator of the Wnt pathway that acts by binding to the protein Vangl1, a previously undrugged target [133].

Diversity-Oriented Synthesis (DOS)

Diversity-Oriented Synthesis (DOS) aims to rapidly generate libraries of complex and structurally diverse small molecules that populate broad regions of chemical space, mimicking the structural complexity of NPs [133] [30].

- **Workflow**: Pluripotent Building Block → Branching Reaction Pathways (e.g., cycloadditions, dihydroxylation) → Diverse Molecular Scaffolds → Library Synthesis → Phenotypic/Target-Agnostic Screening.
- **Application Example**: A DOS library of 242 molecules based on 18 distinct natural product-like scaffolds was synthesized from a solid-supported phosphonate. This led to the discovery of "gemmacin," a novel broad-spectrum antibiotic with potent activity against MRSA and lower cytotoxicity against human cells [39].

Complexity-to-Diversity (CtD)

Complexity-to-Diversity (CtD) leverages readily available NPs as complex starting materials and uses chemoselective reactions to dramatically rearrange their core structures, generating unprecedented scaffolds [133].

- **Workflow**: Readily Available NP (e.g., gibberellic acid, yohimbine) → Chemoselective Reactions (Ring Cleavage, Expansion, Fusion, Rearrangement) → Novel, Diverse Molecular Scaffolds.
- **Application Example**: The diterpene abietic acid was transformed through a series of ring-cleavage and rearrangement reactions into novel scaffolds with high fraction of sp^3 carbons and structural diversity. Libraries derived from such processes have yielded compounds with anti-inflammatory, anti-proliferative, and antibacterial activity [133].

The following diagram illustrates the logical relationship between the NP chemical space and these core exploration strategies.



Detailed Experimental Protocols

This section provides detailed methodologies for key experiments cited in this review, serving as a technical reference for researchers aiming to replicate or build upon these findings.

Systematic Review of NPs Against WHO Priority Pathogens

Objective: To identify and evaluate the efficacy of natural products against antibiotic-resistant "priority pathogens" as defined by the WHO [131].

Methodology:

- **Search Strategy**: A comprehensive literature search is conducted across major databases (PUBMED/MEDLINE, WEB OF SCIENCE, SCOPUS) using customized search strings combining keywords and MeSH terms such as `("natural product*" OR "natural compound*") AND (antibacteri* OR antimicrobial*) AND (MDR OR "multi-drug resistant *")`.
- **Study Selection**: Two independent reviewers screen titles and abstracts against pre-defined inclusion criteria (e.g., original studies, exposure to natural products, outcome of antimicrobial effects against priority pathogens, English language). A third reviewer resolves conflicts.
- **Data Extraction**: A standardized form is used to collect data on authors, publication year, study design, natural product source, plant part used, extraction solvent, target pathogen, and quantitative measures of antimicrobial activity (e.g., MIC, MBC).
- **Quality Assessment**: The methodological rigor of included studies is assessed using tools like the Newcastle–Ottawa Scale for observational studies and the Cochrane Risk of Bias tool for randomized controlled trials.
- **Data Synthesis**: Findings are summarized to identify the most promising NP classes (e.g., alkaloids, flavonoids) and their effectiveness against specific WHO priority pathogens.

Biology-Oriented Synthesis (BIOS) Workflow

Objective: To discover novel bioactive molecules by synthesizing and screening libraries based on simplified NP scaffolds [133] [30].

Methodology:

- **Scaffold Selection**: Use computational algorithms like SCONP or the Scaffold Hunter tool to identify simplified, synthetically feasible sub-structures derived from complex NPs with known bioactivity.
- **Route Design & Validation**: Develop a multistep, often one-pot or solid-phase, synthetic sequence that allows for the efficient production of the target scaffold and its derivatives. The route should incorporate points for diversification.
- **Library Synthesis**: Execute the synthetic plan to produce a focused library of compounds (typically 30-100+ derivatives). Parallel derivatization at specific positions on the scaffold is employed to explore structure-activity relationships (SAR).
- **Biological Screening**: Screen the compound library against a specific biological target or pathway using reporter gene assays, enzymatic assays, or phenotypic screens (e.g., for Wnt or Hedgehog pathway modulation).
- **Hit Validation & Target Identification**: For active compounds (hits), validate activity through dose-response curves (IC_{50}/EC_{50}). Use chemical biology tools like biotinylated analogues for immunoenrichment and pull-down experiments to identify the protein target.

Evaluation of Antimicrobial and Anti-Virulence Activity

Objective: To assess the direct antibacterial and anti-virulence properties of NPs and their derivatives [134].

Methodology:

- **Direct Antimicrobial Activity**:
  - **MIC/MBC Determination**: Use standardized broth microdilution or agar well diffusion methods according to CLSI guidelines to determine the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) against target pathogens (e.g., MRSA, *P. aeruginosa*).
- **Anti-Virulence and Synergy Studies**:
  - **Biofilm Assay**: Treat bacterial cultures with sub-inhibitory concentrations of the NP and quantify biofilm formation using crystal violet staining or other metabolic assays.
  - **Synergy Testing**: Evaluate the combination of NPs with conventional antibiotics using the checkerboard microdilution method to calculate the Fractional Inhibitory Concentration (FIC) index, where FIC ≤ 0.5 indicates synergy.
  - **Mechanistic Studies**: Perform molecular docking to predict interactions with key virulence targets (e.g., *C. albicans* CaCYP51). Use gene expression analysis (qRT-PCR) to measure the downregulation of virulence genes (e.g., hyphae-specific genes in *Candida*).

The Scientist's Toolkit: Essential Reagents and Solutions

The following table details key reagents, materials, and computational tools essential for research in NP-based antimicrobial discovery.

Research Reagent / Solution	Function / Application
Ethanol, Methanol, Ethyl Acetate	Common solvents for the extraction of bioactive compounds (e.g., alkaloids, flavonoids, terpenoids) from plant materials [131].
Broth Microdilution Plates	High-throughput platform for determining Minimum Inhibitory Concentrations (MICs) against bacterial and fungal pathogens [131] [134].
Crystal Violet Stain	Dye used to quantify total biofilm biomass formed by bacteria on abiotic surfaces after treatment with test compounds [134].
Biotinylated Analogues	Chemical probes derived from hit compounds; used for target identification via pull-down assays and immunoenrichment [133].
SCONP / Scaffold Hunter	Computational algorithms for the systematic analysis and simplification of natural product scaffolds to guide BIOS library design [133].
PHASE (Schrödinger)	Software module used for pharmacophore modeling and 3D-QSAR studies to identify crucial structural features for activity and guide molecular design [135] [136].
Solid-Phase Synthesis Resin	Polymeric support (e.g., silyl-polystyrene) used for DOS and other library synthesis, enabling rapid purification and automation [133] [30].

Visualizing Key Signaling Pathways and Mechanisms

Understanding the biological pathways targeted by NPs and the mechanisms of antibiotic resistance is crucial for rational drug design. The diagram below illustrates a key signaling pathway that can be modulated by NP-inspired molecules and the primary mechanisms bacteria use to resist antibiotics.



The exploration of **natural product chemical space** represents a paradigm shift in the fight against antimicrobial resistance. Strategies such as **Biology-Oriented Synthesis**, **Diversity-Oriented Synthesis**, and **Complexity-to-Diversity** provide systematic, rational frameworks to discover novel bioactive molecules that bypass conventional resistance mechanisms [129] [133] [30]. The continued integration of these approaches with advances in synthetic chemistry, computational biology, and chemical genomics is paramount. By leveraging NPs as both inspiration and starting points, the scientific community can reconstruct the antibiotic pipeline, transforming this beacon of hope into a new arsenal of effective therapies to safeguard global health for future generations.

The exploration of natural products for drug discovery represents a journey through an expansive **chemical multiverse**, a term introduced to describe the comprehensive analysis of compound datasets through several distinct chemical spaces, each defined by a different set of chemical representations [137]. Unlike a single, unified chemical space, the chemical multiverse acknowledges that a given set of molecules represented with different descriptors leads to distinct chemical universes, each providing complementary insights into molecular structure and properties [137]. This conceptual framework is particularly relevant to the study of multi-compound extracts from medicinal plants, which are complex, adaptive systems whose therapeutic efficacy often emerges from **synergistic interactions** between their numerous bioactive constituents [138]. The intrinsic complexity of these extracts means that their overall activity cannot be predicted from the activity of isolated compounds alone, as they contain hundreds or even thousands of individual bioactive molecules in varying abundances [138].

Within this chemical multiverse, synergistic interactions between compounds become a vital part of therapeutic efficacy [138]. **Synergy** occurs when the combined effect of compounds is greater than the sum of their individual effects, potentially arising through multiple mechanisms including multi-target effects, enhanced bioavailability, or protection from toxicity [138] [139]. This review explores the theoretical foundations, experimental methodologies, and practical applications of synergistic effects in multi-compound natural extracts, framing this exploration within the broader context of navigating natural product chemical space for drug discovery research.

Theoretical Foundations of Synergistic Interactions

Defining Combination Effects

The interactions between multiple bioactive compounds in natural extracts can be systematically classified based on the observed effect relative to the expected effect from individual components. These classifications provide the vocabulary for describing combination effects:

- **Synergistic Effect**: The combined effect of compounds is greater than the sum of their individual effects [138]. This may also occur when one compound enhances the therapeutic effect of another by regulating its absorption, distribution, metabolism, and excretion, or when individually inactive compounds become active when combined [138].
- **Additive Effect**: The combined effect of two compounds equals the sum of their individual effects [138] [139]. It is crucial to note that additive effects are not simply the sum of effects but are computed from individual effects based on complex mathematical algorithms [138].
- **Antagonistic Effect**: An interaction that results in less than the sum of the effects of the individual compounds [138] [139]. This reduced effect from expectation can be valuable in mitigating toxicity or adverse effects [138].

Table 1: Classification and Definitions of Combination Effects

Interaction Type	Mathematical Relationship	Therapeutic Implication
Synergy	Combined effect > Sum of individual effects	Enables lower doses, reduces adverse effects, enhances efficacy
Additivity	Combined effect = Sum of individual effects	Straightforward dose combination without interaction
Antagonism	Combined effect < Sum of individual effects	May reduce toxicity or undesirable effects

Molecular Mechanisms Underlying Synergy

The superior therapeutic performance of multi-compound extracts compared to isolated constituents can be explained by several fundamental mechanisms through which synergy manifests:

- **Multi-Target Effects (Pharmacodynamic Synergism)**: Different compounds in a mixture simultaneously engage multiple therapeutic targets or pathways relevant to a disease state [139]. This network-level intervention is particularly valuable for complex, multifactorial diseases where single-target approaches often prove insufficient [138].

- **Enhanced Bioavailability (Pharmacokinetic Synergism)**: Certain compounds may improve the absorption, distribution, or metabolic stability of other active compounds in the mixture, thereby increasing their bioavailability and therapeutic concentration [139]. Some natural compounds, while not possessing direct effects themselves, may increase the solubility or inhibit the metabolism of co-administered active compounds [139].

- **Attenuation of Adverse Effects**: Compounds within an extract may interact to reduce the toxicity or side effects associated with individual constituents while maintaining therapeutic efficacy [138]. This protective synergy allows for the use of potentially toxic but highly effective compounds with a improved safety profile [138].



Diagram 1: Fundamental mechanisms of therapeutic synergy in multi-compound extracts

Experimental Design and Methodological Framework

Key Methodologies for Assessing Combination Effects

Rigorous experimental design is essential for accurately identifying and quantifying synergistic interactions in natural product research. Several well-established methodologies enable researchers to distinguish true synergy from simple additive effects:

- **Isobolographic Analysis**: This classical graphical method involves constructing an "isobole" - a line connecting the doses of two individual compounds that each produce the same specified effect level (typically the ED50) [140]. The combined doses that fall below this line indicate

synergy, while points above the line indicate antagonism [140]. The method is based on the concept of **dose equivalence**, where the drug B-equivalent of dose a is calculated as aB/A, leading to the fundamental isobole equation: $a/A + b/B = 1$ [140].

- **Combination Index (CI) Method**: This quantitative approach calculates a combination index to determine the nature of drug interactions [139]. The CI is defined by the equation: $CI = d_1/Dx_1 + d_2/Dx_2$, where d_1 and d_2 are the respective combination doses of drug one and drug two that produce an effect x, and Dx_1 and Dx_2 are the corresponding single doses for drug one and drug two that result in the same effect x [139]. CI values < 1 indicate synergy, CI = 1 indicates additivity, and CI > 1 indicates antagonism [139].
- **Universal Surface Response Analysis**: This statistical method provides a comprehensive estimate of differentiation between synergy, additivity, and antagonism across a range of concentration combinations, offering a more complete interaction profile than single-point measurements [139].

Table 2: Comparison of Major Methodologies for Synergy Detection

Method	Key Principle	Output	Advantages	Limitations
Isobolographic Analysis [140]	Dose equivalence for specified effect level	Graphical representation (isobole)	Intuitive visualization; Clear interpretation	Limited to two compounds at a time; Fixed effect level
Combination Index (CI) [139]	Summation of fractional doses	Numerical index (CI < 1, =1, >1)	Quantitative results; Applicable to multiple compounds	Requires full dose-response curves for each agent
Universal Surface Response [139]	Statistical modeling of response surface	Three-dimensional interaction profile	Comprehensive across concentration ranges	Complex experimental design and analysis

Experimental Workflow for Synergy Screening

A systematic approach to screening synergistic interactions in natural product extracts ensures comprehensive and reproducible results. The following workflow outlines key stages in this process:



Diagram 2: Systematic workflow for screening synergistic interactions

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Essential Research Reagents and Materials for Synergy Studies

Reagent/Material	Function in Synergy Research	Application Notes
Standardized Natural Extracts	Provides consistent, reproducible starting material for combination studies	Chemical fingerprinting recommended; Source and batch documentation critical [138]

Reagent/Material	Function in Synergy Research	Application Notes
Purified Bioactive Compounds	Enables controlled combination studies with known constituents	High purity (>95%) essential for accurate dose-response characterization [138]
Cell-Based Assay Systems	Initial screening platform for combination effects	Include relevant cell lines; Multiple assay endpoints recommended [138] [139]
Reference Standards (e.g., controls, calibration standards)	Ensures analytical validity and experimental consistency	Include positive/negative controls for biological assays [140]
Analytical Grade Solvents	Extraction, purification, and solubilization of compounds	Low UV cutoff for HPLC applications; Mass spectrometry compatibility [138]

Signaling Pathways and Molecular Mechanisms

The therapeutic advantage of multi-compound extracts often derives from their ability to simultaneously modulate multiple interconnected signaling pathways, creating a network response that is difficult to achieve with single compounds. The following diagram illustrates key pathways frequently engaged by synergistic natural product combinations:



Diagram 3: Key signaling pathways modulated by synergistic natural product combinations

Research has demonstrated that medicinal plant extracts target biological systems through the combined action of structurally and functionally diverse active compounds that modulate complex cellular networks [138]. This multi-target approach is particularly evident in antimicrobial applications, where combinations of bioactive compounds can simultaneously disrupt cell membrane integrity, inhibit essential enzymes, and impair cellular energy production, resulting in enhanced efficacy and reduced potential for resistance development [138]. The **polyvalent** nature of these extracts denotes an improved and cooperative effect that cannot be easily attributed to single mechanisms [138].

Applications in Drug Discovery and Development

Antimicrobial Applications

Natural product combinations show significant promise in addressing the growing challenge of antimicrobial resistance. Studies have demonstrated that **whole plant preparations are frequently more effective than isolated compounds** due to synergistic interactions between constituents within them [138]. These combinations can produce enhanced antimicrobial effects through several mechanisms:

- **Multi-target Engagement**: Different compounds in a mixture may simultaneously target cell wall synthesis, membrane integrity, and intracellular processes in microorganisms [138].
- **Resistance Mitigation**: The probability of resistance development decreases significantly when microbes face multiple antimicrobial challenges simultaneously [138].
- **Bioavailability Enhancement**: Certain compounds in mixtures can improve the penetration or activity of co-occurring antimicrobial agents [138].

Research on medicinal plant extracts has revealed that disease resistance is less likely to occur against a combination of bioactive compounds than against single active molecules, highlighting the strategic advantage of multi-component antimicrobial approaches [138].

Chronic Disease Management

Combinations of marine and plant-derived bioactive compounds have demonstrated significant potential for managing chronic non-communicable diseases, including metabolic, inflammatory, and age-related conditions [139]. These combinations work through several synergistic mechanisms:

- **Anti-inflammatory Effects**: Multi-compound extracts can simultaneously inhibit multiple pro-inflammatory pathways (e.g., NF-κB, COX-2, cytokine signaling), resulting in enhanced anti-inflammatory activity compared to single compounds [139].
- **Antioxidant Protection**: Combinations of compounds with complementary antioxidant mechanisms (e.g., free radical scavenging, metal chelation, enzyme induction) provide enhanced protection against oxidative stress [139].
- **Metabolic Regulation**: Bioactive combinations can modulate complex metabolic networks through coordinated effects on multiple signaling pathways (e.g., AMPK, PPARs, insulin signaling) [139].

The development of marine-based functional foods and nutraceuticals represents a particularly promising application, with research showing that combinations of marine bioactive compounds can produce synergistic effects that enhance their preventive and therapeutic potential against chronic diseases [139].

The study of synergistic effects in multi-compound natural extracts represents a paradigm shift in natural product-based drug discovery, moving from reductionist single-compponent approaches to a more holistic systems-level understanding. The **chemical multiverse** concept provides a valuable framework for navigating the complex chemical space of natural products, acknowledging that comprehensive assessment requires multiple complementary representations of chemical structure and properties [137]. This approach aligns with the inherent complexity of natural extracts, whose therapeutic advantages frequently emerge from synergistic interactions between their numerous constituents [138].

Future research in this field should prioritize several key areas: First, developing more sophisticated computational models to predict synergistic interactions based on chemical structures and known biological activities. Second, advancing analytical techniques, particularly metabolomics approaches, to better characterize complex mixtures and identify interaction networks [138]. Third, establishing standardized methodologies and reporting guidelines for synergy research to improve reproducibility and comparability across studies [140] [139]. Finally, increasing efforts to translate in vitro synergy findings to validated clinical outcomes, particularly for complex chronic conditions where multi-target approaches offer distinct advantages over monotherapies [138] [139].

As drug discovery faces increasing challenges with single-target approaches, the strategic exploration of synergistic multi-compound extracts within the natural product chemical multiverse offers a promising path forward. By embracing the complexity of natural extracts rather than attempting to reduce it, researchers can harness the full therapeutic potential of these evolved chemical systems, potentially leading to more effective, safer, and more resistance-resistant therapeutic interventions.

Conclusion

The exploration of the natural product chemical space remains a cornerstone of innovative drug discovery, uniquely positioned to deliver the structural diversity and biological relevance needed to tackle complex diseases. The integration of foundational concepts with advanced methodologies—from AI and high-throughput screening to synthetic biology—is successfully overcoming historical challenges of supply and characterization. The proven track record of NPs, particularly in generating leads for antimicrobial and anticancer agents, validates their continued strategic importance. Future success hinges on a collaborative, multidisciplinary approach that fully embraces technological revolutions, adheres to ethical and regulatory frameworks, and systematically populates the underexplored regions of the biologically relevant chemical space to usher in a new era of therapeutics.

References

- 1. Frontiers | On the biologically relevant chemical : space BioReCS [https://www.frontiersin.org/journals/drug-discovery/articles/10.3389/fddsv.2025.1674289/full]
- 2. as a source for new Chemical - MedChemComm... space drugs [https://pubs.rsc.org/en/content/articlehtml/2010/md/c0md00020e]
- 3. -informed Natural of product to enable... exploration chemical space [https://pubmed.ncbi.nlm.nih.gov/34046620/]
- 4. Relevance of the Trillion-Sized Chemical “eXplore” as a Source... Space [https://pmc.ncbi.nlm.nih.gov/articles/PMC10108389/]
- 5. Chemical space visual navigation in the era of deep learning and Big Data - ScienceDirect [https://www.sciencedirect.com/science/article/pii/S1359644625001059]
- 6. Quantitative Methods in System-Based Drug Discovery | IntechOpen [https://www.intechopen.com/chapters/53186]
- 7. Network-based multi-omics integrative analysis methods in drug ... [https://biodatamining.biomedcentral.com/articles/10.1186/s13040-025-00442-z]
- 8. Exploring chemical space in non-targeted analysis: a proposed ChemSpace tool - PubMed [https://pubmed.ncbi.nlm.nih.gov/36435841/]
- 9. Finding a Better Way to Explore Chemical Space: A Proposed ChemSpace Tool | Research Highlight | PNNL [https://www.pnnl.gov/publications/finding-better-way-explore-chemical-space-proposed-chemspace-tool]
- 10. Frontiers | Core publications in drug discovery and natural product research [https://www.frontiersin.org/journals/natural-products/articles/10.3389/fntpr.2024.1493720/full]
- 11. The pharmaceutical industry and natural : products status... historical [https://link.springer.com/article/10.1007/s11101-014-9367-z]
- 12. Navigating the chemical of space from diverse... natural products [https://link.springer.com/article/10.1007/s11101-025-10198-3]
- 13. Artificial Intelligence in Natural : Product ... Drug Discovery Current [https://pmc.ncbi.nlm.nih.gov/articles/PMC11874025/]

- 14. Lead Optimization in Drug Discovery | Danaher Life Sciences [https://lifesciences.danaher.com/us/en/library/lead-optimization-drug-discovery-guide.html]
- 15. Analog Design Mastery: Chemistry of Natural Products [https://www.numberanalytics.com/blog/analog-design-mastery-chemistry-natural-products]
- 16. Using PCA and TMAP to visualize the space of synthetic vs ... [https://mihir-parekh.medium.com/using-pca-and-tmap-to-visualize-the-space-of-synthetic-vs-natural-molecules-63cd86922fd0]
- 17. Natural products in drug discovery: advances and ... [https://www.nature.com/articles/s41573-020-00114-z]
- 18. Expanding the Range of 'Druggable' Targets with Natural ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC2878877/]
- 19. Time-Dependent Comparison of the Structural Variations ... [https://www.mdpi.com/1422-0067/25/21/11475]
- 20. Unifying principles for the design and evaluation of natural ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC11788825/]
- 21. Art driven by visual representations of chemical space [https://jcheminf.biomedcentral.com/articles/10.1186/s13321-023-00770-4]
- 22. Principal Component Analysis as a Tool for Library Design [https://pmc.ncbi.nlm.nih.gov/articles/PMC4373534/]
- 23. sciencedirect.com/topics/ chemistry / natural - product - chemistry [https://www.sciencedirect.com/topics/chemistry/natural-product-chemistry]
- 24. Charting, navigating, and populating natural product chemical space for drug discovery - PubMed [https://pubmed.ncbi.nlm.nih.gov/22537178/]
- 25. HZI | Natural from products pathways and extreme... underexplored [https://www.helmholtz-hzi.de/en/research/research-groups/details/natural-products-from-underexplored-pathways-and-extreme-environments/]
- 26. Charting biologically relevant chemical space: a structural... [https://pubmed.ncbi.nlm.nih.gov/16301544/]
- 27. 30.1: Classification of Natural - Products LibreTexts Chemistry [https://chem.libretexts.org/Bookshelves/Organic_Chemistry/Basic_Principles_of_Organic_Chemistry_(Roberts_and_Caserio)/30%3A_Natu
- 28. The exploration of macrocycles for... | Nature Reviews Drug Discovery [https://www.nature.com/articles/nrd2590?error=cookies_not_supported&code=b6e8f81c-19d0-46f5-ba2b-9bb3b6129e68]
- 29. with Active Learning and Alchemical... Chemical Space Exploration [https://pmc.ncbi.nlm.nih.gov/articles/PMC9558370/]
- 30. -inspired strategies towards the Natural of product ... discovery novel [https://fjps.springeropen.com/articles/10.1186/s43094-024-00627-z]
- 31. 67 million natural product-like compound database generated via molecular language processing | Scientific Data [https://www.nature.com/articles/s41597-023-02207-x]
- 32. Bridging the Gap Between Natural and Product ... Synthesis Drug [https://pmc.ncbi.nlm.nih.gov/articles/PMC7718793/]
- 33. Accelerating drug discovery with a single carbon atom | ScienceDaily [https://www.sciencedaily.com/releases/2025/05/250506131509.htm]
- 34. The behind-the-scenes chemistry driving drug discovery | St. Jude Research [https://www.stjude.org/research/progress/2025/the-behind-the-scenes-chemistry-driving-drug-discovery.html]
- 35. Artificial Intelligence in Natural Product Drug Discovery: Current Applications and Future Perspectives - PubMed [https://pubmed.ncbi.nlm.nih.gov/39916476/]
- 36. Frontiers | Exploring chemical for “druglike” small molecules in... space [https://www.frontiersin.org/journals/molecular-biosciences/articles/10.3389/fmolb.2025.1553667/full]
- 37. Rapid traversal of vast chemical space using machine learning-guided docking screens | Nature Computational Science [https://www.nature.com/articles/s43588-025-00777-x]
- 38. AI's emerging role in natural product drug discovery | CAS [https://www.cas.org/resources/cas-insights/ais-emerging-role-in-natural-product-drug-discovery]
- 39. AI-Integrated QSAR Modeling for Enhanced Drug Discovery: From Classical Approaches to Deep Learning and Structural Insight [https://www.mdpi.com/1422-0067/26/19/9384]
- 40. Explore drug-like space with deep generative models - PubMed [https://pubmed.ncbi.nlm.nih.gov/36682423/]
- 41. Tackling the issue of confined chemical space with AI-based de novo drug design and molecular optimization - PubMed [https://pubmed.ncbi.nlm.nih.gov/40887314/]
- 42. Top 5 Drug Discovery Trends 2025 Driving Breakthroughs [https://www.pelagobio.com/cetsa-drug-discovery-resources/blog/drug-discovery-trends-2025/]
- 43. Digital Alchemy: The Rise of Machine and Deep Learning in Small-Molecule Drug Discovery - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC12295739/]
- 44. - High of Natural Product and Synthetic... Throughput Screening [https://pmc.ncbi.nlm.nih.gov/articles/PMC10220967/]
- 45. High-Throughput Screening: today’s biochemical and cell-based approaches - ScienceDirect [https://www.sciencedirect.com/science/article/abs/pii/S1359644620303056]
- 46. Cell-based high-throughput screens for the discovery of chemotherapeutic agents - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC3388188/]
- 47. High-throughput screening - Wikipedia [https://en.wikipedia.org/wiki/High-throughput_screening]
- 48. in cancer... | Phenotypic Reviews screening Nature Drug Discovery [https://www.nature.com/articles/nrd4366?error=cookies_not_supported&code=2ade43ce-96a1-4c6a-9534-079a177e6138]

- 49. Novel Chemical Space Exploration via Natural Products - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC2696019/]
- 50. Novel chemical space exploration via natural products - PubMed [https://pubmed.ncbi.nlm.nih.gov/19265440/]
- 51. in Phenotypic : An... | Technology Networks Screening Drug Discovery [https://www.technologynetworks.com/drug-discovery/articles/phenotypic-screening-a-powerful-tool-for-drug-discovery-398572]
- 52. Application of Combination High-Throughput Phenotypic ... Screening [https://pubmed.ncbi.nlm.nih.gov/28510271/]
- 53. Phenotypic screens in antimalarial drug discovery - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC5007148/]
- 54. A phenomics approach for antiviral drug discovery | BMC Biology | Full Text [https://bmcbiol.biomedcentral.com/articles/10.1186/s12915-021-01086-1]
- 55. leveraging big data in plant natural products biosynthesis ... [https://pubmed.ncbi.nlm.nih.gov/40545552/]
- 56. Metabolomics and Genomics in Natural Products Research [https://pmc.ncbi.nlm.nih.gov/articles/PMC8691422/]
- 57. MEANtools integrates multi-omics data to identify ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12327601/]
- 58. Metabolomics and lipidomics strategies in modern drug ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC10543515/]
- 59. Merging metabolomics and genomics provides a catalog of ... [https://gsejournal.biomedcentral.com/articles/10.1186/s12711-025-00960-8]
- 60. Building Natural –Based Product for Drug Discovery... Libraries [https://link.springer.com/article/10.1007/s43450-024-00540-9]
- 61. Fragment Libraries from Large and Novel Synthetic Compounds and... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12044453/]
- 62. | NCCIH Natural Product Libraries [https://www.nccih.nih.gov/grants/natural-product-libraries]
- 63. Pseudo- natural as next-generation scaffolds: redefining the... products [https://pubmed.ncbi.nlm.nih.gov/41215729/]
- 64. Frontiers | Affinity selection mass spectrometry (AS-MS) for prospecting... [https://www.frontiersin.org/journals/natural-products/articles/10.3389/fntpr.2025.1562501/full]
- 65. strategies for In : optimizing Silico compounds... drug discovery natural [https://link.springer.com/article/10.1007/s44371-025-00201-3]
- 66. In silico methods and tools for drug discovery - PubMed [https://pubmed.ncbi.nlm.nih.gov/34520990/]
- 67. Computer-aided Drug Approaches in the Discovery of... Identification [https://pubmed.ncbi.nlm.nih.gov/38698753/]
- 68. Natural product-informed exploration of chemical space to enable bioactive molecular discovery - RSC Medicinal Chemistry (RSC Publishing) [https://pubs.rsc.org/en/content/articlelanding/2021/md/d0md00376j]
- 69. ARTICLE: BIOPTIC B1 Ultra-High-Throughput Virtual ... Screening [https://bioptic.io/science/article-bioptic-b1-lrrk2-jcim-2025]
- 70. Directory of in silico Drug Design tools [https://www.click2drug.org/]
- 71. In silico drug discovery: a machine learning-driven systematic review | Medicinal Chemistry Research [https://link.springer.com/article/10.1007/s00044-024-03260-w]
- 72. — ASE documentation Molecular dynamics [https://ase-lib.org/ase/md.html]
- 73. Settings of Benchmarking ... Parameter Molecular Dynamics [https://link.springer.com/article/10.1007/s40735-024-00891-9]
- 74. High impact technologies for natural products screening [https://pubmed.ncbi.nlm.nih.gov/18084916/]
- 75. NMR-spectroscopic analysis of mixtures: from structure to function - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC3053045/]
- 76. | SpringerLink Natural Product Isolation [https://link.springer.com/protocol/10.1385/1-59259-955-9:1]
- 77. Molecules | Special Issue : Analytical Methods for Characterization and Isolation of Natural Products [https://www.mdpi.com/journal/molecules/special_issues/natural_products_isolation]
- 78. Integrating Chemical and Biological Profiling for the Functional Annotation of Complex Natural Product Mixtures | NCCIH [https://www.nccih.nih.gov/news/events/integrating-chemical-and-biological-profiling-for-the-functional-annotation-of-complex-natural-product-mixtures]
- 79. and Discovery of pharmacologically active plant-derived... resupply [https://pubmed.ncbi.nlm.nih.gov/26281720/]
- 80. Natural Product-like Compound Libraries for screening | Screening Libraries | Life Chemicals [https://lifechemicals.com/screening-libraries/natural-product-like-compound-library]
- 81. for pharmaceutical Synthetic | DDDT biology drug discovery [https://www.dovepress.com/synthetic-biology-for-pharmaceutical-drug-discovery-peer-reviewed-fulltext-article-DDDT]
- 82. Enzyme Discovery via Genome Mining Unlocks Stereodivergence [https://bioengineer.org/enzyme-discovery-via-genome-mining-unlocks-stereodivergence/]
- 83. Compound Libraries for High Throughput/Content Screening | 96-Well [https://www.selleckchem.com/screening/natural-product-library.html]
- 84. Exploring chemical space for “druglike” small molecules in the age of AI - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC11955463/]
- 85. | Aston University Nagoya Protocol [https://www.aston.ac.uk/research/integrity-ethics/nagoya-protocol]
- 86. Biological Diversity (Access And Benefit) Sharing Regulation 2025 [https://visionias.in/current-affairs/monthly-magazine/2025-06-17/environment/biological-diversity-access-and-benefit-sharing-regulation-2025]
- 87. The Nagoya on Protocol and Access - Benefit sharing [https://www.cbd.int/abs]
- 88. Creating and screening natural product libraries - Natural Product Reports (RSC Publishing) DOI:10.1039/C9NP00068B [https://pubs.rsc.org/en/content/articlehtml/2020/np/c9np00068b]
- 89. Recent Advances in Machine-Learning-Based Chemoinformatics: A Comprehensive Review - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC10380192/]

- 90. ChemSAR: an online pipelining platform for molecular SAR modeling | Journal of Cheminformatics | Full Text [https://jcheminf.biomedcentral.com/articles/10.1186/s13321-017-0215-1]
- 91. DeepAutoQSAR | Schrödinger Machine Learning Solutions [https://www.schrodinger.com/platform/products/deepautoqsar/]
- 92. The latest advances with natural products in drug discovery and opportunities for the future: a 2025 update - PubMed [https://pubmed.ncbi.nlm.nih.gov/40391763/]
- 93. Exploring Chemical Space for Drug Discovery Using the Chemical Universe Database - PMC [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3447393/]
- 94. sciencedirect.com/topics/agricultural-and-biological-sciences/ chemotype [https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/chemotype]
- 95. Molecular Diversity Assessment using Chemotypes [https://pubmed.ncbi.nlm.nih.gov/33535959/]
- 96. Chemical Space Exploration with Active Learning and Alchemical Free Energies - PubMed [https://pubmed.ncbi.nlm.nih.gov/36148968/]
- 97. Frontiers | Hybrid Dynamic Pharmacophore Models as Effective Tools... [https://www.frontiersin.org/journals/chemistry/articles/10.3389/fchem.2020.596412/full]
- 98. Contemporary methods for the extraction and isolation of natural ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC10314546/]
- 99. Stress Driven Discovery of Natural From Actinobacteria... Products [https://pmc.ncbi.nlm.nih.gov/articles/PMC8584265/]
- 100. tools in the digital era: Applications and limitations | CoLab ADMET [https://colab.ws/articles/10.1016%2Fbs.apha.2025.01.004]
- 101. Computer-Aided Drug Design Methods - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC5248982/]
- 102. GitHub - DonghaiZHAO-ZJU/ Benchmark - ADMET - 2025 [https://github.com/DonghaiZHAO-ZJU/Benchmark-ADMET-2025]
- 103. Creating and screening natural - PMC product libraries [https://pmc.ncbi.nlm.nih.gov/articles/PMC8494140/]
- 104. Use of Natural as Products Library for Chemical and... Drug Discovery [https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0062839]
- 105. Natural product-derived compounds in clinical trials and drug approvals - PubMed [https://pubmed.ncbi.nlm.nih.gov/40923383/]
- 106. Artemisinin and Its Derivatives as Potential Anticancer Agents | MDPI [https://www.mdpi.com/1420-3049/29/16/3886]
- 107. BIOCELL | Special Issues: Natural -Based Product ... Anticancer Drug [https://www.techscience.com/biocell/special_detail/anticancer_drug_discovery]
- 108. Artemisinins as a novel anti-cancer therapy: Targeting a global cancer pandemic through drug repurposing - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC7564301/]
- 109. -malarial drug; the emerging role of Anti and its derivatives... artemisinin [https://cmjournal.biomedcentral.com/articles/10.1186/s13020-021-00489-0]
- 110. Frontiers | Repurposing Artemisinin and its Derivatives as Anticancer Drugs: A Chance or Challenge? [https://www.frontiersin.org/journals/pharmacology/articles/10.3389/fphar.2021.828856/full]
- 111. -parasitic Drug Anti Exhibits Potent Ivermectin ... Anticancer Activity [https://ar.iiarjournals.org/content/39/9/4837]
- 112. , a potential Ivermectin drug derived from an antiparasitic... anticancer [https://pmc.ncbi.nlm.nih.gov/articles/PMC7505114/]
- 113. Ivermectin reverses the drug resistance in cancer cells through EGFR/ERK/Akt/NF-κB pathway | Journal of Experimental & Clinical Cancer Research | Full Text [https://jeccr.biomedcentral.com/articles/10.1186/s13046-019-1251-7]
- 114. Progress in the discovery and development of anticancer agents from... [https://pubs.rsc.org/en/content/articlelanding/2025/np/d4np00019f]
- 115. Computational-aided design of a library of lactams through a diversity-oriented synthesis strategy - ScienceDirect [https://www.sciencedirect.com/science/article/abs/pii/S0968089620303680]
- 116. Comparative analyses of structural features and scaffold diversity for purchasable compound libraries | Journal of Cheminformatics | Full Text [https://jcheminf.biomedcentral.com/articles/10.1186/s13321-017-0212-4]
- 117. A combinatorial DNA assembly approach to biosynthesis of N - linked ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9990991/]
- 118. Frontiers | Construction of an Escherichia coli chassis for efficient ... [https://www.frontiersin.org/journals/bioengineering-and-biotechnology/articles/10.3389/fbioe.2024.1370685/full]
- 119. Genetic and process engineering strategies for enhanced recombinant... [https://microbialcellfactories.biomedcentral.com/articles/10.1186/s12934-021-01689-x]
- 120. Verification of N - Linked of Proteins Isolated from Glycosylation ... Plant [https://bio-protocol.org/en/bpdetail?id=5444&type=0]
- 121. Antibody–drug conjugates: Recent advances in payloads - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC10547921/]
- 122. Overcoming resistance to antibody-drug conjugates: from mechanistic... [https://jhoonline.biomedcentral.com/articles/10.1186/s13045-025-01752-9]
- 123. Present Scenario and Future Landscape of Payloads for ADCs: Focus on DNA-Interacting Agents - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC11510327/]
- 124. Antibody Drug Conjugate Market Size, Share & Growth [2035] [https://www.rootsanalysis.com/reports/antibody-drug-conjugates-market/270.html]
- 125. of free Challenges concentration analysis in payload ... | ICON plc ADC [https://www.iconplc.com/insights/blog/2025/09/24/challenges-free-payload-concentration-analysis-adc-studies]
- 126. AACR 2025 preview – a surge in dual- payload conjugates | ApexOnco... [https://www.oncologypipeline.com/apexonco/aacr-2025-preview-surge-dual-payload-conjugates]

- 127. ADC Payload Development Services - BOC Sciences [https://adc.bocsci.com/services/adc-payloads-development.html]
- 128. Antibody Drug Conjugates Market Size, Share & Industry Growth Report 2030 [https://www.mordorintelligence.com/industry-reports/antibody-drug-conjugates-market]
- 129. Reconstructing the Antibiotic Pipeline: Natural Alternatives to... [https://pubmed.ncbi.nlm.nih.gov/40867625/]
- 130. Natural Products as Platforms To Overcome Antibiotic Resistance - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC5869711/]
- 131. A systematic review on natural products with antimicrobial potential against WHO's priority pathogens | European Journal of Medical Research | Full Text [https://eurjmedres.biomedcentral.com/articles/10.1186/s40001-025-02717-x]
- 132. Role of natural product diversity in chemical biology - PubMed [https://pubmed.ncbi.nlm.nih.gov/21489856/]
- 133. -informed exploration of Natural to enable... product chemical space [https://pmc.ncbi.nlm.nih.gov/articles/PMC8130614/]
- 134. Antimicrobial and Anti-Infective Activity of Natural Products—Gaining Knowledge from Novel Studies - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC10295247/]
- 135. QSAR and pharmacophore modeling of natural and synthetic antimalarial prodiginines - PubMed [https://pubmed.ncbi.nlm.nih.gov/24010933/]
- 136. Biological activities and 3D QSAR studies of a series of Delisea pulchra (cf. fimbriata) derived natural products - PubMed [https://pubmed.ncbi.nlm.nih.gov/16933872/]
- 137. Multiverse: An Expanded View of Chemical - PMC Chemical Space [https://pmc.ncbi.nlm.nih.gov/articles/PMC9787733/]
- 138. Interactions between Medical Plant-Derived Bioactive Compounds ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9404952/]
- 139. Frontiers | Combination of marine bioactive compounds and extracts ... [https://www.frontiersin.org/journals/nutrition/articles/10.3389/fnut.2022.1047026/full]
- 140. Methods for Assessing Drug Quantitative - PMC Synergism [https://pmc.ncbi.nlm.nih.gov/articles/PMC3379564/]