Central Carbon Metabolism Engineering: Foundational Strategies and Cutting-Edge Applications for Advanced Bioproduction

Generated at: 2026-04-03
Source URL: https://www.metabengsci.com/posts/central-carbon-metabolism-engineering-foundational-strategies-and-cuttingedge-applications-for-advanced-bioproduction

Central Carbon Metabolism Engineering: Foundational Strategies and Cutting-Edge Applications for Advanced Bioproduction

Abstract

This article provides a comprehensive overview of contemporary strategies for engineering central carbon metabolism (CCM) in microbial hosts to enhance the production of pharmaceuticals and high-value chemicals. It explores the fundamental principles of CCM pathways, details advanced methodological tools for pathway manipulation and dynamic control, addresses common optimization challenges and troubleshooting approaches, and discusses validation techniques for assessing engineering success. Tailored for researchers, scientists, and drug development professionals, this review synthesizes recent advances in metabolic engineering, synthetic biology, and systems-level analysis, offering a roadmap for designing efficient microbial cell factories for sustainable biomanufacturing.

Decoding Central Carbon Metabolism: Core Pathways and Fundamental Engineering Principles

Central Carbon Metabolism (CCM) represents the most fundamental metabolic process in living organisms, responsible for maintaining normal cellular growth by managing the breakdown and utilization of carbohydrates, fats, and proteins [1]. This interconnected biochemical network consists primarily of three core pathways: **glycolysis**, the **pentose phosphate pathway (PPP)**, and the **tricarboxylic acid (TCA) cycle** (also known as the citric acid cycle or Krebs cycle) [1] [2]. These pathways collectively generate ATP for cellular activities and provide key intermediates for biosynthetic reactions, making them indispensable for cellular growth and survival [1]. In rapidly proliferating mammalian cells, such as those used in recombinant protein production, CCM plays an essential role in supplying both biosynthetic precursors and energy, with its configuration changing significantly with varying growth rates [3]. The architectural organization of these pathways—including their compartmentalization, regulatory mechanisms, and potential formation of enzyme complexes known as metabolons—enables precise control over metabolic flux in response to cellular demands and environmental conditions [4].

Pathway Architectures and Biochemical Reactions

Glycolysis: The Foundation of Energy Production

Glycolysis serves as the starting point of central carbon metabolism, occurring in the cytoplasm of all cells [1]. This ten-step enzymatic pathway converts glucose into pyruvate through two distinct phases: the energy investment phase and the energy harvesting phase [1]. In the first phase, the cell consumes ATP to phosphorylate glucose, preparing it for cleavage into two three-carbon molecules. The second phase produces ATP and NADH through substrate-level phosphorylation, directly coupled to enzymatic conversion of metabolites [1]. The end product, pyruvate, represents a crucial metabolic intermediate that can be further metabolized through aerobic or anaerobic pathways depending on oxygen availability [1].

Key Regulatory Nodes in Glycolysis:

- **Phosphofructokinase (PFK):** A central regulatory enzyme inhibited by ATP and activated by AMP, reflecting the cell's energy state [1].
- **Glucose-6-phosphate isomerase (PGI):** Exhibits multifaceted roles, with significant correlations to cell cycle progression in rapidly proliferating cells [3].
- **Pyruvate kinase (PK):** Subject to oxidative inhibition under stress conditions, contributing to metabolic flux rerouting [5].

Pentose Phosphate Pathway: Biosynthetic Support and Redox Balance

The pentose phosphate pathway operates as a complementary pathway to glycolysis, serving two primary functions: maintaining cellular redox balance and supporting biosynthesis [1]. The PPP generates NADPH, a key electron donor in anabolic reactions, and ribose-5-phosphate, which is essential for nucleotide synthesis [1]. This pathway can be divided into two interconnected phases: the oxidative phase (which generates NADPH) and the non-oxidative phase (which produces ribose-5-phosphate and can interconvert various sugar phosphates) [1]. Under oxidative stress conditions, carbon flux is rapidly rerouted into the PPP to support NADPH-dependent antioxidant defense mechanisms, a process regulated through complex coordination of multiple enzymatic activities [5].

PPP Flux Regulation:

- **G6PD activity:** Upregulated in response to decreased NADPH/NADP+ ratios during oxidative stress [5].
- **6-phosphogluconate (6PG)-dependent inhibition** of PGI helps redirect flux from glycolysis to PPP [5].
- **Coordination with glycolytic enzymes** through joint inhibition mechanisms enables efficient detoxification across a broad range of stress levels [5].

Tricarboxylic Acid Cycle: Metabolic Hub and Energy Generation

The TCA cycle operates as a central hub for cellular energy production within the mitochondrial matrix [1]. Here, pyruvate (derived from glucose) is converted to acetyl-CoA, which enters the cycle to generate key metabolites including NADH, $FADH_2$, and GTP, all critical for ATP production via oxidative phosphorylation [1]. Beyond its role in energy generation, the TCA cycle provides intermediates that are diverted into biosynthetic pathways; for example, α-ketoglutarate serves as a precursor for glutamate and other amino acids, while oxaloacetate is essential for gluconeogenesis [1]. In photosynthetic organisms, the TCA cycle demonstrates remarkable metabolic plasticity, with major CO_2 fluxes alternating between fixation by the Calvin-Benson-Bassham cycle during light and release by the TCA cycle in darkness [6].

Metabolon Organization in TCA Cycle: Experimental evidence supports the existence of a functional TCA cycle metabolon in plants, with protein-protein interaction studies identifying 158 interactions among 38 mitochondrial proteins [4]. Isotope dilution experiments with isolated potato mitochondria have revealed channelling of citrate and fumarate, suggesting that substrate channelling may enhance pathway efficiency and regulatory control [4]. This supramolecular organization allows reaction intermediates to be isolated from the bulk environment, potentially providing advantages including local metabolite enrichment, isolation of intermediates from competing reactions, protection of unstable intermediates, and sequestration of cytotoxic metabolites [4].

Quantitative Analysis of Metabolic Fluxes

Flux Distribution Under Normal and Stress Conditions

Table 1: Metabolic Flux Redistribution in Response to Oxidative Stress in Human Fibroblasts

Metabolic Pathway/Branch	Baseline Flux (% glucose import)	Oxidative Stress Flux (% glucose import)	Fold Change
oxPPP flux	~20%	~65%	3.25x increase
Lower glycolytic flux	100% (reference)	~33%	3-fold decrease
Nucleotide production	Not specified	Significant increase	Not quantified
Non-oxidative PPP	Not specified	Significant split from oxPPP	Not quantified

Data derived from 13C-labeling analysis using human fibroblast cells exposed to 500μM H_2O_2 [5].

Growth Rate-Dependent Enzymatic Activities in Mammalian Cells

Table 2: Relationship Between Growth Rate and Key Enzymatic Activities in GS-CHO Cell Lines

Metabolic Pathway	Enzymatic Activity Trend with Increasing Growth Rate	Primary Function
TCA Cycle	Elevated activity	Energy supply
Glycolysis	Decreasing tendency	Biosynthetic precursor supply
PPP	Decreasing tendency	Biosynthetic precursor supply
Glucose-6-phosphate isomerase (PGI)	Significant correlation with % of G2/M-phase cells	Multifaceted role in cell cycle progression

Data based on analysis of seven key enzymes from six cell clones of rapidly proliferating glutamine-synthetase Chinese hamster ovary cells [3].

Experimental Methodologies for Central Carbon Metabolism Analysis

Time-Resolved Metabolomics and Isotope Tracing

The investigation of dynamic metabolic transitions requires sophisticated methodologies capable of capturing rapid changes in metabolite levels and flux distributions. One advanced approach involves exposing biological systems to $^{13}CO_2$ for 20 minutes to establish a quasi-steady state before applying an abrupt environmental change (e.g., light-to-dark transition in plant leaves) while maintaining $^{13}CO_2$ feeding [6]. This enables researchers to track carbon movement through metabolic pathways during rapid metabolic reconfigurations.

Protocol: Fast Kill Freeze Clamp Sampling for Time-Resolved Metabolomics

- **Equipment Setup:** Integrate a gas exchange system (e.g., LI-COR 6800) with a fast-freeze mechanism using liquid-nitrogen-cooled copper blocks [6].
- **Environmental Control:** Configure the system to scrub all CO_2 before introducing $^{13}CO_2$ from a regulated source, maintaining precise CO_2 concentration (e.g., 420 ppm) [6].
- **Isotope Labeling:** Equilibrate biological samples under stable conditions before switching to $^{13}CO_2$ for a predetermined period (e.g., 20 minutes) [6].
- **Stimulus Application:** Apply experimental stimulus (e.g., turn off light for dark transition) while maintaining isotope labeling.
- **Rapid Sampling:** Trigger the Fast Kill mechanism at precise time intervals (e.g., 0, 10, 30, 60, 180, 600 seconds) post-stimulus [6].
- **Sample Preservation:** Collect frozen samples in cryogenic storage tubes and maintain at -80°C for subsequent mass spectrometry analysis [6].

This methodology enables resolution of metabolic changes occurring within seconds, with the interval between stimulus application and sample freezing measured at approximately 35 milliseconds [6].

Protein-Protein Interaction Mapping for Metabolon Identification

Identifying and characterizing metabolons requires comprehensive analysis of potential protein-protein interactions among metabolic enzymes. An integrated approach employing multiple complementary techniques provides reliable assessment of these interactions [4].

Protocol: Integrated Metabolon Identification Pipeline

- **Affinity Purification-Mass Spectrometry (AP-MS):**
  - Utilize GFP-tagged affinity purification procedures in cell culture systems
  - Process normalized signal intensities with SAINT software to determine FC-A scores
  - Consider interactions in the top 10% of scores as positive [4]
- **Split-Luciferase (Split-LUC) Assays:**
  - Conduct high-throughput assays in relevant protoplasts or cell systems
  - Quantify relative luminescence units for interaction strength
  - Apply top 10% threshold for positive interactions [4]
- **Yeast-Two-Hybrid (Y2H) Assays:**
  - Perform semiquantitative assays with colony observation at 3, 5, and 8 days
  - Score earlier colony detection as stronger interaction
  - Consider colonies observed at 3 or 5 days as positive [4]
- **Data Integration:**
  - Apply STATIS-like methodology to generate compromise scores
  - Select protein pairs with compromise scores above established thresholds
  - Validate functional interactions through isotope dilution experiments for substrate channelling [4]

13C-Metabolic Flux Analysis (13C-MFA)

Quantitative modeling of metabolic fluxes provides critical insights into pathway operation under different physiological conditions. For central carbon metabolism, particularly the coordination between glycolysis and PPP, 13C-MFA offers a powerful approach for quantifying flux redistribution [5].

Protocol: Stochastic Simulation Algorithm-Based 13C-MFA

- **Experimental Design:**
  - Select appropriate 13C-labeled glucose tracer (e.g., [1-13C], [U-13C])
  - Expose biological system to tracer under controlled conditions
  - Implement precise environmental perturbations (e.g., H_2O_2 exposure for oxidative stress)
- **Data Collection:**
  - Measure mass isotopomer distributions using LC-MS or GC-MS
  - Quantify absolute metabolite concentrations

- Record extracellular flux rates where applicable

- **Flux Determination:**
  - Apply Monte Carlo sampling to determine posterior distribution of flux parameters
  - Use stochastic simulation algorithm to model isotope labeling system
  - Calculate confidence intervals for estimated flux distributions
  - Validate model fit with experimental mass isotopomer data [5]



Figure 1: 13C-Metabolic Flux Analysis Workflow

Advanced Research Applications and Engineering Strategies

Metabolic Engineering of Central Carbon Metabolism

The optimization of central carbon metabolism has emerged as a powerful strategy in metabolic engineering, enabling enhanced production of valuable compounds through rearrangement of global metabolic flux [2]. Engineering approaches focus on either introducing heterologous pathways or optimizing native pathways to increase precursor supply, rebalance energy availability, or adjust redox cofactors.

Key Engineering Strategies:

- **Phosphoketolase (PHK) Pathway Introduction:**
  - Enables direct synthesis of acetyl-CoA from F6P and X5P
  - Increases acetyl-CoA supply for lipid compounds
  - Addresses insufficient erythrose-4-phosphate synthesis in yeast
  - Applications: 25% increase in farnesene production, 19% increase in total lipids in Yarrowia lipolytica [2]

- **ATP:citrate lyase (ACL) Expression:**
  - Directly converts citric acid to acetyl-CoA in TCA cycle
  - Can double mevalonate yield in S. cerevisiae [2]

- **Pyruvate Dehydrogenase (PDH) Pathway Engineering:**
  - Directly converts pyruvate to acetyl-CoA without ATP consumption

- NADP+-dependent modifications in S. cerevisiae resulted in 2-fold increase in acetyl-CoA [2]

- **Regulatory Enzyme Modulation:**
  - Allosteric regulation targets (PFK, PDH)
  - Transcriptional regulation (HIF-1, c-Myc)
  - Post-translational modifications (AMPK, acetylation) [1]

The Scientist's Toolkit: Essential Research Reagents

Table 3: Key Research Reagent Solutions for Central Carbon Metabolism Studies

Reagent/Category	Specific Examples	Function/Application
Stable Isotopes	$^{13}CO_2$ (99+% purity), [1-^{13}C]glucose, [U-^{13}C]glucose	Metabolic flux tracing, quantification of pathway utilization
Mass Spectrometry Standards	Isotopically labeled internal standards for key metabolites	Absolute quantification of metabolite concentrations
Protein Interaction Assay Systems	GFP-tagging vectors, split-luciferase components, yeast-two-hybrid systems	Identification and validation of metabolon formations
Enzyme Activity Assays	Commercial kits for G6PD, PK, PFK, IDH activities	Assessment of metabolic regulation at enzymatic level
Oxidative Stress Inducers	Hydrogen peroxide (H_2O_2) solutions at precise concentrations	Investigation of metabolic adaptation to stress conditions
Genetic Engineering Tools	CRISPR-Cas9 systems, TALEN, ZFN, pathway-specific overexpression/knockout constructs	Targeted modification of central carbon metabolism

Recent Research Advances and Future Perspectives

Recent studies have revealed the remarkable plasticity and dynamic regulation of central carbon metabolism across biological systems. In cyanobacteria, metabolite-level regulation of enzymatic activity controls awakening from metabolic dormancy, demonstrating how allosteric regulation fine-tunes metabolic states [7]. Plant studies have uncovered an exceptionally rapid reconfiguration from CO_2 fixation by the Calvin-Benson-Bassham cycle in light to TCA cycle activation for energy production in darkness, with substantial accompanying changes in amino acid metabolism occurring within seconds of light-dark transition [6].

The emerging understanding of metabolons and substrate channelling has transformed our perspective on metabolic pathway organization. Rather than existing as collections of independent enzymes, central metabolic pathways appear to form functional complexes that enhance metabolic efficiency and regulatory control [4]. This architectural principle likely explains how cells achieve the rapid metabolic transitions necessary for adaptation to changing environmental conditions.

Future research directions will likely focus on harnessing these architectural principles for biotechnological applications. The integration of kinetic modeling with multi-omics data will enable more predictive engineering of central carbon metabolism [5]. Advances in synthetic biology and metabolic engineering are already paving the way for sustainable bioproduction, with engineered microorganisms demonstrating significantly improved conversion efficiencies, such as 91% biodiesel conversion efficiency from lipids and 3-fold increases in butanol yield in engineered Clostridium species [8].

As our understanding of the architecture of central carbon metabolism continues to evolve, so too will our ability to manipulate these fundamental processes for applications ranging from therapeutic development to sustainable bioproduction. The integration of architectural principles with quantitative systems biology approaches promises to unlock new dimensions of metabolic control and engineering potential.

Central carbon metabolism serves as the biochemical core of the cell, governing carbon flux, energy production, and precursor supply for biosynthesis. In metabolic engineering, reprogramming this network is essential for efficient bioproduction. Three nodes—acetyl-Coenzyme A (acetyl-CoA), pyruvate, and phosphoenolpyruvate (PEP)—stand as critical control points due to their unique positions bridging glycolysis, the tricarboxylic acid (TCA) cycle, and anabolic pathways. This whitepaper examines the biochemical roles, quantitative characteristics, and engineering strategies for these key nodes, providing a technical guide for researchers and scientists aiming to optimize microbial cell factories for drug development and bio-based chemical production.

Quantitative Characteristics of Key Metabolic Nodes

The functional importance of acetyl-CoA, pyruvate, and PEP is underscored by their distinct concentration ranges, turnover rates, and connectivity within metabolic networks. Understanding these quantitative parameters is essential for predicting system behavior and designing effective engineering interventions.

Table 1: Quantitative Parameters of Key Metabolic Nodes in E. coli

Metabolite	Typical Intracellular Concentration	Primary Biosynthetic Route	Primary Consuming Pathways
Acetyl-CoA	0.05–1.5 nmol/mg CDW (20–600 μM) [9]	Pyruvate dehydrogenase (Pdh) [9]	TCA cycle, fatty acid biosynthesis, amino acids, polyketides [9]
Pyruvate	Varies with growth phase and conditions	Glycolysis (EMP pathway) [9]	Pyruvate dehydrogenase, pyruvate formate lyase, anaplerotic reactions [9]
PEP	Lower than pyruvate (exact range varies)	Glycolysis (EMP pathway)	Phosphotransferase system (PTS), anaplerotic reactions, aromatics biosynthesis

Table 2: Carbon Source Impact on Acetyl-CoA Pools in E. coli

Carbon Source	Acetyl-CoA Concentration (nmol/mg CDW)	Relative Flux Potential
Glucose	0.82 [9]	High
Glycerol	0.62 [9]	Medium
Succinate	0.37 [9]	Low

Acetyl-CoA: The Central Biosynthetic Precursor

Biochemical Role and Regulation

Acetyl-CoA represents a fundamental two-carbon building block for numerous biosynthetic pathways, serving as the primary precursor for lipids, polyketides, isoprenoids, amino acids, and other valuable bioproducts with applications across biochemical, biofuel, and pharmaceutical industries [9]. This metabolite sits at the convergence of multiple metabolic routes, including glycolysis, fatty acid β-oxidation, and amino acid catabolism [10]. The intracellular concentration of acetyl-CoA is tightly regulated in response to cellular energy status, carbon source availability, and environmental conditions such as salt stress, temperature, pH, and oxygen levels [9].

The primary enzyme responsible for acetyl-CoA biosynthesis from pyruvate under aerobic conditions is the pyruvate dehydrogenase complex (PDH), which catalyzes the oxidative decarboxylation of pyruvate to yield acetyl-CoA, CO_2, and NADH [9]. This massive multi-enzyme complex comprises three subunits: E1 (pyruvate decarboxylase), E2 (dihydrolipoamide acetyltransferase), and E3 (dihydrolipoyl dehydrogenase) [10]. The complex employs five essential coenzymes in its catalytic mechanism: thiamine pyrophosphate (TPP), lipoic acid, coenzyme A (CoA), flavin adenine dinucleotide (FAD), and nicotinamide adenine dinucleotide (NAD) [10]. Under anaerobic conditions, pyruvate formate lyase (Pfl) serves as the major enzyme for acetyl-CoA formation, producing formate as a byproduct [9].

Engineering Strategies for Enhanced Acetyl-CoA Flux

Multiple metabolic engineering strategies have been successfully implemented to overcome native regulatory constraints and enhance acetyl-CoA availability for bioproduction.

3.2.1 Pyruvate Dehydrogenase Optimization Overexpression of the native pyruvate dehydrogenase complex (encoded by *aceE*, *aceF*, and *lpd*) represents a direct approach to increasing acetyl-CoA flux. In E. coli, this strategy has demonstrated a 1.45-fold increase in isoamyl acetate production and a 2-fold elevation in intracellular acetyl-CoA concentration [9]. Fine-tuning PDH expression through plasmids with different copy numbers has also yielded significant improvements in fatty acid production [9]. A particularly innovative approach involves engineering PDH complexes with reduced sensitivity to NADH inhibition. Introduction of an E354K mutation in the E3 subunit (Lpd) created an NADH-insensitive variant that enabled a 5-fold increase in carbon flux through PDH under anaerobic conditions and a 1.6-fold enhancement in butanol production [9].

3.2.2 Enhanced Pyruvate Supply Increasing carbon flux through glycolysis to pyruvate directly drives acetyl-CoA formation. Overexpression of phosphoglycerate kinase (Pgk) and glyceraldehyde-3-phosphate dehydrogenase (GapA) has been shown to increase intracellular acetyl-CoA concentration by nearly 30%, resulting in doubled naringenin production [9]. Alternative glycolytic pathways such as the serine-deamination (SD) pathway and Entner-Doudoroff (ED) pathway have also been successfully engineered to enhance pyruvate flux. Overexpression of *serABC* and *sdaA* (enhancing serine synthesis and deamination) improved acetyl-CoA concentration by 10%, while engineering the *edd-eda* operon and *zwf* gene promoter regions achieved a 3-fold increase in acetyl-CoA concentration, supporting poly-3-hydroxybutyrate (PHB) production at 5.5 g/L [9].

3.2.3 Acetate Reassimilation Pathways Acetate secretion represents a significant loss of acetyl-CoA flux in many microbial systems. Overexpression of acetyl-CoA synthetase (Acs), which catalyzes the ATP-dependent conversion of acetate to acetyl-CoA, can efficiently reassimilate secreted acetate, reducing extracellular acetate from 11 mM to negligible levels and increasing intracellular acetyl-CoA concentration more than 3-fold (reaching 3.5 nmol/mg CDW) [9]. Alternatively, expression of the phosphate acetyltransferase (*pta*) and acetyl-CoA kinase (*ack*) operon has successfully enhanced N-acetylglutamate production by 2-fold [9]. Supplementing culture media with acetate further enhances the effectiveness of these assimilation strategies.

3.2.4 Modular Deregulation Strategies Recent advanced engineering approaches employ comprehensive deregulation of central carbon metabolism modules. A multifaceted strategy encompassing promoter engineering, transcription factor manipulation, biosensor implementation, heterologous enzyme introduction, and mutant enzyme expression has demonstrated remarkable success in Saccharomyces cerevisiae, achieving a 4.7-fold enhancement in 3-hydroxypropionic acid (3-HP) productivity from xylose compared to initially optimized strains [11].



Diagram 1: Metabolic Engineering Strategies for Enhanced Acetyl-CoA Flux

Experimental Protocols for Key Metabolic Engineering Approaches

Protocol: Pyruvate Dehydrogenase Complex Overexpression

Objective: Enhance flux from pyruvate to acetyl-CoA through targeted overexpression of PDH complex genes.

Materials:

- **Plasmids:** High-, medium-, and low-copy number plasmids with inducible promoters (e.g., pTrc, pBAD) [9]
- **Strains:** Production host (e.g., E. coli) with deleted competing pathways as needed
- **Growth Media:** Defined minimal media with target carbon source (e.g., glucose, glycerol)
- **Induction Reagents:** IPTG for lac-based promoters, arabinose for araBAD promoters
- **Analytical Tools:** LC-MS/MS for acetyl-CoA quantification [12], GC-MS for metabolic flux analysis

Methodology:

- Clone PDH operon (*aceE*, *aceF*, *lpd*) into expression vectors of varying copy numbers
- Transform constructs into production host and validate expression via SDS-PAGE
- Cultivate strains in controlled bioreactors with monitoring of growth parameters
- Induce PDH expression during mid-exponential phase
- Quantify intracellular acetyl-CoA concentrations at multiple time points
- Measure target product titers and calculate yield coefficients

Expected Outcomes: 1.5- to 2-fold increase in acetyl-CoA concentration; 1.5- to 2.5-fold improvement in product titers from acetyl-CoA-derived pathways [9].

Protocol: NADH-Insensitive Pyruvate Dehydrogenase Engineering

Objective: Overcome native regulation of PDH by NADH feedback inhibition for improved anaerobic acetyl-CoA production.

Materials:

- **Site-Directed Mutagenesis Kit:** For introducing E354K mutation in *lpd* gene [9]
- **Anaerobic Chamber:** For maintaining oxygen-free cultivation conditions
- **Specialized Media:** Pre-reduced media for anaerobic experiments

Methodology:

- Introduce E354K mutation into *lpd* gene encoding dihydrolipoyl dehydrogenase
- Express mutant PDH complex in production host
- Compare aerobic vs. anaerobic performance of wild-type and mutant PDH
- Quantify NADH sensitivity through enzyme activity assays
- Evaluate impact on product formation under anaerobic conditions

Expected Outcomes: 5-fold increase in carbon flux through PDH under anaerobic conditions; 1.6-fold improvement in anaerobic butanol production [9].

Protocol: Glycolytic Flux Enhancement for Pyruvate Overflow

Objective: Increase glycolytic flux to pyruvate to drive enhanced acetyl-CoA formation.

Materials:

- **Gene Expression Tools:** Constitutive promoters (PJ23119, PTrc-162) for pathway engineering [9]
- **Pathway Enzymes:** Genes for glycolytic enzymes (Pgk, GapA), ED pathway (edd-eda operon), SD pathway (serABC, sdaA)

Methodology:

- Overexpress key glycolytic enzymes (Pgk, GapA) identified through flux balance analysis
- Engineer serine-deamination pathway through *serABC* and *sdaA* overexpression
- Enhance ED pathway flux through promoter replacement of *edd-eda* operon and *zwf*
- Measure intracellular pyruvate and acetyl-CoA concentrations
- Determine flux redistribution using 13C metabolic flux analysis

Expected Outcomes: 10-30% increase in acetyl-CoA concentration; 3-fold enhancement achievable through combinatorial approach [9].

Visualization of Metabolic Network Dynamics

Advanced visualization approaches are essential for interpreting complex metabolic networks and understanding dynamic changes in metabolite pools. The GEM-Vis method enables visualization of time-course metabolomic data within metabolic network maps, representing metabolite concentrations through node fill levels, which allows for intuitive human interpretation of quantitative changes [13]. This approach is particularly valuable for identifying metabolic bottlenecks, understanding regulatory interactions, and observing system responses to genetic interventions.

Regulatory interactions between metabolite pools and reaction steps can be quantified through Regulatory Strength (RS) values, which measure the strength of up- or down-regulation compared to non-inhibited or non-activated states [14]. These values can be visualized on a percentage scale where 100% represents maximal possible inhibition or activation, and 0% indicates absence of regulatory interaction [14].



Diagram 2: Dynamic Visualization Framework for Metabolic Networks

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 3: Essential Research Reagents for Metabolic Engineering Studies

Reagent/Material	Function/Application	Example Use Cases
Acetyl-CoA Synthetase (Acs)	ATP-dependent acetate assimilation to acetyl-CoA	Reversing acetate secretion; increasing acetyl-CoA pools 3-fold [9]
NADH-Insensitive Lpd Mutant (E354K)	Pyruvate dehydrogenase complex component resistant to NADH inhibition	Enhancing anaerobic acetyl-CoA flux 5-fold [9]
Phosphoglycerate Kinase (Pgk)	Glycolytic enzyme catalyzing 1,3-BPG to 3-PG conversion	Increasing glycolytic flux to pyruvate and acetyl-CoA [9]
Constitutive Promoters (PJ23119, PTrc-162)	Transcriptional control elements for pathway engineering	Optimizing expression of ED pathway genes for enhanced pyruvate flux [9]
LC-MS/MS Systems	Quantitative analysis of intracellular metabolites	Absolute quantification of acetyl-CoA and pathway intermediates [12]
13C-Labeled Substrates	Metabolic flux analysis through isotopic tracing	Determining in vivo flux distributions in engineered strains [9]
CRISPR Interference (CRISPRi)	Targeted gene repression without knockout	Fine-tuning competing pathway expression while maintaining viability [9]

Acetyl-CoA, pyruvate, and PEP represent fundamental control points in central carbon metabolism whose engineering is essential for optimizing microbial cell factories. Through targeted strategies including enzyme complex optimization, pathway deregulation, flux enhancement, and dynamic visualization, researchers can significantly enhance carbon flux toward valuable biochemical products. The continued development of sophisticated engineering tools and analytical methods will further advance our ability to reprogram cellular metabolism for pharmaceutical and industrial biotechnology applications.

Natural vs. Synthetic C1 Assimilation Pathways for Carbon Fixation

The escalating concentration of atmospheric carbon dioxide (CO_2) and the environmental imperative to transition toward a circular bioeconomy have catalyzed intense research into one-carbon (C1) assimilation pathways. **C1 compounds**—including CO_2, carbon monoxide (CO), methane (CH_4), methanol, and formate—represent abundant and sustainable feedstocks that can be derived from industrial waste gases, renewable synthesis, or greenhouse gases [15] [16]. Leveraging these compounds for biomanufacturing presents a sustainable alternative to conventional processes that depend on sugar-based feedstocks, which compete with food resources, or fossil fuels [17] [18]. The fundamental challenge lies in engineering efficient biological routes to integrate these simple molecules into **central carbon metabolism (CCM)**, thereby generating the precursors for biomass, fuels, and high-value chemicals.

This field operates at the intersection of synthetic biology and metabolic engineering, exploring two primary strategies: optimizing **natural C1 assimilation pathways** native to autotrophic organisms, and designing **synthetic pathways** implanted into industrially robust, tractable hosts. Natural pathways are a product of evolution but often suffer from kinetic and thermodynamic inefficiencies. In contrast, synthetic pathways are rationally designed for higher theoretical efficiency and better integration with a host's metabolic network, though their implementation faces challenges in functional expression and regulation [18] [19]. The ultimate engineering goal is to rewire central carbon metabolism to create microbial cell factories that can efficiently transform C1 substrates into multi-carbon, value-added products, thereby contributing to carbon neutrality and a sustainable future [17] [15].

Foundational Concepts of C1 Metabolism

Physicochemical and Bioenergetic Considerations

The efficient utilization of C1 compounds is inextricably linked to the principles of energy transduction and redox balance. The **oxidation state** of the carbon atom in a C1 substrate dictates the number of electrons required for its reduction to a target metabolite, profoundly influencing the pathway's thermodynamics and energy demands [15]. For instance, CO_2, the most oxidized form of carbon, requires substantial energy input for reduction, whereas more reduced substrates like methanol can themselves serve as energy sources [15].

A pivotal bioenergetic challenge in C1 metabolism is the **adenosine triphosphate (ATP)** requirement. ATP is essential for cell maintenance, driving thermodynamically unfavorable reactions, and supporting growth. Many native C1 assimilation pathways are ATP-intensive. The Calvin-Benson-Bassham (CBB) cycle, for example, consumes 3 ATP and 2 NADPH per molecule of CO_2 fixed [15] [20]. This high energy demand often creates a **bioenergetic constraint**, limiting the overall carbon flux and yield. Engineering objectives therefore frequently focus on supplementing

ATP supply through strategies such as chemical co-feeding, or coupling metabolism to light (in phototrophs) or electrical energy [15]. The pairing of C1 substrates with target products is also guided by biochemical proximity; synthesizing products that are structurally similar to pathway intermediates (e.g., C2 or C3 metabolites) minimizes the need for further energy-intensive redox reactions and decarboxylation steps, thereby enhancing overall carbon and energy efficiency [15].

The Role of Central Carbon Metabolism

Central carbon metabolism—encompassing glycolysis, the pentose phosphate pathway (PPP), and the tricarboxylic acid (TCA) cycle—serves as the fundamental hub connecting C1 assimilation to all downstream bioproducts. C1 pathways do not operate in isolation; they function to **condense C1 units** into central metabolic intermediates, such as acetyl-CoA, pyruvate, or glyceraldehyde-3-phosphate (GAP) [2] [15]. Once these C2 or C3 building blocks enter the CCM, they can be channeled by the host's endogenous enzymes to generate energy, reducing equivalents (NADPH, NADH), and the precursor molecules for amino acids, lipids, and nucleotides.

Consequently, engineering CCM is a powerful strategy to amplify the flux from C1 substrates toward desired products. Modifications upstream in the CCM can trigger a global rearrangement of metabolic flux, increasing the supply of essential precursors. For example, introducing a heterologous **phosphoketolase (PHK) pathway** creates a shortcut from fructose-6-phosphate and xylulose-5-phosphate to acetyl-CoA, bypassing several steps in the standard glycolysis and PPP. This has been successfully deployed in *Saccharomyces cerevisiae* to increase the supply of acetyl-CoA for producing fuels and the precursor erythrose-4-phosphate for aromatic amino acid synthesis [2]. Thus, the synergistic optimization of both the C1 entry pathway and the host's central metabolic network is critical for achieving high-yield bioproduction.

Natural C1 Assimilation Pathways

Nature has evolved several distinct pathways to assimilate C1 compounds, each with unique biochemistry, energetics, and host distribution. The following table provides a comparative overview of the major natural pathways.

Table 1: Key Characteristics of Natural C1 Assimilation Pathways

Pathway	Primary Substrates	Key Products	Representative Organisms	ATP Consideration	Notable Features
Calvin-Benson-Bassham (CBB) Cycle	CO_2	Glyceraldehyde-3-P (G3P)	Plants, Cyanobacteria, Algae [15]	High (3 ATP/CO_2) [15]	Dominant global pathway; slow RuBisCO kinetics; oxygen sensitivity [15]
Wood-Ljungdahl Pathway (WLP)	CO_2, CO, Formate	Acetyl-CoA	Acetogens (e.g., *Clostridium* spp.) [15]	ATP-limited in gas fermentation [15]	Anaerobic; high theoretical yield; direct C2 product [15]
Serine Cycle	Formaldehyde, CO_2	Acetyl-CoA	Type II Methanotrophs [15]	Requires 2 ATP/acetyl-CoA [15]	Used by methane-oxidizing bacteria; involves TCA cycle intermediates.
Ribulose Monophosphate (RuMP) Cycle	Formaldehyde	Dihydroxyacetone-P (DHAP)	Type I Methanotrophs, *Bacillus* [15]	Lower ATP cost than CBB [15]	Efficient formaldehyde assimilation; linear then cyclic phases.
Reductive Glycine Pathway (rGlyP)	Formate, CO_2, NH_3	Glycine	*Desulfovibrio*, Engineered *S. cerevisiae* [15]	Thermodynamically challenging [15]	Linear and relatively simple structure; promising for engineering [18].

The Calvin-Benson-Bassham (CBB) Cycle

The CBB cycle is the primary photosynthetic carbon fixation pathway on Earth. It is a **reductive pentose phosphate pathway** that operates in the stroma of chloroplasts or the cytoplasm of cyanobacteria. Its key enzyme, **ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)**, catalyzes the carboxylation of ribulose-1,5-bisphosphate (RuBP) with CO_2 to form two molecules of 3-phosphoglycerate (3-PGA) [20]. The 3-PGA is then reduced to glyceraldehyde-3-phosphate (G3P) using ATP and NADPH generated from the light-dependent reactions. Most of the G3P is recycled to regenerate RuBP, while a portion is exported for the synthesis of sugars, starch, and other organic compounds [20].

Despite its evolutionary success, the CBB cycle has significant **engineering limitations**. RuBisCO is notoriously slow and exhibits a competing oxygenase activity that leads to the wasteful process of photorespiration [15] [20]. Furthermore, the cycle's high ATP demand and low carbon fixation efficiency make it less ideal for many industrial bioprocesses. Engineering the CBB cycle into heterotrophic model organisms like *E. coli* has been achieved, but challenges remain in achieving high flux due to kinetic and regulatory incompatibilities [15].

The Wood-Ljungdahl Pathway (WLP)

The WLP is a highly efficient, anaerobic pathway utilized by acetogenic bacteria. It is characterized by its linear, non-cyclic structure and its direct production of the central metabolite **acetyl-CoA** from two one-carbon units [15]. The pathway operates through two parallel branches: the **methyl branch** reduces CO_2 to a methyl group tethered to a tetrahydrofolate carrier, while the **carbonyl branch** reduces CO_2 to carbon monoxide. These two units are then combined with coenzyme A by the acetyl-CoA synthase enzyme to form acetyl-CoA [15].

The primary engineering challenge for the WLP in industrial applications is its **bioenergetic constraint**. Acetogens often face ATP limitation during growth on gaseous substrates like CO_2/CO or syngas due to low substrate solubility and a lack of substrate-level phosphorylation sites in the pathway itself [15]. Overcoming this limitation through strategies such as coupling to ATP-yielding reactions or improving gas mass transfer is a major focus of current research to harness acetogens for the production of biofuels and chemicals beyond acetate.

Synthetic C1 Assimilation Pathways

To overcome the limitations of natural pathways, researchers have designed novel, synthetic routes for C1 assimilation. These pathways are engineered from first principles or by combining enzymes from diverse organisms, with goals such as minimizing ATP consumption, shortening the route to key metabolites, and improving thermodynamic driving force.

Table 2: Overview of Engineered Synthetic C1 Pathways

Pathway Name	Key Substrate	Target Product	Theoretical Advantage	Demonstrated In
Synthetic Acetyl-CoA (SACA) Pathway	CO_2	Acetyl-CoA	Direct synthesis from CO_2 [19]	Enzyme design [19]
Reductive Glycine Pathway (rGlyP)	Formate, CO_2	Glycine, Serine	Linear structure; simplicity [15] [18]	*E. coli*, *S. cerevisiae* [15]
CETCH Cycle	CO_2	Glyoxylate, Malate	In vitro function; oxygen-insensitive [16]	In vitro system [16]
Formolase (Fls) Pathway	Formaldehyde	Dihydroxyacetone, Sugars	Direct C-C bond formation [16] [19]	In vitro cascade [16]
Phosphoketolase (PHK) Pathway	F6P, X5P (from sugars)	Acetyl-CoA	Bypasses standard glycolysis; increases precursor supply [2]	*S. cerevisiae*, *Yarrowia lipolytica* [2]

Design Principles and Implementation

The design of synthetic pathways leverages computational tools to predict feasibility and optimize performance. **Flux Balance Analysis (FBA)** is used to model steady-state metabolic fluxes and identify potential bottlenecks, while **Minimum-Maximum Driving Force (MDF)** analysis helps select pathway variants with the strongest thermodynamic driving forces to ensure high flux [18]. A key design principle is **orthogonality**—creating a pathway with minimal overlap with the host's native regulation to avoid unintended interference and allow for independent control of the carbon flux [18].

Successful implementation requires advanced genetic tools. This includes the use of **carbon source-responsive promoters** (e.g., xylose-inducible promoters) to dynamically control gene expression [21], **biosensors** to monitor intracellular metabolite levels like NADPH, and **protein engineering** to optimize the kinetics and specificity of heterologous enzymes [17] [21]. For example, engineering the first *E. coli* strain to use methanol as a sole carbon source required the implementation of the rGlyP and extensive optimization to overcome metabolic imbalances [16].

Case Study: The Reductive Glycine Pathway (rGlyP)

The rGlyP is a prominent example of a synthetic pathway that has been successfully implemented in model heterotrophs. It is a linear pathway that assimilates formate and CO_2 into central metabolism via glycine and serine. Formate is first condensed with tetrahydrofolate (THF) and then reduced to a methylene group, which is combined with CO_2 and ammonia to form glycine. Two glycine molecules can then be converted into serine, a three-carbon metabolite, which feeds directly into the CCM [15] [18].

The rGlyP is considered promising because it is **non-cyclic** and relatively simple compared to the CBB cycle. It also benefits from using formate, a liquid C1 substrate that is easier to handle than gases and can be produced efficiently via electrochemical CO_2 reduction [18]. However, the pathway faces **thermodynamic hurdles**, particularly in the initial activation of formate, which requires sophisticated engineering to overcome [15].

Comparative Analysis: Natural vs. Synthetic Pathways

The choice between natural and synthetic C1 assimilation pathways involves a multi-factorial trade-off. Natural pathways are biologically proven and can be harnessed in their native hosts, which are often robust and adapted to C1 growth. However, these native hosts are frequently less genetically tractable than established industrial workhorses like *E. coli* and *S. cerevisiae*. Furthermore, natural pathways often feature complex regulation and inherent inefficiencies, such as the slow kinetics of RuBisCO or the ATP limitation of the WLP.

Synthetic pathways offer the potential for **superior efficiency**. They can be designed to be shorter, have higher energy efficiency, and produce key precursors like acetyl-CoA more directly [18] [19]. Their orthogonal nature can also simplify metabolic control. The primary drawback is that they are new to the cell, and their implementation can be a monumental engineering task, requiring the stable expression and coordination of multiple heterologous enzymes, and often resulting in initial imbalances that impede growth [18]. The following diagram synthesizes the logical workflow for selecting and engineering these pathways, from initial assessment to final strain optimization.



Diagram 1: C1 Pathway Selection and Engineering Workflow. This diagram outlines the key decision points and engineering steps in developing a microbial platform for C1-based biomanufacturing, from initial bioprocess definition to final strain optimization and scale-up.

Experimental Protocols for C1 Pathway Engineering

Protocol: Engineering a Heterologous Pathway into a Model Organism

This protocol outlines the key steps for introducing and optimizing a synthetic C1 assimilation pathway, such as the rGlyP or the PHK pathway, into a heterotrophic chassis like *E. coli* or *S. cerevisiae*.

- **Pathway Selection and Design**:
  - Based on the target C1 substrate and desired product, select a candidate pathway. Use computational models (FBA, MDF) to assess theoretical yield, energy demands, and thermodynamic feasibility [18].
  - Design a genetic construct encoding all necessary enzymes. Codon-optimize genes for the host organism. Consider constructing polycistronic operons or using multiple plasmids for expression.
- **Strain Construction**:
  - Transform the host strain with the constructed genetic module.

- For chromosomal integration, use CRISPR-Cas9 or other recombinase systems to insert the pathway genes into designated genomic loci under the control of appropriate promoters [21].

- **Promoter and Expression Optimization**:
  - Characterize a library of promoters with varying strengths and regulatory responses (e.g., constitutive, substrate-inducible) [21]. For example, replace standard promoters with **xylose-responsive promoters** (e.g., pADH2, pSFC1) to dynamically control gene expression during growth on xylose [21].
  - Employ protein engineering (e.g., site-directed mutagenesis) or test heterologous enzyme variants to improve catalytic efficiency and overcome kinetic limitations [17] [16].

- **Metabolic Balancing and Adaptive Laboratory Evolution (ALE)**:
  - Address metabolic imbalances by modulating competing pathways. This may involve knocking out genes that divert key intermediates or overexpressing bottleneck enzymes in the host's central metabolism [2].
  - Subject the engineered strain to ALE under selective pressure (e.g., using the C1 compound as the sole carbon source). Monitor for improved growth and substrate consumption. Sequence evolved strains to identify beneficial mutations [18].

- **Flux Analysis and Validation**:
  - Use **^{13}C-tracer analysis** (e.g., with $^{13}CO_2$ or ^{13}C-formate) to quantify carbon flux through the engineered pathway and central metabolism [22] [18].
  - Measure extracellular metabolites (substrate consumption, product formation) and intracellular cofactor levels (ATP/ADP, NADPH/$NADP^+$) to fully characterize the strain's physiological state [15].

The Scientist's Toolkit: Key Reagents and Solutions

Table 3: Essential Research Reagents for C1 Metabolic Engineering

Reagent / Tool Category	Specific Examples	Function / Application	Reference
C1 Substrates	$^{13}CO_2$, ^{13}C-Methanol, ^{13}C-Formate	Tracer studies for flux analysis; selective pressure during ALE.	[22] [18]
Genetic Parts	Xylose-responsive promoters (pADH2, pSFC1), Constitutive promoters (pTEF1), CRISPR-Cas9 systems	Precision control of gene expression; genome editing.	[21]
Host Chassis	*E. coli*, *S. cerevisiae*, *Cupriavidus necator*, *Pseudomonas putida*	Model organisms with extensive genetic toolkits and known physiology.	[18] [16]
Key Enzymes	Phosphoketolase (PK), Formolase (Fls), RuBisCO, Methanol Dehydrogenase (MDH)	Critical catalysts for synthetic and natural C1 pathways.	[2] [16] [19]
Biosensors	NADPH biosensors, Fatty acyl-CoA biosensors	Real-time monitoring of intracellular metabolite levels for high-throughput screening.	[21]
Analytical Software	Flux Balance Analysis (FBA), Minimum-Maximum Driving Force (MDF)	*In silico* prediction of metabolic flux and pathway thermodynamics.	[18]

The parallel development of natural and synthetic C1 assimilation pathways represents a powerful, dual-pronged strategy for advancing sustainable biomanufacturing. While natural pathways in their native hosts offer immediate, biologically proven systems for specific applications, the future likely belongs to the **rational design and engineering** of synthetic pathways in highly tractable, robust industrial chassis. The integration of **artificial intelligence** for protein design and pathway optimization, coupled with high-throughput genome editing and biosensor-enabled screening, is set to dramatically accelerate the design-build-test-learn cycle [17].

The ultimate success of C1-based technologies will depend not only on biological breakthroughs but also on the integrated consideration of **techno-economic analysis (TEA)** and **life cycle assessment (LCA)** at the earliest stages of research and development [18]. Ensuring that C1 feedstocks are derived from renewable sources and that the overall process is economically viable and environmentally sustainable is paramount. By bridging foundational discoveries in C1 and central carbon metabolism with scalable industrial applications, these engineering strategies hold the key to unlocking a carbon-neutral bioeconomy, turning the problem of greenhouse gases into the foundation for the next generation of chemical production.

Central Carbon Metabolism (CCM), comprising **glycolysis**, the **tricarboxylic acid (TCA) cycle**, and the **pentose phosphate pathway (PPP)**, serves as the most fundamental metabolic process for cellular energy generation and biosynthetic precursor supply [23] [1]. This network not only breaks down carbohydrates to produce **ATP** but also generates and manages critical redox cofactors—**NADH** and **NADPH**—which are essential for

maintaining cellular redox homeostasis and supporting anabolic reactions [24] [1]. The interplay between energy production and redox balance forms a critical foundation for all cellular activities, and its engineering has become a pivotal strategy in metabolic engineering for industrial biotechnology and therapeutic development [23] [25].

The NAD+/NADH redox couple primarily regulates **cellular energy metabolism**, functioning as a key hydride carrier in glycolysis and mitochondrial oxidative phosphorylation [24]. In contrast, the NADP+/NADPH couple primarily maintains **redox balance** and supports **biosynthetic processes** such as fatty acid and nucleic acid synthesis [24] [26]. The balance between these redox couples, together with ATP availability, influences critical cellular decisions including metabolic reprogramming, stress adaptation, and cell fate [24] [27]. Understanding and manipulating these relationships within CCM provides powerful levers for optimizing microbial cell factories for chemical production and addressing pathological conditions associated with metabolic dysregulation [23] [26].

Fundamental Roles of ATP, NADPH, and NADH in CCM

ATP: The Universal Energy Currency

ATP serves as the **primary energy carrier** in cellular systems, coupling energy-releasing catabolic processes with energy-requiring anabolic reactions [1]. In CCM, ATP is generated through two principal mechanisms: **substrate-level phosphorylation** during glycolysis and the TCA cycle, and **oxidative phosphorylation** via the electron transport chain [1]. The ATP/ADP ratio reflects the cellular energy status and allosterically regulates key metabolic enzymes, such as phosphofructokinase in glycolysis, creating feedback loops that maintain energy homeostasis [1].

NADH: The Redox Link to Energy Production

NADH functions as the **central electron carrier** in catabolic processes, collecting reducing equivalents from carbon oxidation in glycolysis and the TCA cycle [24] [28]. These reducing equivalents are subsequently channeled to the mitochondrial electron transport chain, where ATP generation is coupled to NADH reoxidation [1]. The NAD+/NADH ratio represents the **cellular redox state** and influences numerous metabolic pathways, including the flux through the TCA cycle [28]. In rat liver, the total NAD+ + NADH pool is approximately 1 μmole per gram wet weight, about ten times larger than the NADP+ + NADPH pool [28]. The mitochondrial compartment contains 40-70% of total cellular NAD+, highlighting its central role in energy metabolism [28].

NADPH: The Reducing Power for Biosynthesis and Defense

NADPH serves as the **primary reducing agent** for anabolic biosynthesis and oxidative stress defense [24] [1]. The PPP represents a major source of NADPH, generating it through the oxidative phase where glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase catalyze NADP+ reduction [1]. NADPH provides essential reducing power for **lipid synthesis**, **nucleotide precursor production**, and maintenance of the **cellular antioxidant system** through regeneration of reduced glutathione and thioredoxin [24] [1]. The distinct functional specialization of NADH and NADPH, despite their similar structures, allows cells to independently regulate energy production and biosynthetic/defense processes [24].

Table 1: Key Characteristics of Adenine and Nicotinamide Co-factors in CCM

Cofactor	Primary Role	Major Production Pathways	Major Consumption Processes	Cellular Ratio
ATP	Energy transfer & currency	Glycolysis, TCA cycle, OXPHOS	Biosynthesis, active transport, signaling	High ATP/ADP (energy-rich)
NADH	Electron carrier for energy production	Glycolysis, TCA cycle	OXPHOS, redox reactions	High NAD+/NADH (oxidized)
NADPH	Reducing power for biosynthesis & defense	PPP, ME, IDH	Lipid & nucleotide synthesis, antioxidant systems	High NADPH/NADP+ (reduced)

Quantitative Analysis of Energy and Redox Cofactors

The concentrations and ratios of energy and redox cofactors provide critical insights into cellular metabolic states and regulatory control points. Quantitative analyses reveal that **total NAD(H)** pools typically exceed **NADP(H)** pools by approximately tenfold in mammalian cells, with rat liver containing about 1 μmole NAD+ + NADH per gram wet weight compared to much lower NADP+ + NADPH concentrations [28]. Intracellular NAD+ concentrations in animal cells are estimated at **~0.3 mM**, while yeast concentrations range between **1.0-2.0 mM** [28]. These pools are highly compartmentalized, with mitochondria housing **40-70%** of total cellular NAD+ [28].

The **NAD+/NADH ratio** reflects the catabolic redox state and influences flux through dehydrogenase-catalyzed reactions, while the **NADPH/NADP+ ratio** indicates the reductive capacity for anabolism and oxidative defense [24]. Measurements using genetically encoded

biosensors show compartment-specific differences, with nuclear NAD+ estimated at ~110 μM, mitochondrial NAD+ at ~90 μM, and cytosolic NAD+ at ~70 μM in U2OS cells [26]. The free NAD+ concentration in solution is significantly lower than total measurements suggest, as **>80% of NADH fluorescence in mitochondria** originates from protein-bound forms [28].

Table 2: Quantitative Profile of Energy and Redox Cofactors in Cellular Compartments

Parameter	Cytosol	Nucleus	Mitochondria	Overall Cellular
NAD+ concentration	~70 μM [26]	~110 μM [26]	~90 μM [26]	0.3-2.0 mM [28]
NAD+/NADH ratio	Variable by cell type & state	Correlated with cytosol	Independent regulation	Indicator of redox state [28]
NADPH/NADP+ ratio	Maintained in reduced state	Dependent on cytosol	Independent generation	High for anabolic capacity [24]
ATP/ADP ratio	High (~10:1) in energy-rich	Correlated with cytosol	Highest production site	Regulates PFK, PK [1]
Primary functions	Glycolysis, PPP, biosynthesis	Epigenetic regulation, DNA repair	TCA, OXPHOS, FAO	Energy & redox balance [1]

Engineering Strategies for Optimizing Energy and Redox Balance

The introduction of **heterologous metabolic pathways** has emerged as a powerful strategy for rewiring CCM to optimize energy and redox balance. The **phosphoketolase (PHK) pathway**, in particular, has demonstrated significant success in redirecting carbon flux and enhancing precursor supply [23]. This pathway directly converts fructose-6-phosphate and xylulose-5-phosphate to acetyl-phosphate and subsequently to acetyl-CoA, bypassing multiple steps in conventional glycolysis and PPP [23]. In S. cerevisiae, PHK pathway introduction increased farnesene production by 25% through enhanced acetyl-CoA supply and triggered CCM rearrangement [23]. Similarly, PHK expression in Yarrowia lipolytica corrected redox imbalance caused by NADPH overproduction and increased total lipid production by 19% [23].

Other heterologous pathways have shown complementary benefits. The **pyruvate dehydrogenase (PDH) pathway** from E. coli directly converts pyruvate to acetyl-CoA without ATP consumption, potentially conserving energy for other CCM reactions [23]. When modified for NADP+ dependence in S. cerevisiae, this pathway doubled acetyl-CoA levels [23]. ATP:citrate lyase (ACL) from Aspergillus nidulans provides an alternative route to acetyl-CoA directly from citrate in the TCA cycle, doubling mevalonate yield when expressed in S. cerevisiae [23]. These heterologous pathways demonstrate how CCM engineering can selectively enhance carbon flux toward desired products while reconfiguring energy and redox metabolism.

Modular Deregulation of CCM

Recent advances in **modular deregulation strategies** have enabled more systematic engineering of CCM for enhanced product yields. A 2025 study on S. cerevisiae demonstrated a multifaceted approach encompassing **five distinct engineering strategies** to overcome the tight regulation of central metabolism [25]. This included promoter engineering, transcription factor manipulation, biosensor implementation, heterologous enzyme introduction, and mutant enzyme expression [25]. By applying these tools to different CCM modules, researchers achieved a **4.7-fold increase** in 3-hydroxypropionic acid (3-HP) productivity from xylose compared to the initially optimized strain [25].

A key innovation involved characterizing and categorizing promoters based on their response to xylose versus glucose utilization [25]. **Xylose-responsive promoters** driving xylose isomerase expression resulted in faster cellular growth compared to constitutive promoters, demonstrating the advantage of matching regulatory elements with carbon source utilization [25]. This systematic approach to CCM engineering illustrates how coordinated manipulation of multiple regulatory layers can enhance flux through targeted pathways while maintaining cellular viability, effectively optimizing the energy and redox balance for bioproduction.

Redox Cofactor Engineering

Direct manipulation of **redox cofactor availability** represents another strategic approach for optimizing CCM. Introduction of the Deinococcus radiodurans response regulator DR1558 into E. coli altered the expression of CCM-related genes, inducing excess NADPH generation from the PPP and supplying cofactor requirements for PHB biosynthesis [23]. Similarly, engineering NADP+-dependent enzymes into pathways typically utilizing NAD+ can redirect redox flux, as demonstrated with the E. coli PDH pathway modified for NADP+ dependence in S. cerevisiae [23].

The strategic expression of **NADPH-generating enzymes** in the TCA cycle and PPP, coupled with heterologous pathway introduction, enabled engineered Pichia pastoris to produce 23.4 g/L of free fatty acids and 2.0 g/L of fatty alcohols [23]. These examples highlight how targeted manipulation of redox cofactor supply can remove metabolic bottlenecks and enhance production of valuable compounds. The integration of **biosensors for NADPH and fatty acyl-CoA** provides dynamic regulation capabilities, further optimizing the balance between cofactor generation and consumption [25].

Experimental Protocols for Analyzing Energy and Redox States

Metabolomics and Stable Isotope Tracing

Stable isotope tracing combined with **mass isotopologue distribution analysis** provides powerful methodology for investigating CCM flux and redox dynamics [29]. The protocol involves cultivating cells in media containing isotopically labeled substrates (e.g., ^13C-glucose or ^13C-xylose), followed by metabolite extraction and LC-MS analysis. For investigating redox metabolism in fission yeasts, researchers used this approach to demonstrate that S. japonicus operates a fully bifurcated TCA pathway, sustaining amino acid production while optimizing cytosolic NADH oxidation via glycerol-3-phosphate synthesis [29]. This method enables quantification of pathway fluxes through central metabolism, revealing how different organisms manage NADH/NAD+ and NADPH/NADP+ ratios under varying physiological conditions.

The experimental workflow includes: (1) cultivation of cells in minimal media with labeled carbon source; (2) rapid sampling and quenching of metabolism; (3) metabolite extraction using cold methanol/water solutions; (4) LC-MS analysis with appropriate separation columns; (5) computational analysis of mass isotopologue distributions using specialized software; and (6) metabolic flux modeling to infer intracellular reaction rates [29]. This approach revealed that S. japonicus utilizes the PPP as a glycolytic shunt, reducing allosteric inhibition of glycolysis and supporting biomass generation without respiration [29].

Genetically Encoded Biosensor Implementation

Genetically encoded biosensors enable real-time monitoring of energy and redox cofactors in living cells, providing dynamic information about metabolic states. The protocol involves engineering genetic circuits where promoter elements responsive to specific metabolites drive reporter gene expression, or using fluorescent protein fusions that change properties upon metabolite binding [25]. For CCM engineering, researchers have developed biosensors for **NADPH** and **fatty acyl-CoA** to monitor metabolic status and implement dynamic regulation strategies [25].

Implementation typically includes: (1) identification of suitable sensing elements (transcription factors or protein domains) that respond to the target metabolite; (2) fusion of sensing elements with output domains (e.g., fluorescent proteins); (3) validation of sensor response and dynamic range in model systems; (4) integration of sensors into production hosts; and (5) application of sensors for screening or dynamic control of metabolic pathways [25]. In yeast CCM engineering, biosensors enabled identification of optimal pathway expression levels and dynamic regulation of metabolic fluxes to balance redox cofactor supply and demand [25].



CCM Redox Analysis Experimental Workflow

Research Reagent Solutions for CCM Redox Studies

Table 3: Essential Research Reagents for CCM Energy and Redox Studies

Reagent/Category	Specific Examples	Research Applications	Key Functions in CCM Analysis
Isotope-Labeled Substrates	^13C-Glucose, ^13C-Xylose	Metabolic flux analysis [25] [29]	Tracing carbon fate, quantifying pathway fluxes, redox cofactor production
Genetically Encoded Biosensors	NADPH biosensors, Fatty acyl-CoA sensors [25]	Real-time metabolite monitoring	Dynamic measurement of redox cofactors, metabolic status reporting
Fluorescent Reporters	RFP, GFP [25]	Promoter characterization, gene expression	Quantifying promoter strength, pathway activity under different conditions
Heterologous Enzymes	Phosphoketolase (PK), Phosphotransacetylase (PTA) [23]	Pathway engineering	Creating synthetic glycolytic routes, optimizing acetyl-CoA production
Analytical Standards	NAD+, NADH, NADP+, NADPH	HPLC/LC-MS quantification	Absolute quantification of redox cofactors, method calibration

Reagent/Category	Specific Examples	Research Applications	Key Functions in CCM Analysis
Enzyme Assay Kits	NAD/NADH-Glo, NADP/NADPH-Glo	Redox cofactor quantification	Measuring pool sizes and ratios, determining redox states
Specialized Promoters	Xylose-responsive promoters (pADH2, pSFC1) [25]	Metabolic pathway control	Carbon source-responsive expression, dynamic metabolic engineering

Pathway Visualization of Redox Coupling in CCM



Redox Coupling in Central Carbon Metabolism

The intricate interplay between **ATP production**, **NADH oxidation**, and **NADPH generation** within CCM represents a fundamental nexus of cellular metabolism that can be engineered for biotechnological and therapeutic applications. Advances in **heterologous pathway engineering**, **modular deregulation strategies**, and **dynamic biosensor implementation** have dramatically enhanced our ability to manipulate these relationships for improved product yields [23] [25]. The continued development of tools for precise control of gene expression, enzyme activity, and metabolic flux will further accelerate engineering of customized CCM architectures optimized for specific output goals.

Future research directions will likely focus on **enhanced compartmentalization** of metabolic pathways, **orthogonal cofactor systems** to isolate engineered pathways from native regulation, and **machine learning approaches** to predict optimal engineering strategies [25]. As our understanding of the subcellular distribution and dynamics of NAD(H) and NADP(H) pools improves [24] [26], so too will our ability to precisely engineer these systems. The integration of **multi-omics datasets** with advanced computational models promises to unravel the complex regulatory networks governing energy and redox balance, enabling unprecedented control over cellular metabolism for applications ranging from sustainable chemical production to therapeutic interventions for metabolic diseases [26].

Advanced Toolkit for CCM Engineering: From Pathway Design to Dynamic Control

The engineering of **central carbon metabolism (CCM)** represents a frontier in metabolic engineering, enabling the conversion of sustainable feedstocks into valuable biochemicals. Introducing heterologous pathways into microbial chassis allows researchers to overcome native regulatory limitations and thermodynamic inefficiencies, creating optimized systems for industrial bioproduction. This technical guide examines three transformative pathways that have emerged as particularly promising for CCM engineering: the **phosphoketolase (PHK) pathway**, the **reductive glycine pathway (rGlyP)**, and related **synthetic C1 assimilation routes**. These pathways enable enhanced carbon flux direction, improve ATP efficiency, and facilitate the utilization of one-carbon (C1) substrates, addressing critical challenges in sustainable biomanufacturing. As pressure increases to develop carbon-efficient bioprocesses that reduce reliance on sugar-based feedstocks, these engineered pathways offer compelling solutions for a new generation of microbial cell factories [30] [31] [23].

The Phosphoketolase (PHK) Pathway

Pathway Architecture and Thermodynamic Advantages

The PHK pathway operates as a metabolic shortcut that bypasses several steps of the conventional glycolytic pathway, consisting of just two key enzymes: **phosphoketolase (PK)** and **phosphotransacetylase (PTA)**. This minimalist architecture provides significant thermodynamic and stoichiometric advantages over native metabolic routes. The pathway branches at critical nodal metabolites in central carbon metabolism, processing fructose-6-phosphate (F6P) or xylulose-5-phosphate (X5P) to directly produce **acetyl-phosphate (ACP)**, which is subsequently converted to **acetyl-CoA** [23].

From a bioenergetics perspective, the PHK pathway offers superior **ATP efficiency** compared to traditional pathways for acetyl-CoA generation. By circumventing the pyruvate dehydrogenase complex and associated decarboxylation steps, the pathway minimizes carbon loss as CO_2 while conserving energy that would otherwise be expended in metabolic transitions. This makes the PHK pathway particularly valuable for biosynthesis of acetyl-CoA-derived compounds, including lipids, polyhydroxyalkanoates, and various isoprenoids [23].

Key Enzymes and Reaction Mechanisms

Phosphoketolase catalyzes the key bypass reaction that defines this pathway, cleaving F6P or X5P using inorganic phosphate to produce acetyl-phosphate and either erythrose-4-phosphate (from F6P) or glyceraldehyde-3-phosphate (from X5P). This unique phosphorolytic cleavage mechanism differs fundamentally from the hydrolytic reactions of standard glycolysis. **Phosphotransacetylase** then completes the sequence by converting acetyl-phosphate to acetyl-CoA, simultaneously generating ATP and conserving the energy-rich thioester bond [23].

Table 1: Key Enzymes of the PHK Pathway

Enzyme	EC Number	Reaction Catalyzed	Cofactors	Key Features
Phosphoketolase (PK)	EC 4.1.2.9	F6P + Pi → acetyl-P + E4P	Thiamine pyrophosphate	Bypasses multiple glycolytic steps, determines carbon flux split
Phosphoketolase (PK)	EC 4.1.2.9	X5P + Pi → acetyl-P + G3P	Thiamine pyrophosphate	Gateway from PPP to acetyl-CoA production
Phosphotransacetylase (PTA)	EC 2.3.1.8	acetyl-P + CoA → acetyl-CoA + Pi	Coenzyme A	Energy conservation, substrate channeling

Implementation Strategies and Industrial Applications

Successful implementation of the PHK pathway requires careful consideration of host physiology and metabolic context. In *Saccharomyces cerevisiae*, introduction of the PHK pathway has been shown to **redirect carbon flux** from glycolysis toward the pentose phosphate pathway (PPP), significantly increasing the pool of **erythrose-4-phosphate (E4P)**. This precursor enhancement has proven particularly valuable for aromatic compound biosynthesis, with one study demonstrating a remarkable **135-fold increase in tyrosol production** and fed-batch fermentation achieving over **10 g/L of combined tyrosol and salidroside** [23].

Similar strategies have shown promise in oleaginous yeasts such as *Yarrowia lipolytica*, where PHK expression increased total lipid production by approximately **19%** by addressing redox cofactor imbalances. The pathway created a direct route toward NADPH-oxidizing lipid synthesis while simultaneously increasing acetyl-CoA availability. Further optimization through coupling with ATP:citrate lyase expression and NADPH regeneration systems enabled engineered *Pichia pastoris* strains to produce **23.4 g/L of free fatty acids** and **2.0 g/L of fatty alcohols** [23].

The PHK pathway has also demonstrated versatility in supporting production of diverse chemical classes. For protopanaxadiol (a ginsenoside precursor), PHK introduction combined with overexpression of transaldolase and transketolase increased yields to **152.37 mg/L**. For 3-hydroxypropionic acid biosynthesis, PHK pathway implementation increased production by **41.9%** while reducing glycerol byproduct formation by **48.1%**, ultimately achieving an impressive **864.5 mg/L** of 3-HP through additional metabolic engineering [23].



Figure 1: PHK Pathway Integration in Central Carbon Metabolism. The heterologous PHK pathway (blue) creates shortcuts from F6P and X5P to acetyl-CoA, bypassing multiple native metabolic steps (gray). Key intermediates E4P and G3P are highlighted in red.

The Reductive Glycine Pathway (rGlyP)

Pathway Fundamentals and C1 Assimilation Mechanisms

The reductive glycine pathway (rGlyP) has emerged as a particularly efficient synthetic route for **C1 feedstock assimilation**, enabling microbial growth on formate or methanol as sole carbon and energy sources. This linear pathway represents one of the most ATP-efficient mechanisms for C1 incorporation into central metabolism, operating through a series of reversible reactions that ultimately generate glycine and subsequently pyruvate [31].

The core rGlyP module centers on the **glycine cleavage/synthase system** operating in the reductive direction, which assimilates C1 units (as methylene-tetrahydrofolate) with CO_2 and ammonia to form glycine. The pathway then extends through serine biosynthesis, where a second C1 unit is incorporated to generate serine, which is subsequently converted to pyruvate through serine deaminase activity. This elegant two-step C1 assimilation mechanism enables complete carbon backbone synthesis from C1 substrates, supporting both biomass formation and product synthesis [31].

Engineering Formatotrophy and Methylotrophy

Recent breakthroughs have demonstrated the successful implementation of synthetic formatotrophy and methylotrophy in non-native host organisms through rGlyP engineering. In a landmark study, researchers engineered the industrially robust soil bacterium *Pseudomonas putida* to assimilate formate and methanol as sole carbon sources via the rGlyP. Initial strain optimization employed **adaptive laboratory evolution (ALE)** under mixotrophic conditions, leading to substantial reduction in doubling time. Key mutations emerged in both the synthetic pathway genes' promoter regions and within the native genome, highlighting the importance of global regulatory adaptation [30].

The engineering process involved integrating a formate dehydrogenase gene either plasmidically or chromosomally as a mini-Tn5 module, combined with growth-coupled selection. The resulting strain, *P. putida* rG·F, achieved **strict formatotrophic growth** with a doubling time of approximately **28 hours**. Subsequent replacement of the formate dehydrogenase with an engineered methanol dehydrogenase from *Cupriavidus necator*, followed by additional ALE, yielded a fully methylotrophic strain (*P. putida* rG·M) capable of growth on methanol with a doubling time of approximately **24 hours** [30].

Table 2: Performance Metrics of Engineered C1-Assimilating Strains

Host Organism	Pathway	C1 Substrate	Growth Rate (Doubling Time)	Key Engineering Strategy
Pseudomonas putida rG·F	rGlyP	Formate	~28 hours	Formate dehydrogenase integration + ALE
Pseudomonas putida rG·M	rGlyP	Methanol	~24 hours	Methanol dehydrogenase swap + ALE
S. cerevisiae (various)	PHK	Glucose (co-substrate)	N/A	Pathway introduction + precursor enhancement

Comparative Analysis with Natural C1 Assimilation Pathways

The rGlyP offers distinct advantages over natural C1 assimilation pathways such as the Calvin cycle, serine cycle, or ribulose monophosphate pathway. Its **linear architecture** minimizes catalytic steps while maximizing carbon efficiency, and its limited overlap with native metabolic networks in most industrial hosts reduces complications from unintended metabolic cross-talk. Furthermore, the pathway's operation under **microaerobic or anaerobic conditions** provides flexibility for bioreactor configurations and reduces energy demands for aeration [31].

When compared to other synthetic C1 assimilation strategies, the rGlyP demonstrates superior **ATP stoichiometry**, requiring approximately 20-40% less ATP per fixed carbon than competing pathways. This thermodynamic advantage translates directly to improved carbon conversion efficiency and higher theoretical yields for target products. However, challenges remain in achieving sufficient flux through the glycine synthase system, which represents the pathway's kinetic bottleneck in many engineered hosts [31].

Experimental Protocols and Methodologies

Pathway Implementation and Optimization Workflow

Successful implementation of heterologous pathways follows a systematic methodology beginning with careful **codon optimization** of heterologous genes for the target host organism. This initial step is followed by strategic integration into the host genome, typically selecting neutral sites or loci with known high expression. For the PHK pathway, this involves simultaneous introduction of phosphoketolase and phosphotransacetylase genes, often with ribosomal binding site engineering to balance expression levels [23].

Following initial construction, engineers employ **modular pathway optimization** through promoter swapping and RBS tuning to balance expression of individual enzymes. For the rGlyP, this includes careful coordination of formate dehydrogenase or methanol dehydrogenase expression with the glycine cleavage/synthase system components. Advanced strategies incorporate **biosensor-enabled screening** to identify high-flux variants, particularly when engineering C1 assimilation pathways where growth coupling enables direct selection [30] [31].



Figure 2: Experimental Workflow for Heterologous Pathway Implementation. The systematic approach progresses from initial genetic construction (yellow) through strain optimization (green) to final validation (blue).

Analytical Methods and Validation Techniques

Comprehensive validation of functional pathway implementation requires multi-level analytical approaches. **Metabolic flux analysis** using ^{13}C isotopic tracing provides quantitative assessment of carbon routing through engineered pathways, with particular importance for C1 assimilation routes where traditional metrics may be insufficient. For PHK pathway evaluation, key measurements include **intracellular acetyl-CoA pools**, **NADPH/NADP$^+$ ratios**, and **E4P availability** through dedicated sampling and LC-MS quantification [23].

For rGlyP-engineered strains, validation includes demonstration of **substrate-dependent growth** in minimal media with C1 compounds as sole carbon sources, accompanied by quantitative analysis of pathway intermediates (glycine, serine) and verification of labeled carbon incorporation from ^{13}C-formate or ^{13}C-methanol. Deeper mechanistic insights come from **multi-omics analyses**, including transcriptomics to identify adaptive mutations and proteomics to verify enzyme expression and post-translational modifications [30].

The Scientist's Toolkit: Essential Research Reagents

Table 3: Key Research Reagents for Heterologous Pathway Engineering

Reagent/Category	Specific Examples	Function/Application	Technical Considerations
Pathway Enzymes	Phosphoketolase (PK), Phosphotransacetylase (PTA)	PHK pathway implementation	Codon-optimize for host; balance expression levels
C1 Assimilation Enzymes	Formate dehydrogenase, Methanol dehydrogenase	rGlyP implementation for formatotrophy/methylotrophy	Engineer for correct cofactor specificity (NAD+/NADP+)
Expression Plasmids	pET, pRS, pBBR vectors, mini-Tn5 modules	Genetic construction & integration	Select appropriate origin, resistance, promoter strength
Analytical Standards	^{13}C-labeled formate, ^{13}C-methanol, acetyl-CoA, glycine	Metabolic flux analysis, quantification	Use isotope-labeled internal standards for accurate quantification
Selection Markers	Antibiotic resistance, auxotrophic complementation	Strain selection & maintenance	Consider marker recycling for sequential engineering
Culture Media	M9 minimal media, defined C1 substrate media	Functional validation under controlled conditions	Carefully control micronutrient composition for C1 growth

Future Perspectives and Research Directions

The ongoing development of heterologous pathways for CCM engineering continues to evolve toward more sophisticated and integrated approaches. Future research directions include the development of **dynamic regulatory systems** that automatically adjust pathway expression in response to metabolic status, thereby optimizing resource allocation while minimizing burden. For C1 assimilation pathways, major challenges remain in improving **kinetic efficiency** of the rate-limiting glycine synthase reaction, potentially through protein engineering or enzyme mining from exotic microbiomes [31].

The convergence of **machine learning approaches** with structural biology and systems biology promises to accelerate the design-build-test-learn cycle for these complex metabolic systems. Recent advances in predicting protein structures enable more informed enzyme selection and engineering, while pattern recognition in multi-omics data can identify non-intuitive optimization targets. The integration of these computational approaches with high-throughput experimental validation represents the next frontier in heterologous pathway optimization [32].

From an applications perspective, the combination of multiple heterologous pathways in single chassis organisms offers intriguing possibilities for **synergistic carbon utilization**. For example, coupling the PHK pathway's precursor amplification with rGlyP's C1 assimilation capability could enable hybrid metabolism that simultaneously maximizes carbon efficiency and product yield. As the toolkit for robust pathway implementation expands, so too will the range of viable host organisms, moving beyond traditional model systems to industrially proven platforms with innate stress tolerance and superior cultivation characteristics [30] [23].

The precise orchestration of gene expression at both transcriptional and translational levels is a cornerstone of modern metabolic engineering, enabling the rewiring of cellular machinery for bioproduction. This technical guide explores the integrated use of genetic elements—including promoters, ribosome binding sites (RBS), and CRISPR-based tools—for optimizing metabolic pathways, with particular emphasis on engineering central carbon metabolism. We provide a comprehensive analysis of current methodologies, quantitative performance data, detailed experimental protocols, and visual workflows to equip researchers with practical strategies for enhancing microbial cell factory performance. The convergence of these technologies creates a powerful toolkit for overcoming the kinetic and thermodynamic barriers that have traditionally constrained the overproduction of target chemicals in both prokaryotic and eukaryotic chassis.

Central carbon metabolism serves as the fundamental engine of the cell, channeling carbon flux from nutrients into energy, reducing power, and biosynthetic precursors. Engineering this core network is essential for efficient production of biofuels, pharmaceuticals, and commodity chemicals; however, its tight regulatory control poses significant challenges [11]. Transcriptional and translational control mechanisms provide the genetic dials

to rewire this metabolism with the required precision. Promoters initiate transcription with varying strengths and inducibility, ribosome binding sites (RBS) modulate translational efficiency, and CRISPR tools offer programmable regulation of both processes [33] [34].

The third wave of metabolic engineering leverages synthetic biology to design and construct complete metabolic pathways using standardized genetic parts [34]. This paradigm shift enables not only pathway optimization but also the installation of dynamic control systems that automatically balance cell growth with product synthesis. This whitepaper synthesizes current knowledge on these genetic control elements, providing a technical foundation for researchers engineering central carbon metabolism in microbial hosts.

Core Genetic Elements for Expression Control

Promoter Engineering

Promoters, as the initiation point of transcription, represent one of the most critical genetic components for metabolic engineering. Recent advances have expanded the promoter toolkit beyond host-specific elements to include cross-species functionality.

Table 1: Promoter Characteristics and Applications

Promoter Type	Host Organisms	Strength Range	Key Features	Applications
Psh series	*E. coli, B. subtilis, C. glutamicum, S. cerevisiae, P. pastoris*	1.4-1.6x baseline	Cross-species activity; UP element enhanced	Multi-host pathway expression; Chassis screening
UP-modified	*E. coli*	1.6x over Pbs	Binds RNA polymerase α-CTD; -45 to -60 bp upstream	Enhanced protein production; Metabolic pathway tuning
Synthetic bidirectional	*S. cerevisiae*	Varies by sequence	Co-expression of two genes; Library of 168 variants	Taxadiene and β-carotene pathways
Constitutive strong	Various	Varies by host	Minimal sequence; Consistent expression	Standardized metabolic engineering

Cross-species promoters represent a significant innovation, addressing the challenge of screening gene expression constructs across multiple microbial hosts. By integrating consensus motifs from both prokaryotic and eukaryotic systems, researchers have developed Psh promoters that maintain functionality across five different microbial species including both bacteria and yeasts [35]. The incorporation of UP elements—sequences located -45 to -60 bp upstream of the transcriptional start site that bind the RNA polymerase alpha subunit carboxy-terminal domain (α-CTD)—can enhance promoter activity in *E. coli* by up to 1.6-fold [35].

Ribosome Binding Sites and Translational Control

While promoters govern transcription initiation, ribosome binding sites control the efficiency of translation initiation in prokaryotic systems. The RBS sequence influences ribosomal binding affinity through its complementarity to the 16S rRNA, with optimization achieving translational efficiency variations spanning several orders of magnitude.

In eukaryotic systems, where Shine-Dalgarno sequences are absent, Kozak sequences surrounding the start codon perform a analogous function in translation initiation. Engineering these elements enables fine-tuning of protein synthesis rates without altering transcriptional activity or plasmid copy number, making them particularly valuable for balancing multi-enzyme pathways where stoichiometric ratios critically impact flux.

CRISPR-Based Transcriptional and Translational Tools

CRISPR systems have evolved from simple gene-editing tools to versatile platforms for precise gene expression control at both transcriptional and translational levels.

Table 2: CRISPR Tools for Metabolic Engineering

Tool	Mechanism	Key Features	Performance	Applications
dCas9 CRISPRi	Transcriptional repression	Binds DNA, blocks transcription; Genome-wide screens	Strong polar effects on operons	Multiplexed gene knockdowns; Flux balance analysis
dCas13 tlCRISPRi	Translational repression	Binds mRNA, blocks translation; RNA targeting	Up to 6-fold lower polar effects vs dCas9	Independent gene control in operons; Multi-gene regulation

Tool	Mechanism	Key Features	Performance	Applications
CRISPRa	Transcriptional activation	dCas9-effector fusions; Targeted upregulation	Tunable activation levels	Pathway enhancement; Dynamic regulation
Base/Prime editors	DNA sequence alteration	Single-nucleotide changes; DSB-free editing	High precision; Reduced off-target effects	Enzyme engineering; Regulatory element optimization

The distinctive advantage of dCas13-mediated translational repression (tlCRISPRi) lies in its ability to independently regulate individual genes within multi-gene operons, exhibiting up to 6-fold lower polar effects compared to dCas9-based transcriptional repression [36]. This capability is particularly valuable in organisms like *E. coli* where approximately 68% of all genes are organized in operons [36].

Experimental Protocols

Protocol: CRISPR-dCas13 Mediated Translational Repression

Application: Independent gene regulation in multi-gene operons; Fine-tuning metabolic pathways

Materials:

- Plasmid system for dCas13 expression (e.g., p15A vector with J23107 promoter)
- crRNA expression construct (e.g., ColE1 plasmid with J23119 promoter)
- *E. coli* MG1655 or target strain
- Appropriate antibiotics (Carbenicillin, Chloramphenicol, Kanamycin, Spectinomycin)

Method:

- Clone dCas13d and crRNA sequences from source plasmids (e.g., pXR001, pXR002) into appropriate expression vectors [36].
- Transform constructs into *E. coli* target strain using standard chemical competence protocols.
- Plate on LB-agar with appropriate antibiotics based on resistance markers.
- For repression assays, inoculate single colonies in defined medium (e.g., EZ-RDM) with antibiotics.
- Culture in 96-deep-well plates at 37°C with shaking (900 rpm) overnight.
- Measure fluorescence (for reporter systems) or mRNA/protein levels for endogenous targets.
- For kinetic studies, subculture stationary phase cultures to OD600 0.1 and monitor every 30 min for 16 h.

Optimization Parameters:

- Vary dCas13 expression levels through promoter strength modulation
- Test crRNA spacer lengths (22-30 nt) for efficiency
- Evaluate target site position relative to start codon and ribosomal binding site

Protocol: Cross-Species Promoter Characterization

Application: Universal genetic part development; Multi-chassis metabolic engineering

Materials:

- Psh promoter variants
- GFP reporter plasmids with different replication origins
- Microbial hosts: *E. coli* JM109, *B. subtilis* 168, *C. glutamicum*, *S. cerevisiae* CEN.PK2-1C, *P. pastoris* GS115
- Media: LB, YPD, BHI, SD (with appropriate supplements)

Method:

- Construct GFP-expression plasmids via Gibson assembly using standardized syntax.
- Transform into each microbial host using species-specific methods (chemical transformation, electroporation).
- Culture recombinants in 24-deep well plates with appropriate media and selection.
- Grow to mid-exponential phase (OD600 0.5-0.8) under standard conditions for each species.
- Wash cells twice in phosphate-buffered saline before fluorescence measurement.
- Quantify GFP fluorescence using plate reader (excitation 490 nm, emission 530 nm).
- Normalize fluorescence values to cell density (OD600).
- For *P. pastoris* AOX1 promoter controls, use methanol induction (1% v/v) for 24 h after glucose growth.

Analysis:

- Calculate promoter strength as fluorescence/OD600 ratio
- Compare activities across hosts to determine conservation of function
- Use flow cytometry (BD FACSAria III) for single-cell resolution of expression heterogeneity

Protocol: CRISPR-Cas9 Assisted Marker-Free Integration

Application: Rapid strain construction; Multi-locus pathway integration; Promoter characterization

Materials:

- YaliCraft toolkit or similar modular system [37]
- Cas9-gRNA helper plasmid
- Donor DNA with homology arms (40-60 bp)
- Cre-expressing *E. coli* strain for marker excision
- Yeast strain with potential NHEJ pathway deletions

Method:

- Design 20 bp gRNA sequence specific to target genomic locus using prediction tools.
- Assemble gRNA expression cassette via Golden Gate assembly or recombineering with 90-base oligonucleotide.
- Construct donor DNA with 40-60 bp homology arms flanking integration cassette.
- Co-transform Cas9-gRNA helper plasmid and donor DNA into yeast using standard transformation protocol.
- Plate on appropriate selection medium and incubate until colonies appear (2-3 days).
- Screen colonies by colony PCR and sequencing to verify correct integration.
- For marker recovery, passage positive colonies in non-selective medium to allow plasmid loss.
- If marker-free integration fails for essential/growth-impairing genes, switch to marker-based approach using Cre-lox system.

Troubleshooting:

- For slow-growing mutants, use auxotrophic markers for stringent selection
- For multi-copy integration, design gRNAs targeting repetitive genomic regions
- For difficult loci, extend homology arms to 80-100 bp to enhance recombination efficiency

Visualization of Workflows and Mechanisms



Figure 1: Decision Workflow for Selecting CRISPR Tools in Metabolic Engineering. This workflow guides researchers through the process of choosing appropriate CRISPR-based interventions for different metabolic engineering scenarios, from initial objective definition to final validation.



Figure 2: Pyruvate-Responsive Genetic Circuit for Dynamic Metabolic Control. This diagram illustrates the components and logic of a metabolite-responsive biosensor that dynamically regulates gene expression based on pyruvate levels, enabling autonomous balancing of growth and production phases.

The Scientist's Toolkit: Research Reagent Solutions

Table 3: Essential Research Reagents for Transcriptional/Translational Control Experiments

Reagent/Category	Specific Examples	Function/Application	Key Characteristics
CRISPR Plasmid Systems	p15A-dCas13 (J23107 promoter); ColE1-crRNA (J23119 promoter); YaliCraft toolkit	Programmable gene activation/repression; Marker-free integration	Modular design; Species-specific optimization; Episomal or integrated
Microbial Host Strains	*E. coli* MG1655 (reporter studies); *S. cerevisiae* CEN.PK2-1C; *P. pastoris* GS115; *C. glutamicum* ATCC 13032	Metabolic engineering chassis; Pathway implementation	Well-characterized genetics; Industrial relevance; Genetic tractability
Reporter Systems	sfGFP (ex/em: 485/528 nm); mRFP1 (ex/em: 540/600 nm); mTagBFP2 (ex/em: 400/450 nm)	Quantitative promoter/translational efficiency measurement	Different spectral properties; Rapid maturation; High stability
Selection Antibiotics	Carbenicillin (100 μg/mL); Chloramphenicol (25 μg/mL); Kanamycin (30 μg/mL); Geneticin/G418 (600 μg/mL)	Maintenance of episomal plasmids; Selective pressure	Host-specific resistance markers; Concentration optimization required
Analytical Instruments	Plate reader (e.g., Biotek Synergy HTX); Flow cytometer (e.g., BD FACSAria III); LC/MS (e.g., Agilent 6530 LC/Q-TOF)	High-throughput screening; Single-cell analysis; Metabolite quantification	Multi-well format capability; High sensitivity; Quantitative accuracy

Applications in Central Carbon Metabolism Engineering

The integration of transcriptional and translational control tools has demonstrated remarkable success in rewiring central carbon metabolism for enhanced bioproduction. Three key application areas highlight this convergence:

Dynamic Redirection of Carbon Flux

Pyruvate-responsive genetic circuits exemplify the sophisticated dynamic control now possible in eukaryotic chassis. By implementing the *E. coli* PdhR repressor system in *S. cerevisiae*, researchers have created bifunctional circuits that autonomously redirect carbon flux between cell growth and product synthesis phases [38]. This approach addresses the fundamental conflict between biomass accumulation and chemical production by automatically downregulating ethanol synthesis pathways when pyruvate accumulates, simultaneously restoring redox balance and enhancing precursor availability for target compounds like malic acid and 2,3-butanediol.

Multi-Gene Operon Engineering

The combination of dCas9 transcriptional activation (txCRISPRa) with dCas13 translational repression (tlCRISPRi) enables unprecedented precision in regulating individual genes within multi-gene operons [36]. This strategy activates an entire operon transcriptionally while simultaneously repressing translation of specific genes within the same transcriptional unit, effectively customizing the expression stoichiometry of native operons without architectural modifications. This approach has demonstrated efficacy in enhancing biosynthesis of medically relevant human milk oligosaccharides, achieving superior performance compared to transcriptional control alone.

Systems-Level Metabolic Modeling Integration

The implementation of CRISPR-based metabolic engineering is increasingly guided by systems-level modeling approaches. Genome-scale models and flux balance analysis identify optimal gene knockdown targets, while machine learning algorithms improve gRNA efficacy predictions [39]. This model-guided design is particularly valuable for central carbon metabolism engineering, where competing pathway interactions and complex regulation create challenges for intuitive design. The integration of computational prediction with experimental validation creates a virtuous cycle of design-build-test-learn that accelerates strain development.

The integrated toolkit for transcriptional and translational control—spanning promoter engineering, RBS optimization, and CRISPR technologies—provides metabolic engineers with an unprecedented capacity to rewire central carbon metabolism with precision. The quantitative data, experimental protocols, and visualization frameworks presented in this technical guide offer researchers practical resources for implementing these strategies in their microbial engineering programs. As these tools continue to evolve through improvements in cross-species compatibility, dynamic regulation, and model-guided design, they will undoubtedly unlock new frontiers in sustainable bioproduction, enabling the efficient conversion of renewable carbon sources to valuable chemicals, pharmaceuticals, and materials.

Metabolite-responsive biosensors represent a cornerstone of advanced metabolic engineering, enabling real-time monitoring and dynamic control of central carbon metabolism. These genetically encoded circuits allow microbial cell factories to autonomously redistribute metabolic flux, balancing the inherent conflict between cell growth and product synthesis. This technical guide explores the operational principles, design frameworks, and implementation strategies of these sophisticated regulatory tools. By providing detailed methodologies and quantitative performance data, we aim to equip researchers with the knowledge to deploy biosensor-mediated dynamic regulation for enhancing bioproduction efficiency, particularly within engineered central carbon pathways.

Metabolite-responsive biosensors are synthetic genetic circuits that enable cells to sense and respond to intracellular metabolic states, providing a powerful alternative to traditional static control strategies [40]. In metabolic engineering, the efficient production of valuable chemicals is often constrained by the inherent conflict between cell growth and product synthesis [40]. While static engineering strategies like gene knockouts and constitutive pathway overexpression are widely used, they frequently disrupt cellular homeostasis, leading to redox imbalances and toxic intermediate accumulation [40].

Dynamic regulation inspired by natural regulatory networks offers a more sophisticated approach to process optimization [40]. These systems can be externally controlled by light, temperature, or chemical inducers, or they can function autonomously as metabolite-responsive circuits that provide real-time feedback regulation based on the internal state of the cell [40]. This autonomous function is particularly valuable for controlling central carbon metabolism, where metabolite concentrations fluctuate significantly during fermentation processes.

The core function of metabolite-responsive biosensors involves three fundamental components: a sensing mechanism that detects specific intracellular metabolites, a signal transduction system that converts this detection into a regulatory signal, and an output module that modulates gene expression to achieve a desired cellular response [41]. This architecture allows engineers to program cells to make dynamic decisions that optimize metabolic flux toward desired products while maintaining cellular health and viability.

Operational Principles and Molecular Mechanisms

Fundamental Biosensor Architectures

Metabolite-responsive biosensors function through genetically encoded components that detect intracellular metabolites and translate their concentration into programmable gene expression outputs. The most common architecture employs a **repressed-repressor** system, where a metabolite-responsive transcription factor (MRTF) represses gene expression from its cognate promoter in the absence of the target metabolite [42]. When the metabolite is present, it binds to the MRTF, antagonizing its repression activity and allowing gene expression to proceed [42]. This architecture provides the foundation for most dynamic regulation systems in metabolic engineering.

Biosensor performance is quantified through several key parameters:

- **Minimum output**: The baseline expression level in the absence of the target metabolite
- **Maximum output**: The expression level under saturating metabolite concentrations
- **Dynamic range (DR)**: The ratio of maximum to minimum output [42]
- **Sensitivity**: The metabolite concentration required for half-maximal activation

These parameters are crucial for determining how a biosensor will perform in practical applications, and they can be significantly affected by cellular context and growth conditions [42].

Major Biosensor Classes

Transcription Factor-Based Biosensors Metabolite-responsive transcription factors (MRTFs) represent the most extensively utilized class of biosensors. These proteins naturally evolved to interact with various metabolites, causing conformational changes that alter their DNA-binding affinity [41]. For example, the Escherichia coli-derived transcription factor PdhR acts as a pyruvate-responsive repressor by binding to the pdhO site within its target promoter, blocking RNA polymerase recruitment and suppressing downstream gene expression [40]. Pyruvate binding induces a conformational change in PdhR, relieving this repression and allowing transcription to proceed. This mechanism has been successfully exploited in prokaryotes including Bacillus subtilis and E. coli to engineer pyruvate-responsive genetic circuits for enhanced production of various chemicals [40].

CRISPR-Integrated Systems CRISPR-based systems have recently been integrated with traditional biosensor architectures to enhance programmability and scalability. The fusion of transcription factor-based biosensors with CRISPR interference (CRISPRi) systems creates powerful genetic switches capable of precise, signal-dependent transcriptional regulation [43]. For instance, FndCas12a's unique ribonuclease (RNase) activity enables direct processing of CRISPR RNAs (crRNAs) from biosensor-responsive transcripts, creating a seamless link between metabolite sensing and targeted gene repression [43]. These systems address limitations of traditional TF-based systems, including limited scalability, labor-intensive engineering processes, and difficulties in reconfiguring ligand specificity [43].

RNA-Based Biosensors Regulatory RNAs, including riboswitches and engineered riboregulators, offer an alternative to protein-based biosensors. These systems typically function through metabolite-induced structural changes in RNA leader sequences that modulate translation initiation or transcription termination [41]. While not the focus of this guide, RNA-based biosensors provide advantages such as smaller genetic size and faster response times, though they often lack the amplification capability of protein-based systems.

Table 1: Comparison of Major Biosensor Classes

Biosensor Class	Sensing Mechanism	Advantages	Limitations	Example Applications
Transcription Factor-Based	Metabolite binding alters DNA affinity	High sensitivity, natural diversity	Limited scalability, context-dependent	Pyruvate sensing with PdhR [40]
CRISPR-Integrated	TF controls crRNA expression	Programmable, highly specific	Increased genetic complexity	CRISPRi-aided genetic switches [43]
RNA-Based	Metabolite-induced structural changes	Small genetic size, fast response	Limited amplification capability	Riboswitches for coenzyme B12 [41]

Quantitative Biosensor Performance Under varying Growth Conditions

Biosensor performance is intrinsically linked to cellular growth rate, a critical consideration for industrial applications where growth conditions vary significantly. Experimental data reveals that both minimum and maximum biosensor outputs generally decrease with increasing growth rates due to dilution effects from cellular volume expansion [42]. However, the dynamic range (DR) response to growth rate varies significantly between different biosensor systems.

Table 2: Growth Rate Dependence of Biosensor Dynamic Range

Biosensor System	Inducer	DR Trend with Increasing Growth Rate	Key Characteristics
TetR-Ptet	aTc	Increasing	Passive inducer transport [42]
LacI-PLacUV5	IPTG	Increasing	Passive inducer transport [42]
FadR-PAR	Fatty Acids	Decreasing	Active transport via FadD [42]

Research has demonstrated that for TetR and LacI-based biosensors, the minimum output decreases more rapidly than the maximum output with increasing growth rate, resulting in an overall increase in dynamic range [42]. In contrast, the FadR-based fatty acid sensor shows the opposite behavior, with the maximum output decreasing more rapidly than the minimum output, leading to reduced dynamic range at higher growth rates [42]. This difference appears linked to transport mechanisms, as fatty acids require active transport via FadD, whose concentration is growth-dependent, while aTc and IPTG diffuse passively into cells [42].

Kinetic modeling of these systems incorporates dilution effects from cellular growth and accounts for different metabolite transport mechanisms (passive diffusion versus active transport) [42]. These models reveal that the coupling between dynamic range and growth rate sensitivity presents a fundamental trade-off in biosensor design, with implications for deploying biosensors in varying bioprocessing conditions.

Implementation Protocols for Key Systems

Implementing a Pyruvate-Responsive Genetic Circuit in Eukaryotic Chassis

The functional transfer of prokaryotic transcription factors to eukaryotic hosts presents unique challenges due to complex regulatory networks and strict subcellular compartmentalization [40]. The following protocol outlines the implementation of an E. coli PdhR-based pyruvate-responsive system in Saccharomyces cerevisiae:

Strain and Plasmid Construction

- **Strain Selection**: Utilize Pdc-negative S. cerevisiae mutant strains (e.g., TAM strain) that accumulate substantial pyruvate during aerobic cultivation [40].
- **Nuclear Localization**: Engineer a nuclear localization signal (NLS) peptide fused to the N-terminus of PdhR to ensure proper subcellular targeting [40].
- **Promoter Engineering**: Clone the native PdhR-responsive promoter (containing the pdhO operator) upstream of your gene of interest.
- **Reporter Integration**: Implement green fluorescent protein (GFP) as a reporter for characterizing activation levels of the genetic circuit [40].

Culture Conditions and Validation

- **Culture Medium**: Use minimal medium containing 1 g/L $(NH_4)_2SO_4$, 1 g/L KH_2PO_4, 1 g/L K_2HPO_4, 0.2 g/L $MgSO_4 \cdot 7H_2O$, 0.1 g/L leucine, 0.02 g/L histidine, 0.02 g/L methionine, supplemented with glucose [40].
- **Fluorescence Detection**: Monitor GFP fluorescence (excitation: 395 nm, emission: 509 nm) relative to cell density to quantify biosensor activation [40].
- **Circuit Optimization**: Systematically optimize the number of pdhO operator boxes (e.g., 1-4 copies) in the output promoter, as research shows a direct correlation between operator copy number and circuit activation [40].

Application for Metabolic Engineering

- Implement the optimized circuit to dynamically control ethanol synthesis pathways in S. cerevisiae by placing pyruvate decarboxylase genes under PdhR regulation.
- For downstream product synthesis, place pathways for compounds like malic acid or 2,3-butanediol under this dynamic control system to enhance production titers [40].

Establishing a Quorum Sensing-Controlled Type I CRISPRi System

The integration of quorum sensing with CRISPR interference enables autonomous dynamic regulation based on cell density. The following protocol details the implementation in Bacillus subtilis:

System Components and Assembly

- **QS Selection**: Utilize the PhrQ-RapQ-ComA quorum sensing system from B. subtilis [44].
- **CRISPR System**: Employ a type I CRISPR system with cas3 gene deletion for interference without cleavage [44].
- **Vector Construction**: Streamline crRNA vector construction by incorporating direct repeat sequences upstream of spacer regions targeting genes of interest [44].

Optimization and Validation

- **Component Balancing**: Modulate expression levels of PhrQ, RapQ, and ComA components to enhance QS-CRISPRi efficacy [44].
- **Reporter Assays**: Use green fluorescent protein (monitor at excitation 395 nm, emission 509 nm) to quantify repression efficiency relative to cell density (OD_{600}) [44].
- **Fermentation Validation**: Test the system in 5-L fed-batch fermentations without precursor supplementation to evaluate industrial relevance [44].

Application for Metabolic Engineering

- Target key metabolic nodes such as citrate synthase (citZ) in the TCA cycle to balance cell growth and product synthesis for compounds like d-pantothenic acid [44].
- Alternatively, suppress metabolic flux through glycolysis to redirect carbon through the pentose phosphate pathway, enhancing riboflavin production [44].

Experimental Design and Data Analysis Workflows

The implementation of dynamic regulation requires careful experimental design and standardized analysis methods. Below is a generalized workflow for biosensor characterization:

Culture Conditions and Monitoring

- Inoculate transformed strains from single colonies into selective medium and incubate overnight at appropriate temperatures (e.g., 30°C for yeast, 37°C for E. coli/Bacillus) with shaking at 200 rpm [40] [44].
- Subculture overnight cultures (1% v/v inoculum) into fresh medium with appropriate inducers and continue incubation under monitored conditions.
- Measure cell density (OD600) and fluorescence at regular intervals using a plate reader, subtracting background absorbance from blank media controls [43].

Fluorescence Data Analysis

- Calculate fluorescence intensities by subtracting initial fluorescence (0 h) from measurements at specific time points.
- Determine ON-state fluorescence (ΔFluorescenceON) under inducer-present conditions and OFF-state fluorescence (ΔFluorescenceOFF) under uninduced conditions [43].
- Compute dynamic range as the ratio of ON-state to OFF-state fluorescence after normalizing for cell density variations.

Metabolic Flux Validation

- For central carbon metabolism interventions, employ instationary 13C-fluxomics (INST-MFA) to track metabolic rearrangements [45].
- Use 13C-labeled glucose or other carbon sources to trace metabolic fate and quantify flux distributions under biosensor control.
- Correlate flux changes with transcriptomic data to validate intended metabolic remodeling and identify potential compensatory mechanisms.

Research Reagent Solutions

The following table provides key reagents and their applications for implementing dynamic regulation systems:

Table 3: Essential Research Reagents for Biosensor Implementation

Reagent / Genetic Part	Function	Example Application	Key Considerations
Bacterial Transcription Factors (e.g., PdhR, FadR, TetR)	Metabolite sensing and DNA binding	Pyruvate-responsive circuits in eukaryotes [40]	Requires nuclear localization in eukaryotes [40]
Type I/II CRISPR Systems	Programmable transcriptional regulation	CRISPRi for gene repression [44] [43]	Type I shows reduced toxicity compared to Cas9 [44]
Terminator Filters	Reduction of basal transcription	Enhancing dynamic range in genetic switches [43]	Critical for minimizing leaky expression
Nuclear Localization Signal (NLS)	Subcellular targeting in eukaryotes	Functional transfer of prokaryotic TFs [40]	Essential for TF function in eukaryotic nuclei
Quorum Sensing Components (PhrQ, RapQ, ComA)	Cell density-responsive regulation	Autonomous dynamic control in Bacillus subtilis [44]	Enables inducer-free operation
Fluorescent Reporters (GFP, RFP)	Quantitative biosensor characterization	Real-time monitoring of circuit performance [40] [42]	Enables high-throughput screening

Signaling Pathway and Workflow Visualizations



Diagram 1: Integrated biosensor architecture with sensing, transduction, and regulation modules.



Diagram 2: Experimental workflow for biosensor implementation and validation.

Metabolite-responsive biosensors represent a transformative technology for dynamic control of central carbon metabolism, moving beyond static engineering approaches to create self-regulating microbial cell factories. The integration of traditional transcription factor-based systems with modern CRISPR technologies has significantly expanded the programmability and precision of these genetic circuits. However, successful implementation requires careful consideration of cellular context, particularly growth rate dependencies and host-specific adaptations.

As demonstrated through the protocols and data presented in this guide, these systems enable sophisticated metabolic engineering strategies that automatically balance growth and production phases, redirect carbon flux with minimal manual intervention, and enhance overall bioprocess efficiency. Future developments will likely focus on expanding the repertoire of detectable metabolites, improving orthogonality for multiplexed control, and enhancing robustness across varying industrial conditions. Through continued refinement and application, dynamic regulation systems will play an increasingly vital role in sustainable bioproduction and advanced metabolic engineering.

Modular Optimization Strategies for Pathway Engineering

Central carbon metabolism in industrial microorganisms represents a primary target for metabolic engineering, yet its native, tightly regulated nature poses a significant challenge for efforts aimed at increasing metabolic flux toward desired products [21]. Modular deregulation has emerged as a powerful strategic framework to overcome these inherent regulatory constraints. This approach involves partitioning the metabolic network into discrete, manageable functional units that can be independently optimized before reintegration into a high-performance whole system. Within the broader context of central carbon metabolism engineering, modular optimization enables researchers to systematically overcome kinetic bottlenecks, transcriptional limitations, and allosteric control mechanisms that naturally resist flux redistribution. This technical guide examines the core principles, methodologies, and applications of modular optimization strategies, with specific emphasis on recent advances in engineering *Saccharomyces cerevisiae* for efficient utilization of non-glucose carbon sources—a critical capability for sustainable bioproduction from lignocellulosic feedstocks [21] [46].

Core Principles of Modular Optimization

Modular optimization in pathway engineering operates on several foundational principles that distinguish it from traditional monolithic engineering approaches:

- **Functional Encapsulation**: Metabolic pathways are divided into discrete functional modules based on their biochemical transformations, regulatory networks, or contribution to overall system objectives. Common modularizations separate substrate utilization, central metabolic flux, and product synthesis segments [21].
- **Independent Optimizability**: Each module can be engineered, characterized, and optimized with minimal interference to other system components, enabling parallel development and reducing combinatorial complexity during strain construction.
- **Interface Standardization**: Well-defined metabolic intermediates serve as interfaces between modules (e.g., acetyl-CoA connecting central metabolism to product formation pathways), allowing modular exchange and orthogonal optimization [21].
- **Systemic Integration**: Once individually optimized, modules are reintegrated with careful attention to intermodular flux balancing and regulatory harmony to prevent emergent bottlenecks at module interfaces.

In practice, these principles were effectively demonstrated in a recent study engineering *S. cerevisiae* for xylose utilization, where researchers divided the central carbon metabolism into three distinct modules for systematic deregulation: (1) xylose assimilation, (2) central metabolic pathways, and (3) product synthesis modules [21]. This structured approach enabled the implementation of five distinct engineering strategies tailored to each module's specific constraints and requirements.

Key Engineering Strategies and Methodologies

Promoter Engineering for Transcriptional Control

Precise control of gene expression represents a cornerstone of modular optimization, particularly when engineering pathways for non-native substrates like xylose. Recent work has systematically characterized promoter behavior under target cultivation conditions to identify elements with desired expression profiles [21].

Experimental Protocol: Promoter Characterization

- **Transcriptomic Analysis**: Perform RNA-seq on host strains cultivated in target versus reference conditions (e.g., xylose vs. glucose minimal medium) to identify natively regulated promoters.
- **Fluorescence-Based Screening**: Clone selected promoter candidates upstream of fluorescent reporter genes (e.g., RFP, GFP) and integrate into a neutral genomic locus.
- **Quantitative Assessment**: Measure fluorescence intensity across growth phases in target conditions using flow cytometry or plate readers.
- **Categorization**: Classify promoters based on response patterns:
  - Substrate-responsive promoters (e.g., xylose-responsive)
  - Constitutive promoters with stable expression
  - Repressed promoters under target conditions
- **Validation**: Test top candidates by driving expression of pathway genes and assessing functional outcomes (e.g., growth rates, substrate consumption) [21].

This approach identified xylose-responsive promoters such as pADH2 and pSFC1, which when used to control xylose isomerase expression, resulted in faster cellular growth on xylose compared to constitutive promoter pTEF1 [21].

Transcription Factor Manipulation

Master regulatory transcription factors governing carbon metabolism present valuable targets for modular optimization. Engineering strategies include:

- **Overexpression** of activators for target pathways
- **Deletion or knockdown** of repressors that constrain flux
- **Mutagenesis** to alter DNA-binding specificity or effector response
- **Domain swapping** to create chimeric regulators with novel properties

Implementation requires identification of key regulators through chromatin immunoprecipitation, transcriptomics, or genetic screens, followed by precise genetic manipulation using CRISPR/Cas9 or homologous recombination systems.

Heterologous Enzyme Implementation

Introducing non-native enzymes can bypass endogenous regulatory constraints or kinetic limitations:

Experimental Protocol: Heterologous Enzyme Evaluation

- **Candidate Identification**: Mine literature and databases for enzymes with desired catalytic properties (e.g., higher specific activity, reduced regulation).
- **Codon Optimization**: Optimize coding sequences for expression in host organism.
- **Expression Testing**: Evaluate functional expression using plasmid-based systems.
- **Activity Assessment**: Measure in vivo flux through the alternative pathway.
- **Host Adaptation**: Employ directed evolution if necessary to improve performance in host context.

In the xylose utilization case, introduction of xylose isomerase (XI) and xylulokinase (XK) enabled direct conversion of xylose to xylulose-5-phosphate, bypassing the native oxidoreductase pathway [21].

Mutant Enzyme Expression

Protein engineering creates enzyme variants with altered regulatory properties:

- **Allosteric site mutagenesis** to remove feedback inhibition
- **Catalytic site engineering** to improve kinetics
- **Phosphorylation site modification** to eliminate regulatory control
- **Chimeric enzyme construction** to combine beneficial properties

Implementation requires structural knowledge, targeted or random mutagenesis, and high-throughput screening systems to identify beneficial variants.

Biosensor Construction for Metabolic Monitoring

Metabolite-responsive biosensors enable real-time monitoring of metabolic status and dynamic regulation:

Experimental Protocol: Biosensor Implementation

- **Sensing Element Selection**: Choose transcription factors or RNA aptamers that respond to target metabolites.
- **Output Element Connection**: Link sensing element to measurable outputs (e.g., fluorescence).
- **Characterization**: Determine dynamic range, sensitivity, and specificity in host background.
- **Application**: Employ for screening mutant libraries or implementing feedback control circuits.

Recent applications have included NADPH and fatty acyl-CoA biosensors to monitor cofactor balance and metabolic status [21].

Case Study: Engineering Xylose Utilization in S. cerevisiae

A comprehensive example of modular optimization comes from recent work engineering *S. cerevisiae* for efficient xylose conversion to acetyl-CoA-derived products [21]. The metabolic network was divided into three functional modules with specific optimization strategies applied to each.

Table 1: Modular Optimization Strategies for Xylose Utilization in S. cerevisiae

Module	Engineering Target	Strategies Applied	Key Genetic Modifications	Outcome
Module I: Product Synthesis	3-HP production from acetyl-CoA	Heterologous enzyme expression, enzyme splitting, mutant enzymes	Bifunctional MCR from *C. aurantiacus*, split McrN and McrCm with mutation	3-HP production increased to 1.2 g/L on glucose
Module II: Xylose Assimilation	Xylose to xylulose conversion	Promoter engineering, heterologous enzyme implementation	XI and XK expression under xylose-responsive promoters (pADH2, pSFC1)	Improved growth rate on xylose vs. constitutive promoter
Module III: Central Carbon Metabolism	Flux enhancement to acetyl-CoA	Transcription factor manipulation, biosensor construction, mutant enzymes	Multiple TF modulations, metabolic biosensors	4.7-fold increase in 3-HP productivity on xylose

Table 2: Quantitative Assessment of Engineering Strategies in the Case Study

Engineering Component	Baseline Performance	Optimized Performance	Fold Improvement	Assessment Method
3-HP Production (Xylose)	Reference strain	Engineered strain	4.7×	HPLC quantification

Engineering Component	Baseline Performance	Optimized Performance	Fold Improvement	Assessment Method
MCR Configuration	Wild-type MCR	Split McrN + McrCm	~16× (vs. wild-type)	Extracellular 3-HP titer
Promoter Strategy	Constitutive (pTEF1)	Xylose-responsive (pADH2)	Significant growth improvement	Growth rate analysis
Promoter Library Range	Not applicable	42-fold fluorescence range	42×	GFP intensity measurement

The experimental workflow for this comprehensive engineering effort is visualized below:



Essential Research Reagent Solutions

Successful implementation of modular optimization strategies requires specific genetic tools and experimental reagents. The following table catalogs key resources employed in the referenced case study and their functional significance.

Table 3: Essential Research Reagents for Modular Pathway Engineering

Reagent/Category	Specific Examples	Function/Application	Experimental Role
Specialized Promoters	pADH2, pSFC1, pTEF1	Transcriptional control tailored to carbon source	Xylose-responsive expression, constitutive control
Heterologous Enzymes	Xylose isomerase (XI), bifunctional MCR	Pathway bypass, novel functionality	Enable xylose catabolism, 3-HP production
Engineered Enzyme Variants	McrCm (mutant), split MCR components	Enhanced activity, reduced regulation	Improve catalytic efficiency, metabolic flux
Reporter Systems	RFP, GFP	Quantitative promoter characterization	Measure expression strength, dynamics
Biosensor Systems	NADPH biosensor, fatty acyl-CoA sensor	Real-time metabolic monitoring	Dynamic pathway regulation, screening
Analytical Standards	3-HP, xylose, metabolic intermediates	Product quantification and analytics	HPLC calibration, yield determination

Reagent/Category	Specific Examples	Function/Application	Experimental Role
Selection Markers	Antibiotic resistance, auxotrophic markers	Strain selection and maintenance	Selective pressure for genetic elements

Analytical and Validation Methods

Rigorous quantitative analysis is essential for evaluating the success of modular optimization efforts. The referenced study employed multiple analytical approaches to characterize strain performance.

Experimental Protocol: Product Quantification and Strain Validation

- **Culture Conditions**: Grow engineered strains in minimal medium with target carbon source (e.g., 20 g/L xylose) with appropriate controls.
- **Sampling**: Collect samples at regular intervals throughout growth phase for multiple analyses.
- **Substrate Consumption**: Quantify xylose depletion using HPLC with refractive index detection.
- **Product Formation**: Measure 3-HP accumulation via HPLC with comparison to purified standards.
- **Growth Kinetics**: Monitor optical density (OD600) to determine growth rates and biomass yield.
- **Calculations**: Determine key performance metrics including:
  - Volumetric productivity (g/L/h)
  - Specific productivity (g/OD/h)
  - Yield (g product/g substrate)
  - Metabolic flux (mmol/gDCW/h)

The relationship between metabolic engineering strategies and analytical outcomes is visualized below:



Modular optimization represents a paradigm shift in metabolic engineering, moving from sequential gene manipulations to systematic module-level engineering. The case study in engineering xylose utilization in *S. cerevisiae* demonstrates how partitioning metabolic networks into functional units enables targeted application of diverse engineering strategies including promoter engineering, transcription factor manipulation, heterologous enzyme implementation, mutant enzyme expression, and biosensor construction [21]. This approach achieved a 4.7-fold enhancement in 3-HP productivity from xylose, highlighting the power of modular strategies for overcoming the inherent regulatory constraints of central carbon metabolism. As synthetic biology tools continue to advance, particularly in DNA assembly, genome editing, and computational modeling, modular optimization will undoubtedly expand its impact toward more complex metabolic objectives and non-model host organisms. The integration of modular design with emerging capabilities in machine learning and automated strain engineering promises to accelerate development of microbial cell factories for sustainable bioproduction.

Engineering Non-Model Hosts and Polytrophs for Expanded Substrate Use

The expanding demand for sustainable bioprocesses has catalyzed a paradigm shift beyond traditional model microorganisms toward **non-canonical hosts** with unique metabolic capabilities. While model organisms like *Escherichia coli* and *Saccharomyces cerevisiae* benefit from extensive genetic toolkits and well-characterized physiology, they often lack the innate physiological robustness and specialized metabolic features required for efficient utilization of next-generation feedstocks [18] [47]. Engineering **polytrophic microorganisms**—those naturally capable of growing on diverse multi-carbon substrates but not typically on one-carbon (C1) compounds—represents a promising frontier for sustainable bioproduction [18]. These organisms offer untapped potential due to native metabolic properties, enzyme activities, and substrate tolerance that are difficult to engineer from first principles [47].

The imperative for this transition stems from the limitations of sugar-based feedstocks, which compete with food production and agricultural land [18]. C1 molecules such as **methanol, formate, methane, and carbon monoxide** can be derived from or converted to CO_2, enabling a circular carbon economy [18] [48]. Furthermore, non-model hosts such as *Zymomonas mobilis*, *Pseudomonas putida*, and *Cupriavidus necator* possess exceptional industrial characteristics including **toxicity tolerance, broad substrate spectrum, and high product yield** [49] [48]. This

technical guide provides a comprehensive framework for leveraging these organisms to expand substrate utilization capabilities, with emphasis on central carbon metabolism engineering strategies.

Strategic Framework for Host Selection and Engineering

Criteria for Selecting Non-Model Hosts

Selecting an appropriate microbial host requires careful consideration of multiple physiological and metabolic factors aligned with bioprocess requirements:

- **Substrate Utilization Range**: Prioritize organisms with native capability to consume diverse carbon sources, indicating metabolic flexibility [18] [49]
- **Tolerance to Inhibitors and Products**: Assess resilience to feedstock-derived inhibitors (e.g., lignin derivatives) and target products [49]
- **Genetic Accessibility**: Evaluate existing genetic tools and transformation efficiency, though this can be developed [49] [48]
- **Redox and Energy Metabolism**: Consider native cofactor regeneration capabilities and energy conservation mechanisms [18] [50]
- **Pathway Orthogonality**: Prefer hosts where engineered pathways minimally conflict with native metabolic fluxes [18]

Pseudomonas putida KT2440 exemplifies an ideal candidate with native **solvent tolerance, versatile metabolism** encompassing various organic compounds, and a unique cyclic architecture of central carbon metabolism that promotes redox balance [50]. Similarly, *Zymomonas mobilis* possesses exceptional characteristics including **high sugar uptake rate** and ethanol tolerance, though its dominant ethanol production pathway requires engineering for diversion to other products [49].

Bioprocess Considerations Influencing Strain Selection

The bioprocess context must inform host selection and engineering strategies from the outset:

- **Feedstock Characteristics**: Gaseous C1 compounds (CO_2, CO, CH_4) suffer from low solubility, requiring specialized bioreactor designs, while liquid C1 compounds (methanol, formate) circumvent mass transfer constraints [18] [47]
- **Oxygen Requirements**: Anaerobic or micro-oxic conditions preclude obligate aerobes, while aerobic processes favor organisms with efficient oxidative metabolism [18]
- **Downstream Processing**: Cellular properties that simplify product recovery (e.g., secretion capabilities, cell size) should be considered [18]
- **Sustainability Metrics**: Preliminary life cycle assessment (LCA) and techno-economic analysis (TEA) guide economically viable and environmentally sustainable design [18] [49]

Analytical and Computational Tools for Metabolic Investigation

Omics Technologies for Metabolic Characterization

Comprehensive characterization of non-model hosts requires multi-omics approaches to elucidate their metabolic networks:

- **Metabolomics**: Identifies intracellular metabolite pools and their dynamics under different growth conditions [18]
- **Fluxomics**: Quantifies metabolic reaction rates through techniques like ^{13}C-metabolic flux analysis (MFA), revealing carbon flow through central metabolism [49]
- **Transcriptomics**: Uncovers gene expression patterns and regulatory networks [18] [50]
- **Proteomics**: Determines enzyme abundance and post-translational modifications [18]

Integrated omics analysis of *Zymomonas mobilis* under aerobic versus anaerobic conditions revealed differential carbon channeling into acetate and acetoin, guiding metabolic engineering strategies [49]. Similarly, multi-omics profiling of *Pseudomonas putida* exposed to different carbon sources elucidated the HexR regulon and its role in metabolic adaptation [50].

Computational Modeling for Pathway Design

Genome-scale metabolic models (GEMs) are indispensable tools for predicting metabolic behavior and designing engineering strategies:

- **Flux Balance Analysis (FBA)**: Predicts steady-state flux distributions optimizing objectives like biomass formation or product synthesis [18]
- **Enzyme-Constrained Models (ecModels)**: Incorporate enzyme kinetics and proteome limitations for more realistic flux predictions [49]
- **Minimum-Maximum Driving Force (MDF)**: Identifies pathways with highest thermodynamic feasibility [18]
- **Enzyme Cost Minimization (ECM)**: Estimates optimal enzyme and metabolite concentrations minimizing protein investment [18]

The enzyme-constrained model eciZM547 for *Z. mobilis* correctly predicted proteome-limited growth at high glucose uptake rates and accurately simulated carbon distribution between acetate and acetoin, outperforming previous models [49]. Such models are particularly valuable for predicting the behavior of engineered pathways before implementation.

Engineering Strategies for Substrate Expansion

Implementing C1 Assimilation Pathways

Engineering synthetic C1 assimilation in polytrophs requires introducing specialized metabolic routes:

- **Reductive Glycine Pathway (rGlyP)**: Linear pathway with high flux potential for formate and CO_2 assimilation [18] [48]
- **Calvin-Benson-Bassham (CBB) Cycle**: Natural CO_2 fixation pathway implemented synthetically [48]
- **Ribulose Monophosphate (RuMP) Cycle**: Efficient methanol assimilation route [48]
- **Serine Threonine Cycle (STC)**: Synthetic pathway for C1 assimilation [48]

Linear and orthogonal pathways like rGlyP are typically simpler to implement than circular autocatalytic cycles, which require tight control at branch points to prevent intermediate depletion [18]. When engineering *E. coli* for synthetic methylotrophy, implementation of the RuMP cycle required careful balancing of gene expression to prevent metabolic bottlenecks [48].

Overcoming Native Metabolic Dominance

Many non-model hosts possess strong native pathways that compete with engineered routes, requiring specialized strategies:

- **Dominant-Metabolism Compromised Intermediate (DMCI)**: Introducing a mildly toxic but cofactor-imbalanced pathway to weaken native metabolism before engineering the final product pathway [49]
- **Promoter Replacement**: Substituting strong native promoters with inducible or weaker versions to reduce flux through competitive pathways [49]
- **CRISPR-Mediated Genome Editing**: Precisely deleting or modifying key genes in native dominant pathways [49]

The DMCI strategy successfully addressed the dominant ethanol production pathway in *Z. mobilis* by first introducing a 2,3-butanediol pathway creating cofactor imbalance, then engineering the resulting intermediate chassis for high-yield D-lactate production (≥140 g/L from glucose) [49].

Regulatory Network Engineering

Modifying transcriptional regulatory networks is essential for optimizing flux through engineered pathways:

- **Transcription Factor Manipulation**: Deleting or overexpressing regulatory genes to reprogram cellular metabolism [11] [50]
- **Promoter Engineering**: Designing synthetic promoters with desired strength and regulation [11]
- **Biosensor Implementation**: Developing metabolite-responsive regulators for dynamic pathway control [11]

In *P. putida*, HexR was found to exert extensive control beyond glycolysis, repressing glycolytic genes while activating gluconeogenic and glyoxylate shunt genes during growth on acetate [50]. Understanding such regulatory landscapes enables more effective engineering strategies.

Experimental Protocols and Workflows

Standardized Engineering Workflow for Non-Model Hosts

Implementing a systematic engineering pipeline is crucial for success with non-model organisms:



Systematic Engineering Workflow for Non-Model Hosts

Genetic Tool Development Protocol

Establishing genetic manipulation capabilities is foundational for engineering non-model hosts:

- **Transformation System Establishment**
  - Test electroporation, chemical transformation, and conjugation protocols
  - Identify functional selectable markers (antibiotic resistance, auxotrophic complementation)
  - Determine copy number and stability of plasmid vectors
- **CRISPR Tool Adaptation**
  - Identify compatible CRISPR-Cas systems (Type I, II, or V)
  - Optimize guide RNA design and expression
  - Establish repair mechanisms (homologous recombination, MMEJ, NHEJ)
- **Parts Characterization**
  - Isolate and characterize native promoters, RBS, terminators
  - Develop modular plasmid systems with standardized parts
  - Validate part functionality across growth conditions

For *Z. mobilis*, researchers developed efficient genome-editing tools based on heterologous CRISPR-Cas12a and endogenous Type I-F CRISPR-Cas systems with associated repair pathways like microhomology-mediated end joining (MMEJ) [49].

Metabolic Flux Analysis Protocol

Quantifying metabolic fluxes provides critical insights for engineering central carbon metabolism:

- **^{13}C-Labeling Experiment**

- Grow cells on ^{13}C-labeled substrate (e.g., [1-^{13}C]glucose)
- Harvest cells during mid-exponential phase
- Quench metabolism rapidly (cold methanol)
- Extract intracellular metabolites

- **Mass Spectrometry Analysis**
  - Analyze mass isotopomer distributions of metabolic intermediates
  - Determine labeling patterns in proteinogenic amino acids
  - Quantify fractional enrichment

- **Flux Calculation**
  - Use computational software (13CFLUX, INCA)
  - Incorporate stoichiometric constraints from metabolic model
  - Iteratively fit simulated and measured labeling patterns
  - Calculate net and exchange fluxes

13C-MFA in *Z. mobilis* ZM4 under aerobic conditions revealed significant carbon flux into acetate and acetoin, differing from anaerobic conditions and informing engineering strategies [49].

Performance Metrics and Case Studies

Quantitative Performance of Engineered Strains

Recent advances in engineering non-model hosts have yielded impressive results across various organisms and products:

Table 1: Performance Metrics of Engineered Non-Model Hosts

Host Organism	Engineering Strategy	Product	Titer (g/L)	Yield (g/g)	Substrate	Citation
Zymomonas mobilis	DMCI strategy	D-lactate	140.92	0.97	Glucose	[49]
Zymomonas mobilis	DMCI strategy	D-lactate	104.6	>0.97	Corncob residue hydrolysate	[49]
Pseudomonas putida	HexR regulon engineering	Muconic acid	Not specified	Improved	Acetate	[50]
E. coli (synthetic)	rGlyP implementation	Biomass	Not specified	Not specified	Formate/CO_2	[48]

Techno-Economic and Sustainability Assessment

Integrating economic and environmental analyses early in the engineering process is crucial for commercial viability:

- **Techno-Economic Analysis (TEA)**: Estimates production costs and identifies economic bottlenecks [18] [49]
- **Life Cycle Assessment (LCA)**: Evaluates environmental impacts across the entire process [18] [49]
- **Carbon Footprint Calculation**: Assesses greenhouse gas emissions compared to conventional processes [18]

TEA and LCA of D-lactate production from corncob residue hydrolysate using engineered *Z. mobilis* demonstrated both commercial feasibility and significant greenhouse gas reduction potential [49]. Such analyses should guide target selection and process optimization to ensure sustainability goals are met.

Research Reagent Solutions

Essential research tools and reagents for engineering non-model hosts:

Table 2: Key Research Reagents for Engineering Non-Model Hosts

Reagent/Tool	Function	Examples/Specifications
CRISPR-Cas Systems	Genome editing	Cas9, Cas12a, endogenous Type I-F systems [49]
Metabolic Models	Pathway prediction & analysis	iZM547, eciZM547 [49]
Omics Platforms	Systems-level characterization	RNA-seq, proteomics, metabolomics platforms [18] [49]
Flux Analysis Tools	Metabolic flux quantification	13C-MFA, flux balance analysis [18] [49]

Reagent/Tool	Function	Examples/Specifications
Genetic Parts	Pathway regulation	Native C1-inducible promoters, RBS libraries [18] [11]
Non-canonical Cofactors	Pathway insulation	Engineered cofactors to isolate synthetic pathways [51]

Visualization of Metabolic Engineering Strategies

Central Carbon Metabolism Engineering

Engineering central carbon metabolism in non-model hosts requires careful consideration of native pathway architecture and potential conflicts:



Metabolic Engineering Strategy Overview

Engineering non-model hosts and polytrophs for expanded substrate use represents a transformative approach for sustainable biomanufacturing. The strategies outlined in this technical guide—from host selection and metabolic characterization to pathway implementation and regulatory engineering—provide a roadmap for leveraging microbial diversity for industrial applications. Future advances will likely come from several emerging areas:

- **Automated Strain Engineering**: High-throughput platforms integrating design, construction, and testing cycles [50]
- **Machine Learning Applications**: Predictive models for part performance, enzyme design, and metabolic behavior [50]
- **Non-canonical Cofactor Engineering**: Orthogonal cofactor systems to insulate engineered pathways from native metabolism [51]
- **Consortium Engineering**: Multi-species systems distributing metabolic tasks across specialized organisms

As genetic toolboxes expand for diverse microorganisms, the synthetic biology community is poised to unlock the immense potential of non-model hosts, ultimately enabling bioprocesses that are both economically competitive and environmentally sustainable.

Overcoming Metabolic Hurdles: Balancing Flux, Cofactors, and Cellular Burden

Identifying and Alleviating Metabolic Bottlenecks and Flux Imbalances

Metabolic bottlenecks and flux imbalances are critical constraints in metabolic engineering that limit the efficient channeling of carbon through biosynthetic pathways, ultimately reducing product yield and productivity. These constraints arise from kinetic, thermodynamic, and regulatory limitations within the intricate network of cellular metabolism [52]. In the specific context of central carbon metabolism engineering, which serves as the core gateway for carbon distribution to various downstream pathways, identifying and alleviating these limitations is paramount for achieving industrial-scale production of bio-based chemicals and pharmaceuticals.

The tightly regulated nature of central carbon metabolism in production hosts like *Saccharomyces cerevisiae* presents significant challenges for engineering efforts aimed at increasing flux through targeted pathways [11]. This technical guide provides a comprehensive framework for identifying flux limitations using advanced computational and experimental approaches, with specialized methodologies for implementing targeted strategies to alleviate these constraints within central carbon metabolism engineering frameworks.

Theoretical Foundations: From Stoichiometric Models to Flux Imbalances

Flux Balance Analysis and the Steady-State Assumption

Constraint-based modeling, particularly Flux Balance Analysis (FBA), has emerged as a fundamental framework for studying metabolic networks at genome scale. FBA relies on the steady-state assumption, where intracellular metabolites are assumed to not accumulate or deplete over time, mathematically represented as:

$$\mathbf{S}\mathbf{v} = 0$$

where **S** is the stoichiometric matrix and **v** is the vector of metabolic fluxes [52]. This formulation enables prediction of reaction fluxes that satisfy mass conservation constraints while optimizing cellular objectives, typically biomass production for growing cells. FBA has demonstrated remarkable agreement with experimental data for wild-type cells in rich media [52].

Flux Imbalance Analysis and Shadow Prices

Flux Imbalance Analysis (FIA) extends FBA by examining the biological significance of deviations from the steady-state assumption. Mathematically, this sensitivity analysis is captured through the dual problem of the linear programming formulation, with corresponding variables known as **shadow prices** (λ_i) [53] [54].

Shadow prices quantify the change in the cellular growth rate upon a unit change in the steady-state constraint of a metabolite (b_i). Metabolites with **negative shadow prices** are identified as growth-limiting, indicating that allowing additional outflow would increase biomass production [54]. This theoretical framework provides a direct link between stoichiometric modeling and metabolite control analysis.

Table 1: Interpretation of Shadow Prices in Metabolic Models

Shadow Price Value	Biological Interpretation	Metabolite Status
Negative ($\lambda < 0$)	Metabolite is growth-limiting	Allowing depletion increases growth
Zero ($\lambda = 0$)	Metabolite non-limiting	No effect on growth upon depletion
Positive ($\lambda > 0$)	Metabolite accumulation beneficial	Increasing concentration boosts growth

Computational Methodologies for Identifying Bottlenecks

Constraint-Based Modeling Approaches

Constraint-based modeling provides multiple methodologies for identifying flux bottlenecks:

Flux Balance Analysis (FBA): Identifies optimal flux distributions maximizing cellular objectives. Reactions operating at maximum capacity in FBA solutions represent potential bottlenecks [52].

Flux Variability Analysis (FVA): Determines the range of possible fluxes for each reaction while maintaining near-optimal growth (typically ≥90% of maximum). Reactions with narrow flux ranges indicate tight metabolic control points [52].

Flux Imbalance Analysis (FIA): Utilizes shadow prices to identify metabolites whose imbalance most significantly impacts cellular growth. Metabolites with highly negative shadow prices represent prime targets for engineering [53] [54].

Network Analysis of Metabolic Flux

Advanced graph-based approaches provide systems-level analysis of metabolic bottlenecks. The **Mass Flow Graph (MFG)** framework constructs directed graphs where edges represent metabolite flow from source to target reactions, weighted by flux values obtained from FBA [55].

This methodology captures environment-dependent rerouting of metabolic flows and identifies critical choke-point reactions through network centrality measures. Unlike structural graphs, MFGs incorporate directional information and minimize bias from pool metabolites, providing more biologically relevant bottleneck identification [55].



Experimental Methodologies for Validation

Metabolomics for Experimental Flux Determination

Time-dependent metabolomics provides experimental validation of computational predictions. The protocol involves:

Sample Collection and Quenching: Rapid sampling of cultures followed by immediate quenching in cold methanol (-40°C) to arrest metabolic activity.

Metabolite Extraction: Using optimized solvent systems (e.g., methanol:acetonitrile:water 40:40:20) for comprehensive metabolite recovery.

LC-MS/MS Analysis: Reversed-phase or HILIC chromatography coupled to high-resolution mass spectrometry for metabolite separation and quantification.

Data Analysis: Tracking metabolite pool sizes over time following environmental perturbations. Metabolites with negative shadow prices exhibit lower temporal variation, confirming their tight regulation [53] [54].

Chemostat Cultivation for Growth Limitation Studies

Controlled nutrient limitation in chemostats enables systematic investigation of metabolic bottlenecks:

Culture Conditions: Maintain cells at steady-state growth under defined nutrient limitations (C, N, P, S).

Metabolite Analysis: Quantify intracellular metabolite pools under each limitation condition.

Correlation Analysis: Relate metabolite shadow prices to measured degrees of growth limitation. Experimental data from *Saccharomyces cerevisiae* demonstrates significant anticorrelation between shadow prices and growth limitation [53].

Table 2: Experimental Techniques for Bottleneck Identification

Method	Key Measurements	Bottleneck Indicators	Applications
Time-Resolved Metabolomics	Metabolite concentration changes over time	Low temporal variation in key metabolites	Transient response to perturbations
Chemostat Cultivation	Intracellular metabolite levels under nutrient limitation	Correlation with shadow prices	Nutrient-specific limitations
Fluxomics (13C-MFA)	Metabolic flux rates through pathways	Reactions operating at capacity	In vivo flux determination
Enzyme Activity Assays	Vmax, Km of enzymes	Low catalytic efficiency	Kinetic limitations

Engineering Strategies for Alleviating Bottlenecks

Modular Deregulation of Central Carbon Metabolism

A multifaceted engineering approach targeting multiple regulatory layers simultaneously has proven effective for alleviating flux bottlenecks. In a case study focusing on xylose utilization in *S. cerevisiae*, implementation of five complementary strategies achieved a 4.7-fold enhancement in 3-hydroxypropionic acid productivity [11]:

Promoter Engineering: Replacement of native promoters with constitutive or inducible variants to optimize enzyme expression levels.

Transcription Factor Manipulation: Engineering of regulatory proteins to overcome transcriptional repression.

Biosensor Implementation: Development of metabolite-responsive sensors for high-throughput screening of optimized strains.

Heterologous Enzyme Expression: Introduction of non-native enzymes with superior kinetic properties or regulation.

Mutant Enzyme Expression: Utilization of enzyme variants with alleviated allosteric regulation or improved catalytic efficiency.

Cofactor Engineering and Energy Supply Optimization

Cofactor imbalances frequently create thermodynamic bottlenecks in engineered pathways:

NAD/NADH Balancing: Expression of transhydrogenases or NADH-consuming pathways to maintain redox balance.

ATP Supply Enhancement: Optimization of ATP-generating pathways to support energy-intensive biosynthesis.

Cofactor Specificity Switching: Engineering enzymes to utilize more abundant or sustainable cofactor pools.

Pathway Analysis and Comparison Algorithms

Metabolic pathway comparison enables identification of evolutionary solutions to flux bottlenecks across organisms. Low-cost algorithms for pairwise comparison include:

Graph Transformation Approach: Conversion of 2D pathway graphs to 1D linear structures using breadth-first traversal, followed by sequence alignment techniques (global, local, or semi-global alignment) to quantify similarity [56].

Differentiation by Pairs Method: Calculation of numerical differentiation values based on the ratio of differences to total reactions, providing intuitive homology assessment [56].

These algorithms facilitate comparative analysis of bottleneck solutions in native versus engineered pathways, guiding strategic interventions.



The Scientist's Toolkit: Research Reagent Solutions

Table 3: Essential Research Reagents for Metabolic Bottleneck Analysis

Reagent/Resource	Function	Application Examples
Genome-Scale Metabolic Models (e.g., iML1515, iMM904)	Constraint-based flux prediction	FBA, FVA, FIA simulations
LP Solvers (e.g., COBRA, Gurobi, CPLEX)	Linear programming optimization	Solving FBA problems
Metabolomics Standards (isotope-labeled internal standards)	Quantitative mass spectrometry	Absolute metabolite quantification
Pathway Databases (KEGG, MetaCyc, BioCyc)	Reference metabolic pathways	Pathway comparison and analysis
CRISPR Interference/Activation Systems	Targeted gene regulation	Transcription factor manipulation
Metabolite Biosensors (Transcription factor-based)	High-throughput screening	Isolation of optimized variants
Heterologous Enzyme Libraries	Alternative pathway implementation	Bypassing native regulation

Integration with Broader Engineering Strategies

Successful alleviation of metabolic bottlenecks requires integration with broader central carbon metabolism engineering frameworks:

Energy Supply Augmentation: Coordination with ATP-generating pathways to support flux increases.

Redox Balancing: Implementation of systems to maintain cofactor homeostasis under high flux conditions.

Precursor Optimization: Engineering of upstream pathways to ensure adequate supply of building blocks.

Regulatory Network Editing: CRISPR-based modification of transcriptional regulators controlling central carbon metabolism.

As demonstrated in recent advances in microbial carbon dioxide fixation, overcoming kinetic and thermodynamic barriers through directed evolution of enzymes and pathway design represents the next frontier in metabolic bottleneck engineering [57].

The systematic identification and alleviation of metabolic bottlenecks through integrated computational and experimental approaches provides a powerful framework for optimizing central carbon metabolism in engineered strains. The continued development of more sophisticated analytical methods, combined with modular engineering strategies targeting multiple regulatory layers simultaneously, will further enhance our ability to design efficient microbial cell factories for pharmaceutical and industrial applications.

Managing Cofactor Regeneration and Redox Imbalances

Cofactor regeneration and redox balance are critical determinants of efficiency in engineered microbial cell factories. Imbalances in cofactors such as NAD(P)H/NAD(P)+ and ATP/ADP can lead to metabolic bottlenecks, reduced cell viability, and suboptimal production of target compounds. This technical guide examines advanced strategies for managing redox homeostasis, drawing upon recent developments in cofactor engineering, pathway design, and computational modeling. Framed within the broader context of central carbon metabolism engineering, this review provides researchers with experimental protocols, key reagent solutions, and visualization tools to overcome redox-related limitations in bioproduction pipelines.

Central carbon metabolism serves as the cornerstone of microbial biosynthesis, channeling carbon substrates toward energy generation, redox maintenance, and biomass formation. **Cofactors are crucial chemicals** that maintain cellular redox balance and drive synthetic and catabolic reactions, participating in approximately 1,500 enzymatic reactions in microbial metabolism [58]. The intracellular redox state is primarily determined by the ratio of NADH/NAD+ rather than that of NADPH/NADP+ [59]. When **redox imbalance occurs**—typically when glycolytic NADH production exceeds cellular oxidation capacity—it triggers overflow metabolism toward reduced byproducts like ethanol and glycerol, resulting in carbon and energy waste, cellular damage, and metabolic arrest [59].

Engineering redox balance through cofactor systems has emerged as a powerful approach to maximize carbon flux toward target metabolites while maintaining thermodynamic feasibility [60]. This guide synthesizes recent advances in cofactor engineering strategies, experimental methodologies, and computational tools to address redox imbalances in engineered microbial systems for pharmaceutical and chemical production.

Core Principles of Cofactor Balance

Key Cofactor Systems and Metabolic Functions

Table 1: Primary Cofactors in Microbial Metabolism and Their Roles

Cofactor	Primary Functions	Key Metabolic Pathways	Cellular Locations
ATP/ADP	Energy transfer, Metabolic regulation	Glycolysis, TCA cycle, Oxidative phosphorylation	Cytoplasm, Mitochondria

Cofactor	Primary Functions	Key Metabolic Pathways	Cellular Locations
NADH/NAD+	Catabolic redox carrier, Electron donor for respiration	Glycolysis, TCA cycle, Fermentation	Cytoplasm, Mitochondria
NADPH/NADP+	Anabolic reducing power, Antioxidant defense	Pentose phosphate pathway, Fatty acid synthesis	Cytoplasm
Acetyl-CoA	Carbon carrier, Metabolic precursor, Protein modification	Glycolysis, TCA cycle, Fatty acid metabolism	Cytoplasm, Mitochondria

The **ATP/NADPH demand ratio** varies significantly across biological processes. For instance, the Calvin-Benson-Bassham cycle requires an ATP/NADPH ratio of approximately 1.5, while photosynthetic electron transport typically provides only 1.28, creating inherent imbalance that cells must compensate for through alternative pathways [61]. Understanding these **inherent mismatches** between cofactor supply and demand is essential for designing effective engineering strategies.

Quantitative Analysis of Cofactor Demands

Table 2: Cofactor Turnover Rates and ATP/NADPH Ratios in Synechocystis Under Different Production Scenarios

Metabolic Objective	Growth Condition	NADPH Turnover (mmol/gDW/h)	ATP Turnover (mmol/gDW/h)	ATP/NADPH Ratio
Biomass Formation	Autotrophic	3.87-5.49	7.24-8.61	2.11
Biomass Formation	Mixotrophic	3.87-5.49	7.24-8.61	19.49
Isobutene Production	Autotrophic	3.87-5.49	7.24-8.61	<2.11
Isoprene Production	Mixotrophic	3.87-5.49	7.24-8.61	<19.49
1-Undecene Production	Autotrophic	3.87-5.49	7.24-8.61	<2.11

Flux-sum analysis reveals that **NADPH preference** is pronounced in cyanobacteria under autotrophic conditions, with NADPH turnover rates (3.87-5.49 mmol/gDW/h) significantly exceeding NADH turnover (0.01-0.49 mmol/gDW/h) [61]. This preference must be considered when engineering heterologous pathways that may have different cofactor specificities. The **interconnected nature** of cofactor systems means that perturbations to one pool inevitably affect others, necessitating systems-level approaches to cofactor engineering.

Engineering Strategies for Redox Balance

Cofactor Regeneration Systems

Enzyme-based regeneration approaches provide separate reactions to replenish active cofactor forms in situ. For example, the water-forming NADH oxidase (NOX) from *Streptococcus pneumoniae* and alternative oxidase (AOX1) from *Histoplasma capsulatum* have been successfully employed in *Saccharomyces cerevisiae* to alleviate redox imbalance [59]. These systems **oxidize excess NADH** without ATP generation, directly addressing NADH/NAD+ imbalance while reducing byproduct formation.

The **phosphite dehydrogenase (PtxD)**-based NADH regeneration system represents an advanced cofactor engineering approach that decouples cofactor supply from central carbon metabolism. In E. coli, this system enabled a 41.3 mol% lactate fraction in P(3HB-co-LA) copolymer biosynthesis while reducing acetate byproduct formation to 1.58 g/L [62]. This **flexible regulation** demonstrates how orthogonal cofactor regeneration systems can enhance production without disrupting bacterial growth.

Pathway Engineering for Cofactor Supply

Redirecting carbon flux through NADPH-generating pathways represents a fundamental strategy for enhancing reducing power. In Yarrowia lipolytica engineered for β-carotene production, pushing flux through the **pentose phosphate pathway** and introducing NADP+-dependent glyceraldehyde-3-phosphate dehydrogenase proved more effective than the phosphoketolase-phosphotransacetylase (PK-PTA) pathway [63]. This approach increased β-carotene production to 809.2 mg/L by enhancing NADPH regeneration capacity.

The **synthetic two-step isopentenol utilization pathway** (IUP) demonstrates how bypassing native regulation can enhance cofactor efficiency. When implemented in Y. lipolytica for β-carotene production, this approach ultimately achieved remarkable titers of 39.5 g/L in fed-batch fermentation [63], highlighting the potential of **non-canonical routes** for cofactor optimization.

Transhydrogenation and Cofactor Interconversion

Microbial transhydrogenases facilitate interconversion between NADH and NADPH pools, providing flexibility in redox balancing. In E. coli, the membrane-bound transhydrogenase PntAB and soluble transhydrogenase UdhA (SthA) play complementary roles in maintaining cofactor homeostasis [64]. Engineering the expression and activity of these enzymes can **modulate NADPH availability**—a critical parameter for anabolic pathways.

Studies have demonstrated that manipulating transhydrogenase expression can significantly impact product profiles. For example, enhancing UdhA activity in E. coli increased product yield and NADPH availability, while PntAB overexpression had contrasting effects [64]. These findings emphasize the need for **context-specific optimization** of transhydrogenase expression based on the redox demands of engineered pathways.

Experimental Protocols and Methodologies

Assessing Intracellular Redox Status

Protocol: Quantifying NADH/NAD+ Ratios in S. cerevisiae

- **Culture Preparation**: Grow engineered yeast strains in appropriate selective medium at 30°C with shaking at 250 rpm until mid-exponential phase (OD600 ≈ 0.8-1.0).
- **Metabolite Extraction**: Rapidly harvest 10 mL culture by vacuum filtration and immediately immerse cell pellet in 5 mL of pre-chilled 80% methanol (-80°C). Agitate vigorously for 10 minutes at -20°C.
- **Sample Processing**: Centrifuge at 15,000 × g for 15 minutes at 4°C. Collect supernatant and evaporate methanol under nitrogen stream. Resuspend dried metabolites in 200 µL assay buffer.
- **Enzymatic Assay**: Use NADH/NAD+ quantification kits following manufacturer protocols. Measure absorbance at 565 nm for formazan product derived from NADH-dependent reduction of tetrazolium salts.
- **Data Normalization**: Normalize cofactor concentrations to total cellular protein content determined by Bradford assay.

This protocol successfully detected a 78.0% reduction in free NADH/NAD+ ratio following NOX implementation in 7-dehydrocholesterol-producing S. cerevisiae, correlating with reduced glycerol and ethanol accumulation [59].

Implementing Cofactor Regeneration Systems

Protocol: Integrating Heterologous NADH Oxidase for Redox Rebalancing

- **Genetic Construct Design**: Codon-optimize nox gene from Streptococcus pneumoniae for target host. Clone under control of constitutive promoter (e.g., TEF1p) in integration vector.
- **Strain Transformation**: Introduce construct into production host using appropriate method (e.g., lithium acetate transformation for yeast, electroporation for bacteria).
- **Screening and Validation**: Select transformants on appropriate antibiotic media. Verify integration by colony PCR and Sanger sequencing.
- **Phenotypic Characterization**: Assess redox rebalancing by measuring glycerol and ethanol production, growth rates, and target compound synthesis in shake flask cultures.
- **Bioreactor Validation**: Scale optimal strains to controlled bioreactors with monitoring of dissolved oxygen, pH, and carbon source utilization.

This methodology enabled a 74.4% increase in 7-dehydrocholesterol production in S. cerevisiae, demonstrating the efficacy of enzymatic cofactor regeneration strategies [59].

Computational Modeling and Design Tools

Stoichiometric Modeling for Redox Balance Analysis

Flux Balance Analysis (FBA) and related constraint-based approaches enable prediction of intracellular flux distributions that maintain cofactor balance under steady-state assumptions. These methods have revealed that the **rate of NAD(P)H regeneration**, rather than ATP regeneration, primarily controls the ATP/NADPH ratio and thereby bioproduction in cyanobacteria [61].

Phenotypic phase plane analysis of alkene-overproducing Synechocystis strains demonstrated the inherent trade-off between growth and product synthesis, with maximal alkene production rates resulting in zero biomass accumulation and vice versa [61]. Such analyses help identify **optimal operating points** that balance growth and production objectives.

Flux-Sum Analysis for Cofactor Turnover Assessment

Flux-sum analysis quantifies metabolite turnover rates, providing insights into cofactor usage patterns. This approach has shown that **ATP pool sizes** remain relatively constant (narrow range) across different production scenarios, while NADPH levels exhibit larger variations [61]. This information guides engineering strategies focused on the most limiting cofactors.

Visualization of Cofactor Engineering Strategies



Cofactor Engineering and Regeneration Pathways

The Scientist's Toolkit: Research Reagent Solutions

Table 3: Essential Research Reagents for Cofactor Engineering Studies

Reagent/Component	Function/Application	Example Source/Sequence
Water-forming NADH Oxidase (NOX)	Oxidizes NADH to NAD+ without reactive oxygen species production	*Streptococcus pneumoniae* (GenBank: AF014458) [59]
Alternative Oxidase (AOX1)	Mitochondrial oxidase that accepts electrons from ubiquinol	*Histoplasma capsulatum* (GenBank: AF133236) [59]
Phosphite Dehydrogenase (PtxD)	NAD+ regeneration using phosphite as electron donor	Codon-optimized for target host [62]
Transhydrogenases (PntAB, UdhA)	Interconvert NADH and NADPH pools	*E. coli* native enzymes [64]
NADP+-dependent GAPDH	Shifts glycolytic flux toward NADPH generation	*Clostridium acetobutylicum* GapN (AAK81579.1) [63]
Phosphoketolase (PK)	Bypasses NADPH-limiting steps in pentose phosphate pathway	*Clostridium acetobutylicum* (NP_347971.1) [63]

Effective management of cofactor regeneration and redox imbalances represents a critical frontier in metabolic engineering for pharmaceutical and chemical production. The strategies outlined in this guide—including enzyme-based cofactor regeneration, pathway rebalancing, and computational modeling—provide researchers with a comprehensive toolkit for optimizing microbial cell factories. As the field advances, the integration of **non-canonical redox cofactors** and dynamic regulation systems will further enhance our ability to control metabolic flux toward desired products while maintaining cellular fitness. The continued refinement of these approaches promises to accelerate the development of sustainable bioprocesses for drug development and chemical synthesis.

Strategies for Reducing By-Product Formation (e.g., Acetate, Ethanol)

In the pursuit of microbial cell factories for efficient bio-production, central carbon metabolism (CCM) serves as the fundamental network generating energy, reducing power, and precursor metabolites for biosynthesis. However, native metabolic fluxes often prioritize cellular growth and homeostasis, leading to the formation of undesirable by-products such as acetate and ethanol, which constrain the yield of target compounds. Framed within broader research on CCM engineering strategies, this technical guide synthesizes current metabolic engineering approaches for optimizing carbon flux. We detail methods to systematically rewire microbial metabolism, redirecting carbon from wasteful by-products toward valuable biochemicals and biofuels, thereby enhancing process economics and sustainability in pharmaceutical and industrial biotechnology.

Core Metabolic Engineering Strategies

Pathway Manipulation and Gene Knockouts

A direct strategy for reducing by-product formation involves the deletion of genes encoding enzymes responsible for diverting carbon toward undesirable compounds. This approach forces carbon flux toward the desired pathways.

- **Competitive Pathway Disruption**: In *E. coli*, a significant amount of carbon is often lost to acetate and lactate under anaerobic conditions. Deleting the genes for **acetate kinase (ackA)** and **lactate dehydrogenase (ldhA)** effectively blocks these secretion routes. In one study, this double knockout in an ethyl acetate-producing strain increased the ethyl acetate titer from 2.7 mM to 9.1 mM, demonstrating a more efficient channeling of carbon [65].
- **Redox Metabolism Engineering**: In *Saccharomyces cerevisiae*, glycerol formation is essential for re-oxidizing surplus NADH under anaerobic conditions. To reduce this carbon loss, **GPD2**, the isoenzyme primarily involved in redox balancing (as opposed to the osmoregulatory Gpd1), is a key deletion target. Knocking out **GPD2** in engineered yeast strains has been shown to reduce glycerol yield by over 90%, significantly improving ethanol and other target product yields [66] [67].

Introducing non-native pathways can create more efficient, direct routes for carbon conversion, bypassing native nodes that lead to by-product formation.

- **The Phosphoketolase (PHK) Pathway**: This heterologous pathway, involving phosphoketolase (PK) and phosphotransacetylase (PTA), provides a shortcut from glycolysis and the pentose phosphate pathway to acetyl-CoA. It minimizes carbon loss and addresses redox imbalances.
  - **Application**: In *S. cerevisiae*, introducing the PHK pathway shifted glycolytic flux, increasing the supply of the precursor erythrose-4-phosphate (E4P) and enabling a yield of 12.5 g/L of *p*-hydroxycinnamic acid. It has also been used to increase the synthesis of acetyl-CoA-derived compounds like fatty acids and 3-hydroxypropionic acid (3-HP) [23].
- **Reductive Acetyl-CoA Pathway**: The introduction of a heterologous **acetylating acetaldehyde dehydrogenase (A-ALD)** enables the NADH-dependent reduction of acetyl-CoA to acetaldehyde, which is then reduced to ethanol. This pathway can functionally replace glycerol formation as a redox sink. When applied in *S. cerevisiae*, this strategy allowed the strain to consume acetate and reduce glycerol formation, thereby increasing the ethanol yield by up to 13% in high-osmolarity cultures [67].

Optimization of Central Carbon Metabolism (CCM) Flux

Global rearrangement of the central metabolic network is often necessary to support high fluxes through engineered pathways.

- **The PRK-RuBisCO Bypass**: This strategy involves introducing the Calvin-cycle enzymes **phosphoribulokinase (PRK)** and **ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)**. It bypasses the oxidative step in glycolysis, reducing NADH formation and eliminating the need for glycerol as a redox sink. An engineered *S. cerevisiae* strain with this pathway showed a 96% lower glycerol yield and a 10% higher ethanol yield on glucose compared to a non-engineered reference strain [66].
- **Modular Deregulation of CCM**: For non-conventional carbon sources like xylose, a systematic approach to deregulate CCM is required. This can involve:
  - **Promoter Engineering**: Using carbon source-responsive promoters (e.g., xylose-responsive promoters) to dynamically control the expression of key catabolic genes.
  - **Enzyme Replacement**: Swapping endogenous, highly regulated enzymes with less-regulated heterologous counterparts.
  - **Biosensor Implementation**: Employing biosensors to monitor and regulate intracellular metabolite levels like NADPH [21].

Table 1: Summary of Key Metabolic Engineering Strategies and Outcomes

Strategy	Host Organism	Target By-Product	Engineering Action	Key Outcome
Competitive Pathway Disruption	*E. coli*	Acetate, Lactate	Deletion of *ackA* and *ldhA* genes	Increased ethyl acetate titer to 9.1 mM [65]
Redox Metabolism Engineering	*S. cerevisiae*	Glycerol	Deletion of *GPD2*	>90% reduction in glycerol yield; higher ethanol yield [66]
Heterologous PHK Pathway	*S. cerevisiae*	---	Expression of phosphoketolase and phosphotransacetylase	12.5 g/L yield of *p*-hydroxycinnamic acid; increased acetyl-CoA supply [23]
A-ALD Pathway	*S. cerevisiae*	Glycerol, Acetate	Expression of acetylating acetaldehyde dehydrogenase (A-ALD)	13% higher ethanol yield in high-osmolarity culture [67]

Strategy	Host Organism	Target By-Product	Engineering Action	Key Outcome
PRK-RuBisCO Bypass	*S. cerevisiae*	Glycerol	Expression of PRK and RuBisCO	96% lower glycerol yield; 10% higher ethanol yield [66]
Multi-Modular CCM Optimization	*S. cerevisiae*	---	Promoter engineering, TF modulation, heterologous enzymes	4.7-fold increase in 3-HP productivity from xylose [21]

Detailed Experimental Protocols

Protocol: Anaerobic Production of Ethyl Acetate in EngineeredE. coli

This protocol details the construction and cultivation of an *E. coli* strain engineered for anaerobic ethyl acetate production, with minimized acetate and lactate by-product formation [65].

- **Strain Construction**:
  - Start with *E. coli* BW25113 (DE3) as the base strain.
  - Use a CRISPR/Cas9 or lambda Red recombinase system to delete the **acetate kinase (*ackA*)** and **lactate dehydrogenase (*ldhA*)** genes to block major by-product pathways.
  - Introduce a plasmid containing the gene for **alcohol acetyltransferase (AAT)**, such as *eat1* from *Kluyveromyces marxianus* or *Wickerhamomyces anomalus*, under the control of an inducible promoter (e.g., LacI/T7 or XylS/Pm).
- **Anaerobic Cultivation**:
  - **Medium**: Use a defined minimal medium (e.g., M9) or a rich medium (e.g., LB) supplemented with an appropriate carbon source, typically **20 g/L glucose**.
  - **Culture Conditions**: Inoculate serum bottles or bioreactors with the engineered strain. Flush the headspace with an inert gas (e.g., N_2 or Argon) to establish and maintain anaerobic conditions.
  - **Induction**: Add the inducer molecule (e.g., IPTG for the LacI/T7 promoter) at the early exponential growth phase to trigger AAT expression. Optimize inducer concentration (e.g., 0.1-1.0 mM IPTG) to balance enzyme activity and cell fitness.
  - **Process Optimization**: For bioreactor studies, implement **continuous gas stripping** (e.g., with N_2) to remove volatile ethyl acetate from the culture broth, mitigating product inhibition and shifting the reaction equilibrium toward ester synthesis [65].
- **Analytical Methods**:
  - **Product Quantification**: Analyze culture supernatants using **Gas Chromatography (GC)** equipped with a flame ionization detector (FID) or a mass spectrometer (GC-MS) to quantify ethyl acetate, ethanol, and residual acetate.
  - **Metabolite Analysis**: Use **High-Performance Liquid Chromatography (HPLC)** to measure concentrations of glucose, organic acids (lactate, formate), and other extracellular metabolites.

Protocol: Replacing Glycerol Production with Acetate Reduction inS. cerevisiae

This protocol describes the engineering of a yeast strain to reduce acetate to ethanol, thereby replacing glycerol formation as the primary mechanism for NADH re-oxidation [67].

- **Strain Engineering**:
  - Use a *S. cerevisiae* CEN.PK lineage strain as the host.
  - Delete the *GPD2* gene to disrupt the primary redox-balancing glycerol pathway.
  - Express a heterologous **acetylating acetaldehyde dehydrogenase (A-ALD)**, such as *eutE* from *E. coli*, from a strong, constitutive yeast promoter.
  - To prevent an ATP-wasting substrate cycle, consider deleting the major cytosolic aldehyde dehydrogenase gene *ALD6*.
- **Anaerobic Bioreactor Cultivation**:
  - **Medium**: Use a synthetic defined medium with **50 g/L glucose** and supplement with **5-10 mmol/L acetate** to serve as the electron acceptor.
  - **Bioreactor Setup**: Cultivate in a stirred-tank bioreactor with controlled **anaerobic conditions**. Maintain constant temperature (30°C) and pH (5.0). Sparge the medium with an N_2/CO_2 mixture (e.g., 95/5 vol/vol) to ensure anaerobiosis and remove CO_2.
  - **Sampling**: Take periodic samples to monitor optical density (cell growth), substrate consumption (glucose, acetate), and product formation (ethanol, glycerol).

- **Analytical Methods**:
  - **Extracellular Metabolites**: Use HPLC to quantify glycerol, ethanol, acetate, and glucose concentrations in the culture supernatant.
  - **Yield Calculations**: Calculate the yields of ethanol and glycerol on glucose (C-mol product / C-mol glucose) to evaluate the efficiency of carbon redistribution.

Visualization of Engineering Strategies

The following diagrams illustrate the logical relationships and metabolic fluxes of the core engineering strategies discussed.

Metabolic Engineering Workflow for By-product Reduction

This diagram outlines the decision-making process for selecting and implementing strategies to reduce by-product formation in microbial cell factories.



Engineered Pathways in Central Carbon Metabolism

This diagram maps key engineering strategies onto the central carbon metabolism of a typical microorganism like *E. coli* or *S. cerevisiae*, showing how they reroute carbon flux.



The Scientist's Toolkit: Key Reagents and Materials

Table 2: Essential Research Reagents for Metabolic Engineering to Reduce By-Products

Reagent / Material	Function / Application	Examples & Notes
CRISPR/Cas9 System	For precise gene knockouts (e.g., *ackA*, *ldhA*, *GPD2*) and integration of heterologous genes.	Enables rapid, multiplexed genome editing in both *E. coli* and *S. cerevisiae*.
Inducible Promoter Systems	To control the timing and level of expression of heterologous enzymes, optimizing metabolic flux and reducing toxicity.	**LacI/T7** and **XylS/Pm** systems are widely used for fine-tuning expression in *E. coli* [65].
Heterologous Enzyme Genes	Key catalytic components for implementing new pathways.	**A-ALD (*eutE*)**: For acetate reduction. **Eat1**: For ethyl acetate synthesis. **PK/PTA**: For the PHK pathway [65] [67] [23].
Anaerobic Bioreactor	Provides controlled conditions (temperature, pH, anaerobiosis) for evaluating engineered strains under process-relevant conditions.	Essential for validating redox-balancing strategies that only function in the absence of oxygen [65] [67].
Gas Chromatography (GC)	Analytical instrument for quantifying volatile products and by-products.	Critical for measuring compounds like **ethyl acetate, ethanol, and acetate** in fermentation broth [65].
HPLC System	For quantifying substrates (e.g., glucose) and non-volatile metabolites (e.g., glycerol, organic acids) in culture supernatants.	Standard method for tracking carbon utilization and by-product formation.

Addressing Thermodynamic and Kinetic Barriers in Synthetic Pathways

The engineering of central carbon metabolism (CCM) in microbial hosts represents a cornerstone of modern metabolic engineering, enabling the sustainable bioproduction of fuels, chemicals, and pharmaceuticals. However, the inherent thermodynamic and kinetic constraints of metabolic pathways often limit carbon flux and final product yields, presenting significant challenges for industrial applications. **Thermodynamic barriers** determine the fundamental feasibility and directionality of biochemical reactions, while **kinetic barriers** control the rates at which these reactions proceed. Overcoming these limitations requires a systematic approach that integrates mechanistic understanding with advanced engineering strategies. Within the broader context of CCM engineering research, addressing these barriers is paramount for redirecting metabolic flux from core pathways—such as glycolysis, the tricarboxylic acid (TCA) cycle, and the pentose phosphate pathway (PPP)—toward target compounds. This

whitepaper provides an in-depth technical guide to the latest methodologies for identifying, analyzing, and overcoming thermodynamic and kinetic bottlenecks in synthetic pathways, with a particular emphasis on applications within central carbon metabolism.

Thermodynamic Barriers in Metabolic Pathways

Fundamental Principles and Analytical Frameworks

The thermodynamic landscape of a metabolic pathway is primarily governed by the **Gibbs free energy change (ΔrG)** of its constituent reactions. This parameter indicates the thermodynamic feasibility of a reaction under physiological conditions. The relationship is defined as $\Delta rG = \Delta rG^\circ + RT \ln Q$, where ΔrG° is the **standard Gibbs free energy change**, R is the universal gas constant, T is temperature, and Q is the reaction quotient [68]. Reactions with substantially negative ΔrG values are considered **thermodynamically favorable** and often act as key flux-control points in pathways like glycolysis, where reactions catalyzed by hexokinase (HK), phosphofructokinase (PFK), and pyruvate kinase (PK) exhibit this property [68].

The **Minimum-Maximum Driving Force (MDF)** computational framework has emerged as a powerful tool for analyzing pathway thermodynamics. MDF identifies metabolic routes with the highest thermodynamic driving forces, enabling researchers to select and design pathways that are inherently more efficient [18]. At the genome scale, reactions with substantial negative ΔrG values have been identified as **Thermodynamic Driver Reactions (TDRs)**, which exhibit distinctive network topological properties and higher variability in associated enzyme abundance, consistent with their potential roles in controlling metabolic fluxes [68].

Table 1: Key Thermodynamic Parameters and Their Metabolic Significance

Parameter	Definition	Metabolic Significance	Analysis Method
ΔrG° (Standard Gibbs Free Energy Change)	Energy change under standard conditions (1 M concentrations, pH 7.0)	Reflects intrinsic thermodynamic stability difference between products and reactants	Group Contribution Method, Component Contribution Method, dGbyG ML Model [68]
ΔrG (Reaction Gibbs Free Energy Change)	Energy change under physiological metabolite concentrations	Determines actual thermodynamic feasibility and directionality of metabolic reactions	13C-MFA, GC-MS, LC-MS with thermodynamic modeling [68]
Thermodynamic Driving Force	Degree to which a reaction is from equilibrium	Impacts flux control; larger driving forces often indicate rate-limiting steps	MDF (Minimum-Maximum Driving Force) analysis [18]
Reaction Affinity	Negative of ΔrG	Quantifies the driving force for reaction flux; higher affinity promotes forward flux	Thermodynamic flux balance analysis [68]

Strategies for Overcoming Thermodynamic Barriers

Several advanced strategies have been developed to overcome thermodynamic limitations in synthetic pathways:

- **Pathway Bifurcation and Compartmentalization**: Non-model organisms like *Schizosaccharomyces japonicus* demonstrate natural optimization of central carbon metabolism through strategies such as running a **fully bifurcated TCA pathway** to sustain amino acid production without respiration. This organism also optimizes cytosolic NADH oxidation via **glycerol-3-phosphate synthesis**, maintaining redox balance under fermentative conditions [29].
- **Intermediate Phase Engineering**: Inspired by principles in materials science, this approach involves stabilizing **metastable intermediate phases** that serve as thermodynamic templates during pathway reactions. These intermediates possess lower energy barriers ($\Delta G_2 < \Delta G_1$) for nucleation, enabling controlled transformation to the final stable metabolic products [69].
- **Energy Ratchet Mechanisms**: Molecular engineering approaches can create systems where energy inputs (e.g., light) periodically modulate asymmetric energy landscapes. This technique, observed in light-driven molecular pumps, rectifies Brownian motion through **coupled kinetic and thermodynamic control**, enabling directional transport against concentration gradients [70].

Kinetic Barriers in Pathway Engineering

Kinetic barriers in synthetic pathways primarily arise from **suboptimal enzyme kinetics**, **allosteric regulation**, and **metabolic imbalances**. In central carbon metabolism, tightly regulated enzymes often resist engineering attempts to alter flux balance, particularly when introducing non-native carbon sources like xylose [21]. Key kinetic limitations include:

- **Inefficient enzyme kinetics** with C1 substrates (CO_2, methane, methanol) [17]

- **Allosteric inhibition** of glycolytic enzymes by downstream metabolites [29]
- **Rate-limiting steps** in canonical pathways such as glycolysis and TCA cycle [68]
- **Cofactor imbalances** resulting from heterologous pathway introduction [2]

Advanced monitoring techniques employ **biosensors to detect intracellular metabolites** like NADPH and fatty acyl-CoA in real-time, enabling identification of kinetic bottlenecks during pathway operation [21].

Engineering Solutions for Kinetic Optimization

Enzyme and Pathway Engineering

- **Heterologous Enzyme Expression**: Introduction of non-native enzymes with superior kinetic properties can bypass native regulatory mechanisms. For example, the **phosphoketolase (PHK) pathway** provides a thermodynamically favorable route to acetyl-CoA by directly converting fructose-6-phosphate (F6P) and xylulose-5-phosphate (X5P) to acetyl-phosphate and subsequently to acetyl-CoA, enhancing flux toward acetyl-CoA-derived products [2]. In *S. cerevisiae*, this strategy increased 3-hydroxypropionic acid (3-HP) production by 41.9% while reducing glycerol byproduct formation by 48.1% [2].

- **Enzyme Fusion and Scaffolding**: Aggregating catalytic components into fusion proteins can enhance substrate channeling, though this approach requires careful optimization. In 3-HP production pathways, aggregating McrN and McrCm components showed variable effectiveness depending on the fusion architecture, indicating that physical proximity alone doesn't guarantee improved kinetics [21].

- **Directed Evolution and Mutagenesis**: Engineering mutant enzymes with improved catalytic efficiency or reduced allosteric inhibition. For instance, expression of a mutant acetyl-CoA synthetase (ACS^SEL641P) enhanced acetyl-CoA synthesis rates in yeast strains [2].

Regulatory and Expression Optimization

- **Promoter Engineering**: Systematic evaluation and implementation of carbon source-responsive promoters enable dynamic control of gene expression. In xylose utilization, **xylose-responsive promoters** (e.g., pADH2, pSFC1) driving xylose isomerase expression resulted in faster growth compared to constitutive promoters, demonstrating the kinetic advantage of regulon-controlled expression [21].

- **Transcription Factor Manipulation**: Modulating the expression of global regulators can reshape metabolic networks. Introducing the *Deinococcus radiodurans* response regulator DR1558 into *E. coli* enhanced expression of CCM genes and improved NADPH generation from PPP to meet cofactor demands during PHB biosynthesis [2].

Table 2: Kinetic Engineering Strategies for Central Carbon Metabolism

Engineering Strategy	Mechanism of Action	Application Example	Performance Outcome
Heterologous Pathway Introduction	Provides thermodynamically favorable bypass routes	PHK pathway in *S. cerevisiae* for acetyl-CoA production	25% increase in farnesene yield; 41.9% increase in 3-HP production [2]
Promoter Engineering	Enables carbon source-responsive expression control	Xylose-responsive promoters for xylose assimilation genes	Faster growth on xylose compared to constitutive promoters [21]
Enzyme Abundance Optimization	Alleviates rate-limiting steps through increased catalyst concentration	Multicopy integration of transaldolase 1 (Tal1) and transketolase 1 (Tkl1)	Increased protopanaxadiol yield to 152.37 mg/L in yeast [2]
Allosteric Regulation Removal	Eliminates feedback inhibition of key enzymes	Modification of amino acid sites on proteins to reduce inhibition	Enhanced flux through central carbon metabolism [21]
Cofactor Balancing	Matches cofactor supply with pathway demand	Expression of NADP+-dependent PDH pathway in yeast	2-fold increase in acetyl-CoA levels [2]

Integrated Experimental Methodologies

Protocol for Thermodynamic Analysis of Metabolic Reactions

Objective: Determine the thermodynamic feasibility of reactions in a synthetic pathway under physiological conditions.

Materials:

- **dGbyG Machine Learning Model**: Graph neural network-based predictor for ΔrG° of metabolic reactions from molecular structures [68]
- **Stable Isotope Tracers** (e.g., 13C-labeled substrates): For metabolic flux analysis [29]
- **Mass Spectrometry Systems**: GC-MS or LC-MS for metabolite quantification [68]

- **eQuilibrator 3.0 Database**: Repository for Gibbs free energy predictions [68]

Procedure:

- **Pathway Identification**: Define all enzymatic reactions in the target synthetic pathway.
- **ΔrG° Prediction**: Calculate standard Gibbs free energy changes using dGbyG or eQuilibrator database [68].
- **Metabolite Concentration Measurement**: Cultivate cells under relevant conditions, perform rapid quenching, extract metabolites, and quantify absolute concentrations using LC-MS/MS with internal standards.
- **ΔrG Calculation**: Compute actual reaction Gibbs free energy using the formula $\Delta rG = \Delta rG^\circ + RT \ln Q$ with measured metabolite concentrations.
- **Thermodynamic Driving Force Assessment**: Identify reactions with ΔrG values close to zero (reversible) or highly negative (irreversible) using MDF analysis [18].
- **Flux Control Coefficient Determination**: Correlate ΔrG values with enzyme abundances and flux control coefficients to identify thermodynamic drivers [68].

Protocol for Kinetic Pathway Optimization

Objective: Enhance kinetic flux through synthetic pathways by addressing rate-limiting steps.

Materials:

- **Carbon Source-Responsive Promoters**: Libraries of promoters with characterized expression profiles on target substrates [21]
- **Biosensor Systems**: Genetic constructs for real-time monitoring of metabolites (e.g., NADPH, acyl-CoAs) [21]
- **CRISPR-Cas9 Tools**: For precise genome editing and regulatory element manipulation [71]
- **Omic Analysis Platforms**: RNA-seq, proteomics, and metabolomics for systems-level analysis [21]

Procedure:

- **Promoter Characterization**: Evaluate promoter strength and regulation under target cultivation conditions using fluorescent reporters (e.g., RFP, GFP) [21].
- **Pathway Module Expression Tuning**: Assemble pathway genes with promoters of appropriate strengths to balance expression levels and minimize metabolic burden.
- **Biosensor Integration**: Implement metabolite biosensors to monitor pathway intermediate levels and identify kinetic bottlenecks in real-time [21].
- **Enzyme Engineering**: Apply directed evolution or structure-based mutagenesis to improve catalytic efficiency, reduce inhibition, or alter cofactor specificity of rate-limiting enzymes.
- **Dynamic Regulation**: Implement feedback-controlled expression systems that automatically adjust enzyme levels based on metabolite concentrations.
- **Fed-Batch Fermentation Validation**: Evaluate kinetic performance in controlled bioreactors with carefully designed feeding strategies to maintain optimal substrate concentrations.

Computational and Modeling Approaches

Thermodynamic and Kinetic Modeling Tools

Advanced computational frameworks are essential for predicting and optimizing pathway performance:

- **Graph Neural Networks (GNNs)**: The dGbyG model applies GNNs to predict ΔrG° values directly from molecular structures, outperforming traditional group contribution methods in both accuracy and coverage of genome-scale metabolic reactions [68].

- **Flux Balance Analysis (FBA)**: Constrains-based modeling approach that predicts steady-state flux distributions optimizing cellular objectives (e.g., biomass formation, product yield) [18].

- **Enzyme Cost Minimization (ECM)**: Estimates optimal enzyme and metabolite concentrations supporting desired flux distributions while minimizing cellular protein investment [18].

- **Two-Step (2S) Nucleation Models**: Describe non-equilibrium processes where metastable intermediate phases with lower energy barriers form first, providing frameworks for understanding intermediate-driven pathway optimization [69].

Artificial Intelligence and Machine Learning Applications

- **Machine Learning-Guided Protein Design**: AI models accelerate the engineering of enzymes with improved kinetics and thermodynamics for C1 assimilation [17].

- **Generative AI for Pathway Design**: Predicts optimal metabolic routes and identifies suitable non-model chassis organisms with native beneficial properties [18].
- **Large Language Models**: Assist in predicting molecular interactions and optimizing crystallization pathways in metabolic engineering contexts [69].

The Scientist's Toolkit: Essential Research Reagents

Table 3: Key Research Reagent Solutions for Thermodynamic and Kinetic Studies

Reagent/Category	Function/Application	Example Specifics
Stable Isotope Tracers	Metabolic flux analysis (MFA)	13C-labeled substrates (e.g., 13C-glucose, 13C-xylose) for quantifying pathway fluxes [29]
Biosensor Systems	Real-time monitoring of metabolites	Genetic constructs for detecting NADPH, fatty acyl-CoA, or other pathway intermediates [21]
Carbon Source-Responsive Promoters	Tunable gene expression control	Xylose-responsive promoters (pADH2, pSFC1) for optimized expression during xylose utilization [21]
Heterologous Enzyme Kits	Pathway optimization and bypass	Phosphoketolase (PK) and phosphotransacetylase (PTA) for PHK pathway implementation [2]
CRISPR-Cas9 Systems	Genome editing and regulation	Tools for gene knockouts, promoter replacements, and transcription factor manipulation [71]
Thermodynamic Databases	ΔG° prediction and validation	eQuilibrator 3.0 database with Component Contribution method for ~5,000 human metabolic reactions [68]
Machine Learning Models	Prediction of thermodynamic parameters	dGbyG GNN-based model for standard Gibbs free energy prediction [68]

Visualizing Experimental Workflows and Pathway Relationships

Diagram 1: Integrated workflow for addressing thermodynamic and kinetic barriers in synthetic pathway engineering. The methodology combines computational predictions with experimental validation through an iterative design-build-test-learn cycle.



Diagram 2: Metabolic pathway engineering with heterologous bypass routes. The PHK pathway creates a thermodynamically favorable shortcut from central carbon metabolism intermediates to acetyl-CoA, enhancing flux toward target products while addressing kinetic and thermodynamic bottlenecks in native routes.

Addressing thermodynamic and kinetic barriers represents a critical frontier in synthetic pathway engineering within central carbon metabolism. The integrated methodologies presented in this technical guide—spanning advanced computational modeling, sophisticated experimental protocols, and

cutting-edge analytical techniques—provide researchers with a comprehensive toolkit for overcoming these fundamental limitations. The continuing convergence of **artificial intelligence** with traditional metabolic engineering, particularly through GNN-based thermodynamic prediction and machine learning-guided enzyme design, promises to further accelerate the development of efficient microbial cell factories. By systematically applying these principles to engineer both native and synthetic pathways, researchers can achieve unprecedented control over metabolic fluxes, enabling the sustainable production of valuable chemicals, pharmaceuticals, and fuels from diverse carbon sources while contributing to a circular carbon economy.

Integrated Multi-Omics Approaches for Systems-Level Optimization

The pursuit of systems-level optimization in biological engineering, particularly within central carbon metabolism (CCM), necessitates a holistic understanding of interconnected molecular layers. Integrated multi-omics approaches provide a powerful framework to achieve this by simultaneously interrogating genomes, transcriptomes, proteomes, and metabolomes. This technical guide delineates the methodologies, computational tools, and experimental protocols for deploying multi-omics to elucidate and engineer the complex regulatory networks of CCM. By framing these techniques within the context of CCM engineering strategies, we provide researchers and drug development professionals with a blueprint for enhancing the production of biofuels, pharmaceuticals, and other valuable compounds through targeted metabolic optimization.

Central carbon metabolism, comprising glycolysis, the tricarboxylic acid (TCA) cycle, and the pentose phosphate pathway (PPP), serves as the fundamental engine for cellular energy production and precursor generation [23] [72]. Its optimization is a central goal in metabolic engineering for industrial and therapeutic applications. However, traditional single-omics studies provide limited insights into the complex, multi-layered regulation of these pathways. Multi-omics integration addresses this by combining diverse datasets to reconstruct intricate biological pathways and networks, enabling a systems-level perspective [73].

The need for this integration is particularly evident in precision medicine and advanced bioproduction. For instance, in microbial engineering, the introduction of heterologous pathways like the phosphoketolase (PHK) pathway can rewire CCM, increasing precursor supply and correcting redox imbalances to boost yields of compounds like fatty acids and farnesene [23]. Such interventions create ripple effects across the entire metabolic network, effects that can only be fully comprehended and optimized through multi-omics analysis. This guide details the frameworks and methods for such an integrated analysis, providing a pathway to deconvolute CCM complexity for superior engineering outcomes.

Computational Frameworks for Data Integration

The integration of heterogeneous, high-dimensional omics data presents significant computational challenges, including variations in data scales, nomenclature, and the inherent noise of each technology [73]. Several powerful frameworks have been developed to meet these challenges, enabling researchers to extract robust, biologically meaningful signals.

Reference Materials and Ratio-Based Profiling

A foundational step for reliable integration is the use of multi-omics reference materials to control for technical variability. The Quartet Project provides suites of publicly available reference materials (DNA, RNA, protein, metabolites) derived from immortalized cell lines of a family quartet [74]. These materials provide a built-in ground truth defined by genetic relationships and the central dogma of molecular biology.

A paradigm shift from absolute to **ratio-based profiling** is recommended to mitigate irreproducibility. This approach involves scaling the absolute feature values of study samples relative to those of a concurrently measured common reference sample (e.g., one of the Quartet samples) on a feature-by-feature basis [74]. This method produces highly reproducible and comparable data across batches, laboratories, and technology platforms, forming a robust foundation for all subsequent integration and analysis.

Tools and Workflows for Multi-Omics Analysis

A range of computational tools facilitates the exploration and integration of multi-omics data, many designed for accessibility to non-bioinformaticians.

- **MiBiOmics:** This web-based application allows for the visualization, exploration, and integration of up to three omics datasets. It implements ordination techniques like Principal Component Analysis (PCA) and network-based approaches, including Weighted Gene Correlation Network Analysis (WGCNA). A key feature is its **multi-WGCNA** protocol, which identifies associations between modules (groups of highly correlated features) from different omics layers and links them to external phenotypic traits, effectively reducing dimensionality to pinpoint robust, multi-omics biomarkers [75].
- **iNetModels:** This platform is an interactive database and visualization tool for pre-computed Multi-Omics Biological Networks (MOBNs) and tissue-specific Gene Co-expression Networks (GCNs). It allows users to query features of interest and visualize their associations across other omics layers, such as clinical chemistry, proteomics, metabolomics, and microbiome data from the same individuals [76].
- **Metabolic-Informed Neural Networks (MINN):** For a more mechanistic integration, MINN represents a hybrid approach that embeds Genome-Scale Metabolic Models (GEMs) within a neural network architecture. This framework utilizes multi-omics data to predict metabolic

fluxes in *E. coli*, combining the pattern recognition strength of machine learning with the structured biochemical knowledge of GEMs, thereby enhancing prediction accuracy and interpretability [77].

Table 1: Key Computational Tools for Multi-Omics Integration

Tool	Type	Key Function	Applicability to CCM
Quartet Reference Materials [74]	Reference Suite	Provides ground truth for data QC and integration	Enables precise measurement of CCM enzyme expression and metabolite levels
MiBiOmics [75]	Web Application / Workflow	Multi-omics exploration via ordination and network analysis (multi-WGCNA)	Identifies CCM-related gene/protein modules and links them to metabolic output traits
iNetModels [76]	Database & Visualization Platform	Interactive exploration of pre-computed multi-omics association networks	Allows query of CCM enzymes/metabolites to find cross-omics regulators
MINN [77]	Hybrid ML-Mechanistic Model	Integrates omics data with GEMs to predict metabolic flux	Predicts flux through glycolytic, TCA, or PPP pathways from omics input

Experimental Protocols for Multi-Omics in CCM Engineering

Applying multi-omics to CCM optimization requires a structured experimental workflow, from perturbation to data integration. The following protocols outline a standardized pipeline.

Protocol 1: Systems-Level Analysis of an Engineered CCM Pathway

This protocol is designed to assess the global impact of a genetic perturbation in the CCM, such as the introduction of a heterologous pathway.

- **Strain Engineering and Cultivation:**
  - Introduce the genetic modification of interest (e.g., heterologous PHK pathway from *Aspergillus nidulans* [23]) into the host chassis (e.g., *S. cerevisiae* or *E. coli*).
  - Cultivate the engineered strain and an unmodified control in biological triplicates under defined conditions (e.g., minimal glucose medium).
  - Harvest cells at mid-exponential growth phase by rapid centrifugation. Snap-freeze cell pellets in liquid nitrogen and store at -80°C.
- **Multi-Omics Sample Preparation:**
  - **Genomics:** Extract genomic DNA using a commercial kit. Verify the successful integration of the heterologous pathway via whole-genome sequencing or PCR.
  - **Transcriptomics:** Extract total RNA using a hot phenol protocol or commercial kit. Assess RNA integrity (RIN > 8.0). Prepare libraries for RNA sequencing (RNA-seq) using a standardized kit (e.g., Illumina).
  - **Proteomics:** Lyse cells and perform protein extraction. Digest proteins with trypsin. Analyze peptides using liquid chromatography-tandem mass spectrometry (LC-MS/MS) on a high-resolution instrument.
  - **Metabolomics:** Quench metabolism rapidly (e.g., cold methanol). Extract intracellular metabolites. Analyze polar metabolites via GC-MS or LC-MS and lipids via LC-MS.
- **Data Generation and Horizontal Integration:**
  - Process each omics dataset with established pipelines (e.g., RNA-seq alignment and counts, proteomics搜谱 and quantification, metabolomics peak picking and annotation).
  - Perform **ratio-based profiling** using a common reference sample (e.g., the unmodified control strain or a Quartet reference) to generate comparable log-ratio values for all features [74].
  - Conduct within-omics differential analysis (e.g., DESeq2 for RNA-seq, limma for proteomics) to identify significantly altered features.
- **Vertical Integration and Systems Analysis:**
  - Use a tool like **MiBiOmics** to perform an integrated analysis [75].
  - Upload the normalized and filtered datasets (genomics variant as a trait, transcriptomics, proteomics, metabolomics) along with an annotation table including the "engineering status" (e.g., PHK+ vs. control) and key performance metrics (e.g., product titer, growth rate).
  - Perform a **multi-WGCNA** analysis to identify modules of highly correlated genes, proteins, and metabolites that are significantly associated with the engineered CCM state.
  - Extract features from the most significant multi-omics module for pathway enrichment analysis (e.g., KEGG, GO) to identify affected biological processes beyond the immediate CCM.



Multi-Omics Experimental Workflow

Protocol 2: Multi-Omics for Dynamic Flux Analysis

This protocol focuses on capturing time-resolved changes in CCM flux in response to a nutrient shift or other dynamic perturbation.

- **Dynamic Perturbation and Sampling:**
  - Grow cells in a controlled bioreactor under steady-state conditions.
  - Introduce a defined perturbation (e.g., carbon source shift from glucose to xylose, pulse of a stressor).
  - Collect samples at multiple time points (e.g., 0, 5, 15, 30, 60, 120 minutes) post-perturbation for multi-omics analysis as described in Protocol 1.
- **Data Integration with Metabolic Models:**
  - Generate time-series data for transcriptome, proteome, and metabolome.
  - Integrate the data into a **Metabolic-Informed Neural Network (MINN)** [77].
  - The MINN framework uses the omics data as input and the embedded Genome-Scale Metabolic Model (GEM) to constrain the output, predicting dynamic changes in metabolic flux for reactions in glycolysis, TCA, and PPP.
  - Validate key flux predictions experimentally (e.g., using 13C metabolic flux analysis).

The Scientist's Toolkit: Essential Reagents and Materials

Successful execution of multi-omics studies relies on a suite of reliable reagents, materials, and computational resources.

Table 2: Research Reagent Solutions for Multi-Omics Studies

Item	Function / Description	Example Use in CCM Context
Quartet Reference Materials [74]	Certified reference materials (DNA, RNA, protein, metabolites) for inter-laboratory calibration and QC.	Normalizing sample measurements for CCM-related enzymes (e.g., phosphofructokinase) and metabolites (e.g., acetyl-CoA).

Item	Function / Description	Example Use in CCM Context
RNA-seq Library Prep Kit	For converting extracted RNA into sequencing-ready libraries.	Profiling transcriptional changes in genes encoding CCM enzymes (e.g., *TAL1*, *TKL1* in PPP [23]) in engineered strains.
LC-MS/MS Grade Solvents	High-purity solvents for proteomic and metabolomic sample preparation and separation.	Ensuring reproducible identification and quantification of CCM metabolites and proteins.
Trypsin, Proteomics Grade	Enzyme for digesting proteins into peptides for LC-MS/MS analysis.	Digesting cellular proteome to monitor abundance of CCM enzymes like pyruvate dehydrogenase [23].
Genome-Scale Metabolic Model (GEM)	A computational model of an organism's metabolism.	Providing a mechanistic framework for integrating omics data and predicting flux (e.g., using MINN [77]).
WGCNA R Package [75]	A widely used R package for weighted correlation network analysis.	Constructing co-expression networks to find modules of genes/proteins associated with high-flux CCM states.

Visualization and Interpretation of Integrated Data

The final step involves translating integrated data into actionable biological insights, particularly regarding the structure and regulation of the CCM network.

Constructing an Integrated CCM Network

The results from vertical integration, such as the multi-WGCNA in MiBiOmics, can be used to build a refined model of CCM.

- **Identify Key Multi-Omics Modules:** Pinpoint modules that are significantly associated with both the engineering trait (e.g., PHK expression) and key performance indicators (e.g., farnesene yield) [23].
- **Map Modules to CCM Pathways:** Overlay the features (genes, proteins, metabolites) from these key modules onto the canonical CCM map. This highlights which specific pathway steps are most strongly co-regulated with the desired outcome.
- **Infer Regulatory Links:** Identify transcription factors or signaling proteins within the modules that may act as novel regulators of CCM flux.
- **Pinpoint New Targets:** Metabolites within the module that show strong cross-omics correlations can indicate potential bottlenecks or off-target effects, revealing new candidates for engineering (e.g., downregulating a competing pathway).



Multi-Omics View of an Engineered CCM

This diagram illustrates how multi-omics data layers provide a systems view of CCM after introducing a heterologous PHK pathway. Genomics confirms the presence of the new gene, transcriptomics and proteomics show the cellular response in related pathways (e.g., upregulation of PPP enzymes TAL1/TKL1), and metabolomics captures the resulting changes in metabolite pools (e.g., NADPH, Citrate), revealing the interconnected network effects of the engineering strategy [23].

Measuring Success: Analytical Frameworks and Host Performance Benchmarks

Metabolic Flux Analysis (MFA) and Flux Balance Analysis (FBA)

Metabolic Flux Analysis (MFA) and Flux Balance Analysis (FBA) represent cornerstone methodologies in systems biology for quantifying and predicting metabolic phenotypes. As computational and experimental frameworks, they enable researchers to decipher the complex flow of metabolites through biochemical networks, providing critical insights into cellular physiology. Within the strategic context of central carbon metabolism (CCM) engineering, these flux analysis techniques have become indispensable for rational strain design, bioprocess optimization, and therapeutic target identification. This technical guide comprehensively examines the fundamental principles, methodological workflows, and practical applications of MFA and FBA, with particular emphasis on their transformative role in redirecting carbon flux for enhanced production of valuable biochemicals and pharmaceuticals.

Metabolic flux analysis refers to a suite of computational and experimental techniques designed to quantify the rates of metabolic reactions through biochemical pathways in living cells. As the functional endpoint of cellular regulation, metabolic fluxes provide a direct link between genetic makeup and phenotypic expression, offering unique insights into metabolic network operation that cannot be obtained from transcriptomic or proteomic data alone [78]. The term "metabolic flux" was first used in the 1940s, but it was not until the 21st century that researchers began using the term "fluxomics" to describe the comprehensive analysis of metabolic fluxes [78]. Flux analysis methodologies have since become fundamental tools for understanding how genetic modifications and environmental perturbations reshape cellular metabolism.

Two primary approaches dominate the field: **Flux Balance Analysis (FBA)**, a constraint-based modeling approach that predicts flux distributions using optimization principles, and **13C-Metabolic Flux Analysis (13C-MFA)**, an experimental approach that employs isotopic tracers to quantify intracellular fluxes [78] [79]. While FBA generates hypotheses about metabolic capabilities through in silico simulation, 13C-MFA provides experimental validation and precise quantification of in vivo metabolic fluxes. When applied to central carbon metabolism engineering—which encompasses glycolysis, the pentose phosphate pathway, and the tricarboxylic acid cycle—these complementary techniques enable systematic redesign of microbial chassis for optimized biochemical production [23].

Fundamental Principles

Flux Balance Analysis (FBA)

Flux Balance Analysis operates on the fundamental principle of mass conservation within metabolic networks. The core mathematical framework represents metabolism through a stoichiometric matrix **S** of dimensions $m \times n$, where m represents metabolites and n represents biochemical reactions [79]. Each element Sij corresponds to the stoichiometric coefficient of metabolite i in reaction j. Under the assumption of steady-state metabolite concentrations, the system is described by the mass balance equation:

$$\mathbf{Sv} = 0$$

where **v** is the vector of metabolic fluxes [79]. This equation constrains the solution space such that the production and consumption of each metabolite must balance.

FBA further refines the solution space by incorporating capacity constraints through upper and lower bounds on reaction fluxes:

$$\alpha i \leq vi \leq \beta i$$

The final element involves defining a biological objective function **Z = cTv**, which represents a linear combination of fluxes that the cell is presumed to optimize [79]. For microbial systems, this is typically biomass production, simulating cellular growth. The complete optimization problem becomes:

$$\text{Maximize } Z = cTv, \text{ subject to } Sv = 0 \text{ and } \alpha i \leq vi \leq \beta i$$

This linear programming problem can be solved efficiently even for genome-scale metabolic models, enabling rapid prediction of phenotypic behavior under various genetic and environmental conditions [79].

Metabolic Flux Analysis (MFA)

In contrast to FBA, 13C-MFA employs experimental data from isotope labeling experiments to determine intracellular fluxes. The fundamental principle involves tracking the fate of 13C-labeled atoms from specific substrates as they propagate through metabolic networks [78] [80]. When cells

are cultivated with 13C-labeled substrates (e.g., [1,2-13C]glucose or [U-13C]glucose), the labeled carbon atoms incorporate into metabolic intermediates in patterns that reflect the activities of different pathways [78].

The key assumption in classical 13C-MFA is that the system is at both metabolic and isotopic steady state, meaning metabolite concentrations and isotope distributions remain constant over time [78] [80]. Under these conditions, the relationship between net fluxes and isotopic labeling patterns can be described by algebraic equations. Mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy are used to measure the isotopic labeling distributions of intracellular metabolites, providing the experimental data used to compute the most statistically likely flux map [78].

Table 1: Comparison of Major Flux Analysis Techniques

Method	Tracer Required	Metabolic Steady State	Isotopic Steady State	Key Applications
FBA	No	Yes	Not applicable	Genome-scale prediction, growth simulation, gene knockout studies [79]
13C-MFA	Yes	Yes	Yes	Precise quantification of central carbon metabolism fluxes [78]
13C-INST-MFA	Yes	Yes	No	Systems with slow isotopic labeling, autotrophic organisms [78] [80]
DMFA	Optional	No	Not necessarily	Fermentation processes, dynamic metabolic shifts [78]

Methodological Workflows

Workflow for 13C-Metabolic Flux Analysis

The experimental workflow for 13C-MFA involves multiple critical stages that must be carefully optimized to ensure accurate flux determination:

1. Experimental Design and Tracer Selection: The process begins with selection of appropriate 13C-labeled substrates based on the metabolic pathways of interest. Common choices include [1,2-13C]glucose, [1,6-13C]glucose, or uniformly labeled [U-13C]glucose for central carbon metabolism studies [78]. The labeling pattern is chosen to maximize discrimination between alternative metabolic routes.

2. Cell Cultivation and Labeling: Cells are first pre-cultured in unlabeled medium until metabolic steady state is achieved. The medium is then replaced with labeled substrate solution, and cultivation continues until isotopic steady state is reached—typically requiring several hours for microbial systems but potentially days for mammalian cells [78].

3. Metabolite Quenching and Extraction: Metabolism is rapidly quenched using cold methanol to preserve in vivo flux states. Intracellular metabolites are then extracted using methanol/water or chloroform/methanol solvent systems [80].

4. Analytical Measurement: Isotopic labeling patterns are quantified using either Mass Spectrometry (MS; ~62.6% of studies) or Nuclear Magnetic Resonance (NMR; ~35.6% of studies) [78]. MS offers higher sensitivity, while NMR provides positional labeling information without requiring fragmentation.

5. Computational Flux Estimation: The measured labeling data is integrated with a stoichiometric model to compute the most probable flux distribution. This typically involves minimizing the difference between experimentally observed and computationally simulated labeling patterns using specialized software platforms [78].



Workflow for Flux Balance Analysis

The computational workflow for FBA involves constructing and interrogating genome-scale metabolic models:

1. Network Reconstruction: The process begins with compiling a comprehensive list of metabolic reactions, their stoichiometries, and gene-protein-reaction associations from biochemical databases and literature [79].

2. Stoichiometric Matrix Formation: The metabolic network is represented mathematically as a stoichiometric matrix **S**, where rows correspond to metabolites and columns to reactions [79].

3. Constraint Definition: Physiologically relevant constraints are applied, including substrate uptake rates, thermodynamic constraints (reversibility/irreversibility), and enzyme capacity constraints [79].

4. Objective Function Selection: An appropriate biological objective is defined, typically biomass formation for predicting growth phenotypes or product synthesis for biotechnological applications [79].

5. Linear Programming Solution: The constrained optimization problem is solved using linear programming algorithms to identify the flux distribution that maximizes the objective function [79].

6. Model Validation and Refinement: Predictions are compared with experimental data, and the model is refined iteratively to improve its predictive capability [79].

Table 2: Essential Research Reagents and Tools for Flux Analysis

Category	Specific Examples	Function/Purpose
Isotopic Tracers	[1,2-13C]glucose, [U-13C]glucose, 13C-NaHCO3	Carbon source for 13C-MFA; enables tracking of metabolic fate [78]
Analytical Instruments	LC-MS, GC-MS, NMR spectrometers	Quantification of metabolite concentrations and isotopic labeling patterns [78]
Computational Tools	COBRA Toolbox, 13CFLUX2, INCA, OpenFLUX	Software platforms for flux calculation and simulation [78] [79]
Biological Materials	Microbial chassis (E. coli, S. cerevisiae), Cell culture lines	Model organisms/systems for metabolic engineering studies [23]

Advanced Methodological Developments

Extensions to Core Methodologies

The fundamental frameworks of FBA and MFA have spawned numerous advanced methodologies that address specific limitations and expand application scope:

Isotopically Non-Stationary MFA (INST-MFA) addresses the limitation of traditional 13C-MFA by analyzing transient isotopic labeling patterns before the system reaches isotopic steady state [78] [80]. This approach is particularly valuable for systems with slow isotopic labeling dynamics and enables more rapid flux determination. INST-MFA requires solving ordinary differential equations rather than algebraic balance equations, substantially increasing computational complexity [78].

Dynamic FBA (dFBA) extends FBA to non-steady-state conditions by incorporating dynamic changes in extracellular substrate concentrations and integrating them with the metabolic model [81]. This enables simulation of batch and fed-batch fermentation processes where metabolic fluxes change over time.

Thermodynamics-based MFA (TMFA) incorporates thermodynamic constraints, including Gibbs free energy changes of reactions, to eliminate thermodynamically infeasible flux solutions [80]. This approach provides more physiologically realistic predictions and helps identify thermodynamic bottleneck reactions.

Two-Stage FBA for Drug Target Identification employs sequential linear programming models to first identify pathological metabolic states and then determine medication states with minimal side effects [82]. This approach has shown promise for identifying potential drug targets in metabolic disorders.

Recent Computational Innovations

Recent methodological advances focus on integrating multiple data types and improving prediction accuracy:

TIObjFind represents a novel framework that integrates Metabolic Pathway Analysis (MPA) with FBA to identify context-specific objective functions [81]. By calculating "Coefficients of Importance" that quantify each reaction's contribution to cellular objectives under different conditions, this approach better captures metabolic adaptations to environmental changes.

NEXT-FBA implements a hybrid stoichiometric/data-driven approach that incorporates high-throughput omics data to improve intracellular flux predictions [83]. This methodology helps bridge the gap between metabolic potential predicted by FBA and actual metabolic states realized in biological systems.

ObjFind Framework identifies optimal objective functions for FBA by maximizing a weighted sum of fluxes while minimizing deviations from experimental flux data [81]. This approach helps address the challenge of selecting appropriate biological objectives for different physiological states.



Applications in Central Carbon Metabolism Engineering

Metabolic Engineering Strategies

Central carbon metabolism serves as the fundamental engine of cellular metabolism, generating energy, reducing equivalents, and precursor metabolites for biosynthesis. Engineering CCM has consequently become a cornerstone strategy for optimizing microbial cell factories:

Heterologous Pathway Introduction represents a powerful approach for rewiring central metabolism. The **phosphoketolase (PHK) pathway** has been successfully introduced into *S. cerevisiae* to create a shortcut from fructose-6-phosphate and xylulose-5-phosphate to acetyl-CoA, bypassing several steps of conventional glycolysis [23]. This pathway modification increases acetyl-CoA precursor supply and redirects carbon flux toward acetyl-CoA-derived products, resulting in 25% increase in farnesene production and significant improvements in free fatty acid synthesis [23].

Pyruvate Dehydrogenase (PDH) Bypass engineering addresses the compartmentalization limitations in eukaryotic systems. Expression of an NADP+-dependent pyruvate dehydrogenase complex from *E. coli* in yeast cytosol created a direct route from pyruvate to acetyl-CoA without ATP consumption, doubling intracellular acetyl-CoA levels [23].

Modular Deregulation Strategies enable coordinated optimization of complex metabolic networks. Recent work in *S. cerevisiae* demonstrated that dividing central carbon metabolism into discrete modules (xylose uptake, central metabolic pathways, and product synthesis modules) with systematic deregulation using promoter engineering, transcription factor manipulation, and heterologous enzyme expression resulted in a 4.7-fold increase in 3-hydroxypropionic acid productivity from xylose [21].

Pharmaceutical and Biomedical Applications

Flux analysis methodologies have proven invaluable in pharmaceutical research and development:

Drug Target Identification through two-stage FBA has successfully identified enzyme targets for metabolic disorders such as hyperuricemia [82]. By comparing flux distributions in pathological and healthy states while minimizing side effects, this approach pinpoints strategic intervention points for therapeutic development.

Metabolic Bottleneck Identification in pathogenic organisms enables development of selective antimicrobials. FBA-based essentiality analysis of *Mycobacterium tuberculosis* and *Plasmodium falciparum* metabolic networks has identified enzymes crucial for pathogen survival but non-essential in human hosts, revealing promising drug targets [82].

Toxicology Assessment using 13C-MFA provides insights into drug-induced metabolic perturbations. By quantifying changes in metabolic flux patterns in response to drug candidates, researchers can predict mechanism-based toxicity early in drug development [78].

Table 3: Representative Applications of Flux Analysis in Metabolic Engineering

Application Domain	Specific Example	Engineering Strategy	Outcome
Biofuel Production	Ethanol from xylose in yeast	FBA-guided strain optimization	Enhanced xylose utilization and ethanol yield [80]
Natural Product Synthesis	Aromatic amino acid derivatives	PHK pathway introduction	12.5 g/L p-hydroxycinnamic acid, 154.9 mg/g glucose yield [23]
Therapeutic Compound Production	Ginsenoside precursor PPD	PHK pathway + Tal1/Tkl1 overexpression	152.37 mg/L protopanaxadiol [23]
Bulk Chemical Production	3-Hydroxypropionic acid	Modular CCM deregulation	4.7-fold productivity increase on xylose [21]

Experimental Protocols

Protocol for 13C-Flux Analysis in Microbial Systems

Materials and Reagents

- 13C-labeled substrate (e.g., [U-13C]glucose)
- Defined mineral medium
- Quenching solution (60% aqueous methanol, -40°C)
- Extraction solvent (methanol:chloroform:water, 5:2:2)
- Internal standards for MS analysis

Procedure

- **Pre-culture Preparation**: Inoculate cells from frozen stock into unlabeled medium and grow until mid-exponential phase.
- **Labeling Experiment**: Harvest cells and resuspend in fresh medium containing 13C-labeled substrate at the same concentration as pre-culture. Use adequate culture volume for subsequent metabolite analysis (typically 50-100 mL).
- **Metabolic Quenching**: At isotopic steady state (typically 2-3 generations for microbes), rapidly transfer culture to cold quenching solution (-40°C) at 1:1 ratio. Immediately centrifuge at -20°C.
- **Metabolite Extraction**: Resuspend cell pellet in cold extraction solvent. Agitate vigorously for 1 hour at 4°C. Centrifuge to remove cell debris and transfer supernatant to fresh tubes.
- **Sample Analysis**: Derivatize metabolites for GC-MS analysis or prepare samples for LC-MS/MS. For NMR-based flux analysis, concentrate extracts and reconstitute in appropriate deuterated solvent.
- **Data Processing**: Correct raw mass isotopomer distributions for natural isotope abundances. Integrate with stoichiometric model for flux calculation using specialized software (e.g., 13CFLUX2, INCA).

Critical Considerations

- Ensure metabolic steady state by maintaining constant growth conditions
- Verify isotopic steady state through time-course sampling

- Include appropriate controls and replicates for statistical analysis
- Optimize quenching protocol for specific microbial strain

Protocol for FBA-Based Metabolic Engineering

Computational Resources

- COBRA Toolbox for MATLAB
- Genome-scale metabolic model (e.g., iML1515 for E. coli, iTO977 for S. cerevisiae)
- Linear programming solver (e.g., Gurobi, CPLEX)

Procedure

- **Model Reconstruction**: Curate genome-scale metabolic model from databases (KEGG, BioCyc) and literature evidence.
- **Constraint Definition**: Set substrate uptake constraints based on experimental measurements. Define product secretion and biomass composition constraints.
- **Objective Selection**: Implement appropriate objective function (e.g., biomass production, ATP maximization, or product synthesis).
- **Flux Prediction**: Solve linear programming problem to obtain optimal flux distribution.
- **Gene Deletion Analysis**: Identify essential genes and synthetic lethal pairs through systematic in silico knockouts.
- **Robustness Analysis**: Determine sensitivity of objective function to changes in key reaction fluxes.
- **Experimental Validation**: Compare predictions with experimental growth phenotypes or flux measurements.

Implementation Notes

- Use parsimonious FBA (pFBA) to eliminate thermodynamically infeasible cycles
- Perform flux variability analysis to identify alternate optimal solutions
- Integrate transcriptomic data to create context-specific models
- Employ OptKnock algorithm for metabolic engineering strain design

Metabolic Flux Analysis and Flux Balance Analysis have evolved from specialized methodologies to essential tools in the metabolic engineering toolkit. The integration of these complementary approaches—FBA providing predictive capability and design guidance, and MFA delivering experimental validation and quantitative insight—has dramatically accelerated progress in central carbon metabolism engineering. As computational power increases and analytical techniques advance, flux analysis methodologies will continue to expand their impact, enabling more sophisticated redesign of microbial factories for sustainable chemical production and providing increasingly powerful approaches for therapeutic target identification. The ongoing development of hybrid approaches that combine stoichiometric modeling with machine learning and multi-omics data integration promises to further enhance our ability to understand and engineer metabolic systems for biomedical and industrial applications.

Techno-Economic Analysis (TEA) and Life Cycle Assessment (LCA)

Techno-Economic Analysis (TEA) and Life Cycle Assessment (LCA) have emerged as critical methodological frameworks for evaluating the economic viability and environmental impacts of emerging biotechnologies, particularly within metabolic engineering and carbon dioxide removal (CDR) research. These analytical tools provide systematic approaches for assessing novel technologies during development phases, enabling researchers to identify potential bottlenecks, guide optimization efforts, and make informed decisions about scale-up potential. Within central carbon metabolism engineering, TEA and LCA serve as essential bridges between laboratory-scale innovation and industrial implementation, offering quantitative metrics for comparing alternative metabolic pathways, host organisms, and process configurations.

The integration of TEA and LCA at early research stages is especially crucial for technologies targeting climate change mitigation, such as carbon capture and utilization (CCU) and one-carbon (C1) biomanufacturing platforms [84]. These assessments provide comprehensive evaluations that extend beyond technical performance to encompass economic feasibility and environmental sustainability, ensuring that research investments align with long-term climate goals and circular economy principles. As the field advances, harmonized application of TEA and LCA methodologies enables direct comparison across diverse technological approaches, from engineering microbial chassis for C1 assimilation to deploying large-scale ocean-based carbon dioxide removal strategies [85] [18].

Fundamental Principles and Methodologies

Techno-Economic Analysis (TEA) Framework

Techno-Economic Analysis provides a systematic methodology for evaluating the economic feasibility of technological processes by integrating process design, engineering performance, and financial modeling. For metabolic engineering applications, TEA typically follows a structured approach beginning with process synthesis and simulation, followed by capital and operating cost estimation, and culminating in financial metrics calculation [85].

The TEA framework applied to carbon metabolism engineering projects typically encompasses several critical components. First, **process modeling** involves creating detailed mass and energy balances for the proposed biomanufacturing pathway, accounting for all inputs (feedstocks, nutrients, energy) and outputs (target products, byproducts, waste streams). Second, **capital expenditure (CAPEX)** estimation covers costs for bioreactors, downstream processing equipment, utilities infrastructure, and buildings. Third, **operating expenditure (OPEX)** includes raw materials, labor, maintenance, utilities, and waste disposal. Finally, **financial analysis** integrates these costs to calculate key performance indicators such as minimum selling price (MSP), return on investment (ROI), and payback period [86].

For novel carbon dioxide removal approaches like ocean iron fertilization (OIF), specialized TEA frameworks have been developed that incorporate technology readiness levels (TRL) and learning curves. These frameworks distinguish between first-of-a-kind (FOAK) costs for initial deployments and nth-of-a-kind (NOAK) costs for mature implementations, accounting for potential cost reductions through technological learning and scale optimization [85].

Life Cycle Assessment (LCA) Framework

Life Cycle Assessment provides a comprehensive methodology for evaluating the environmental impacts of products, processes, or services throughout their complete life cycle, from raw material extraction to end-of-life disposal or recycling. The standardized LCA framework, as defined by ISO 14040 and 14044 standards, comprises four interdependent phases [84] [87].

The **goal and scope definition** phase establishes the study's objectives, system boundaries, functional unit, and intended application. For central carbon metabolism engineering, this typically involves defining whether the assessment will focus solely on climate impacts (carbon footprint) or include multiple environmental impact categories such as eutrophication, acidification, and resource depletion. The **life cycle inventory (LCI)** phase involves compiling quantitative data on all material and energy inputs and environmental releases associated with the defined system. The **life cycle impact assessment (LCIA)** phase translates inventory data into potential environmental impacts using characterization factors that relate emissions to their environmental effects. Finally, the **interpretation** phase evaluates results, checks consistency with goal and scope, and provides conclusions and recommendations [84] [87].

Table 1: Key LCA Impact Categories Relevant to Carbon Metabolism Engineering

Impact Category	Description	Common Characterization Factors
Global Warming Potential (GWP)	Contribution to climate change through greenhouse gas emissions	CO_2 equivalents (CO_2eq) over 100-year timeframe
Acidification Potential	Potential to release protons (H^+) to terrestrial and aquatic environments	SO_2 equivalents
Eutrophication Potential	Nutrient over-enrichment of ecosystems	PO_4^{3-} equivalents
Abiotic Resource Depletion	Consumption of non-renewable resources	Antimony equivalents
Land Use	Impacts associated with occupation and transformation of land	Various metrics including soil organic carbon changes

Harmonization of TEA and LCA

The integration of TEA and LCA methodologies creates a powerful decision-support tool for evaluating sustainable technologies, enabling simultaneous assessment of economic viability and environmental performance. Harmonization efforts address challenges in system boundary alignment, data consistency, and interpretation of combined results [84].

Key considerations for TEA-LCA integration include **temporal alignment** (ensuring both assessments reference the same technology maturity stage), **system boundary consistency** (including equivalent processes and life cycle stages), and **functional unit harmonization** (using consistent bases for comparison). For carbon metabolism engineering, this often involves expressing both economic and environmental metrics per unit of target product (e.g., $/kg product and kg CO_2eq/kg product) [84].

Emerging frameworks also incorporate **technology learning curves (TLCs)** into prospective TEA and LCA to forecast how both economic and environmental performance might improve as technologies mature and benefit from cumulative experience and scale [84]. This approach is particularly valuable for early-stage metabolic engineering projects where significant optimization is anticipated between laboratory demonstration and commercial implementation.

Application in Central Carbon Metabolism Engineering

Metabolic Engineering of C1 Assimilation Pathways

The engineering of central carbon metabolism for one-carbon (C1) compound assimilation represents a rapidly advancing frontier in biotechnology, with significant implications for sustainable manufacturing and carbon circularity. TEA and LCA play crucial roles in guiding pathway selection, host engineering, and process development for these platforms [18].

Non-model microorganisms offer particular promise for C1-based biomanufacturing due to native metabolic properties, enzyme activities, and substrate tolerance that may be difficult to engineer into conventional chassis organisms. For example, *Pseudomonas putida* KT2440 possesses inherent qualities that establish it as a premier synthetic biology chassis, including efficient growth with minimal maintenance requirements and natural capability to harness diverse substrates including acetate, crude glycerol, and lignin-derived aromatic compounds [50]. Engineering these organisms for enhanced C1 assimilation involves implementing standardized workflows that encompass substrate and target product evaluation, fermentation parameters, bioreactor selection, and downstream processing—all informed by preliminary TEA and LCA [18].

Key metabolic engineering strategies for improving C1 assimilation efficiency include **flux balance analysis (FBA)** to predict steady-state flux distributions optimizing objectives like biomass formation, **enzyme cost minimization (ECM)** to estimate optimal enzyme and metabolite concentrations supporting desired flux distributions, and **minimum-maximum driving force (MDF)** models to identify pathways with highest thermodynamic driving forces [18]. These computational approaches, when coupled with TEA and LCA, enable researchers to prioritize engineering targets that simultaneously enhance both economic and environmental performance.



Diagram 1: C1 Biomanufacturing Development Workflow

Techno-Economic Challenges in C1 Biomanufacturing

Despite significant technical advances, C1 biomanufacturing faces substantial economic barriers that must be addressed to achieve commercial viability. Comprehensive TEA studies reveal several critical challenges [86].

Low carbon conversion efficiency represents a fundamental economic constraint, with C1 feedstock-to-chemical conversion rates frequently below 10%—significantly lower than conventional petrochemical routes. This inefficiency drives increased capital expenditures by requiring larger-scale infrastructure to compensate for productivity losses. Fermentation-related equipment typically accounts for the largest share of capital costs, often exceeding 90% of total equipment expenses in biological C1 conversion processes [86].

Feedstock cost and variability present additional economic hurdles. Unlike centralized petroleum refining with consistent supply chains, C1 resources are often decentralized with significant regional variations in availability, composition, and cost. For example, waste methane sources range from less than one ton per day at wastewater treatment plants to 31 tons per day at landfills, creating disparate economies of scale. Furthermore, feedstock costs typically dominate operating expenditures, accounting for more than 57% of OPEX in many C1 conversion processes [86].

Table 2: Economic Comparison of Carbon Utilization Technologies

Technology	Cost Range (per tCO_2)	Key Cost Drivers	Technology Readiness
Ocean Iron Fertilization	$25-$53,000 (sensitivity range) [85]	MRV processes, nutrient robbing, carbon export efficiency	Early deployment (TRL 6-7)
Indirect Carbonation (mining waste)	$500-$1,800 (with acid recycling) [88]	Acid-base recycling, carbonate precipitation, critical mineral recovery	Pilot scale (TRL 5-6)

Technology	Cost Range (per tCO_2)	Key Cost Drivers	Technology Readiness
C1 Biomanufacturing (3-HP production)	Varies by pathway	Feedstock cost, bioreactor capacity, carbon yield	Lab to pilot scale (TRL 3-5)
Waste-derived Activated Carbon	Sensitivity to variable costs	Activation chemicals, renewable energy integration	Commercial (TRL 9)

Life Cycle Assessment of Carbon Metabolism Strategies

LCA provides critical insights into the environmental performance of engineered carbon metabolism systems, revealing trade-offs and improvement opportunities that may not be apparent from technical or economic metrics alone [89] [87].

For C1 biomanufacturing platforms, LCA studies consistently highlight the importance of **feedstock sourcing** and **energy inputs** in determining overall environmental impacts. C1 substrates derived from industrial waste streams (e.g., steel mill off-gases) typically offer superior environmental performance compared to purpose-produced C1 feedstocks, due to avoided emissions from waste management and allocation of burdens across multiple products. Similarly, integrating renewable energy sources can reduce life cycle emissions by up to 72%, as demonstrated in activated carbon production for CO_2 capture [89].

The **choice of functional unit** significantly influences LCA results and interpretation. While mass-based functional units (e.g., kg CO_2eq per kg product) are commonly used, monetary units (e.g., kg CO_2eq per $ value) may provide complementary perspectives, particularly for differentiated products with significant price variations. This distinction is evident in agricultural LCAs, where grass-finished beef demonstrates different environmental performance relative to conventional beef when evaluated using economic versus mass-based functional units [87].

Experimental and Computational Methodologies

Integrated TEA/LCA Workflow for Metabolic Engineering

Implementing harmonized TEA and LCA for central carbon metabolism engineering requires systematic methodologies that span experimental and computational approaches. The following workflow outlines key stages for comprehensive technology assessment [18] [86].

Stage 1: Goal Definition and Scoping

- Define assessment objectives and decision context
- Establish system boundaries (cradle-to-gate vs. cradle-to-grave)
- Select functional unit(s) aligned with TEA and LCA
- Identify benchmark technologies for comparison

Stage 2: Inventory Analysis

- Compile experimental data on pathway performance (titer, rate, yield)
- Quantify material and energy inputs for unit operations
- Document equipment specifications and operating parameters
- Characterize waste streams and emissions

Stage 3: Impact Assessment

- Calculate techno-economic metrics (CAPEX, OPEX, MSP)
- Quantify environmental impacts using LCIA methods
- Conduct uncertainty and sensitivity analyses
- Perform comparative analysis against benchmarks

Stage 4: Interpretation and Integration

- Identify critical technical and economic bottlenecks
- Propose design modifications to improve performance
- Evaluate trade-offs between economic and environmental objectives
- Formulate recommendations for research prioritization

Protocol for Prospective TEA of Novel CDR Approaches

Ocean iron fertilization and other marine carbon dioxide removal (mCDR) strategies require specialized TEA methodologies that account for unique monitoring, reporting, and verification (MRV) requirements and environmental uncertainties. The following protocol adapts conventional TEA frameworks to these challenges [85].

Process Decomposition and Modeling

- Divide the OIF process into discrete stages: material production, transport to ocean, CDR operation, and MRV
- Develop mass and energy balances for each stage
- Model carbon sequestration accounting for export efficiency, ventilation losses, and equivalent carbon losses from N_2O production

Cost Element Characterization

- Estimate capital costs including specialized vessels, mixing equipment, storage tanks, and monitoring systems
- Quantify operating costs: iron source, labor, vessel operation, maintenance, and MRV implementation
- Include administrative, regulatory compliance, and verification expenses often excluded from preliminary assessments

Uncertainty and Sensitivity Analysis

- Identify high-impact parameters through local sensitivity analysis (one-at-a-time variation)
- Assess cost sensitivity to oceanographic parameters (export efficiency, nutrient robbing) and engineering parameters (equipment costs)
- Develop probabilistic cost models using Monte Carlo simulation to characterize uncertainty ranges

Learning Rate Application

- Apply technology learning curves to estimate cost reductions from FOAK to NOAK deployments
- Differentiate between potentially reducible costs (engineering, operations) and fixed costs (MRV, verification)
- Contextualize results relative to other CDR approaches using harmonized assessment guidelines



Diagram 2: Integrated TEA/LCA Methodology

Essential Research Reagents and Tools

Research Reagent Solutions for Metabolic Engineering

Advanced metabolic engineering of central carbon metabolism requires specialized reagents and tools for pathway construction, host engineering, and performance characterization. The following table details essential research materials and their applications in developing C1 assimilation platforms [18] [57] [50].

Table 3: Essential Research Reagents for Carbon Metabolism Engineering

Reagent/Tool Category	Specific Examples	Function in Metabolic Engineering
Molecular Cloning Tools	CRISPR-Cas9 systems, SacB counter-selection vectors, Gateway cloning systems	Genome editing, pathway integration, multi-gene assembly

Reagent/Tool Category	Specific Examples	Function in Metabolic Engineering
Analytical Standards	^{30}C-labeled substrates, metabolite standards, internal standards	Metabolic flux analysis, quantification of pathway intermediates
Culture Media Components	M9 minimal medium, defined nutrient mixes, selective antibiotics	Controlled cultivation conditions, mutant selection, phenotypic characterization
Enzyme Assay Kits	NADP/NADPH quantification, metabolic enzyme activity assays, ATP determination	Pathway validation, kinetic parameter determination, bottleneck identification
Omics Analysis Platforms	RNA sequencing kits, proteomics sample preparation, metabolomics extraction	Systems-level analysis of engineered strains, regulatory network mapping

Computational Tools for TEA and LCA

Computational modeling represents an essential component of metabolic engineering TEA and LCA, enabling prediction of system performance and identification of optimization targets prior to resource-intensive experimental implementation [18] [84].

Metabolic Modeling Platforms include flux balance analysis (FBA) software such as COBRA Toolbox and RAVEN, which enable constraint-based modeling of metabolic networks. These tools predict theoretical yield maxima, identify essential genes, and suggest knockout targets for growth-coupled production. **Process Simulation Software** like Aspen Plus and SuperPro Designer facilitate detailed process modeling, equipment sizing, and energy integration—providing critical data for capital and operating cost estimation. **LCA Software** including openLCA, SimaPro, and the GREET model provide databases and calculation methods for environmental impact assessment across multiple categories [86] [84].

Integrated computational workflows combine these tools to enable multi-objective optimization of metabolic engineering strategies. For example, flux balance analysis can identify pathway configurations that maximize carbon conversion efficiency, with results subsequently fed into process simulation to estimate energy requirements, which then inform both TEA and LCA calculations. This integrated approach allows researchers to rapidly evaluate design alternatives and focus experimental efforts on the most promising configurations [18] [86].

Techno-Economic Analysis and Life Cycle Assessment provide indispensable frameworks for guiding the development of sustainable biotechnologies based on engineered central carbon metabolism. As metabolic engineering capabilities advance, enabling increasingly sophisticated manipulation of C1 assimilation pathways and carbon fixation strategies, the integration of TEA and LCA at early research stages becomes ever more critical for ensuring that laboratory innovations translate to economically viable and environmentally beneficial industrial processes.

The continued harmonization of TEA and LCA methodologies, development of prospective assessment frameworks that incorporate technology learning, and creation of standardized benchmarking approaches will enhance our ability to objectively evaluate and compare emerging carbon management technologies. Furthermore, the expanding application of these analytical methods to novel carbon dioxide removal strategies—from ocean iron fertilization to microbial C1 biomanufacturing—will provide essential insights for prioritizing research investments and designing policies that effectively address the urgent challenge of climate change while advancing toward a circular carbon economy.

Comparative Performance of Model vs. Non-Model Chassis Organisms

The selection of microbial chassis organisms is a foundational decision in metabolic engineering and synthetic biology. For decades, the field has been dominated by a narrow set of model organisms, primarily *Escherichia coli* and *Saccharomyces cerevisiae*, valued for their genetic tractability and well-characterized physiology. However, the expanding scope of biotechnological applications—ranging from sustainable biomanufacturing to therapeutic development—has exposed the limitations of these traditional workhorses. This whitepaper provides a comparative analysis of model and non-model chassis organisms, focusing on their performance within the critical context of central carbon metabolism (CCM) engineering. By examining innate physiological advantages, engineering challenges, and emerging strategies for domesticating novel microbes, this review aims to guide researchers in making informed chassis selection decisions for advanced metabolic engineering applications.

The historical bias toward model organisms in synthetic biology has been a significant self-imposed design constraint [90]. While these chassis have been invaluable for proof-of-concept systems and foundational genetic tool development, they may not represent the optimal platform for many industrial applications. The concept of **Broad-Host-Range (BHR) synthetic biology** has emerged to challenge this paradigm, advocating for the strategic selection of microbial hosts based on innate functional traits rather than defaulting to traditional models [90].

Central carbon metabolism serves as the core engine of the cell, governing the flow of carbon and energy from nutrients to biomass and products. Engineering CCM is therefore a powerful strategy for redirecting metabolic flux toward desired compounds [2]. The suitability of a chassis for CCM engineering depends on a complex interplay of factors, including its native metabolic network architecture, regulatory systems, and physiological constraints. This review systematically compares the capabilities of model and non-model organisms as chassis for CCM engineering, providing a framework for rational chassis selection in metabolic engineering projects.

Comparative Analysis: Physiological and Metabolic Attributes

The performance of a microbial chassis is determined by a combination of its innate physiological attributes and the ease with which it can be genetically engineered. The table below summarizes key comparative characteristics of model and non-model organisms.

Table 1: Comparative Attributes of Model and Non-Model Chassis Organisms

Attribute	Model Organisms (e.g., *E. coli*, *S. cerevisiae*)	Non-Model Organisms (e.g., *Z. mobilis*, *C. necator*, *H. bluephagenesis*)
Genetic Toolkits	Extensive, standardized, and optimized [90]	Often limited; requires development for each new host [49] [91]
Physiological Knowledge	Deeply characterized with comprehensive omics datasets [92]	Often treated as "black boxes"; information can be scarce [18]
Native CCM Features	Standard pathways (e.g., glycolysis, TCA); well-understood regulation [2]	Diverse and specialized (e.g., ED pathway in *Z. mobilis*) [49] [93]
Biotechnological Niches	General-purpose chassis; proven for many products	Specialists for specific substrates/products; often superior in harsh conditions [90] [91]
Typical Engineering Workflow	Streamlined DBTL cycles	Challenging; focus on initial tool development and host characterization [18]
Regulatory & Safety	Generally recognized as safe (GRAS) status for some	Varies; may require new safety approvals [91]

The "Chassis Effect" on Engineered Functions

A critical consideration in chassis selection is the **"chassis effect"**—where identical genetic constructs exhibit different behaviors across host organisms due to host-construct interactions [90]. These interactions arise from:

- **Resource competition** for finite cellular resources like ribosomes, RNA polymerase, and metabolites [90].
- **Regulatory crosstalk** between introduced genetic circuits and the host's native regulatory networks [90].
- **Metabolic burden** and growth feedback, which can alter host physiology and circuit performance unpredictably [90].

Systematic comparisons have shown that host selection significantly influences key performance parameters of genetic devices, including output signal strength, response time, and stability [90]. This positions the host chassis not merely as a passive platform but as an active **tuning module** that can be leveraged to optimize system performance [90].

Central Carbon Metabolism Engineering Strategies

Engineering central carbon metabolism is a primary lever for altering metabolic flux. The strategies and their implementation differ significantly between model and non-model hosts.

Engineering Approaches and Outcomes

Table 2: CCM Engineering Strategies and Representative Outcomes

Engineering Strategy	Description	Example Organism	Key Result	Citation
Heterologous Pathway Introduction	Adding non-native pathways to create shortcuts or bypass bottlenecks.	*S. cerevisiae*	Introduction of the phosphoketolase (PHK) pathway increased acetyl-CoA supply, boosting fatty acid ester production to ~5100 g/CDW. [2]	
Dominant Pathway Compromise	Attenuating native, high-flux pathways to redirect carbon.	*Z. mobilis*	Weakening the dominant ethanol pathway enabled redirection of carbon to D-lactate, achieving >140 g/L. [49]	
Enzyme & Regulatory Optimization	Modifying expression of key CCM enzymes or transcription factors.	*E. coli*	Expression of a response regulator (DR1558) improved NADPH supply and enhanced PHB biosynthesis. [2]	

Engineering Strategy	Description	Example Organism	Key Result	Citation
Genome Reduction	Removing non-essential genes to reduce metabolic burden and improve predictability.	*E. coli*, *B. subtilis*	Deletion of mobile genetic elements increased recombinant protein production by 20-25% and genomic stability. [91]	

Model-Based Metabolic Design Constraints

Computational models are indispensable for guiding CCM engineering. The constraints used in these models directly impact the feasibility of their predictions.

Table 3: Key Constraints for Model-Based Metabolic Design

Constraint Category	Specific Constraints	Application in Kinetic Models	Application in Stoichiometric Models
General (Universal)	Mass conservation, Energy balance, Steady-state assumption, Thermodynamic constraints	Yes	Yes (Foundation of FBA)
Organism-Level	Total enzyme activity, Homeostatic constraint (limits on metabolite concentration changes), Cytotoxic metabolite levels	Yes	Limited
Experiment-Level	Substrate uptake rates, Oxygen availability, Cell size/surface/volume ratios	Yes (if dynamic)	Yes

The integration of **enzyme constraints** into Genome-Scale Metabolic Models (GEMs) has been a significant advance. For example, the development of eciZM547 for *Zymomonas mobilis* provided more accurate predictions of metabolic flux and growth by accounting for the proteomic cost of enzyme synthesis, unlike its predecessor iZM516 [49].

Experimental Workflows and Protocols

The practical workflow for engineering CCM varies significantly between established model organisms and emerging non-model chassis.

A Generalized Workflow for Engineering Non-Model Chassis

The following diagram illustrates a structured workflow for developing and engineering a non-model organism, integrating tools from synthetic biology and systems biology.



Diagram 1: Non-model chassis development and engineering workflow. VR: Vector Resources; TEA: Techno-Economic Analysis; LCA: Life Cycle Assessment.

Protocol: Absolute Quantification of Central Carbon Metabolites

A critical step in characterizing chassis performance is the precise measurement of intracellular metabolite concentrations. The following protocol, adapted from quantitative mass spectrometry workflows, is essential for validating CCM models [92].

1. Sampling and Quenching:

- Rapidly culture samples (e.g., via rapid filtration or cold methanol quenching) to instantly halt metabolic activity.
- Critical Parameters: Speed and temperature are paramount to prevent metabolite turnover.

2. Metabolite Extraction:

- Use a combination of cold organic solvents (e.g., methanol:acetonitrile:water) for comprehensive extraction of hydrophilic metabolites.
- Include internal standards, preferably **U-13C-labeled metabolite extracts** or other isotopologues, for **isotope dilution mass spectrometry (IDMS)** to correct for matrix effects and ionization efficiency [92].

3. Chromatographic Separation and MS Detection:

- Employ multiple complementary LC-MS/MS methods to cover the physicochemical diversity of the central carbon metabolome:
  - **Reversed-Phase (RP) LC:** For moderately hydrophobic metabolites.
  - **Hydrophilic Interaction Liquid Chromatography (HILIC):** For polar metabolites.
  - **Ion Chromatography (IC):** For superior resolution of phosphorylated sugars and organic acids [92].
- Use tandem mass spectrometry (MS/MS) in selected reaction monitoring (SRM) mode for high selectivity and sensitivity.

4. Data Processing and Quantification:

- Use the response ratio (analyte vs. internal standard) for quantification.
- Interpolate absolute concentrations from calibration curves of pure analytical standards.
- Normalize to cell density, cell dry weight (DW), and cell volume to report final **intracellular concentrations in µmol/gDW or mM** [92].

The Scientist's Toolkit: Essential Research Reagents

Successful CCM engineering relies on a suite of specialized reagents and tools. The following table details key solutions for chassis evaluation and engineering.

Table 4: Key Research Reagent Solutions for CCM Engineering

Reagent / Tool Category	Specific Examples	Function & Application
Genetic Toolkits	SEVA (Standard European Vector Architecture) plasmids, Broad-host-range CRISPR systems [90]	Enable standardized genetic manipulation across diverse bacterial hosts.
Quantitative Metabolomics Standards	U-13C-labeled S. cerevisiae extract, Commercially available 13C/15N isotopologues [92]	Essential for absolute quantification of intracellular metabolites via isotope dilution.
Analytical Standards	Purified metabolites for glycolysis, PPP, TCA cycle, amino acids, nucleotides [92]	Used to create calibration curves for precise concentration determination in LC-MS/MS.
Enzyme Kinetic Parameters	kcat values from databases (e.g., AutoPACMEN, DLkcat) [49]	Parameterize enzyme-constrained metabolic models (ecModels) for improved predictions.
Specialized Culture Media	Mineral and rich media for physiological comparison; C1 substrates (methanol, formate) [18] [92]	Assess chassis performance and substrate utilization range under different conditions.

The dichotomy between model and non-model chassis organisms is artificial and increasingly obsolete. The future of metabolic engineering lies in a **portfolio approach**, where the chassis is selected as a tunable design parameter based on the specific requirements of the application [90]. Model organisms will remain powerful platforms for fundamental research and for products where their physiology is adequate. However, for applications demanding specialized capabilities—such as the utilization of one-carbon feedstocks, operation under extreme conditions, or the synthesis of complex natural products—non-model organisms often provide a superior and more efficient starting point [18] [91].

The ongoing development of **broad-host-range synthetic biology tools** [90], more sophisticated **enzyme-constrained metabolic models** [49], and streamlined **genome reduction techniques** [91] is systematically lowering the barrier to domesticating non-model microbes. Integrating techno-economic analysis (TEA) and life cycle assessment (LCA) at the initial design stage will further ensure that the engineered chassis not only achieves high performance in the laboratory but also drives sustainable and economically viable bioprocesses [18] [49]. By moving beyond traditional models and strategically leveraging microbial diversity, researchers can unlock new possibilities for a sustainable bio-based economy.

The engineering of **central carbon metabolism (CCM)** is a cornerstone of modern metabolic engineering, providing the foundational precursors, energy, and redox cofactors essential for biosynthetic pathways. Optimization of CCM—encompassing glycolysis, the tricarboxylic acid (TCA) cycle, and the pentose phosphate pathway (PPP)—enables global rearrangement of metabolic flux to support the high-yield production of valuable compounds. This whitepaper benchmarks advanced engineering strategies through three seminal case studies: the anti-malarial drug **artemisinin**, diverse **isoprenoid** platforms, and the platform chemical **3-hydroxypropionic acid (3-HP)**. Each case demonstrates the critical interplay between precursor availability, cofactor balancing, and pathway localization, framed within the context of CCM engineering strategies. The insights derived provide a template for optimizing the production of high-value natural products and bulk chemicals in microbial and plant-based systems.

Artemisinin: Engineering Isoprenoid Flux in Multiple Chassis

Artemisinin, a potent sesquiterpene lactone anti-malarial, is synthesized from isoprenoid precursors supplied by the **methylerythritol phosphate (MEP)** pathway in plants or the **mevalonate (MVA)** pathway in engineered microbes. Its biosynthesis highlights the critical need to augment the supply of the universal five-carbon isoprenoid building blocks, **isopentenyl diphosphate (IPP)** and **dimethylallyl diphosphate (DMAPP)**, and their downstream conjugate, **farnesyl pyrophosphate (FPP)**.

Key Metabolic Engineering Strategies

- **Enhancing Precursor Supply**: In *Artemisia annua*, overexpression of genes from the native MVA pathway, such as *HMGR* (the rate-limiting step), and the MEP pathway, such as *HDR*, significantly increases artemisinin yield [94]. In microbial chassis, the entire MVA pathway is often heterologously expressed to bolster IPP and DMAPP flux [95].
- **Upregulating Committed Pathway Steps**: Overexpression of **amorpha-4,11-diene synthase (ADS)**, which cyclizes FPP to amorphadiene, and the cytochrome P450 enzyme **CYP71AV1**, which oxidizes amorphadiene to artemisinic acid, are proven effective strategies [94].

- **Blocking Competitive Pathways**: Engineering chassis to reduce carbon flux toward sterol synthesis or other sesquiterpenes increases the precursor pool available for artemisinin pathway [2].
- **Host and Pathway Compartmentalization**: Compartmentalizing the artemisinin pathway in plant chloroplasts or engineering peroxisomal flux in yeast optimizes substrate channeling and mitigates metabolic crosstalk [96].
- **Advanced Chassis Engineering**:
  - In *Bacillus subtilis*, fusion of amorphadiene synthase to **Green Fluorescent Protein (GFP)** enhanced enzyme stability and expression. Coupled with MEP pathway and FPPS overexpression, this yielded **416 mg/L** of amorphadiene [97].
  - **Dual-acting immuno-antibiotics (DAIAs)** represent a novel strategy where IspH inhibitors in the MEP pathway directly kill pathogens like *Acinetobacter baumannii* and *Klebsiella pneumoniae* while simultaneously activating human Vγ9Vδ2 T-cells for a synergistic immune response [98].

Table 1: Benchmarking Artemisinin and Precursor Production in Engineered Chassis

Chassis Organism	Engineering Strategy	Key Enzymes/Pathways Targeted	Maximum Reported Titer	Citation
Artemisia annua (Plant)	Overexpression of pathway genes; Increasing GST density	AaADS, AaCYP71AV1, AaCPR, AaDBR2, HMGR, FPS	0.1-1% of Dry Weight (native)	[94]
Saccharomyces cerevisiae (Yeast)	Heterologous MVA pathway; ADS & CYP71AV1 expression	MVA pathway, ADS, CYP71AV1	Not Specified	[95]
Escherichia coli (Bacteria)	Heterologous MVA pathway; ADS expression	MVA pathway, ADS	>100 mg/L (amorphadiene)	[95]
Bacillus subtilis (Bacteria)	GFP-ADS fusion; MEP & FPPS overexpression	MEP pathway, FPPS, GFP-ADS	416 mg/L (amorphadiene)	[97]

Experimental Protocol: High-Titer Amorphadiene Production inB. subtilis

Objective: To achieve high-level production of amorphadiene in *Bacillus subtilis* through integrated pathway engineering and medium optimization [97].

Methodology:

- **Strain Construction:**
  - Genetically fuse amorphadiene synthase (ADS) with Green Fluorescent Protein (GFP) to enhance protein expression and stability.
  - Establish a co-expression system with a synthetic operon containing key MEP pathway genes (*dxs*, *ispD*, etc.).
  - Overexpress farnesyl pyrophosphate synthase (FPPS) to ensure sufficient FPP substrate for ADS.
  - Combinatorially integrate these genetic modifications (GFP-ADS, FPPS, plasmid-borne MEP operon) into the *B. subtilis* chromosome.
- **Fermentation and Analysis:**
  - Inoculate engineered strains in shake flasks with a defined production medium.
  - Conduct systematic medium optimization using **Experimental Design** methodologies (e.g., Response Surface Methodology) to identify critical components. Key supplements identified include **pyruvate** (a MEP pathway precursor) and **K_2HPO_4** (for pH buffering and phosphorus supply).
  - Extract amorphadiene from the culture broth using an organic solvent (e.g., dodecane) overlay.
  - Quantify amorphadiene titers using **Gas Chromatography-Mass Spectrometry (GC-MS)**.

Isoprenoid Platforms: Diversifying Biosynthesis beyond the Isoprene Rule

Isoprenoids represent the largest class of natural products. Traditional production relies on the condensation of C5 IPP and DMAPP, conforming to the "isoprene rule." Recent advances focus on diversifying these platforms by utilizing atypical carbon substrates and creating non-canonical isoprenoid analogs.

Key Metabolic Engineering Strategies

- **Carbon Source Switching**: To reduce substrate costs, engineering efforts have enabled isoprenoid production from C1 sources (methanol, CO_2) in methylotrophic *Pichia pastoris* or cyanobacteria, and from aromatic compounds in *Pseudomonas putida* [95].
- **Generating Isoprenoid Analogs**: A novel yeast-based platform bypasses endogenous IPP/DMAPP synthesis. It uses the kinase pair **AtFKI/AtIPK** to phosphorylate exogenous **prenol-like alcohols** (e.g., C6) into non-canonical DMAPP/IPP analogs. These are incorporated by promiscuous terpene synthases to produce isoprenoids with extended carbon skeletons (e.g., C11 ethyllinalool) [99].

- **Pathway Selection and Balancing**: The choice between the MVA (cytosolic) and MEP (plastid) pathways is critical. Hybrid strategies, such as introducing the MVA pathway in *E. coli*, have successfully increased flux. Fine-tuning using CRISPRi to balance gene expression within these pathways is a key optimization step [95].
- **Non-Model Chassis Development**: Oleaginous yeasts like *Yarrowia lipolytica* and *Rhodosporidium toruloides* are emerging chassis due to their high innate acetyl-CoA flux and tolerance to lipophilic compounds [95].

Table 2: Isoprenoid Production Platforms and Engineering Strategies

Platform / Strategy	Chassis Organism	Carbon Source	Key Innovation	Application Example
Standard MVA/MEP	*E. coli, S. cerevisiae*	Glucose, Glycerol	Pathway balancing; CRISPRi tuning	High-titer production of bisabolene, limonene [95]
C1 Metabolism	*P. pastoris*, Cyanobacteria	Methanol, CO_2	Lower cost substrate; Sustainable feedstock	Isoprenoid production from waste streams [95]
Non-Canonical Analog Production	*S. cerevisiae*	Glucose + C6 alcohol	AtFKI/AtIPK kinase pathway	Synthesis of ethyllinalool and cannabinoid analogs [99]
Oleaginous Yeast	*Y. lipolytica, R. toruloides*	Various	Native high acetyl-CoA flux	Production of terpenoids from lipids [95]

Experimental Protocol: Biosynthesis of Isoprenoid Analogs in Yeast

Objective: To produce non-canonical isoprenoid analogs in *Saccharomyces cerevisiae* using an engineered pathway that bypasses native IPP and DMAPP synthesis [99].

Methodology:

- **Strain Engineering:**
  - Clone and express the kinase pair **AtFKI** (*Arabidopsis thaliana* farnesol kinase) and **AtIPK** (*A. thaliana* isopentenyl phosphate kinase) in a suitable yeast strain (e.g., EGY48).
  - Introduce genes for the downstream terpene biosynthetic pathway (e.g., linalool synthase for monoterpenes, cannabinoid acid synthase for cannabinoids).
- **Feeding and Fermentation:**
  - Supplement the culture medium with the desired non-canonical alcohol (e.g., prenol for C5, C6 alcohols for C6 analogs).
  - Induce expression of the heterologous kinases and synthases.
- **Product Analysis:**
  - For volatile compounds (e.g., linalool analogs), use **Headspace Solid-Phase Microextraction (HS-SPME)** coupled with **GC-MS** for identification and quantification.
  - For non-volatile compounds (e.g., cannabinoid analogs), employ Liquid Chromatography-Mass Spectrometry (LC-MS).

3-HP Production: Optimizing the Malonyl-CoA Pathway

3-Hydroxypropionic acid (3-HP) is a key platform chemical for acrylic acid and biodegradable plastics. Its biosynthesis primarily relies on the malonyl-CoA pathway, making the optimization of malonyl-CoA and NADPH supply from CCM a primary engineering target.

Key Metabolic Engineering Strategies

- **Enhancing Malonyl-CoA Supply**: This is achieved by overexpressing **acetyl-CoA carboxylase (ACC1)**, a rate-limiting enzyme. A common strategy uses an ACC1 mutant (S659A/S1157A) to bypass phosphorylation-based inhibition [100].
- **Introducing Efficient Reductases**: The bifunctional **malonyl-CoA reductase (MCR) from *Chloroflexus aurantiacus*** is expressed to convert malonyl-CoA to 3-HP. Splitting MCR into two functional domains (MCR-N and MCR-C) and expressing them separately significantly improves activity and 3-HP yield [101] [100].
- **Boosting Cofactor Regeneration**: Overexpression of genes like cytosolic NADH kinase (*cPOSSc*) increases NADPH availability, which is consumed by MCR [101].
- **Blocking Competing Pathways**: Knocking out genes in the glyoxylate cycle (*CIT2*, *MLS1*) and 3-HP degradation pathways (*MMSDH*, *HPDH*) channels flux toward 3-HP accumulation [100].
- **Host Engineering and Pathway Introduction**:
  - In *Pichia pastoris*, expressing *mcrCa* and engineering for precursor availability achieved a titer of **24.75 g/L** and a volumetric productivity of **0.54 g/L/h** in fed-batch culture [101].

- In *Yarrowia lipolytica*, introducing MCR, *ACSL641P*, and overexpressing *ACC1* while knocking out degradation pathways yielded **16.23 g/L** 3-HP in a fed-batch bioreactor [100].

Table 3: Benchmarking 3-HP Production in Yeast Chassis via the Malonyl-CoA Pathway

Chassis Organism	Engineering Strategy	Key Genetic Modifications	Max Titer (Fed-Batch)	Volumetric Productivity (g/L/h)	Citation
Pichia pastoris	MCR expression; Precursor optimization	*mcrCa*; NADPH & Malonyl-CoA supply genes	24.75 g/L	0.54	[101]
Yarrowia lipolytica	MCR expression; Blocking degradation	*MCR-N/CCa*, *ACC1*, *ACSL641P*, Δ*MMSDH*, Δ*HPDH*	16.23 g/L	Not Specified	[100]
Saccharomyces cerevisiae	Dynamic sensor-regulator system	FapR-based malonyl-CoA sensor; *PMP1*, *TPL1* overexpression	1.2 g/L (shake flask)	Not Specified	[100]

Experimental Protocol: High-Level 3-HP Production inP. pastoris

Objective: To engineer *Pichia pastoris* for efficient production of 3-HP from glycerol via the malonyl-CoA pathway [101].

Methodology:

- **Base Strain Construction:**
  - Integrate the gene encoding the bifunctional **malonyl-CoA reductase from *Chloroflexus aurantiacus (mcrCa)*** into the *P. pastoris* genome under a strong constitutive promoter (e.g., GAP promoter).
- **Strain Optimization:**
  - **Protein Engineering:** Split the *mcrCa* gene and express the N-terminal (MCR-N) and C-terminal (MCR-C) domains separately to improve catalytic efficiency.
  - **Metabolic Engineering:** Implement further genetic modifications to increase the supply of the MCR substrates, malonyl-CoA and NADPH. This can involve overexpressing native *ACC1* and enzymes from the PPP.
- **Fed-Batch Fermentation:**
  - Cultivate the engineered strain in a bioreactor with a defined minimal medium using glycerol as the sole carbon source.
  - Initiate a fed-batch phase with a controlled glycerol feed rate to maintain growth and production while avoiding overflow metabolism.
  - Maintain the pH below the pKa of 3-HP (4.5) to facilitate product retention in its anionic form and simplify downstream processing.
- **Analytics:**
  - Monitor cell density (OD600), glycerol consumption, and 3-HP production.
  - Quantify 3-HP concentration in the culture supernatant using **High-Performance Liquid Chromatography (HPLC)**.

Pathway Visualization and Logical Workflows

Central Carbon Metabolism and Product Synthesis

Diagram 1: CCM routes to 3-HP, Artemisinin, and Isoprenoids. Key nodes and pathways are color-coded: CCM intermediates (white), 3-HP pathway (red), isoprenoid/artemisinin pathway (green). The diagram shows how glycolysis and TCA cycle feed acetyl-CoA and malonyl-CoA for 3-HP, while the MEP and MVA pathways generate IPP/DMAPP for artemisinin. Abbreviations: ACC, acetyl-CoA carboxylase; MCR, malonyl-CoA reductase; MEP, methylerythritol phosphate pathway; MVA, mevalonate pathway; IspH, key MEP pathway enzyme; ADS, amorphadiene synthase.

Experimental Workflow for Microbial Metabolic Engineering



Diagram 2: Generalized metabolic engineering workflow. The process is iterative, starting with target identification and moving through host selection, genetic construction, cultivation, and analytical validation. Systems-level analysis informs further rational or dynamic engineering to optimize production, creating a feedback loop for continuous strain improvement.

The Scientist's Toolkit: Essential Research Reagents and Materials

Table 4: Key Reagents and Materials for Metabolic Engineering in Featured Case Studies

Reagent / Material	Function / Application	Example Use Case(s)
Malonyl-CoA Reductase (MCR)	Key enzyme in malonyl-CoA pathway; converts malonyl-CoA to 3-HP via malonate semialdehyde.	3-HP production in *P. pastoris* [101] and *Y. lipolytica* [100].
Amorpha-4,11-diene Synthase (ADS)	Terpene synthase; catalyzes the cyclization of FPP to amorphadiene, the first committed step in artemisinin biosynthesis.	Artemisinin precursor production in *E. coli*, yeast [95], and *B. subtilis* [97].
IspH Inhibitors (e.g., C23 analogs)	Small molecule inhibitors of the MEP pathway enzyme IspH; act as dual-acting immuno-antibiotics.	Antibacterial therapy against multidrug-resistant Gram-negative pathogens [98].
Acetyl-CoA Carboxylase (ACC1)	Catalyzes the ATP-dependent carboxylation of acetyl-CoA to malonyl-CoA; a rate-limiting step in fatty acid and 3-HP biosynthesis.	Engineered in *S. cerevisiae* and *Y. lipolytica* to boost malonyl-CoA supply for 3-HP [100].
AtFKI / AtIPK Kinase Pair	Enables phosphorylation of non-canonical prenol-like alcohols to form non-C5 IPP/DMAPP analogs.	Biosynthesis of isoprenoid analogs (e.g., ethyllinalool) in *S. cerevisiae* [99].
GFP Fusion Protein	Used as a solubility and expression tag to enhance the stability and production of recombinant proteins.	Improved expression and activity of ADS in *B. subtilis* [97].
FapR-based Malonyl-CoA Sensor	Transcription factor-based biosensor for detecting intracellular malonyl-CoA levels.	Dynamic regulation and high-throughput screening of engineered *S. cerevisiae* for 3-HP production [100].

The scaling of fermentation processes from laboratory to industrial scale is a critical, complex, and costly endeavor in biomanufacturing. Despite advances in metabolic engineering and synthetic biology that have streamlined the development of high-yielding microbial chassis, the **transition to**

large-scale production remains a significant bottleneck. This process can take **3–10 years** and cost between **$100 million to $1 billion**, often failing to replicate laboratory performance in an industrial setting [102]. A primary reason for this failure is the **disconnect between strain development and production realities**; while laboratory-scale fermenters offer homogeneous and tightly controlled conditions, industrial-scale bioreactors introduce **physical and chemical gradients** that can negatively impact microbial physiology and productivity [103] [102]. Viewing scale-up through the lens of **central carbon metabolism (CCM)** is crucial, as the fluxes through these core pathways are highly sensitive to environmental fluctuations. Successful scale-up, therefore, requires not only robust microbial strains but also a deep understanding of how large-scale bioreactor hydrodynamics interact with cellular physiology [102]. This guide outlines the key challenges, scale-down experimental strategies, and modeling tools essential for bridging this lab-to-factory gap.

Key Challenges in Fermentation Scale-Up

Scaling a fermentation process introduces fundamental changes in the physical and chemical environment experienced by the microbial cells. Understanding these challenges is the first step toward designing a successful scale-up strategy.

Bioreactor Heterogeneity and Gradients

In contrast to the homogeneous conditions of lab-scale vessels, industrial-scale fermenters exhibit significant spatial and temporal variations.

- **Mixing and Mass Transfer Limitations**: As bioreactor volume increases, achieving uniform mixing becomes more difficult. Parameters like **mixing time** increase substantially, leading to heterogeneous zones. Cells circulate through varying conditions of **dissolved oxygen (DO)**, **substrate concentration**, and **pH** [102]. For example, in an aerobic process, oxygen concentrations can be high at the sparger bottom and low at the top, with gradients in between. Similarly, nutrient concentrations are often highest near the feed point and lowest in other regions [103].
- **Impact on Physiology and Yield**: These gradients force microorganisms to experience **dynamic and often suboptimal conditions**, which can alter their metabolic flux. Cells may shift their CCM in response to feast-famine cycles, leading to **reduced overall yield** and the **production of unwanted by-products** like acetate in *E. coli* [102] [104]. This unpredictability makes it difficult to know the optimal time to stop the fermentation, complicating production planning [103].

Table 1: Key Physical-Chemical Parameter Changes During Scale-Up

Parameter	Laboratory Scale (e.g., 10 L)	Industrial Scale (e.g., 10,000 L)	Impact on Process
Mixing Time	Seconds to a few minutes	Several minutes to hours	Creates substrate, pH, and gas gradients [102]
Oxygen Transfer Rate (OTR)	Easily maintained high	Can become a major bottleneck	Limits growth and productivity in aerobic fermentations [105]
Heat Transfer	Rapid cooling/heating	Slow cooling/heating (several hours)	Critical for processes requiring precise temperature control or rapid termination [103]
Shear Stress	Generally low and uniform	Can be very high near impellers	Can damage sensitive cells or alter morphology [102] [105]

Physiological Responses and Metabolic Challenges

The non-ideal environment of a large bioreactor directly challenges the tightly regulated central carbon metabolism of production strains.

- **Overflow Metabolism**: A common issue in large-scale fermentation is the onset of overflow metabolism, where high, transient local glucose concentrations trigger cells to produce waste metabolites, most notably **acetate** in *E. coli*. Acetate is a major problem as it can **inhibit cell growth and recombinant protein production**, even at low concentrations (e.g., 0.5 g/L), and represents a significant carbon loss [104].
- **Carbon Catabolite Regulation and Energetics**: The dynamic environment can disrupt the energy and redox balance (ATP, NADH, NADPH) of the cell. Engineering strategies that modify CCM to enhance precursor supply, such as introducing the **phosphoketolase (PHK) pathway** to increase acetyl-CoA, can be successful in lab settings. However, their performance in large-scale, gradient-prone environments is not guaranteed and must be validated [23].

A Framework for Successful Scale-Up: "Scaling-Down"

The traditional approach of sequential scale-up through progressively larger vessels is slow, expensive, and risks late-stage failure. A more effective strategy is "**scaling-down**," where the constraints and heterogeneous conditions of the production bioreactor are mimicked and studied at a small, manageable scale [103].

Scale-Down Experimental Methodologies

Scale-down systems are designed to replicate, in a controlled manner, the fluctuating conditions cells experience in a large tank.

- **System Design and Principles**: These systems typically consist of one or more small fermenters linked to devices that create environmental oscillations. A common setup involves a **stirred tank reactor (STR)** connected to a **plug flow reactor (PFR)**. The circulation time and conditions within the PFR (e.g., oxygen limitation, substrate excess) can be tuned to mimic the mixing time and heterogeneity of the large-scale plant [102].
- **Protocol: Two-Compartment Scale-Down System**
  - **Apparatus Setup**: Connect a standard laboratory fermenter (STR, e.g., 1-5 L) to a long, narrow tube (PFR) using a peristaltic pump to control the circulation loop. The PFR can be equipped with ports for substrate feeding and gas monitoring.
  - **Process Configuration**: Operate the STR under constant, well-controlled conditions. The PFR is designed to create a specific stressor—for example, by making it anaerobic to mimic an oxygen gradient or by injecting a concentrated substrate feed to create a high-glucose zone.
  - **Cell Cultivation**: Inoculate the system and allow the culture to circulate between the STR and PFR. The **circulation time** is set to match the **mixing time** of the target industrial bioreactor.
  - **Sampling and Analysis**: Sample from both the STR and the PFR outlet. Analyze for metabolites (e.g., glucose, acetate), dissolved oxygen, and cell density. Use techniques like **rapid sampling and quenching** to measure short-term metabolic responses (e.g., changes in ATP, NADPH, and central metabolite pools) [102].
  - **Physiological Assessment**: Compare results with a control fermentation run in a single, homogeneous STR. Key performance indicators (KPIs) include **final product titer, yield, productivity, and by-product formation**. Transcriptomic and metabolomic analyses can reveal how the CCM is being reconfigured in response to the oscillations [102].

Application: Testing CCM-Engineered Strains

Scale-down systems are invaluable for vetting engineered strains. A strain engineered with a heterologous PHK pathway might show a 25% increase in farnesene yield in a lab flask [23]. However, when tested in a scale-down system that mimics substrate gradients, its performance might drop if the pathway enzymes are sensitive to transient substrate shortages or if the pathway disrupts the cell's redox balance under dynamic conditions. This early failure allows engineers to either re-design the strain for robustness or adjust the process conditions before costly large-scale runs.

Model-Based Tools for Scale-Up Prediction

Mathematical models are powerful tools for interpreting scale-down data and predicting large-scale performance.

Kinetic and Constraint-Based Modeling

- **Unstructured Kinetic Models**: These models describe the fermentation process at a macro level, focusing on the relationships between cell growth, substrate consumption, and product formation. They are particularly useful for predicting overall bioreactor dynamics and optimizing feeding strategies in fed-batch processes [106].

Table 2: Common Unstructured Kinetic Models for Fermentation Processes

Model Name	Mathematical Expression	Application Context
Monod	$\mu = \mu_{max} \frac{S}{K_S + S}$	Microbial growth limited by a single substrate [106]
Haldane–Andrew	$\mu = \mu_{max} \frac{S}{K_S + S + S^2/K_i}$	Growth with substrate inhibition at high concentrations [106]
Contois	$\mu = \mu_{max} \frac{S}{K_S X + S}$	High-cell-density cultures where growth is limited by mass transfer [106]
Luedeking–Piret	$\frac{dP}{dt} = \alpha \frac{dX}{dt} + \beta X$	Product formation (P) that is both growth- and non-growth-associated [106]

- **Constraint-Based Modeling (CBM)**: CBM, such as Genome-Scale Metabolic Models (GEMs), simulates the flow of metabolites through the entire metabolic network. It is used to predict how genetic modifications or environmental changes alter carbon flux, helping to identify potential bottlenecks in CCM during scale-up [106].

Integrated Modeling: CFD and Biological Models

The most advanced approach for scale-up prediction involves coupling biological models with **Computational Fluid Dynamics (CFD)**.

- **Principle**: CFD models simulate the complex flow, mixing, and mass transfer phenomena inside a large bioreactor. By coupling a CFD model with a kinetic or CBM model, it is possible to predict how spatial variations in the environment will impact the local physiology of cells and the overall fermentation performance [106].

- **Workflow**:
  - A **CFD model** of the industrial bioreactor is developed to map environmental parameters (e.g., substrate, DO) throughout the vessel over time.
  - The resulting data on environmental heterogeneity are fed into a **kinetic or CBM** that describes the cell's physiological response.
  - The integrated model predicts key outcomes like overall product titer and by-product formation, allowing for virtual optimization of the large-scale process before implementation [106].

The following diagram illustrates the logical workflow and integration of these model-based tools for rational scale-up.



Model-Based Scale-Up Prediction Workflow

The Scientist's Toolkit: Key Reagents and Solutions

The following table details essential reagents and materials used in the development and scale-up of fermentation processes, particularly those involving central carbon metabolism engineering.

Table 3: Key Research Reagent Solutions for Fermentation Scale-Up

Reagent/Material	Function	Example Application
Xylose-Responsive Promoters	Genetically control gene expression specifically when xylose is the carbon source.	Enhancing xylose utilization in *S. cerevisiae* by driving expression of xylose isomerase (XI), leading to faster growth [21].
Phosphoketolase (PK) Pathway Enzymes	Provide a heterologous route to convert fructose-6-P and xylulose-5-P directly to acetyl-CoA.	Increasing acetyl-CoA precursor supply for products like fatty acids, farnesene, or 3-HP; can re-route carbon flux in CCM [23].
Biosensors (NADPH, Fatty Acyl-CoA)	Monitor intracellular metabolite levels in real-time, linking metabolic state to a measurable output (e.g., fluorescence).	Dynamic monitoring of redox cofactor status during scale-down experiments to identify metabolic bottlenecks [21].
Scale-Down Bioreactor Systems	Physically simulate the heterogeneous conditions (O_2, substrate gradients) of large-scale production tanks at lab scale.	Pre-validating strain and process robustness by exposing cultures to oscillating environmental conditions [103] [102].

The successful transition of a fermentation process from the laboratory to an industrial plant is a multifaceted challenge that hinges on more than just the intrinsic productivity of the microbial strain. It requires a proactive approach that integrates an understanding of **large-scale bioreactor hydrodynamics** with the **physiology and metabolism** of the production organism. By employing **scale-down methodologies** and **advanced modeling techniques**, researchers can de-risk the scale-up process. Testing strains under simulated industrial conditions and using predictive models allows for the identification and rectification of potential failures early in the development timeline. Ultimately, a rational scale-up strategy that considers the constraints of central carbon metabolism within the complex flow field of a production bioreactor is indispensable for achieving commercially viable biomanufacturing.

Conclusion

Engineering central carbon metabolism is a powerful, multifaceted approach to reprogramming microbial factories for efficient bioproduction. Success hinges on integrating foundational knowledge of metabolic pathways with advanced tools for dynamic control and modular design, while systematically addressing flux imbalances and cofactor requirements through iterative optimization. Future progress will be driven by the development of more efficient enzymes and pathways, the application of AI-driven design, and the creation of electro-microbial hybrid systems. For biomedical research, these advances promise to establish more robust and sustainable platforms for producing complex pharmaceuticals, drug precursors, and other high-value therapeutic compounds, ultimately accelerating the transition to a circular bioeconomy.

References

- 1. Overview of Central Carbon Metabolism [https://metabolomics.creative-proteomics.com/resource/central-carbon-metabolism-analysis-pathways-applications.htm]
- 2. Advances in the optimization of central carbon metabolism in ... [https://microbialcellfactories.biomedcentral.com/articles/10.1186/s12934-023-02090-6]
- 3. Understanding central carbon metabolism of rapidly ... [https://pubmed.ncbi.nlm.nih.gov/26108557/]
- 4. Protein-protein interactions and metabolite channelling in ... [https://www.nature.com/articles/ncomms15212]
- 5. Quantitative modeling of pentose phosphate pathway ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9287814/]
- 6. Time-resolved targeted metabolomics shows an abrupt switch ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12479584/]
- 7. Metabolite-level regulation of enzymatic activity controls ... [https://www.sciencedirect.com/science/article/pii/S0960982224015185]
- 8. Synthetic biology and metabolic engineering paving the way ... [https://pubs.rsc.org/en/content/articlehtml/2025/ya/d5ya00118h]
- 9. Metabolic Engineering Design Strategies for Increasing Acetyl ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC7240943/]
- 10. 6.2: Citric Acid Cycle & Related Pathways [https://bio.libretexts.org/Bookshelves/Biochemistry/Book%3A_Biochemistry_Free_For_All_(Ahern_Rajagopal_and_Tan)/06%3A_Metabol
- 11. Li X, et al. (2025) | SGD [https://yeastgenome.org/reference/S100000585]
- 12. Effective data visualization strategies in untargeted ... [https://pubs.rsc.org/en/content/articlehtml/2025/np/d4np00039k]
- 13. Visualizing metabolic network dynamics through time-series ... [https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-020-3415-z]
- 14. Visualizing regulatory interactions in metabolic networks - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC2216428/]
- 15. Bringing Carbon to Life via One-Carbon Metabolism - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC11972661/]
- 16. Industrial applicability of enzymatic and whole-cell ... [https://www.nature.com/articles/s41467-025-60777-3]
- 17. Advancing Synthetic Biology for Sustainable One-Carbon ... [https://www.sciencedirect.com/science/article/pii/S2950155525000655]
- 18. A blueprint for designing the next-generation of synthetic C ... [https://www.nature.com/articles/s41467-025-64483-y]
- 19. C1 photochemotrophy – rethinking one-carbon metabolism ... [https://www.sciencedirect.com/science/article/abs/pii/S0167779925000034]
- 20. Calvin cycle [https://en.wikipedia.org/wiki/Calvin_cycle]
- 21. Modular deregulation of central carbon metabolism for ... [https://www.nature.com/articles/s41467-025-59966-x]
- 22. The 'photosynthetic C1 pathway' links carbon assimilation ... [https://www.nature.com/articles/s42003-024-07142-0]
- 23. Advances in the optimization of central carbon metabolism in ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC10122336/]
- 24. NAD(H) and NADP(H) Redox Couples and Cellular Energy ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC5737637/]
- 25. Modular deregulation of central carbon metabolism for ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12084563/]
- 26. NAD+ metabolism: pathophysiologic mechanisms and ... [https://www.nature.com/articles/s41392-020-00311-7]
- 27. coordination of redox homeostasis and carbon flux ... [https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2025.1637328/full]
- 28. Nicotinamide adenine dinucleotide [https://en.wikipedia.org/wiki/Nicotinamide_adenine_dinucleotide]
- 29. Optimization of energy production and central carbon ... [https://www.sciencedirect.com/science/article/pii/S0960982223005286]
- 30. Synthetic C1 metabolism in Pseudomonas putida enables ... [https://pubmed.ncbi.nlm.nih.gov/40824035/]
- 31. Engineering the Reductive Glycine Pathway: A Promising ... [https://pubmed.ncbi.nlm.nih.gov/35364693/]
- 32. 2025 Plant Metabolic Engineering Conference GRC [https://www.grc.org/plant-metabolic-engineering-conference/2025/]
- 33. Engineering Genetic Elements for Microbial Protein ... [https://www.sciencedirect.com/science/article/pii/S2405805X2500170X]
- 34. Hierarchical metabolic engineering for rewiring cellular ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12514951/]
- 35. Engineering artificial cross-species promoters with different ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC11366860/]
- 36. CRISPR-Cas tools for simultaneous transcription & ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC11109947/]
- 37. A DNA assembly toolkit to unlock the CRISPR/Cas9 ... [https://www.nature.com/articles/s42003-023-05202-5]
- 38. Pyruvate-responsive genetic circuit biosensors for ... [https://www.sciencedirect.com/science/article/abs/pii/S1385894725116537]
- 39. Systems-Level Modeling for CRISPR-Based Metabolic ... [https://pubmed.ncbi.nlm.nih.gov/39119666/]
- 40. Pyruvate-responsive genetic circuit biosensors for ... [https://www.sciencedirect.com/science/article/pii/S1385894725116537]
- 41. Applications and advances of metabolite biosensors for ... [https://www.sciencedirect.com/science/article/abs/pii/S1096717615000774]
- 42. The Growth Dependent Design Constraints of Transcription ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9994378/]
- 43. Modular and signal-responsive transcriptional regulation ... [https://jbioleng.biomedcentral.com/articles/10.1186/s13036-025-00526-8]
- 44. A quorum sensing-controlled type I CRISPRi toolkit for ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12276013/]
- 45. Central transcriptional regulator controls photosynthetic ... [https://www.nature.com/articles/s41467-024-49090-7]
- 46. Synthetic Biology and Engineering - Volume 3/Issue 2 ... [https://www.sciepublish.com/journals/sbe/roll/3/2]
- 47. A blueprint for designing the next-generation of synthetic C1 ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12494876/]
- 48. Synergistic investigation of natural and synthetic C1-trophic ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC10590403/]
- 49. Paradigm of engineering recalcitrant non-model ... [https://www.nature.com/articles/s41467-024-54897-5]
- 50. Redefining HexR regulatory landscape in Pseudomonas ... [https://www.sciencedirect.com/science/article/abs/pii/S1096717625001879]
- 51. Engineering Noncanonical Cofactors To Expand Cellular ... [https://pubmed.ncbi.nlm.nih.gov/40923492/]
- 52. Metabolic Modelling as a Framework for Metabolomics Data ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC7465778/]

- 53. Flux imbalance analysis and the sensitivity of cellular growth ... [https://pubmed.ncbi.nlm.nih.gov/24009492/]
- 54. Flux Imbalance Analysis and the Sensitivity of Cellular Growth ... [https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1003195]
- 55. Flux-dependent graphs for metabolic networks [https://www.nature.com/articles/s41540-018-0067-y]
- 56. Low-Cost Algorithms for Metabolic Pathway Pairwise ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC8883897/]
- 57. Recent advancements in microbial carbon dioxide fixation [https://pubs.rsc.org/en/content/articlehtml/2025/gc/d5gc03349g]
- 58. Frontiers | Application of cofactors in the regulation of microbial... [https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1145784/full]
- 59. Alleviating Redox Imbalance Enhances... | PLOS One [https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0130840]
- 60. through Engineering systems redox balance cofactor [https://pubmed.ncbi.nlm.nih.gov/24794722/]
- 61. Optimal energy and redox in the cyanobacterium... metabolism [https://www.nature.com/articles/s41540-023-00307-3?error=cookies_not_supported]
- 62. with Phosphite Dehydrogenase... | Newswise Cofactor Engineering [https://www.newswise.com/articles/cofactor-engineering-with-phosphite-dehydrogenase-enables-flexible-regulation-of-lactate-based-copolymer-biosynthesis-in-e-coli]
- 63. Yarrowia lipolytica for the production of β-carotene by... Engineering [https://jbioleng.biomedcentral.com/articles/10.1186/s13036-025-00476-1]
- 64. [PDF] Redox in industrial... | Semantic Scholar cofactor engineering [https://www.semanticscholar.org/paper/Redox-cofactor-engineering-in-industrial-recent-and-Liu-Li/171203714710e47353ef59fb2024f906a487409c]
- 65. Multilevel optimisation of anaerobic ethyl acetate production in ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC7137189/]
- 66. Optimizing the balance between heterologous acetate [https://pmc.ncbi.nlm.nih.gov/articles/PMC10647013/]
- 67. Metabolic engineering strategies for optimizing acetate ... [https://biotechnologyforbiofuels.biomedcentral.com/articles/10.1186/s13068-017-0791-3]
- 68. Unraveling principles of thermodynamics for genome-scale ... [https://www.sciencedirect.com/science/article/abs/pii/S2405471225002261]
- 69. Unraveling the thermodynamic origins and kinetic ... [https://www.sciencedirect.com/science/article/abs/pii/S2590238525001870]
- 70. Light-driven molecular pumps - RSC Publishing [https://pubs.rsc.org/en/content/articlehtml/2025/ob/d5ob01154j]
- 71. Recent advancements in microbial carbon dioxide fixation [https://pubs.rsc.org/en/content/articlelanding/2025/gc/d5gc03349g]
- 72. What is Central Carbon Metabolism? [https://www.metwarebio.com/what-is-central-carbon-metabolism/]
- 73. Multi-Omics in Biological System: Harnessing the Power of ... [https://www.intechopen.com/chapters/1214067]
- 74. Multi-omics data integration using ratio-based quantitative ... [https://www.nature.com/articles/s41587-023-01934-1]
- 75. MiBiOmics: an interactive web application for multi-omics data ... [https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-020-03921-8]
- 76. an interactive visualization and database of multi-omics data [https://pmc.ncbi.nlm.nih.gov/articles/PMC8262747/]
- 77. MINN: A metabolic-informed neural network for integrating ... [https://www.sciencedirect.com/science/article/pii/S2001037025003265]
- 78. Metabolic flux analysis: a comprehensive review on sample ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9449821/]
- 79. What is flux balance analysis? - PMC [https://pmc.ncbi.nlm.nih.gov/articles/PMC3108565/]
- 80. Metabolic flux analysis [https://en.wikipedia.org/wiki/Metabolic_flux_analysis]
- 81. Optimization-based framework with flux balance analysis (FBA ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12578352/]
- 82. Two-stage flux balance analysis of metabolic networks for ... [https://bmcsystbiol.biomedcentral.com/articles/10.1186/1752-0509-5-S1-S11]
- 83. NEXT-FBA: A hybrid stoichiometric/data-driven approach to ... [https://www.sciencedirect.com/science/article/pii/S1096717625000461]
- 84. Editorial: Harmonizing life cycle analysis (LCA) and techno ... [https://www.frontiersin.org/journals/climate/articles/10.3389/fclim.2023.1204840/full]
- 85. Frontiers | Techno - economic of ocean iron fertilization analysis [https://www.frontiersin.org/journals/climate/articles/10.3389/fclim.2025.1509367/full]
- 86. Economic and sustainable revolution to facilitate one- ... [https://www.nature.com/articles/s41467-025-60247-w]
- 87. Comparative Techno-Economic and Carbon Footprint ... [https://www.mdpi.com/2077-0472/15/5/472]
- 88. - Techno of indirect carbonation processes for... economic analysis [https://pubs.rsc.org/en/content/articlelanding/2025/ya/d4ya00567h]
- 89. Circular Carbon Capture: Comparative Life Cycle and... [https://pubmed.ncbi.nlm.nih.gov/41275387/]
- 90. Broad-Host-Range Synthetic Biology: Rethinking Microbial ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC12538584/]
- 91. Promising non-model microbial cell factories obtained by ... [https://www.frontiersin.org/journals/bioengineering-and-biotechnology/articles/10.3389/fbioe.2024.1427248/full]
- 92. Absolute Quantification of the Central Carbon Metabolome ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC7073941/]
- 93. Modeling central metabolism and energy biosynthesis across ... [https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-016-2887-8]
- 94. Advanced metabolic engineering strategies for increasing ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC10898619/]
- 95. Diversifying Isoprenoid Platforms via Atypical Carbon ... [https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2021.791089/full]
- 96. Compartmentalized Metabolic Engineering for Artemisinin ... [https://www.sciencedirect.com/science/article/pii/S1674205216302222]

- 97. High level production of amorphadiene using Bacillus ... [https://www.sciencedirect.com/science/article/pii/S1871678420301850]
- 98. IspH inhibitors kill Gram-negative bacteria and mobilize ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC8776033/]
- 99. Systematic biotechnological production of isoprenoid analogs ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC11873216/]
- 100. Engineering 3-Hydroxypropionic Acid Production from ... [https://www.mdpi.com/2309-608X/9/5/573]
- 101. Benchmarking recombinant Pichia pastoris for 3 ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC8313290/]
- 102. Understanding the scale-up of fermentation processes from ... [https://www.sciencedirect.com/science/article/abs/pii/S1004954120306984]
- 103. How to scale fermentation processes from lab to factory [https://www.foodnavigator.com/Article/2025/06/17/how-to-scale-fermentation-processes-from-lab-to-factory/]
- 104. Recent advances in engineering the central carbon ... [https://microbialcellfactories.biomedcentral.com/articles/10.1186/1475-2859-11-50]
- 105. Scaling Up: From Laboratory to Industrial-Scale Fermentation [https://www.linkedin.com/pulse/scaling-up-from-laboratory-industrial-scale-fermentation-44q2c]
- 106. Optimization and Scale-Up of Fermentation Processes Driven ... [https://pmc.ncbi.nlm.nih.gov/articles/PMC9495923/]