

Lab Report: Characterization and Analysis of Oil-Based Mixtures

Introduction

In this report, we detail the testing and analysis of various oil-based mixtures using advanced laboratory instruments. Each mixture was tested to assess its properties and composition. The carried-out experiments include conductivity measurements, chromatographic analysis, optical density readings, and viscosity evaluations, among others. Below is a comprehensive presentation of our findings.

Materials and Methods

The samples were meticulously prepared, incorporating different oils with additives such as gums and vitamins. The procedures were conducted using the following instruments:

Table 1: Instrumentation and Sample Data

	Instrument	Primary Component	Secondary Component	Additive 1	Additive 2	Measurement	Unit
Conductivity Meter	CM-215	Jobba Oil	Gum	nan	nan	1450.0	uS/cm
Gas Chromatograph	GC-2010	Coconut Oil	Gum	Glycerin	nan	350.0	ppm
UV-Vis Spectrophotometer	UV-2600	Jobba Oil	Vitamin E	nan	nan	1.8	Abs
HPLC System	HPLC-9000	Jobba Oil	nan	nan	nan	0.45	mg/L
Microplate Reader	MRX	Coconut Oil	Cetyl Alcohol	Vitamin E	nan	2.3	OD
Viscometer	VS-300	Almond Oil	Beeswax	Vitamin E	nan	7000.12	cP
Viscometer	VS-300	Almond Oil	Cetyl Alcohol	nan	nan	7218.31	cP

Observations

Conductivity Analysis:The presence of gummy substances in the Jojoba Oil solution resulted in significant ionic content, indicated by the conductivity value of 1450 uS/cm. High ionic strength might suggest dissolution of salts or similar components which could be impurities or intentional additives.

Chromatographic Behavior:Coconut Oil mixed with Gum and Glycerin displayed a peak concentration of aromatic

compounds at 350 ppm using Gas Chromatograph GC-2010. This reading implies a moderate level of glycerol esters, potentially affecting the hydrophobicity of the mixture.

Optical Characteristics and Absorption:As seen with the UV-Vis Spectrophotometer UV-2600, the inclusion of Vitamin E in Jojoba Oil marked an absorption peak of 1.8 Abs, indicative of the broad absorption spectrum of tocopherols.

Viscosity Insights:Almond Oil combined with Beeswax and Vitamin E measured a notable viscosity of 7000.12 cP. This formulation is considerably thicker compared to the Almond Oil and Cetyl Alcohol mixture, which registered at 7218.31 cP. The presence of cetyl alcohol typically enhances fluidity due to its lubricating properties.

Table 2: Summary of Results

Sample	Ionic Strength (uS/cm)	Aromatic Compounds (ppm)	Absorption (Abs)	Viscosity (cP)	Observations
Jojoba Oil + Gum	1450.0	nan	nan	nan	High ionic content
Coconut Oil + Gum + Glycerin	nan	350.0	nan	nan	Moderate level of glycerol esters
Jojoba Oil + Vitamin E	nan	nan	1.8	nan	Tocopherol absorption spectrum
Almond Oil + Beeswax + Vitamin E	nan	nan	nan	7000.12	High viscosity due to wax additives
Almond Oil + Cetyl Alcohol	nan	nan	nan	7218.31	Enhanced fluidity with cetyl alcohol

Conclusion

In drawing the final analysis, each mixture of oil and supplementary materials exhibited distinct physical and chemical properties, profoundly influenced by its constituents. While some displayed notable ionic and aromatic features, others highlighted unique viscosity profiles. These findings underscore the necessity of careful formulation to achieve desired material characteristics for specific applications. Further studies might explore the implications of these properties on broader industrial and cosmetic applications.

Random Note: Despite the occasional deviation in measurements due to environmental variations, the results are largely consistent, reflecting the inherent stability of the samples under investigation.