

F-test Summary Tables

2024 transfer-function polynomial scan

Central and forward categories

Candidates: 0x0, 0x1, 0x2, 1x0, 1x1, 1x2, 2x1, 2x2

Numerical values are reproduced directly from the central and forward F-test CSV summaries.

CMS Work in Progress

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
0x0	0x1	1	2	165	6.319	0.0129	p < 0.05
0x0	0x2	1	3	165	0	1.0000	not significant
0x0	1x0	1	2	165	11.79	7.55e-04	p < 0.05
0x0	1x1	1	3	165	0	1.0000	not significant
0x0	1x2	1	4	165	0.6624	0.5763	not significant
0x0	2x1	1	4	165	0	1.0000	not significant
0x0	2x2	1	5	165	9.535	6.12e-07	p < 0.05
0x1	0x2	2	3	165	0	1.0000	not significant
0x1	1x1	2	3	165	0	1.0000	not significant
0x1	1x2	2	4	165	0	1.0000	not significant
0x1	2x1	2	4	165	0	1.0000	not significant
0x1	2x2	2	5	165	10.25	3.28e-06	p < 0.05

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
1x0	0x2	2	3	165	0	1.0000	not significant
0x2	1x2	3	4	165	184.7	< numerical precision	p < 0.05
0x2	2x1	3	4	165	0	1.0000	not significant
0x2	2x2	3	5	165	130.1	< numerical precision	p < 0.05
1x0	1x1	2	3	165	0	1.0000	not significant
1x0	1x2	2	4	165	0	1.0000	not significant
1x0	2x1	2	4	165	0	1.0000	not significant
1x0	2x2	2	5	165	8.258	3.84e-05	p < 0.05
1x1	1x2	3	4	165	63.17	5.03e-13	p < 0.05
1x1	2x1	3	4	165	0	1.0000	not significant
1x1	2x2	3	5	165	56.26	< numerical precision	p < 0.05
1x2	2x2	4	5	165	35.72	1.43e-08	p < 0.05

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
2x1	2x2	4	5	165	511.9	< numerical precision	p < 0.05

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
0x0	0x1	1	2	165	0.1779	0.6737	not significant
0x0	0x2	1	3	165	0	1.0000	not significant
0x0	1x0	1	2	165	0	1.0000	not significant
0x0	1x1	1	3	165	0	1.0000	not significant
0x0	1x2	1	4	165	0	1.0000	not significant
0x0	2x1	1	4	165	3.228	0.0241	p < 0.05
0x0	2x2	1	5	165	0	1.0000	not significant
0x1	0x2	2	3	165	0	1.0000	not significant
0x1	1x1	2	3	165	0	1.0000	not significant
0x1	1x2	2	4	165	0	1.0000	not significant
0x1	2x1	2	4	165	4.748	0.0099	p < 0.05
0x1	2x2	2	5	165	0	1.0000	not significant

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
1x0	0x2	2	3	165	0	1.0000	not significant
0x2	1x2	3	4	165	0	1.0000	not significant
0x2	2x1	3	4	165	315.5	< numerical precision	p < 0.05
0x2	2x2	3	5	165	0	1.0000	not significant
1x0	1x1	2	3	165	0	1.0000	not significant
1x0	1x2	2	4	165	0	1.0000	not significant
1x0	2x1	2	4	165	157.7	3.70e-14	p < 0.05
1x0	2x2	2	5	165	0	1.0000	not significant
1x1	1x2	3	4	165	0	1.0000	not significant
1x1	2x1	3	4	165	327.6	< numerical precision	p < 0.05
1x1	2x2	3	5	165	0	1.0000	not significant
1x2	2x2	4	5	165	17.8	4.10e-05	p < 0.05

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.

Simple	Complex	p1	p2	Bins	F statistic	p-value	Pairwise result
2x1	2x2	4	5	165	0	1.0000	not significant

Red rows have $p < 0.05$. This is a pairwise statistical comparison, not by itself a final model-selection rule.