

# Parabola Question

EE24BTECH11040 - Mandara Hosur

## Question:

Given that the length of the major axis of a conic is 16 units and the coordinates of the foci are  $(0, \pm 6)$ , find the equation of the conic.

## Solution:

2 foci exist for this conic. Therefore, it must be an ellipse or a hyperbola.

The general equation of a conic with directrix  $\mathbf{n}^\top \mathbf{x} = c$  can be written as

$$g(\mathbf{x}) = \mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f$$

Let focus  $\mathbf{F}_1$  be  $(0, 6)$  and focus  $\mathbf{F}_2$  be  $(0, -6)$

$$\mathbf{u} = \frac{\mathbf{F}_1 + \mathbf{F}_2}{2} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ and } \mathbf{n} \equiv \mathbf{F}_1 - \mathbf{F}_2 \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Given that  $e$  represents eccentricity of the conic,  $\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 1 - e^2 \end{pmatrix}$

$$\text{Length of major axis} = 16 = 2\sqrt{\frac{|f|}{|1 - e^2|}}$$

Rearranging this equation gives

$$8\sqrt{|1 - e^2|} = \sqrt{|f|} \quad (0.1)$$

$$\text{Now, } \mathbf{F} = \frac{ce^2 \mathbf{n} - \mathbf{u}}{\lambda_2} \text{ and } \lambda_2 = \|\mathbf{n}\|^2.$$

$$\Rightarrow \mathbf{F} = \frac{ce^2 \mathbf{n} - \mathbf{u}}{\|\mathbf{n}\|^2}$$

Substituting values of  $\mathbf{F}_1$ ,  $\mathbf{F}_2$ ,  $\mathbf{n}$ , and  $\mathbf{u}$  and simplifying reduces the above equation to

$$\pm ce^2 = 6 \quad (0.2)$$

Consider another equation

$$c = \frac{e\mathbf{u}^\top \mathbf{n} \pm \sqrt{e^2 (\mathbf{u}^\top \mathbf{n})^2 - \lambda_2 (e^2 - 1) (\|\mathbf{u}\|^2 - \lambda_2 f)}}{\lambda_2 e (e^2 - 1)}$$

Substituting known values gives

$$c = \pm \frac{1}{e} \sqrt{\frac{|f|}{e^2 - 1}} \quad (0.3)$$

We have 3 equations and 3 variables ( $c$ ,  $f$ ,  $e$ ).

Upon solving, we get  $e = \frac{3}{4}$ ,  $c = \pm \frac{32}{3}$ , and  $|f| = 28$

However,  $f$  can only have one value (either  $+28$  or  $-28$ ).

To find this, assume some point  $(0, \alpha)$  to lie on the major axis of the conic. Substituting this point in the general conic equation  $g(\mathbf{x}) = 0$  along with values of  $\mathbf{V}$  and  $\mathbf{u}$  gives

$$\frac{7\alpha^2}{16} + f = 0$$

This implies that  $f$  must be negative, and therefore  $f = -28$ .

The equation of the conic thus becomes

$$g(\mathbf{x}) = \mathbf{x}^\top \begin{pmatrix} 1 & 0 \\ 0 & \frac{7}{16} \end{pmatrix} \mathbf{x} - 28$$




Fig. 0.1: Plot of conic