

# Computational Methods of Optimization

## First Midterm(7th Dec, 2020)

Time: 60 minutes

### **Instructions**

- Answer all questions
- See upload instructions in the form

In the following, assume that  $f$  is a  $\mathcal{C}^1$  function defined from  $\mathbb{R}^d \rightarrow \mathbb{R}$  unless otherwise mentioned. Let  $\mathbf{I} = [e_1, \dots, e_d]$  be a  $d \times d$  matrix with  $e_j$  be the  $j$ th column. Also  $\mathbf{x} = [x_1, x_2, \dots, x_d]^\top \in \mathbb{R}^d$  and  $\|\mathbf{x}\| = \sqrt{\mathbf{x}^\top \mathbf{x}}$ . Set of real symmetric  $d \times d$  matrices will be denoted by  $\mathcal{S}_d$ .  $[n]$  will denote the set  $\{1, 2, \dots, n\}$

1. (10 points) Please indicate True(T) or False(F) in the space given after each question. All questions carry equal marks

- (a) Let  $a < b$  where  $a, b \in \mathbb{R}$  and  $h : [a, b] \rightarrow \mathbb{R}$  be differentiable and satisfies  $h(a) = h(b)$ . Then  $h$  has a critical point in  $(a, b)$ . \_\_\_\_
- (b) Suppose the function defined in the previous question satisfies  $|h(x) - h(y)| \leq 1|x - y|$  for all  $x, y \in (a, b)$ . There could exist a point in  $(a, b)$  such that  $h'(x) \geq 2$ , where  $h'(x)$  is derivative of  $h$  at  $x$ . \_\_\_\_
- (c) If  $f$  is a coercive function then the global minimum must lie at one of the critical points. Recall that a critical point is a point,  $\mathbf{x}$ , such that  $\nabla f(\mathbf{x}) = 0$ . \_\_\_\_
- (d) Consider  $g : \mathbb{R} \rightarrow \mathbb{R}, g(u) = u^2 - \frac{1}{3}u^3$ . The function has a global minimum. \_\_\_\_
- (e) The local maximum of  $g$  (defined in the previous question) is at  $u = 0$ . \_\_\_\_

2. Let  $f : S \subset \mathbb{R}^d \rightarrow \mathbb{R} \in \mathcal{C}^2$  function. Let  $H(\mathbf{x})$  be the Hessian of  $f$  with eigenvectors denoted by columns of  $U \in \mathbb{R}^{d \times d}$

- (a) (2 points) Consider  $S = \{\mathbf{x} | \mathbf{x} = \mathbf{x}^* + \mathbf{U}\mathbf{v}, \mathbf{v} \in \mathbb{R}^d\}$ .  $\bar{S}$ , the complement of the set  $S$ , is not empty. True or False. Justify with reasons.

- (b) (4 points) The Hessian,  $H(\mathbf{x}^*)$  at a stationary point  $\mathbf{x}^*$  has one eigenvalue 0 and the rest positive. For any  $\mathbf{x} \in S$  with  $\mathbf{v} \neq 0$   $f(\mathbf{x}) > f(\mathbf{x}^*)$  is true. Prove or disprove

- (c) (4 points) Suppose  $H(\mathbf{x}^*)$  has negative and positive eigenvalues. Construct two distinct points,  $\mathbf{x}^1$  and  $\mathbf{x}^2$  in terms of  $\mathbf{U}$  so that

$$f(\mathbf{x}^1) < f(\mathbf{x}) < f(\mathbf{x}^2)$$

3. Let  $f(x) = \sqrt{1 + x^2}, x \in \mathbb{R}$ .

- (a) (2 points) Find the  $x^*$ , global minimum of the problem

- (b) (4 points) Let  $x^{(k)}$  be the output of the  $k$ th iteration of Newton's method applied to  $f(x)$ . Find a function  $g$  such that

$$|x^{(k+1)} - x^*| \leq |g(x^{(k)} - x^*)|$$

- (c) (4 points) Using the above relationship find largest  $a$  so that the Newton's method is most effective for any  $x^{(0)} \in (x^* - a, x^* + a)$ .

4. Consider minimizing the function

$$f(\mathbf{x}) = \frac{1}{2} \mathbf{x}^\top Q \mathbf{x} - \mathbf{b}^\top \mathbf{x}$$

over  $\mathbf{x} \in \mathbb{R}^d$  with  $Q \in \mathcal{S}_d^+$ ,  $\mathbf{b} \in \mathbb{R}^d$  using the steepest descent iterates,

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} - \alpha_k \nabla f(\mathbf{x}^{(k)})$$

executed with exact step-size selection strategy. Let  $f^* = f(\mathbf{x}^*)$  be the global minimum. Let  $g(\mathbf{y}) = f(A\mathbf{x})$  where  $A$  is a  $d \times d$  matrix. Consider applying steepest descent procedure to  $g$  i.e.

$$\mathbf{y}^{(k+1)} = \mathbf{y}^{(k)} - \alpha_k \nabla g(\mathbf{y}^{(k)})$$

and  $g^*$  is the global minimum of  $g$  attained at  $\mathbf{y}^*$ .

(a) (4 points) State the Hessian of  $f$  and  $g$ . Derive the relationship between  $\mathbf{x}^*$  and  $\mathbf{y}^*$

(b) (3 points) What is the convergence rate of the steepest descent procedure for  $g$ ?

(c) (3 points) What is the best value of  $A$ ?

5. Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a  $C^{(2)}$  function. Consider the Quasi-newton update

**a.**  $s^{(k)} = -G^{(k)} \nabla f(\mathbf{x}^{(k)})$ ,

**b.**  $\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k s^{(k)}$

**c.**  $G^{(k+1)} = G^{(k)} + A^{(k)} E A^{(k)\top}$

where  $\gamma_k = \nabla f(\mathbf{x}^{(k+1)}) - \nabla f(\mathbf{x}^{(k)})$ ,  $\delta_k = \mathbf{x}^{(k+1)} - \mathbf{x}^{(k)}$ , and  $A^{(k)} = [\delta_k \quad G^{(k)} \gamma^{(k)}]$  and  $E$  is a matrix.

(a) (1 point) What is your name

(b) (4 points) Find  $E$  corresponding to DFP updates.

(c) (5 points) Show that for exact line search DFP updates yield positive definite matrices.