

E1 222 Stochastic Models and Applications

P.S. Sastry
sastry@iisc.ac.in

Reference Material

- ▶ V.K. Rohatgi and A.K.Md.E. Saleh, An Introduction to probability and Statistics, Wiley, 2nd edition, 2018
- ▶ S.Ross, 'Introduction to Probability Models', Elsevier, 12th edition, 2019.

Reference Material

- ▶ V.K. Rohatgi and A.K.Md.E. Saleh, An Introduction to probability and Statistics, Wiley, 2nd edition, 2018
- ▶ S.Ross, 'Introduction to Probability Models', Elsevier, 12th edition, 2019.
- ▶ P G Hoel, S Port and C Stone, Introduction to Probability Theory, 1971.
- ▶ P G Hoel, S Port and C Stone, Introduction to Stochastic Processes, 1971.

Course Prerequisites / Background needed

- ▶ Calculus
 - ▶ continuity, differentiability, derivatives, functions of several variables, partial derivatives, integration, multiple integrals or integration over \mathbb{R}^n , convergence of sequences and series, Taylor series

Course Prerequisites / Background needed

- ▶ Calculus
 - ▶ continuity, differentiability, derivatives, functions of several variables, partial derivatives, integration, multiple integrals or integration over \mathbb{R}^n , convergence of sequences and series, Taylor series
- ▶ Matrix theory
 - ▶ vector spaces, linear independence, linear transformations, matrices, rank, determinant, eigen values and eigen vectors

Course Prerequisites / Background needed

- ▶ Calculus
 - ▶ continuity, differentiability, derivatives, functions of several variables, partial derivatives, integration, multiple integrals or integration over \mathbb{R}^n , convergence of sequences and series, Taylor series
- ▶ Matrix theory
 - ▶ vector spaces, linear independence, linear transformations, matrices, rank, determinant, eigen values and eigen vectors
- ▶ In addition, knowledge of basic probability is assumed. I assume all students are familiar with the following:
Random experiment, sample space, events, conditional probability, independent events, simple combinatorial probability computations

Course Prerequisites / Background needed

- ▶ Calculus
 - ▶ continuity, differentiability, derivatives, functions of several variables, partial derivatives, integration, multiple integrals or integration over \mathbb{R}^n , convergence of sequences and series, Taylor series
- ▶ Matrix theory
 - ▶ vector spaces, linear independence, linear transformations, matrices, rank, determinant, eigen values and eigen vectors
- ▶ In addition, knowledge of basic probability is assumed. I assume all students are familiar with the following:
Random experiment, sample space, events, conditional probability, independent events, simple combinatorial probability computations

But we would review the basic probability in the first two classes.

Course grading

- ▶ Mid-Term Tests and Assignments: 70%
Final Exam: 30%
- ▶ Three mid-term tests for 20 marks each. We will have 2-3 assignments for 10 marks. (Tentative)

Course grading

- ▶ Mid-Term Tests and Assignments: 70%
Final Exam: 30%
- ▶ Three mid-term tests for 20 marks each. We will have 2-3 assignments for 10 marks. (Tentative)
- ▶ Please remember this is essentially a Maths course

Probability Theory

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena – “individually not predictable but have a lot of regularity at a population level”

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena – “individually not predictable but have a lot of regularity at a population level”
- ▶ E.g., Recommender systems are useful for Amazon or Netflix because at a population level customer behaviour can be predicted well.

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena – “individually not predictable but have a lot of regularity at a population level”
- ▶ E.g., Recommender systems are useful for Amazon or Netflix because at a population level customer behaviour can be predicted well.
- ▶ Example random phenomena: Tossing a coin, rolling a dice etc – familiar to you all

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena – “individually not predictable but have a lot of regularity at a population level”
- ▶ E.g., Recommender systems are useful for Amazon or Netflix because at a population level customer behaviour can be predicted well.
- ▶ Example random phenomena: Tossing a coin, rolling a dice etc – familiar to you all
- ▶ It is also useful in many engineering systems, e.g., for taking care of noise.

Probability Theory

- ▶ Probability Theory – branch of mathematics that deals with modeling and analysis of random phenomena.
- ▶ Random Phenomena – “individually not predictable but have a lot of regularity at a population level”
- ▶ E.g., Recommender systems are useful for Amazon or Netflix because at a population level customer behaviour can be predicted well.
- ▶ Example random phenomena: Tossing a coin, rolling a dice etc – familiar to you all
- ▶ It is also useful in many engineering systems, e.g., for taking care of noise.
- ▶ Probability theory is also needed for Statistics that deals with making inferences from data.

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)
 - ▶ Policies for decision making under uncertainty

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)
 - ▶ Policies for decision making under uncertainty
 - ▶ Pattern Recognition, prediction from data
 - ▶

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)
 - ▶ Policies for decision making under uncertainty
 - ▶ Pattern Recognition, prediction from data
 - ▶
- ▶ We may use probability models for analysing algorithms. (e.g., average case complexity of algorithms)

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)
 - ▶ Policies for decision making under uncertainty
 - ▶ Pattern Recognition, prediction from data
 - ▶
- ▶ We may use probability models for analysing algorithms. (e.g., average case complexity of algorithms)
- ▶ We may deliberately introduce randomness in an algorithm (e.g., ALOHA protocol, Primality testing)
- ▶

- ▶ In many engineering problems one needs to deal with random inputs where probability models are useful
 - ▶ Dealing with dynamical systems subjected to noise (e.g., Kalman filter)
 - ▶ Policies for decision making under uncertainty
 - ▶ Pattern Recognition, prediction from data
 - ▶
- ▶ We may use probability models for analysing algorithms. (e.g., average case complexity of algorithms)
- ▶ We may deliberately introduce randomness in an algorithm (e.g., ALOHA protocol, Primality testing)
- ▶

This is only a 'sample' of possible application scenarios!

Review of basic probability

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

We use the following Notation:

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

We use the following Notation:

- ▶ Sample space – Ω

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

We use the following Notation:

- ▶ Sample space – Ω

Elements of Ω are the outcomes of the random experiment

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

We use the following Notation:

► Sample space – Ω

Elements of Ω are the outcomes of the random experiment

We write $\Omega = \{\omega_1, \omega_2, \dots\}$ when it is countable

Review of basic probability

We assume all of you are familiar with the terms:

*random experiment, outcomes of random experiment,
sample space, events etc.*

We use the following Notation:

- ▶ Sample space – Ω

Elements of Ω are the outcomes of the random experiment

We write $\Omega = \{\omega_1, \omega_2, \dots\}$ when it is countable

- ▶ An event is, by definition, a subset of Ω

Review of basic probability

We assume all of you are familiar with the terms:

random experiment, outcomes of random experiment, sample space, events etc.

We use the following Notation:

- ▶ Sample space – Ω
Elements of Ω are the outcomes of the random experiment
We write $\Omega = \{\omega_1, \omega_2, \dots\}$ when it is countable
- ▶ An event is, by definition, a subset of Ω
- ▶ Set of all possible events – $\mathcal{F} \subset 2^\Omega$ (power set of Ω)
Each event is a subset of Ω

Review of basic probability

We assume all of you are familiar with the terms:

random experiment, outcomes of random experiment, sample space, events etc.

We use the following Notation:

- ▶ Sample space – Ω

Elements of Ω are the outcomes of the random experiment

We write $\Omega = \{\omega_1, \omega_2, \dots\}$ when it is countable

- ▶ An event is, by definition, a subset of Ω
- ▶ Set of all possible events – $\mathcal{F} \subset 2^\Omega$ (power set of Ω)

Each event is a subset of Ω

For now, we take $\mathcal{F} = 2^\Omega$ (power set of Ω)

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathbb{R}$ satisfying

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathbb{R}$ satisfying

A1 Non-negativity: $P(A) \geq 0, \forall A \in \mathcal{F}$

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathbb{R}$ satisfying

A1 Non-negativity: $P(A) \geq 0, \forall A \in \mathcal{F}$

A2 Normalization: $P(\Omega) = 1$

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathfrak{R}$ satisfying

A1 Non-negativity: $P(A) \geq 0, \forall A \in \mathcal{F}$

A2 Normalization: $P(\Omega) = 1$

A3 σ -additivity: If $A_1, A_2, \dots \in \mathcal{F}$ satisfy $A_i \cap A_j = \emptyset, \forall i \neq j$ then

$$P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$$

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathfrak{R}$ satisfying

A1 Non-negativity: $P(A) \geq 0, \forall A \in \mathcal{F}$

A2 Normalization: $P(\Omega) = 1$

A3 σ -additivity: If $A_1, A_2, \dots \in \mathcal{F}$ satisfy $A_i \cap A_j = \phi, \forall i \neq j$ then

$$P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$$

Events satisfying $A_i \cap A_j = \phi, \forall i \neq j$ are said to be **mutually exclusive**

Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

Formally, $P : \mathcal{F} \rightarrow \mathfrak{R}$ satisfying

A1 Non-negativity: $P(A) \geq 0, \forall A \in \mathcal{F}$

A2 Normalization: $P(\Omega) = 1$

A3 σ -additivity: If $A_1, A_2, \dots \in \mathcal{F}$ satisfy $A_i \cap A_j = \phi, \forall i \neq j$ then

$$P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$$

Events satisfying $A_i \cap A_j = \phi, \forall i \neq j$ are said to be **mutually exclusive**

(Ω, \mathcal{F}, P) is called the **Probability Space**

Probability axioms

$P : \mathcal{F} \rightarrow \mathbb{R}$, $\mathcal{F} \subset 2^\Omega$ (Events are subsets of Ω)

Probability axioms

$P : \mathcal{F} \rightarrow \mathbb{R}$, $\mathcal{F} \subset 2^\Omega$ (Events are subsets of Ω)

A1 $P(A) \geq 0$, $\forall A \in \mathcal{F}$

A2 $P(\Omega) = 1$

A3 If $A_i \cap A_j = \emptyset$, $\forall i \neq j$ then $P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$

Probability axioms

$P : \mathcal{F} \rightarrow \mathbb{R}$, $\mathcal{F} \subset 2^\Omega$ (Events are subsets of Ω)

A1 $P(A) \geq 0$, $\forall A \in \mathcal{F}$

A2 $P(\Omega) = 1$

A3 If $A_i \cap A_j = \emptyset$, $\forall i \neq j$ then $P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$

► For these axioms to make sense, we are assuming

(i). $\Omega \in \mathcal{F}$ and (ii). $A_1, A_2, \dots \in \mathcal{F} \Rightarrow (\cup_i A_i) \in \mathcal{F}$

When $\mathcal{F} = 2^\Omega$ this is true.

Probability axioms

$P : \mathcal{F} \rightarrow \mathbb{R}$, $\mathcal{F} \subset 2^\Omega$ (Events are subsets of Ω)

A1 $P(A) \geq 0$, $\forall A \in \mathcal{F}$

A2 $P(\Omega) = 1$

A3 If $A_i \cap A_j = \emptyset$, $\forall i \neq j$ then $P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$

► For these axioms to make sense, we are assuming

(i). $\Omega \in \mathcal{F}$ and (ii). $A_1, A_2, \dots \in \mathcal{F} \Rightarrow (\cup_i A_i) \in \mathcal{F}$

When $\mathcal{F} = 2^\Omega$ this is true.

► Note that axiom A3 is about countable unions.

Probability axioms

$P : \mathcal{F} \rightarrow \mathfrak{R}$, $\mathcal{F} \subset 2^\Omega$ (Events are subsets of Ω)

A1 $P(A) \geq 0$, $\forall A \in \mathcal{F}$

A2 $P(\Omega) = 1$

A3 If $A_i \cap A_j = \emptyset$, $\forall i \neq j$ then $P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$

► For these axioms to make sense, we are assuming

(i). $\Omega \in \mathcal{F}$ and (ii). $A_1, A_2, \dots \in \mathcal{F} \Rightarrow (\cup_i A_i) \in \mathcal{F}$

When $\mathcal{F} = 2^\Omega$ this is true.

► Note that axiom A3 is about countable unions.

► As defined, the co-domain of the function P is \mathfrak{R} .

However, the axioms imply that it takes values in $[0, 1]$.

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A))$$

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A)) = P(A) + P(B - A)$$

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A)) = P(A) + P(B - A) \geq P(A)$$

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A)) = P(A) + P(B - A) \geq P(A)$$

This also shows $P(B - A) = P(B) - P(A)$ when $A \subset B$.

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A)) = P(A) + P(B - A) \geq P(A)$$

This also shows $P(B - A) = P(B) - P(A)$ when $A \subset B$.

- ▶ If A is an event, then $A \subset \Omega$ and hence
 $P(A) \leq P(\Omega) = 1$.

Simple consequences of the axioms

- ▶ Notation: A^c is complement of A .
 $C = A + B$ implies A, B are mutually exclusive and C is their union.
- ▶ Let $A \subset B$ be events. Then $B = A + (B - A)$.
Now we can show $P(A) \leq P(B)$:

$$P(B) = P(A + (B - A)) = P(A) + P(B - A) \geq P(A)$$

This also shows $P(B - A) = P(B) - P(A)$ when $A \subset B$.

- ▶ If A is an event, then $A \subset \Omega$ and hence $P(A) \leq P(\Omega) = 1$.
- ▶ We can show $P(A^c) = 1 - P(A)$ as

$$1 = P(\Omega) = P(A + A^c) = P(A) + P(A^c)$$

- ▶ There are many such properties (I assume familiar to you) that can be derived from the axioms.

- ▶ There are many such properties (I assume familiar to you) that can be derived from the axioms.
- ▶ Here are a few important ones. (Proof is left to you as an exercise!)

- ▶ There are many such properties (I assume familiar to you) that can be derived from the axioms.
- ▶ Here are a few important ones. (Proof is left to you as an exercise!)

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- ▶ There are many such properties (I assume familiar to you) that can be derived from the axioms.
- ▶ Here are a few important ones. (Proof is left to you as an exercise!)

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\begin{aligned}
 P(U_{i=1}^n A_i) &= \sum_i P(A_i) - \sum_i \sum_{j>i} P(A_i \cap A_j) \\
 &+ \sum_i \sum_{j>i} \sum_{k>j} P(A_i \cap A_j \cap A_k) - \cdots + (-1)^{n+1} P(\cap_i A_i)
 \end{aligned}$$

Known as inclusion-exclusion formula

Case of finite Ω – Example

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$.

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\})$$

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\})$$

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event, A .

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event, A .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event, A .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.
- ▶ Thus, ‘equally likely’ is one way of specifying the probability function (in case of finite Ω).

Case of finite Ω – Example

- ▶ Let $\Omega = \{\omega_1, \dots, \omega_n\}$, $\mathcal{F} = 2^\Omega$, and P is specified through ‘equally likely’ assumption.
- ▶ That is, $P(\{\omega_i\}) = \frac{1}{n}$. (Note the notation)
- ▶ Suppose $A = \{\omega_1, \omega_2, \omega_3\}$. Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event, A .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.
- ▶ Thus, ‘equally likely’ is one way of specifying the probability function (in case of finite Ω).
- ▶ An obvious point worth remembering: specifying P for singleton events fixes it for all other events.