

## E2 212: Homework - 7

### 1 Topics

- Variational characterization of eigenvalues

### 2 Problems

1. Let  $\mathbf{A} \in \mathbb{C}^{n \times n}$  be a Hermitian matrix, let  $\mathbf{x} \in \mathbb{C}^n$  be a given nonzero vector, and let  $\alpha \triangleq \mathbf{x}^H \mathbf{A} \mathbf{x} / \mathbf{x}^H \mathbf{x}$ . Show that there exists at least one eigenvalue of  $\mathbf{A}$  in the interval  $(-\infty, \alpha]$  and at least one in  $[\alpha, \infty)$ .
2. If  $\mathbf{A} \in \mathbb{C}^{n \times n}$  be a Hermitian matrix. Show that the following three optimization problems all have the same solution:

(a)  $\max_{\mathbf{x}^H \mathbf{x}=1} \mathbf{x}^H \mathbf{A} \mathbf{x}$

(b)  $\max_{\mathbf{x} \neq 0} \frac{\mathbf{x}^H \mathbf{A} \mathbf{x}}{\mathbf{x}^H \mathbf{x}}$

(c)  $\min_{\mathbf{x}^H \mathbf{A} \mathbf{x}=1} \mathbf{x}^H \mathbf{x}$

3. Show that, if  $\lambda_i$  is any eigenvalue of  $\mathbf{A} \in \mathbb{C}^{n \times n}$  (not necessarily Hermitian), then, one has the bounds

$$\min_{\mathbf{x} \neq 0} \left| \frac{\mathbf{x}^H \mathbf{A} \mathbf{x}}{\mathbf{x}^H \mathbf{x}} \right| \leq |\lambda_i| \leq \max_{\mathbf{x} \neq 0} \left| \frac{\mathbf{x}^H \mathbf{A} \mathbf{x}}{\mathbf{x}^H \mathbf{x}} \right|, i = 1, 2, \dots, n. \quad (1)$$

4. If  $\mathbf{A} \in \mathbb{C}^{n \times n}$  is Hermitian and if  $\mathbf{x}^H \mathbf{A} \mathbf{x} \geq 0$  for all  $\mathbf{x}$  in a  $k$ -dimensional subspace, then prove that  $\mathbf{A}$  has at least  $k$  nonnegative eigenvalues.
5. Let  $\mathbf{A} \in \mathbb{C}^{n \times n}$  and  $\mathbf{B} \in \mathbb{C}^{n \times n}$  be two Hermitian matrices. Prove that

$$|\lambda_k(\mathbf{A} + \mathbf{B}) - \lambda_k(\mathbf{A})| \leq \rho(\mathbf{B})$$

for all  $k = 1, 2, \dots, n$ , where  $\rho(\mathbf{B})$  is the spectral radius of  $\mathbf{B}$ , and where the eigenvalues of  $\mathbf{A}$  and  $\mathbf{A} + \mathbf{B}$  are arranged, as usual, in increasing order.

6. If  $\mathbf{A} \in \mathbb{C}^{n \times n}$  and  $\mathbf{B} \in \mathbb{C}^{n \times n}$  are Hermitian matrices with eigenvalues arranged in increasing order, and if  $1 \leq k \leq n$ , show that

$$\lambda_k(\mathbf{A} + \mathbf{B}) \leq \min\{\lambda_i(\mathbf{A}) + \lambda_j(\mathbf{B}) : i + j = k + n\}.$$

7. Explain why the Courant-Fischer theorem discussed in class is equivalent to

$$\lambda_k = \min_{\{\mathcal{S} : \dim \mathcal{S} = k\}} \max_{\{\mathbf{x} : 0 \neq \mathbf{x} \in \mathcal{S}\}} \frac{\mathbf{x}^H \mathbf{A} \mathbf{x}}{\mathbf{x}^H \mathbf{x}}$$

and

$$\lambda_k = \max_{\{\mathcal{S} : \dim \mathcal{S} = n - k + 1\}} \min_{\{\mathbf{x} : 0 \neq \mathbf{x} \in \mathcal{S}\}} \frac{\mathbf{x}^H \mathbf{A} \mathbf{x}}{\mathbf{x}^H \mathbf{x}}$$

where  $\mathbf{A} \in \mathbb{C}^{n \times n}$  is a Hermitian matrix with eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$ ,  $k \in \{1, \dots, n\}$  and  $\mathcal{S}$  denotes a subspace of  $\mathbb{C}^n$ .

8. Let  $\mathbf{A} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ . What are the eigenvalues of  $\mathbf{A}$ ? What is  $\max\{\mathbf{x}^T \mathbf{A} \mathbf{x} / \mathbf{x}^T \mathbf{x} : 0 \neq \mathbf{x} \in \mathbb{R}^2\}$ ? Does this contradict the Rayleigh-Ritz theorem?
9. Let  $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n}$  be Hermitian. Use Weyl's theorem to show that  $\lambda_1(\mathbf{B}) \leq \lambda_i(\mathbf{A} + \mathbf{B}) - \lambda_i(\mathbf{A}) \leq \lambda_n(\mathbf{B})$ . Hence, conclude that  $|\lambda_i(\mathbf{A} + \mathbf{B}) - \lambda_i(\mathbf{A})| \leq \rho(\mathbf{B})$  for all  $i = 1, \dots, n$ . This is an example of a *perturbation theorem* for the eigenvalues of a Hermitian matrix.
10. Let  $\lambda, a \in \mathbb{R}$ ,  $\mathbf{y} \in \mathbb{C}^n$ , and  $\mathbf{A} = \begin{bmatrix} \lambda \mathbf{I}_n & \mathbf{y} \\ \mathbf{y}^H & a \end{bmatrix} \in \mathbb{C}^{(n+1) \times (n+1)}$ . Use the Cauchy interlacing theorem to show that  $\lambda$  is an eigenvalue of  $\mathbf{A}$  with multiplicity at least  $n - 1$ . What are the other two eigenvalues?