

## E2-212 MATRIX THEORY: ASSIGNMENT 11

**Question 1.** Let  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$ , and  $\mathbf{A} \in \mathbb{C}^{n \times n}$ . Let  $\mathbf{B} = \mathbf{I} - \mathbf{A}$  and  $\mathbf{x}_0 \in \mathbb{C}^n$  be an arbitrary vector. Given  $\mathbf{A}$  and  $\mathbf{y}$ , we wish to solve the linear system of equations  $\mathbf{A}\mathbf{x} = \mathbf{y}$  as follows: (7 points)

For  $i = 1, 2, 3, \dots$ ,

Do :  $\mathbf{x}_i = \mathbf{B}\mathbf{x}_{i-1} + \mathbf{y}$

- (a) Let  $\varepsilon_i = \mathbf{x}_i - \mathbf{x}$  be the error in the  $i$ th iteration. Show that  $\varepsilon_i = \mathbf{B}^i(\mathbf{x}_0 - \mathbf{x})$ .
- (b) Conclude that if  $\rho(\mathbf{B}) < 1$ , then this algorithm works (in the sense that  $\mathbf{x}_i \rightarrow \mathbf{x}$  as  $i \rightarrow \infty$ ).
- (c) Use the Gersgorin theorem to show that a hermitian  $\mathbf{A}$  must be strictly diagonally dominant and  $\|\mathbf{A}\|_\infty < 2$  to ensure that this algorithm works.

**Question 2.** Let  $\mathbf{y}$  and  $\mathbf{x}$  be unit  $\ell_2$ -norm left and right eigenvectors respectively of  $\mathbf{C} \in \mathbb{C}^{n \times n}$  corresponding to a simple eigenvalue  $\lambda$ . Let  $S(\lambda) \triangleq |\mathbf{y}^H \mathbf{x}|$ . Prove that  $S(\lambda) \neq 0$ . (3 points)  
 (Hint for proof by contradiction: Show that  $\mathbf{x} \in \mathcal{R}(\mathbf{C} - \lambda \mathbf{I})$ . Also, there can exist only 1 linearly independent generalized eigenvector corresponding to simple eigenvalues.)