

# Recap: Markov Chain

- ▶ Let  $X_n$ ,  $n = 0, 1, \dots$  be a sequence of discrete random variables taking values in  $S$ .

# Recap: Markov Chain

- ▶ Let  $X_n$ ,  $n = 0, 1, \dots$  be a sequence of discrete random variables taking values in  $S$ .
- ▶ We say it is a Markov chain if

$$Pr[X_{n+1} = x_{n+1} | X_n = x_n, X_{n-1} = x_{n-1} \cdots X_0 = x_0] = Pr[X_{n+1} = x_{n+1} | X_n = x_n],$$

# Recap: Markov Chain

- ▶ Let  $X_n$ ,  $n = 0, 1, \dots$  be a sequence of discrete random variables taking values in  $S$ .
- ▶ We say it is a Markov chain if

$$Pr[X_{n+1} = x_{n+1} | X_n = x_n, X_{n-1} = x_{n-1} \cdots X_0 = x_0] = Pr[X_{n+1} = x_{n+1} | X_n = x_n],$$

- ▶ We can write it as

$$f_{X_{n+1}|X_n, \dots, X_0}(x_{n+1} | x_n, \dots, x_0) = f_{X_{n+1}|X_n}(x_{n+1} | x_n), \quad \forall x_i$$

# Recap: Markov Chain

- ▶ Let  $X_n$ ,  $n = 0, 1, \dots$  be a sequence of discrete random variables taking values in  $S$ .
- ▶ We say it is a Markov chain if

$$Pr[X_{n+1} = x_{n+1} | X_n = x_n, X_{n-1} = x_{n-1} \dots X_0 = x_0] = Pr[X_{n+1} = x_{n+1} | X_n = x_n],$$

- ▶ We can write it as

$$f_{X_{n+1}|X_n, \dots, X_0}(x_{n+1} | x_n, \dots, x_0) = f_{X_{n+1}|X_n}(x_{n+1} | x_n), \quad \forall x_i$$

- ▶ For a Markov chain, given the current state, the future evolution is independent of the history of how you reached the current state

# Recap: Transition Probabilities

- ▶ Transition probabilities:  $P(x, y) = \Pr[X_{n+1} = y | X_n = x]$

# Recap: Transition Probabilities

- ▶ Transition probabilities:  $P(x, y) = Pr[X_{n+1} = y | X_n = x]$   
Chain is homogeneous:  
 $Pr[X_{n+1} = y | X_n = x] = Pr[X_1 = y | X_0 = x], \forall n$

# Recap: Transition Probabilities

- ▶ Transition probabilities:  $P(x, y) = Pr[X_{n+1} = y | X_n = x]$   
Chain is homogeneous:  
 $Pr[X_{n+1} = y | X_n = x] = Pr[X_1 = y | X_0 = x], \forall n$
- ▶ Initial probabilities  $\pi_0(x) = Pr[X_0 = x]$

# Recap: Transition Probabilities

- ▶ Transition probabilities:  $P(x, y) = Pr[X_{n+1} = y | X_n = x]$   
Chain is homogeneous:  
 $Pr[X_{n+1} = y | X_n = x] = Pr[X_1 = y | X_0 = x], \forall n$
- ▶ Initial probabilities  $\pi_0(x) = Pr[X_0 = x]$
- ▶ Similarly,  $\pi_n(x) = Pr[X_n = x]$



# Recap: Chapman-Kolmogorov Equations

- ▶  $n$ -step transition probabilities:

$$P^n(x, y) = \Pr[X_n = y | X_0 = x]$$

# Recap: Chapman-Kolmogorov Equations

- ▶  $n$ -step transition probabilities:

$$P^n(x, y) = \Pr[X_n = y | X_0 = x]$$

- ▶ These satisfy Chapman-Kolmogorov equations:

$$P^{m+n}(x, y) = \sum_z P^m(x, z) P^n(z, y)$$

# Recap: Chapman-Kolmogorov Equations

- ▶  $n$ -step transition probabilities:

$$P^n(x, y) = \Pr[X_n = y | X_0 = x]$$

- ▶ These satisfy Chapman-Kolmogorov equations:

$$P^{m+n}(x, y) = \sum_z P^m(x, z) P^n(z, y)$$

- ▶ For a finite chain, the  $n$ -step transition probability matrix is  $n$ -fold product of the transition probability matrix

## Recap: transient and recurrent states

- ▶ Hitting time for  $y$ :  $T_y = \min\{n > 0 : X_n = y\}$

## Recap: transient and recurrent states

- ▶ Hitting time for  $y$ :  $T_y = \min\{n > 0 : X_n = y\}$
- ▶  $\rho_{xy} = P_x(T_y < \infty)$ .

# Recap: transient and recurrent states

- ▶ Hitting time for  $y$ :  $T_y = \min\{n > 0 : X_n = y\}$
- ▶  $\rho_{xy} = P_x(T_y < \infty)$ .
- ▶ A state  $y$  is called transient if  $\rho_{yy} < 1$ ; it is called recurrent if  $\rho_{yy} = 1$ .

# Recap: transient and recurrent states

- ▶ Hitting time for  $y$ :  $T_y = \min\{n > 0 : X_n = y\}$
- ▶  $\rho_{xy} = P_x(T_y < \infty)$ .
- ▶ A state  $y$  is called transient if  $\rho_{yy} < 1$ ; it is called recurrent if  $\rho_{yy} = 1$ .
- ▶  $N(y)$  – total number of visits to  $y$

# Recap: transient and recurrent states

- ▶ Hitting time for  $y$ :  $T_y = \min\{n > 0 : X_n = y\}$
- ▶  $\rho_{xy} = P_x(T_y < \infty)$ .
- ▶ A state  $y$  is called transient if  $\rho_{yy} < 1$ ; it is called recurrent if  $\rho_{yy} = 1$ .
- ▶  $N(y)$  – total number of visits to  $y$
- ▶  $G(x, y) = E_x[N(y)]$



# Recap

**Theorem:**

# Recap

## Theorem:

(i). Let  $y$  be transient. Then

$$P_x(N(y) < \infty) = 1, \forall x \text{ and } G(x, y) = \frac{\rho_{xy}}{1 - \rho_{yy}} < \infty, \forall x$$

# Recap

## Theorem:

(i). Let  $y$  be transient. Then

$$P_x(N(y) < \infty) = 1, \forall x \text{ and } G(x, y) = \frac{\rho_{xy}}{1 - \rho_{yy}} < \infty, \forall x$$

(ii) Let  $y$  be recurrent. Then

$$P_y[N(y) = \infty] = 1, \text{ and } G(y, y) = E_y[N(y)] = \infty$$

# Recap

## Theorem:

(i). Let  $y$  be transient. Then

$$P_x(N(y) < \infty) = 1, \forall x \text{ and } G(x, y) = \frac{\rho_{xy}}{1 - \rho_{yy}} < \infty, \forall x$$

(ii) Let  $y$  be recurrent. Then

$$P_y[N(y) = \infty] = 1, \text{ and } G(y, y) = E_y[N(y)] = \infty$$

$$P_x[N(y) = \infty] = \rho_{xy}, \text{ and } G(x, y) = \begin{cases} 0 & \text{if } \rho_{xy} = 0 \\ \infty & \text{if } \rho_{xy} > 0 \end{cases}$$

# Recap

- ▶ Transient states are visited only finitely many times while recurrent states are visited infinitely often

# Recap

- ▶ Transient states are visited only finitely many times while recurrent states are visited infinitely often
- ▶ A finite chain should have at least one recurrent state

# Recap

- ▶ Transient states are visited only finitely many times while recurrent states are visited infinitely often
- ▶ A finite chain should have at least one recurrent state
- ▶ We say,  $x$  leads to  $y$  if  $\rho_{xy} > 0$

# Recap

- ▶ Transient states are visited only finitely many times while recurrent states are visited infinitely often
- ▶ A finite chain should have at least one recurrent state
- ▶ We say,  $x$  leads to  $y$  if  $\rho_{xy} > 0$

**Theorem:** If  $x$  is recurrent and  $x$  leads to  $y$  then  $y$  is recurrent and  $\rho_{xy} = \rho_{yx} = 1$ .



# Recap: closed and irreducible sets

- ▶ A set of states,  $C \subset S$  is said to be irreducible if  $x$  leads to  $y$  for all  $x, y \in C$

# Recap: closed and irreducible sets

- ▶ A set of states,  $C \subset S$  is said to be irreducible if  $x$  leads to  $y$  for all  $x, y \in C$
- ▶ An irreducible set is also called a communicating class

# Recap: closed and irreducible sets

- ▶ A set of states,  $C \subset S$  is said to be irreducible if  $x$  leads to  $y$  for all  $x, y \in C$
- ▶ An irreducible set is also called a communicating class
- ▶ A set of states,  $C \subset S$ , is said to be closed if  $x \in C$ ,  $y \notin C$  implies  $\rho_{xy} = 0$ .

# Recap: closed and irreducible sets

- ▶ A set of states,  $C \subset S$  is said to be irreducible if  $x$  leads to  $y$  for all  $x, y \in C$
- ▶ An irreducible set is also called a communicating class
- ▶ A set of states,  $C \subset S$ , is said to be closed if  $x \in C$ ,  $y \notin C$  implies  $\rho_{xy} = 0$ .
- ▶ Once the chain visits a state in a closed set, it cannot leave that set.

# Recap: Partition of state space

- ▶  $S = S_T + S_R$ , transient and recurrent states

# Recap: Partition of state space

- ▶  $S = S_T + S_R$ , transient and recurrent states and

$$S_R = C_1 + C_2 + \dots$$

where  $C_i$  are closed and irreducible

# Recap: Partition of state space

- ▶  $S = S_T + S_R$ , transient and recurrent states and

$$S_R = C_1 + C_2 + \dots$$

where  $C_i$  are closed and irreducible

- ▶ We can calculate absorption probabilities for  $C_i$  using

$$\rho_C(x) = \sum_{y \in C} P(x, y) + \sum_{y \in S_T} P(x, y) \rho_C(y)$$

# Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$



## Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ For finite chains,  $P^T \pi = \pi$

# Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ For finite chains,  $P^T \pi = \pi$
- ▶ When  $\pi$  is stationary distribution,  
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$

# Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ For finite chains,  $P^T \pi = \pi$
- ▶ When  $\pi$  is stationary distribution,  
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If  $\pi_n = \pi, \forall n$  then  $\pi$  is a stationary distribution

# Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ For finite chains,  $P^T \pi = \pi$
- ▶ When  $\pi$  is stationary distribution,  
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If  $\pi_n = \pi, \forall n$  then  $\pi$  is a stationary distribution
- ▶ For a finite chain, a stationary distribution always exists.

# Recap: Stationary distribution

- ▶  $\pi$  is said to be a stationary distribution for the Markov chain with transition probabilities  $P$  if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ For finite chains,  $P^T \pi = \pi$
- ▶ When  $\pi$  is stationary distribution,  
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If  $\pi_n = \pi, \forall n$  then  $\pi$  is a stationary distribution
- ▶ For a finite chain, a stationary distribution always exists.
- ▶ The stationary distribution, when it exists, is related to expected fraction of time spent in different states.

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

$$G_n(x, y) \triangleq E_x[N_n(y)]$$



- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

$$G_n(x, y) \triangleq E_x[N_n(y)] = \sum_{m=1}^n E_x[I_y(X_m)]$$

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

$$G_n(x, y) \triangleq E_x[N_n(y)] = \sum_{m=1}^n E_x[I_y(X_m)] = \sum_{m=1}^n P^m(x, y)$$

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

$$G_n(x, y) \triangleq E_x[N_n(y)] = \sum_{m=1}^n E_x[I_y(X_m)] = \sum_{m=1}^n P^m(x, y)$$

- ▶ Expected fraction of time spent in  $y$  till  $n$  is

$$\frac{G_n(x, y)}{n} = \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ Let  $I_y(X_n)$  be indicator of  $[X_n = y]$
- ▶ Number of visits to  $y$  till  $n$ :  $N_n(y) = \sum_{m=1}^n I_y(X_m)$

$$G_n(x, y) \triangleq E_x[N_n(y)] = \sum_{m=1}^n E_x[I_y(X_m)] = \sum_{m=1}^n P^m(x, y)$$

- ▶ Expected fraction of time spent in  $y$  till  $n$  is

$$\frac{G_n(x, y)}{n} = \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ We will first establish a limit for the above as  $n \rightarrow \infty$

- ▶ Suppose  $y$  is transient. Then we have

- Suppose  $y$  is transient. Then we have

$$\lim_{n \rightarrow \infty} N_n(y) = N(y)$$

- Suppose  $y$  is transient. Then we have

$$\lim_{n \rightarrow \infty} N_n(y) = N(y)$$

and  $Pr[N(y) < \infty] = 1$   $\lim_{n \rightarrow \infty} G_n(x, y) = G(x, y) < \infty$

► Suppose  $y$  is transient. Then we have

$$\lim_{n \rightarrow \infty} N_n(y) = N(y)$$

$$\text{and } Pr[N(y) < \infty] = 1 \quad \lim_{n \rightarrow \infty} G_n(x, y) = G(x, y) < \infty$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = 0 \text{ (w.p.1)} \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = 0$$



- Suppose  $y$  is transient. Then we have

$$\lim_{n \rightarrow \infty} N_n(y) = N(y)$$

$$\text{and } Pr[N(y) < \infty] = 1 \quad \lim_{n \rightarrow \infty} G_n(x, y) = G(x, y) < \infty$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = 0 \text{ (w.p.1)} \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = 0$$

- The expected fraction of time spent in a transient state is zero.

- Suppose  $y$  is transient. Then we have

$$\lim_{n \rightarrow \infty} N_n(y) = N(y)$$

$$\text{and } Pr[N(y) < \infty] = 1 \quad \lim_{n \rightarrow \infty} G_n(x, y) = G(x, y) < \infty$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = 0 \text{ (w.p.1)} \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = 0$$

- The expected fraction of time spent in a transient state is zero.
- This is intuitively obvious

- ▶ Now, let  $y$  be recurrent

- ▶ Now, let  $y$  be recurrent
- ▶ Then,  $P_y[T_y < \infty] = 1$

- ▶ Now, let  $y$  be recurrent
- ▶ Then,  $P_y[T_y < \infty] = 1$
- ▶ Define  $m_y = E_y[T_y]$

- ▶ Now, let  $y$  be recurrent
- ▶ Then,  $P_y[T_y < \infty] = 1$
- ▶ Define  $m_y = E_y[T_y]$
- ▶  $m_y$  is mean return time to  $y$

- ▶ Now, let  $y$  be recurrent
- ▶ Then,  $P_y[T_y < \infty] = 1$
- ▶ Define  $m_y = E_y[T_y]$
- ▶  $m_y$  is mean return time to  $y$
- ▶ We will show that  $\frac{N_n(y)}{n}$  converges to  $\frac{1}{m_y}$  if the chain starts in  $y$ .

- ▶ Now, let  $y$  be recurrent
- ▶ Then,  $P_y[T_y < \infty] = 1$
- ▶ Define  $m_y = E_y[T_y]$
- ▶  $m_y$  is mean return time to  $y$
- ▶ We will show that  $\frac{N_n(y)}{n}$  converges to  $\frac{1}{m_y}$  if the chain starts in  $y$ .
- ▶ Convergence would be with probability one.



- ▶ Consider a chain started in  $y$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$
- ▶ Also,  $T_y^r = W_y^1 + \cdots + W_y^r$

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$
- ▶ Also,  $T_y^r = W_y^1 + \cdots + W_y^r$
- ▶  $W_y^r$  are the “waiting times”

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$
- ▶ Also,  $T_y^r = W_y^1 + \cdots + W_y^r$
- ▶  $W_y^r$  are the “waiting times”
- ▶ By Markovian property we should expect them to be iid



- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$
- ▶ Also,  $T_y^r = W_y^1 + \cdots + W_y^r$
- ▶  $W_y^r$  are the “waiting times”
- ▶ By Markovian property we should expect them to be iid
- ▶ We will prove this.

- ▶ Consider a chain started in  $y$
- ▶ let  $T_y^r$  be time of  $r^{th}$  visit to  $y$ ,  $r \geq 1$

$$T_y^r = \min\{n \geq 1 : N_n(y) = r\}$$

- ▶ Define  $W_y^1 = T_y^1 = T_y$  and  $W_y^r = T_y^r - T_y^{r-1}$ ,  $r > 1$
- ▶ Note that  $E_y[W_y^1] = E_y[T_y] = m_y$
- ▶ Also,  $T_y^r = W_y^1 + \cdots + W_y^r$
- ▶  $W_y^r$  are the “waiting times”
- ▶ By Markovian property we should expect them to be iid
- ▶ We will prove this.
- ▶ Then  $T_y^r/r$  converges to  $m_y$  by law of large umbers

► We have

► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

- We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

- Using the Markovian property, we get

► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

► Using the Markovian property, we get

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid X_{k_1+k_2} = y]$$

► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

► Using the Markovian property, we get

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid X_{k_1+k_2} = y]$$

$$= Pr[X_j \neq y, 1 \leq j \leq k_3 - 1, X_{k_3} = y \mid X_0 = y]$$



► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

► Using the Markovian property, we get

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid X_{k_1+k_2} = y]$$

$$= Pr[X_j \neq y, 1 \leq j \leq k_3 - 1, X_{k_3} = y \mid X_0 = y]$$

$$= P_y[W_y^1 = k_3]$$

► We have

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid B]$$

where  $B = [X_{k_1+k_2} = y, X_{k_1} = y, X_j \neq y, j < k_1 + k_2, j \neq k_1]$

► Using the Markovian property, we get

$$Pr[W_y^3 = k_3 | W_y^2 = k_2, W_y^1 = k_1] =$$

$$Pr[X_{k_1+k_2+j} \neq y, 1 \leq j \leq k_3 - 1, X_{k_1+k_2+k_3} = y \mid X_{k_1+k_2} = y]$$

$$= Pr[X_j \neq y, 1 \leq j \leq k_3 - 1, X_{k_3} = y \mid X_0 = y]$$

$$= P_y[W_y^1 = k_3]$$

► In general, we get

$$Pr[W_y^r = k_r \mid W_y^{r-1} = k_{r-1}, \dots, W_y^1 = k_1] = P_y[W_y^1 = k_r]$$

- ▶ This shows the waiting time are iid

- This shows the waiting time are iid

$$P_y[W_y^2 = k_2] = \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1]$$

- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \end{aligned}$$

- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] \end{aligned}$$

- This shows the waiting time are iid

$$\begin{aligned}P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\&= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\&= P_y[W_y^1 = k_2] \\&\Rightarrow \text{identically distributed}\end{aligned}$$

- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] \\ \Rightarrow &\text{ identically distributed} \end{aligned}$$

$$P_y[W_y^2 = k_2, W_y^1 = k_1] = P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1]$$



- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] \\ \Rightarrow &\text{ identically distributed} \end{aligned}$$

$$\begin{aligned} P_y[W_y^2 = k_2, W_y^1 = k_1] &= P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \end{aligned}$$

- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] \\ \Rightarrow &\text{ identically distributed} \end{aligned}$$

$$\begin{aligned} P_y[W_y^2 = k_2, W_y^1 = k_1] &= P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^2 = k_2] P_y[W_y^1 = k_1] \end{aligned}$$

- This shows the waiting time are iid

$$\begin{aligned} P_y[W_y^2 = k_2] &= \sum_{k_1} P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= \sum_{k_1} P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] \\ \Rightarrow &\text{ identically distributed} \end{aligned}$$

$$\begin{aligned} P_y[W_y^2 = k_2, W_y^1 = k_1] &= P_y[W_y^2 = k_2 \mid W_y^1 = k_1] P_y[W_y^1 = k_1] \\ &= P_y[W_y^1 = k_2] P_y[W_y^1 = k_1] \\ &= P_y[W_y^2 = k_2] P_y[W_y^1 = k_1] \\ \Rightarrow &\text{ independent} \end{aligned}$$

- ▶ We have shown  $W_y^r$ ,  $r = 1, 2, \dots$  are iid

- ▶ We have shown  $W_y^r$ ,  $r = 1, 2, \dots$  are iid
- ▶ Since  $E[W_y^1] = m_y$ , by strong law of large numbers,

$$\lim_{k \rightarrow \infty} \frac{T_y^k}{k} = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{r=1}^k W_y^r = m_y, \quad (w.p.1)$$

- ▶ We have shown  $W_y^r$ ,  $r = 1, 2, \dots$  are iid
- ▶ Since  $E[W_y^1] = m_y$ , by strong law of large numbers,

$$\lim_{k \rightarrow \infty} \frac{T_y^k}{k} = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{r=1}^k W_y^r = m_y, \quad (w.p.1)$$

- ▶ Note that this is true even if  $m_y = \infty$

- For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- $N_n(y)$  is the number of visits to  $y$  till time step  $n$



- ▶ For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- ▶  $N_n(y)$  is the number of visits to  $y$  till time step  $n$
- ▶ Suppose  $N_{50}(y) = 8$

- For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- $N_n(y)$  is the number of visits to  $y$  till time step  $n$
- Suppose  $N_{50}(y) = 8$  – Visited  $y$  8 times till time 50.

- For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- $N_n(y)$  is the number of visits to  $y$  till time step  $n$
- Suppose  $N_{50}(y) = 8$  – Visited  $y$  8 times till time 50.
- So, the 8<sup>th</sup> visit occurred at or before time 50.

- ▶ For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- ▶  $N_n(y)$  is the number of visits to  $y$  till time step  $n$
- ▶ Suppose  $N_{50}(y) = 8$  – Visited  $y$  8 times till time 50.
- ▶ So, the  $8^{th}$  visit occurred at or before time 50.
- ▶ The  $9^{th}$  visit has not occurred till 50.

- For all  $n$  such that  $N_n(y) \geq 1$ , we have

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

- $N_n(y)$  is the number of visits to  $y$  till time step  $n$
- Suppose  $N_{50}(y) = 8$  – Visited  $y$  8 times till time 50.
- So, the  $8^{th}$  visit occurred at or before time 50.
- The  $9^{th}$  visit has not occurred till 50.
- So, time of  $9^{th}$  visit is beyond 50.

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

► Now we have

$$\frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} < \frac{T_y^{N_n(y)+1}}{N_n(y)}$$

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

► Now we have

$$\frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} < \frac{T_y^{N_n(y)+1}}{N_n(y)}$$

► We know that



$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

► Now we have

$$\frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} < \frac{T_y^{N_n(y)+1}}{N_n(y)}$$

► We know that

► As  $n \rightarrow \infty$ ,  $N_n(y) \rightarrow \infty$ , *w.p.1*

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

► Now we have

$$\frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} < \frac{T_y^{N_n(y)+1}}{N_n(y)}$$

► We know that

- As  $n \rightarrow \infty$ ,  $N_n(y) \rightarrow \infty$ , *w.p.1*
- As  $n \rightarrow \infty$ ,  $\frac{T_y^n}{n} \rightarrow m_y$ , *w.p.1*

$$T_y^{N_n(y)} \leq n < T_y^{N_n(y)+1}$$

► Now we have

$$\frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} < \frac{T_y^{N_n(y)+1}}{N_n(y)}$$

► We know that

► As  $n \rightarrow \infty$ ,  $N_n(y) \rightarrow \infty$ , *w.p.1*

► As  $n \rightarrow \infty$ ,  $\frac{T_y^{N_n(y)}}{N_n(y)} \rightarrow m_y$ , *w.p.1*

► Hence we get

$$\lim_{n \rightarrow \infty} \frac{n}{N_n(y)} = m_y, \quad w.p.1$$

or

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{m_y}, \quad w.p.1$$

- ▶ All this is true if the chain started in  $y$ .

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Since  $0 \leq \frac{N_n(y)}{n} \leq 1$ , almost sure convergence implies convergence in mean

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Since  $0 \leq \frac{N_n(y)}{n} \leq 1$ , almost sure convergence implies convergence in mean

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} E_x \left[ \frac{N_n(y)}{n} \right]$$



- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Since  $0 \leq \frac{N_n(y)}{n} \leq 1$ , almost sure convergence implies convergence in mean

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} E_x \left[ \frac{N_n(y)}{n} \right] = \lim_{n \rightarrow \infty} \frac{P_x[T_y < \infty]}{m_y}$$

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Since  $0 \leq \frac{N_n(y)}{n} \leq 1$ , almost sure convergence implies convergence in mean

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} E_x \left[ \frac{N_n(y)}{n} \right] = \lim_{n \rightarrow \infty} \frac{P_x[T_y < \infty]}{m_y} = \frac{\rho_{xy}}{m_y}$$

- ▶ All this is true if the chain started in  $y$ .
- ▶ That means it is true if the chain visits  $y$  once.
- ▶ So, we get

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Since  $0 \leq \frac{N_n(y)}{n} \leq 1$ , almost sure convergence implies convergence in mean

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} E_x \left[ \frac{N_n(y)}{n} \right] = \lim_{n \rightarrow \infty} \frac{P_x[T_y < \infty]}{m_y} = \frac{\rho_{xy}}{m_y}$$

- ▶ The fraction of time spent in each recurrent state is inversely proportional to the mean recurrence time

► Thus we have proved the following theorem

- ▶ Thus we have proved the following theorem
- ▶ **Theorem:**  
Let  $y$  be recurrent. Then

► Thus we have proved the following theorem

► **Theorem:**

Let  $y$  be recurrent. Then

1

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

- ▶ Thus we have proved the following theorem

- ▶ **Theorem:**

Let  $y$  be recurrent. Then

1

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

2

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \frac{\rho_{xy}}{m_y}$$

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.



- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .  
Though  $P_y[T_y < \infty] = 1$ , we can have  $E_y[T_y] = \infty$ .

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .  
Though  $P_y[T_y < \infty] = 1$ , we can have  $E_y[T_y] = \infty$ .
- ▶ What if  $m_y = \infty$ ,  $\forall y$ ?

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .  
Though  $P_y[T_y < \infty] = 1$ , we can have  $E_y[T_y] = \infty$ .
- ▶ What if  $m_y = \infty$ ,  $\forall y$ ?
- ▶ Does not seem reasonable for a finite chain.

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .  
Though  $P_y[T_y < \infty] = 1$ , we can have  $E_y[T_y] = \infty$ .
- ▶ What if  $m_y = \infty$ ,  $\forall y$ ?
- ▶ Does not seem reasonable for a finite chain.
- ▶ But for infinite chains??

- ▶ The limiting fraction of time spent in a state is inversely proportional to  $m_y$ , the mean return time.
- ▶ Intuitively, the stationary probability of a state could be the limiting fraction of time spent in that state.
- ▶ Thus  $\pi(y) = \frac{1}{m_y}$  is a good candidate for stationary distribution.
- ▶ We first note that we can have  $m_y = \infty$ .  
Though  $P_y[T_y < \infty] = 1$ , we can have  $E_y[T_y] = \infty$ .
- ▶ What if  $m_y = \infty$ ,  $\forall y$ ?
- ▶ Does not seem reasonable for a finite chain.
- ▶ But for infinite chains??
- ▶ Let us characterize  $y$  for which  $m_y = \infty$



- ▶ A recurrent state  $y$  is called **null recurrent** if  $m_y = \infty$ .

- ▶ A recurrent state  $y$  is called **null recurrent** if  $m_y = \infty$ .
- ▶  $y$  is called **positive recurrent** if  $m_y < \infty$

- ▶ A recurrent state  $y$  is called **null recurrent** if  $m_y = \infty$ .
- ▶  $y$  is called **positive recurrent** if  $m_y < \infty$
- ▶ We earlier saw that the fraction of time spent in a transient state is zero.

- ▶ A recurrent state  $y$  is called **null recurrent** if  $m_y = \infty$ .
- ▶  $y$  is called **positive recurrent** if  $m_y < \infty$
- ▶ We earlier saw that the fraction of time spent in a transient state is zero.
- ▶ Suppose  $y$  is null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{m_y} = 0$$

- ▶ A recurrent state  $y$  is called **null recurrent** if  $m_y = \infty$ .
- ▶  $y$  is called **positive recurrent** if  $m_y < \infty$
- ▶ We earlier saw that the fraction of time spent in a transient state is zero.
- ▶ Suppose  $y$  is null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{m_y} = 0$$

- ▶ Thus the limiting fraction of time spent by the chain in transient and null recurrent states is zero.

- **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

- **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

**Proof**

- ▶ **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

**Proof**

- ▶ Since  $x$  is recurrent and  $x$  leads to  $y$  we know  $\exists n_0, n_1$  s.t.  $P^{n_0}(x, y) > 0$ ,  $P^{n_1}(y, x) > 0$  and



- ▶ **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

**Proof**

- ▶ Since  $x$  is recurrent and  $x$  leads to  $y$  we know  $\exists n_0, n_1$  s.t.  $P^{n_0}(x, y) > 0$ ,  $P^{n_1}(y, x) > 0$  and

$$P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x)P^m(x, x)P^{n_0}(x, y), \quad \forall m$$

- **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

**Proof**

- Since  $x$  is recurrent and  $x$  leads to  $y$  we know  $\exists n_0, n_1$  s.t.  $P^{n_0}(x, y) > 0$ ,  $P^{n_1}(y, x) > 0$  and

$$P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x)P^m(x, x)P^{n_0}(x, y), \quad \forall m$$

Summing the above for  $m = 1, 2, \dots, n$  and dividing by  $n$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

- **Theorem:** Let  $x$  be positive recurrent and let  $x$  lead to  $y$ . Then  $y$  is positive recurrent.

**Proof**

- Since  $x$  is recurrent and  $x$  leads to  $y$  we know  $\exists n_0, n_1$  s.t.  $P^{n_0}(x, y) > 0$ ,  $P^{n_1}(y, x) > 0$  and

$$P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) P^m(x, x) P^{n_0}(x, y), \quad \forall m$$

Summing the above for  $m = 1, 2, \dots, n$  and dividing by  $n$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

If we now let  $n \rightarrow \infty$ , the RHS goes to  $P^{n_1}(y, x) \frac{1}{m_x} P^{n_0}(x, y) > 0$ .

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) = \frac{1}{n} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y)$$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\begin{aligned} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) &= \frac{1}{n} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \\ &= \frac{n_1 + n + n_0}{n} \frac{1}{n_1 + n + n_0} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \end{aligned}$$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\begin{aligned} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) &= \frac{1}{n} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \\ &= \frac{n_1 + n + n_0}{n} \frac{1}{n_1 + n + n_0} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \end{aligned}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) = \frac{1}{m_y}$$



$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \frac{1}{n} \sum_{m=1}^n P^m(x, x) P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\begin{aligned} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) &= \frac{1}{n} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \\ &= \frac{n_1 + n + n_0}{n} \frac{1}{n_1 + n + n_0} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \end{aligned}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) = \frac{1}{m_y}$$

$$\Rightarrow \frac{1}{m_y} \geq P^{n_1}(y, x) \frac{1}{m_x} P^{n_0}(x, y) > 0$$

$$\frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) \geq P^{n_1}(y, x) \quad \frac{1}{n} \sum_{m=1}^n P^m(x, x) \quad P^{n_0}(x, y), \quad \forall n$$

► We can write the *LHS* of above as

$$\begin{aligned} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) &= \frac{1}{n} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \\ &= \frac{n_1+n+n_0}{n} \frac{1}{n_1+n+n_0} \sum_{m=1}^{n_1+n+n_0} P^m(y, y) - \frac{1}{n} \sum_{m=1}^{n_1+n_0} P^m(y, y) \end{aligned}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^{n_1+m+n_0}(y, y) = \frac{1}{m_y}$$

$$\Rightarrow \frac{1}{m_y} \geq P^{n_1}(y, x) \frac{1}{m_x} P^{n_0}(x, y) > 0$$

which implies  $y$  is positive recurrent

- ▶ Thus, in a closed irreducible set of recurrent states, if one state is positive recurrent then all are positive recurrent.

- ▶ Thus, in a closed irreducible set of recurrent states, if one state is positive recurrent then all are positive recurrent.
- ▶ Hence, in the partition:  $S_R = C_1 + C_2 + \dots$ , each  $C_i$  is either wholly positive recurrent or wholly null recurrent.

- ▶ Thus, in a closed irreducible set of recurrent states, if one state is positive recurrent then all are positive recurrent.
- ▶ Hence, in the partition:  $S_R = C_1 + C_2 + \dots$ , each  $C_i$  is either wholly positive recurrent or wholly null recurrent.
- ▶ We next show that a finite chain cannot have any null recurrent states.

- ▶ Let  $C$  be a finite closed set of recurrent states.

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$



- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \quad \forall m, \quad \forall x \in C.$

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \forall m, \forall x \in C$ .
- ▶ Thus we get

$$1 = \frac{1}{n} \sum_{m=1}^n \sum_{y \in C} P^m(x, y)$$

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \forall m, \forall x \in C$ .
- ▶ Thus we get

$$1 = \frac{1}{n} \sum_{m=1}^n \sum_{y \in C} P^m(x, y) = \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y), \quad \forall n$$

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \forall m, \forall x \in C$ .
- ▶ Thus we get

$$1 = \frac{1}{n} \sum_{m=1}^n \sum_{y \in C} P^m(x, y) = \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y), \quad \forall n$$

$$\Rightarrow 1 = \lim_{n \rightarrow \infty} \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \forall m, \forall x \in C$ .
- ▶ Thus we get

$$1 = \frac{1}{n} \sum_{m=1}^n \sum_{y \in C} P^m(x, y) = \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y), \quad \forall n$$

$$\Rightarrow 1 = \lim_{n \rightarrow \infty} \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = \sum_{y \in C} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0$$

- ▶ Let  $C$  be a finite closed set of recurrent states.
- ▶ Suppose all states in  $C$  are null recurrent. Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0, \quad \forall x, y \in C$$

- ▶ Since  $C$  is closed,  $\sum_{y \in C} P^m(x, y) = 1, \forall m, \forall x \in C$ .
- ▶ Thus we get

$$1 = \frac{1}{n} \sum_{m=1}^n \sum_{y \in C} P^m(x, y) = \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y), \quad \forall n$$

$$\Rightarrow 1 = \lim_{n \rightarrow \infty} \sum_{y \in C} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = \sum_{y \in C} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0$$

where we could take the limit inside the sum because  $C$  is finite.

- ▶ If  $C$  is a finite closed set of recurrent states then all states in it cannot be null recurrent.

- ▶ If  $C$  is a finite closed set of recurrent states then all states in it cannot be null recurrent.
- ▶ Actually what we showed is that any closed finite set must have at least one positive recurrent state.




- ▶ If  $C$  is a finite closed set of recurrent states then all states in it cannot be null recurrent.
- ▶ Actually what we showed is that any closed finite set must have at least one positive recurrent state.
- ▶ Hence, in a finite chain, every closed irreducible set of recurrent states contains only positive recurrent states.

- ▶ If  $C$  is a finite closed set of recurrent states then all states in it cannot be null recurrent.
- ▶ Actually what we showed is that any closed finite set must have at least one positive recurrent state.
- ▶ Hence, in a finite chain, every closed irreducible set of recurrent states contains only positive recurrent states.
- ▶ Hence, a finite chain cannot have a null recurrent state.


# Example of null recurrent chain

- Consider the chain with state space  $\{0, 1, \dots\}$  given by



# Example of null recurrent chain


- Consider the chain with state space  $\{0, 1, \dots\}$  given by



- Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ .

## Example of null recurrent chain

- Consider the chain with state space  $\{0, 1, \dots\}$  given by




- Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have

$$P_0[T_0 = j+1] = q(j)$$

## Example of null recurrent chain

- ▶ Consider the chain with state space  $\{0, 1, \dots\}$  given by




- ▶ Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have

$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_o[T_0 = j]$$

# Example of null recurrent chain

- Consider the chain with state space  $\{0, 1, \dots\}$  given by




- Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have

$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_0[T_0 = j] = \sum_{j=2}^{\infty} j q(j-1)$$

(Note that  $P_0[T_0 = 1] = 0$ )

# Example of null recurrent chain

- ▶ Consider the chain with state space  $\{0, 1, \dots\}$  given by



- ▶ Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have

$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_0[T_0 = j] = \sum_{j=2}^{\infty} j q(j-1)$$


(Note that  $P_0[T_0 = 1] = 0$ )

- ▶ So,  $m_0 = \infty$  if the  $q(\cdot)$  distribution has infinite expectation.



# Example of null recurrent chain

- ▶ Consider the chain with state space  $\{0, 1, \dots\}$  given by



- ▶ Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have


$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_0[T_0 = j] = \sum_{j=2}^{\infty} j q(j-1)$$

(Note that  $P_0[T_0 = 1] = 0$ )

- ▶ So,  $m_0 = \infty$  if the  $q(\cdot)$  distribution has infinite expectation. For example, if  $q(k) = \frac{c}{k^2}$

# Example of null recurrent chain

- ▶ Consider the chain with state space  $\{0, 1, \dots\}$  given by



- ▶ Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have


$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_0[T_0 = j] = \sum_{j=2}^{\infty} j q(j-1)$$

(Note that  $P_0[T_0 = 1] = 0$ )

- ▶ So,  $m_0 = \infty$  if the  $q(\cdot)$  distribution has infinite expectation. For example, if  $q(k) = \frac{c}{k^2}$
- ▶ Then state 0 is null recurrent.

## Example of null recurrent chain

- Consider the chain with state space  $\{0, 1, \dots\}$  given by



- Here,  $q(k) \geq 0, \forall k$  and  $\sum_{k=1}^{\infty} q(k) = 1$ . We have

$$P_0[T_0 = j+1] = q(j) \Rightarrow m_0 = \sum_{j=2}^{\infty} j P_0[T_0 = j] = \sum_{j=2}^{\infty} j q(j-1)$$

(Note that  $P_0[T_0 = 1] = 0$ )

- So,  $m_0 = \infty$  if the  $q(\cdot)$  distribution has infinite expectation. For example, if  $q(k) = \frac{c}{k^2}$
- Then state 0 is null recurrent. Implies chain is null recurrent

- ▶ Suppose  $\pi$  is a stationary distribution.

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

$$\pi(y) = \sum_x \pi(x) P^m(x, y) \quad \forall m$$

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

$$\pi(y) = \sum_x \pi(x) P^m(x, y) \quad \forall m$$

$$\Rightarrow \pi(y) = \frac{1}{n} \sum_{m=1}^n \sum_x \pi(x) P^m(x, y)$$



- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

$$\pi(y) = \sum_x \pi(x) P^m(x, y) \quad \forall m$$

$$\Rightarrow \pi(y) = \frac{1}{n} \sum_{m=1}^n \sum_x \pi(x) P^m(x, y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

$$\pi(y) = \sum_x \pi(x) P^m(x, y) \quad \forall m$$

$$\Rightarrow \pi(y) = \frac{1}{n} \sum_{m=1}^n \sum_x \pi(x) P^m(x, y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

$$\Rightarrow \pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ Suppose  $\pi$  is a stationary distribution.
- ▶ Then  $\pi(y) = 0$  if  $y$  is transient or null recurrent
- ▶ We prove this as follows

$$\pi(y) = \sum_x \pi(x) P^m(x, y) \quad \forall m$$

$$\Rightarrow \pi(y) = \frac{1}{n} \sum_{m=1}^n \sum_x \pi(x) P^m(x, y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

$$\Rightarrow \pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- ▶ The proof is complete if we can take the limit inside the sum

► **Bounded Convergence Theorem:** Suppose  $a(x) \geq 0$ ,  $\forall x \in S$  and  $\sum_x a(x) < \infty$ . Let  $b_n(x)$ ,  $x \in S$  be such that  $|b_n(x)| \leq K$ ,  $\forall x, n$  and suppose  $\lim_{n \rightarrow \infty} b_n(x) = b(x)$ ,  $\forall x \in S$ . Then

$$\lim_{n \rightarrow \infty} \sum_{x \in S} a(x) b_n(x) = \sum_{x \in S} a(x) \lim_{n \rightarrow \infty} b_n(x) = \sum_{x \in S} a(x) b(x)$$

- **Bounded Convergence Theorem:** Suppose  $a(x) \geq 0$ ,  $\forall x \in S$  and  $\sum_x a(x) < \infty$ . Let  $b_n(x)$ ,  $x \in S$  be such that  $|b_n(x)| \leq K$ ,  $\forall x, n$  and suppose  $\lim_{n \rightarrow \infty} b_n(x) = b(x)$ ,  $\forall x \in S$ . Then

$$\lim_{n \rightarrow \infty} \sum_{x \in S} a(x) b_n(x) = \sum_{x \in S} a(x) \lim_{n \rightarrow \infty} b_n(x) = \sum_{x \in S} a(x) b(x)$$

- We had

$$\pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- **Bounded Convergence Theorem:** Suppose  $a(x) \geq 0$ ,  $\forall x \in S$  and  $\sum_x a(x) < \infty$ . Let  $b_n(x)$ ,  $x \in S$  be such that  $|b_n(x)| \leq K$ ,  $\forall x, n$  and suppose  $\lim_{n \rightarrow \infty} b_n(x) = b(x)$ ,  $\forall x \in S$ . Then

$$\lim_{n \rightarrow \infty} \sum_{x \in S} a(x) b_n(x) = \sum_{x \in S} a(x) \lim_{n \rightarrow \infty} b_n(x) = \sum_{x \in S} a(x) b(x)$$

- We had

$$\pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- We have

$$\pi(x) \geq 0; \quad \sum_x \pi(x) = 1; \quad 0 \leq \frac{1}{n} \sum_{m=1}^n P^m(x, y) \leq 1, \forall x$$

- **Bounded Convergence Theorem:** Suppose  $a(x) \geq 0$ ,  $\forall x \in S$  and  $\sum_x a(x) < \infty$ . Let  $b_n(x)$ ,  $x \in S$  be such that  $|b_n(x)| \leq K$ ,  $\forall x, n$  and suppose  $\lim_{n \rightarrow \infty} b_n(x) = b(x)$ ,  $\forall x \in S$ . Then

$$\lim_{n \rightarrow \infty} \sum_{x \in S} a(x) b_n(x) = \sum_{x \in S} a(x) \lim_{n \rightarrow \infty} b_n(x) = \sum_{x \in S} a(x) b(x)$$

- We had

$$\pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- We have

$$\pi(x) \geq 0; \quad \sum_x \pi(x) = 1; \quad 0 \leq \frac{1}{n} \sum_{m=1}^n P^m(x, y) \leq 1, \forall x$$

- Hence, if  $y$  is transient or null recurrent, then

$$\pi(y) = \sum_x \pi(x) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = 0$$

- ▶ In any stationary distribution  $\pi$ , we would have  $\pi(y) = 0$  if  $y$  is transient or null recurrent.



- ▶ In any stationary distribution  $\pi$ , we would have  $\pi(y) = 0$  if  $y$  is transient or null recurrent.
- ▶ Hence an irreducible transient or null recurrent chain would not have a stationary distribution.

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x)P(x, y)$ . Then

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x)P(x, y)$ . Then

$$\pi(y) = \sum_x \pi(x) P^m(x, y), \forall m$$

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x)P(x, y)$ . Then

$$\pi(y) = \sum_x \pi(x) P^m(x, y), \forall m$$

$$\Rightarrow \pi(y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y), \forall n$$

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x)P(x, y)$ . Then

$$\pi(y) = \sum_x \pi(x) P^m(x, y), \forall m$$

$$\Rightarrow \pi(y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y), \forall n$$

$$\Rightarrow \pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x)P(x, y)$ . Then

$$\pi(y) = \sum_x \pi(x) P^m(x, y), \forall m$$

$$\Rightarrow \pi(y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y), \forall n$$

$$\Rightarrow \pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

$$\Rightarrow \pi(y) = \sum_x \pi(x) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

- **Theorem** An irreducible positive recurrent chain has a unique stationary distribution given by

$$\pi(y) = \frac{1}{m_y}, \quad \forall y \in S$$

- Suppose  $\exists \pi$  such that  $\pi(y) = \sum_x \pi(x) P(x, y)$ . Then

$$\pi(y) = \sum_x \pi(x) P^m(x, y), \quad \forall m$$

$$\Rightarrow \pi(y) = \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y), \quad \forall n$$

$$\Rightarrow \pi(y) = \lim_{n \rightarrow \infty} \sum_x \pi(x) \frac{1}{n} \sum_{m=1}^n P^m(x, y)$$

$$\begin{aligned} \Rightarrow \pi(y) &= \sum_x \pi(x) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) \\ &= \sum_x \pi(x) \frac{1}{m_y} = \frac{1}{m_y} \end{aligned}$$



- To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .

- ▶ To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .
- ▶ We also need to show  $\frac{1}{m_y} = \sum_x \frac{1}{m_x} P(x, y)$

- ▶ To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .
- ▶ We also need to show  $\frac{1}{m_y} = \sum_x \frac{1}{m_x} P(x, y)$
- ▶ We skip these steps in the proof.

- ▶ To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .
- ▶ We also need to show  $\frac{1}{m_y} = \sum_x \frac{1}{m_x} P(x, y)$
- ▶ We skip these steps in the proof.
- ▶ The theorem shows that an irreducible positive recurrent chain has a unique stationary distribution

- ▶ To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .
- ▶ We also need to show  $\frac{1}{m_y} = \sum_x \frac{1}{m_x} P(x, y)$
- ▶ We skip these steps in the proof.
- ▶ The theorem shows that an irreducible positive recurrent chain has a unique stationary distribution
- ▶ Corollary: An irreducible chain has a stationary distribution if and only if it is positive recurrent

- ▶ To complete the proof, we need to show  $\sum_y \frac{1}{m_y} = 1$ .
- ▶ We also need to show  $\frac{1}{m_y} = \sum_x \frac{1}{m_x} P(x, y)$
- ▶ We skip these steps in the proof.
- ▶ The theorem shows that an irreducible positive recurrent chain has a unique stationary distribution
- ▶ Corollary: An irreducible chain has a stationary distribution if and only if it is positive recurrent
- ▶ An irreducible finite chain has a unique stationary distribution

- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$

- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$  (easily verified)



- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$  (easily verified)
- ▶ Let  $C$  be a closed irreducible set of positive recurrent states.

- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$  (easily verified)
- ▶ Let  $C$  be a closed irreducible set of positive recurrent states.

Then there is a unique stationary distribution  $\pi$  that satisfies  $\pi(y) = 0, \forall y \notin C$ .

- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$  (easily verified)
- ▶ Let  $C$  be a closed irreducible set of positive recurrent states.  
Then there is a unique stationary distribution  $\pi$  that satisfies  $\pi(y) = 0, \forall y \notin C$ .
- ▶ Any other stationary distribution of the chain is a convex combination of the stationary distributions concentrated on each of the closed irreducible sets of positive recurrent states.

- ▶ If  $\pi^1$  and  $\pi^2$  are stationary distributions, then so is  $\alpha\pi^1 + (1 - \alpha)\pi^2$  (easily verified)
- ▶ Let  $C$  be a closed irreducible set of positive recurrent states.  
Then there is a unique stationary distribution  $\pi$  that satisfies  $\pi(y) = 0, \forall y \notin C$ .
- ▶ Any other stationary distribution of the chain is a convex combination of the stationary distributions concentrated on each of the closed irreducible sets of positive recurrent states.
- ▶ This answers all questions about existence and uniqueness of stationary distributions

- ▶ Consider an irreducible positive recurrent chain.

- ▶ Consider an irreducible positive recurrent chain.
- ▶ It has a unique stationary distribution and  $\frac{1}{n} \sum_{m=1}^n P^m(x, y)$  converges to  $\pi(y)$ .

- ▶ Consider an irreducible positive recurrent chain.
- ▶ It has a unique stationary distribution and  $\frac{1}{n} \sum_{m=1}^n P^m(x, y)$  converges to  $\pi(y)$ .
- ▶ The next question is convergence of  $\pi_n$

$$\lim_{n \rightarrow \infty} \pi_n(y) = \lim_{n \rightarrow \infty} \sum_x \pi_0(x) P^n(x, y) = ?$$

- ▶ Consider an irreducible positive recurrent chain.
- ▶ It has a unique stationary distribution and  $\frac{1}{n} \sum_{m=1}^n P^m(x, y)$  converges to  $\pi(y)$ .
- ▶ The next question is convergence of  $\pi_n$

$$\lim_{n \rightarrow \infty} \pi_n(y) = \lim_{n \rightarrow \infty} \sum_x \pi_0(x) P^n(x, y) = ?$$

- ▶ If  $P^n(x, y)$  converges to  $g(y)$  then that would be the stationary distribution and  $\pi_n$  converges to it



- ▶ Consider an irreducible positive recurrent chain.
- ▶ It has a unique stationary distribution and  $\frac{1}{n} \sum_{m=1}^n P^m(x, y)$  converges to  $\pi(y)$ .
- ▶ The next question is convergence of  $\pi_n$

$$\lim_{n \rightarrow \infty} \pi_n(y) = \lim_{n \rightarrow \infty} \sum_x \pi_0(x) P^n(x, y) = ?$$

- ▶ If  $P^n(x, y)$  converges to  $g(y)$  then that would be the stationary distribution and  $\pi_n$  converges to it
- ▶ But,  $\frac{1}{n} \sum_{m=1}^n a_m$  may have a limit though  $\lim_{n \rightarrow \infty} a_n$  may not exist.

- ▶ Consider an irreducible positive recurrent chain.
- ▶ It has a unique stationary distribution and  $\frac{1}{n} \sum_{m=1}^n P^m(x, y)$  converges to  $\pi(y)$ .
- ▶ The next question is convergence of  $\pi_n$

$$\lim_{n \rightarrow \infty} \pi_n(y) = \lim_{n \rightarrow \infty} \sum_x \pi_0(x) P^n(x, y) = ?$$

- ▶ If  $P^n(x, y)$  converges to  $g(y)$  then that would be the stationary distribution and  $\pi_n$  converges to it
- ▶ But,  $\frac{1}{n} \sum_{m=1}^n a_m$  may have a limit though  $\lim_{n \rightarrow \infty} a_n$  may not exist.  
For example,  $a_n = (-1)^n$

- Consider a chain with transition probabilities

$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- Consider a chain with transition probabilities

$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


- One can show  $\pi^T = [\frac{1}{8} \ \frac{3}{8} \ \frac{3}{8} \ \frac{1}{8}]$

- Consider a chain with transition probabilities

$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


- One can show  $\pi^T = [\frac{1}{8} \ \frac{3}{8} \ \frac{3}{8} \ \frac{1}{8}]$
- However,  $P^n$  goes to different limits based on whether  $n$  is even or odd

- The chain is the following



$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$


- The chain is the following



$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- We can return to a state only after even number of time steps

- ▶ The chain is the following




$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- ▶ We can return to a state only after even number of time steps
- ▶ That is why  $P^n$  does not go to a limit



- ▶ The chain is the following



$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1/3 & 0 & 2/3 & 0 \\ 0 & 2/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- ▶ We can return to a state only after even number of time steps
- ▶ That is why  $P^n$  does not go to a limit
- ▶ Such a chain is called a periodic chain

- We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$

- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$
- ▶ If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$

- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$
- ▶ If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- ▶ Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .

- We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- If  $P(x, x) > 0$  then  $d_x = 1$
- If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .
- For any  $n$  s.t.  $P^n(y, y) > 0$ , we get  $P^{n_1+n+n_2}(x, x) > 0$

- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$
- ▶ If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- ▶ Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .
- ▶ For any  $n$  s.t.  $P^n(y, y) > 0$ , we get  $P^{n_1+n+n_2}(x, x) > 0$
- ▶ Hence,  $d_x$  divides  $n$  for all  $n$  s.t.  $P^n(y, y) > 0 \Rightarrow d_x \leq d_y$

- We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- If  $P(x, x) > 0$  then  $d_x = 1$
- If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .
- For any  $n$  s.t.  $P^n(y, y) > 0$ , we get  $P^{n_1+n+n_2}(x, x) > 0$
- Hence,  $d_x$  divides  $n$  for all  $n$  s.t.  $P^n(y, y) > 0 \Rightarrow d_x \leq d_y$
- Similarly,  $d_y \leq d_x$  and hence  $d_y = d_x$



- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$
- ▶ If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- ▶ Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .
- ▶ For any  $n$  s.t.  $P^n(y, y) > 0$ , we get  $P^{n_1+n+n_2}(x, x) > 0$
- ▶ Hence,  $d_x$  divides  $n$  for all  $n$  s.t.  $P^n(y, y) > 0 \Rightarrow d_x \leq d_y$
- ▶ Similarly,  $d_y \leq d_x$  and hence  $d_y = d_x$
- ▶ All states in an irreducible chain have the same period.

- ▶ We define period of a state  $x$  as

$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

- ▶ If  $P(x, x) > 0$  then  $d_x = 1$
- ▶ If  $x$  leads to  $y$  and  $y$  leads to  $x$ , then  $d_x = d_y$
- ▶ Let  $P^{n_1}(x, y) > 0$ ,  $P^{n_2}(y, x) > 0$ . Then  
 $P^{n_1+n_2}(x, x) > 0 \Rightarrow d_x$  divides  $n_1 + n_2$ .
- ▶ For any  $n$  s.t.  $P^n(y, y) > 0$ , we get  $P^{n_1+n+n_2}(x, x) > 0$
- ▶ Hence,  $d_x$  divides  $n$  for all  $n$  s.t.  $P^n(y, y) > 0 \Rightarrow d_x \leq d_y$
- ▶ Similarly,  $d_y \leq d_x$  and hence  $d_y = d_x$
- ▶ All states in an irreducible chain have the same period.
- ▶ If the period is 1 then chain is called aperiodic

- ▶ The extra condition we need for convergence of  $\pi_n$  is aperiodicity

- ▶ The extra condition we need for convergence of  $\pi_n$  is aperiodicity
- ▶ For an aperiodic, irreducible, positive recurrent chain, there is a unique stationary distribution and  $\pi_n$  converges to it irrespective of what  $\pi_0$  is.

- ▶ The extra condition we need for convergence of  $\pi_n$  is aperiodicity
- ▶ For an aperiodic, irreducible, positive recurrent chain, there is a unique stationary distribution and  $\pi_n$  converges to it irrespective of what  $\pi_0$  is.
- ▶ An aperiodic, irreducible, positive recurrent chain is called an ergodic chain