

# Recap: Expectation

## Recap: Expectation

- ▶ Let  $X$  be a discrete rv with  $X \in \{x_1, x_2, \dots\}$ . Then

$$E[X] = \sum_i x_i f_X(x_i)$$

## Recap: Expectation

- ▶ Let  $X$  be a discrete rv with  $X \in \{x_1, x_2, \dots\}$ . Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If  $X$  is a continuous random variable with pdf,  $f_X$ ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

# Recap: Expectation

- ▶ Let  $X$  be a discrete rv with  $X \in \{x_1, x_2, \dots\}$ . Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If  $X$  is a continuous random variable with pdf,  $f_X$ ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

# Recap: Expectation

- ▶ Let  $X$  be a discrete rv with  $X \in \{x_1, x_2, \dots\}$ . Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If  $X$  is a continuous random variable with pdf,  $f_X$ ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

- ▶ We take the expectation to exist when the sum or integral above is absolutely convergent

# Recap: Expectation

- ▶ Let  $X$  be a discrete rv with  $X \in \{x_1, x_2, \dots\}$ . Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If  $X$  is a continuous random variable with pdf,  $f_X$ ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

- ▶ We take the expectation to exist when the sum or integral above is absolutely convergent
- ▶ Note that expectation is defined for all random variables

## Recap: Expectation of a function of a random variable

- ▶ Let  $X$  be a rv and let  $Y = g(X)$ . Then,

# Recap: Expectation of a function of a random variable

- ▶ Let  $X$  be a rv and let  $Y = g(X)$ . Then,
- ▶  $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$



# Recap: Expectation of a function of a random variable

- ▶ Let  $X$  be a rv and let  $Y = g(X)$ . Then,
- ▶  $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$
- ▶ That is, if  $X$  is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

# Recap: Expectation of a function of a random variable

- ▶ Let  $X$  be a rv and let  $Y = g(X)$ . Then,
- ▶  $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$
- ▶ That is, if  $X$  is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

- ▶ If  $X$  and  $Y$  are continuous

$$EY = \int y f_Y(y) \, dy = \int g(x) f_X(x) \, dx$$

# Recap: Expectation of a function of a random variable

- ▶ Let  $X$  be a rv and let  $Y = g(X)$ . Then,
- ▶  $EY = \int y dF_Y(y) = \int g(x) dF_X(x)$
- ▶ That is, if  $X$  is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

- ▶ If  $X$  and  $Y$  are continuous

$$EY = \int y f_Y(y) dy = \int g(x) f_X(x) dx$$

- ▶ This is true for all rv's.

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- If  $X \geq 0$  then  $EX \geq 0$

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If  $X \geq 0$  then  $EX \geq 0$
- ▶  $E[b] = b$  where  $b$  is a constant

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If  $X \geq 0$  then  $EX \geq 0$
- ▶  $E[b] = b$  where  $b$  is a constant
- ▶  $E[ag(X)] = aE[g(X)]$  where  $a$  is a constant

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If  $X \geq 0$  then  $EX \geq 0$
- ▶  $E[b] = b$  where  $b$  is a constant
- ▶  $E[ag(X)] = aE[g(X)]$  where  $a$  is a constant
- ▶  $E[aX + b] = aE[X] + b$  where  $a, b$  are constants.



# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If  $X \geq 0$  then  $EX \geq 0$
- ▶  $E[b] = b$  where  $b$  is a constant
- ▶  $E[ag(X)] = aE[g(X)]$  where  $a$  is a constant
- ▶  $E[aX + b] = aE[X] + b$  where  $a, b$  are constants.
- ▶  $E[ag_1(X) + bg_2(X)] = aE[g_1(X)] + bE[g_2(X)]$

# Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If  $X \geq 0$  then  $EX \geq 0$
- ▶  $E[b] = b$  where  $b$  is a constant
- ▶  $E[ag(X)] = aE[g(X)]$  where  $a$  is a constant
- ▶  $E[aX + b] = aE[X] + b$  where  $a, b$  are constants.
- ▶  $E[ag_1(X) + bg_2(X)] = aE[g_1(X)] + bE[g_2(X)]$
- ▶  $E[(X - c)^2] \geq E[(X - EX)^2], \forall c$

## Recap: Variance of random variable

$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

## Recap: Variance of random variable

$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

- Properties of Variance:

# Recap: Variance of random variable

$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

- ▶ Properties of Variance:

- ▶  $\text{Var}(X) \geq 0$

# Recap: Variance of random variable

$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

- ▶ Properties of Variance:
  - ▶  $\text{Var}(X) \geq 0$
  - ▶  $\text{Var}(X + c) = \text{Var}(X)$

# Recap: Variance of random variable

$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

► Properties of Variance:

- $\text{Var}(X) \geq 0$
- $\text{Var}(X + c) = \text{Var}(X)$
- $\text{Var}(cX) = c^2 \text{Var}(X)$

# Recap: Moments of a random variable

- ▶ The  $k^{th}$  (order) moment of  $X$  is

$$m_k = E[X^k] = \int x^k dF_X(x)$$



# Recap: Moments of a random variable

- ▶ The  $k^{th}$  (order) moment of  $X$  is

$$m_k = E[X^k] = \int x^k dF_X(x)$$

- ▶ The  $k^{th}$  central moment of  $X$  is

$$s_k = E[(X - EX)^k] = \int (x - EX)^k dF_X(x)$$

# Recap: Moments of a random variable

- ▶ The  $k^{th}$  (order) moment of  $X$  is

$$m_k = E[X^k] = \int x^k dF_X(x)$$

- ▶ The  $k^{th}$  central moment of  $X$  is

$$s_k = E[(X - EX)^k] = \int (x - EX)^k dF_X(x)$$

- ▶ If moment of order  $k$  is finite then so is moment of order  $s$  for  $s < k$ .

# Moment generating function

- ▶ The moment generating function (mgf) of rv  $X$ ,  $M_X : \Re \rightarrow \Re$ , is defined by

$$M_X(t) = Ee^{tX}$$

# Moment generating function

- ▶ The moment generating function (mgf) of rv  $X$ ,  $M_X : \Re \rightarrow \Re$ , is defined by

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

# Moment generating function

- ▶ The moment generating function (mgf) of rv  $X$ ,  $M_X : \Re \rightarrow \Re$ , is defined by

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

- ▶ We say the mgf exists if  $E[e^{tX}] < \infty$  for  $t$  in some interval around zero

# Moment generating function

- ▶ The moment generating function (mgf) of rv  $X$ ,  $M_X : \Re \rightarrow \Re$ , is defined by

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

- ▶ We say the mgf exists if  $E[e^{tX}] < \infty$  for  $t$  in some interval around zero
- ▶ The mgf may not exist for some random variables.

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.



- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.
- ▶ Then we can get the moments of  $X$  by successive differentiation of  $M_X(t)$ .

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.
- ▶ Then we can get the moments of  $X$  by successive differentiation of  $M_X(t)$ .

$$\left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} E[e^{tX}] \right|_{t=0}$$

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.
- ▶ Then we can get the moments of  $X$  by successive differentiation of  $M_X(t)$ .

$$\left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} E[e^{tX}] \right|_{t=0} = E[Xe^{tX}] \Big|_{t=0}$$

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.
- ▶ Then we can get the moments of  $X$  by successive differentiation of  $M_X(t)$ .

$$\left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} E[e^{tX}] \right|_{t=0} = E[Xe^{tX}]|_{t=0} = EX$$

- ▶ The mgf of  $X$  is:  $M_X(t) = E[e^{tX}]$ .
- ▶ If  $M_X(t)$  exists (for  $t \in [-a, a]$  for some  $a > 0$ ) then all its derivatives also exist.
- ▶ Then we can get the moments of  $X$  by successive differentiation of  $M_X(t)$ .

$$\left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} E[e^{tX}] \right|_{t=0} = E[Xe^{tX}]|_{t=0} = EX$$

- ▶ In general

$$\left. \frac{d^k M_X(t)}{dt^k} \right|_{t=0} = E[X^k]$$

- ▶ We can easily see this by expanding  $e^{tX}$  in Taylor series:

- ▶ We can easily see this by expanding  $e^{tX}$  in Taylor series:

$$M_X(t) = Ee^{tX} = E \left[ 1 + \frac{tX}{1!} + \frac{t^2 X^2}{2!} + \frac{t^3 X^3}{3!} + \frac{t^4 X^4}{4!} + \dots \right]$$

- We can easily see this by expanding  $e^{tX}$  in Taylor series:

$$\begin{aligned}M_X(t) &= Ee^{tX} = E \left[ 1 + \frac{tX}{1!} + \frac{t^2X^2}{2!} + \frac{t^3X^3}{3!} + \frac{t^4X^4}{4!} + \dots \right] \\&= 1 + \frac{t}{1!}EX + \frac{t^2}{2!}EX^2 + \frac{t^3}{3!}EX^3 + \frac{t^4}{4!}EX^4 + \dots\end{aligned}$$



- ▶ We can easily see this by expanding  $e^{tX}$  in Taylor series:

$$\begin{aligned}M_X(t) &= Ee^{tX} = E \left[ 1 + \frac{tX}{1!} + \frac{t^2X^2}{2!} + \frac{t^3X^3}{3!} + \frac{t^4X^4}{4!} + \dots \right] \\&= 1 + \frac{t}{1!}EX + \frac{t^2}{2!}EX^2 + \frac{t^3}{3!}EX^3 + \frac{t^4}{4!}EX^4 + \dots\end{aligned}$$

- ▶ Now we can do term-wise differentiation.

- ▶ We can easily see this by expanding  $e^{tX}$  in Taylor series:

$$\begin{aligned}M_X(t) &= Ee^{tX} = E \left[ 1 + \frac{tX}{1!} + \frac{t^2 X^2}{2!} + \frac{t^3 X^3}{3!} + \frac{t^4 X^4}{4!} + \dots \right] \\&= 1 + \frac{t}{1!} EX + \frac{t^2}{2!} EX^2 + \frac{t^3}{3!} EX^3 + \frac{t^4}{4!} EX^4 + \dots\end{aligned}$$

- ▶ Now we can do term-wise differentiation. For example

$$\frac{d^3 M_X(t)}{dt^3} = 0 + 0 + 0 + \frac{3 * 2 * 1 * t^0}{3!} EX^3 + \frac{4 * 3 * 2 * t}{4!} EX^4 + \dots$$

- ▶ We can easily see this by expanding  $e^{tX}$  in Taylor series:

$$\begin{aligned} M_X(t) &= Ee^{tX} = E \left[ 1 + \frac{tX}{1!} + \frac{t^2 X^2}{2!} + \frac{t^3 X^3}{3!} + \frac{t^4 X^4}{4!} + \dots \right] \\ &= 1 + \frac{t}{1!} EX + \frac{t^2}{2!} EX^2 + \frac{t^3}{3!} EX^3 + \frac{t^4}{4!} EX^4 + \dots \end{aligned}$$

- ▶ Now we can do term-wise differentiation. For example

$$\frac{d^3 M_X(t)}{dt^3} = 0 + 0 + 0 + \frac{3 * 2 * 1 * t^0}{3!} EX^3 + \frac{4 * 3 * 2 * t}{4!} EX^4 + \dots$$

- ▶ Hence we get

$$\left. \frac{d^3 M_X(t)}{dt^3} \right|_{t=0} = E[X^3]$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$M_X(t) = E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda}$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \end{aligned}$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} \end{aligned}$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$



## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

► Now, by differentiating it we can find  $EX$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

► Now, by differentiating it we can find  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0}$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

► Now, by differentiating it we can find  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = e^{\lambda(e^t - 1)} \lambda e^t \Big|_{t=0}$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

► Now, by differentiating it we can find  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = e^{\lambda(e^t - 1)} \lambda e^t \Big|_{t=0} = \lambda$$

## Example – Moment generating function for Poisson

►  $f_X(k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad k = 0, 1, \dots$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{k=0}^{\infty} e^{tk} \frac{\lambda^k}{k!} e^{-\lambda} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{1}{k!} (\lambda e^t)^k \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

► Now, by differentiating it we can find  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = e^{\lambda(e^t - 1)} \lambda e^t \Big|_{t=0} = \lambda$$

(Exercise: Differentiate it twice to find  $EX^2$  and hence show that variance is  $\lambda$ ).

## mgf of exponential rv

- ▶  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$M_X(t) = E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx$$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \int_0^\infty e^{tx} \lambda e^{-\lambda x} dx \\ &= \int_0^\infty \lambda e^{-x(\lambda-t)} dx \end{aligned}$$



## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \int_0^\infty e^{tx} \lambda e^{-\lambda x} dx \\ &= \int_0^\infty \lambda e^{-x(\lambda-t)} dx \end{aligned}$$

This is finite if  $t < \lambda$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$M_X(t) = E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx$$

$$= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx$$

This is finite if  $t < \lambda$

$$= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty}$$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$M_X(t) = E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx$$

$$= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx$$

This is finite if  $t < \lambda$

$$= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty}$$

$$= \frac{\lambda}{\lambda-t}, \quad t < \lambda$$

## mgf of exponential rv

- ▶  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx \\ &= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx \end{aligned}$$

This is finite if  $t < \lambda$

$$\begin{aligned} &= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty} \\ &= \frac{\lambda}{\lambda-t}, \quad t < \lambda \end{aligned}$$

- ▶ We can use this to compute  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0}$$

## mgf of exponential rv

- ▶  $f_X(x) = \lambda e^{-\lambda x}$ ,  $x > 0$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx \\ &= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx \end{aligned}$$

This is finite if  $t < \lambda$

$$\begin{aligned} &= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty} \\ &= \frac{\lambda}{\lambda-t}, \quad t < \lambda \end{aligned}$$

- ▶ We can use this to compute  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} \left( \frac{\lambda}{\lambda-t} \right) \right|_{t=0}$$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$M_X(t) = E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx$$

$$= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx$$

This is finite if  $t < \lambda$

$$= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty}$$

$$= \frac{\lambda}{\lambda-t}, \quad t < \lambda$$

► We can use this to compute  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} \left( \frac{\lambda}{\lambda-t} \right) \right|_{t=0} = \left. \frac{\lambda}{(\lambda-t)^2} \right|_{t=0}$$

## mgf of exponential rv

►  $f_X(x) = \lambda e^{-\lambda x}, x > 0$

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx \\ &= \int_0^{\infty} \lambda e^{-x(\lambda-t)} dx \end{aligned}$$

This is finite if  $t < \lambda$

$$\begin{aligned} &= \left. \frac{\lambda e^{-x(\lambda-t)}}{-(\lambda-t)} \right|_0^{\infty} \\ &= \frac{\lambda}{\lambda-t}, \quad t < \lambda \end{aligned}$$

► We can use this to compute  $EX$

$$EX = \left. \frac{dM_X(t)}{dt} \right|_{t=0} = \left. \frac{d}{dt} \left( \frac{\lambda}{\lambda-t} \right) \right|_{t=0} = \left. \frac{\lambda}{(\lambda-t)^2} \right|_{t=0} = \frac{1}{\lambda}$$

- For mgf to exist we need  $E[e^{tX}] < \infty$  for  $t \in [-a, a]$  for some  $a > 0$ .



- ▶ For mgf to exist we need  $E[e^{tX}] < \infty$  for  $t \in [-a, a]$  for some  $a > 0$ .
- ▶ If  $M_X(t)$  exists then all moments of  $X$  are finite.

- ▶ For mgf to exist we need  $E[e^{tX}] < \infty$  for  $t \in [-a, a]$  for some  $a > 0$ .
- ▶ If  $M_X(t)$  exists then all moments of  $X$  are finite.
- ▶ However, all moments may be finite but the mgf may not exist.

- ▶ For mgf to exist we need  $E[e^{tX}] < \infty$  for  $t \in [-a, a]$  for some  $a > 0$ .
- ▶ If  $M_X(t)$  exists then all moments of  $X$  are finite.
- ▶ However, all moments may be finite but the mgf may not exist.
- ▶ When mgf exists, it uniquely determines the df

- ▶ For mgf to exist we need  $E[e^{tX}] < \infty$  for  $t \in [-a, a]$  for some  $a > 0$ .
- ▶ If  $M_X(t)$  exists then all moments of  $X$  are finite.
- ▶ However, all moments may be finite but the mgf may not exist.
- ▶ When mgf exists, it uniquely determines the df
- ▶ We are not saying moments uniquely determine the distribution; we are saying mgf uniquely determines the distribution

# Characteristic Function

- ▶ The characteristic function of  $X$  is defined by

$$\phi_X(t) = E[e^{itX}] = \int e^{itx} dF_X(x) \quad (i = \sqrt{-1})$$

# Characteristic Function

- ▶ The characteristic function of  $X$  is defined by

$$\phi_X(t) = E[e^{itX}] = \int e^{itx} dF_X(x) \quad (i = \sqrt{-1})$$

- ▶ If  $X$  is continuous rv,

$$\phi_X(t) = E[e^{itX}] = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx$$

# Characteristic Function

- ▶ The characteristic function of  $X$  is defined by

$$\phi_X(t) = E[e^{itX}] = \int e^{itx} dF_X(x) \quad (i = \sqrt{-1})$$

- ▶ If  $X$  is continuous rv,

$$\phi_X(t) = E[e^{itX}] = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx$$

- ▶ Characteristic function always exists because  $|e^{itx}| = 1, \forall t, x$

# Characteristic Function

- ▶ The characteristic function of  $X$  is defined by

$$\phi_X(t) = E[e^{itX}] = \int e^{itx} dF_X(x) \quad (i = \sqrt{-1})$$

- ▶ If  $X$  is continuous rv,

$$\phi_X(t) = E[e^{itX}] = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx$$

- ▶ Characteristic function always exists because  
 $|e^{itx}| = 1, \forall t, x$

- ▶ For example,

$$\left| \int_{-\infty}^{\infty} e^{itx} f_X(x) dx \right| \leq \int_{-\infty}^{\infty} |e^{itx}| |f_X(x)| dx = \int_{-\infty}^{\infty} f_X(x) dx = 1$$



# Characteristic Function

- ▶ The characteristic function of  $X$  is defined by

$$\phi_X(t) = E[e^{itX}] = \int e^{itx} dF_X(x) \quad (i = \sqrt{-1})$$

- ▶ If  $X$  is continuous rv,

$$\phi_X(t) = E[e^{itX}] = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx$$

- ▶ Characteristic function always exists because  $|e^{itx}| = 1, \forall t, x$

- ▶ For example,

$$\left| \int_{-\infty}^{\infty} e^{itx} f_X(x) dx \right| \leq \int_{-\infty}^{\infty} |e^{itx}| |f_X(x)| dx = \int_{-\infty}^{\infty} f_X(x) dx = 1$$

- ▶ We would consider  $\phi_X$  later in the course

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k, \quad s \in \mathbb{R}$$

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k)s^k, \quad s \in \mathbb{R}$$

- ▶ This infinite sum converges (absolutely) for  $|s| \leq 1$ .

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k, \quad s \in \mathbb{R}$$

- ▶ This infinite sum converges (absolutely) for  $|s| \leq 1$ .
- ▶ We have

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k)s^k, \quad s \in \mathbb{R}$$

- ▶ This infinite sum converges (absolutely) for  $|s| \leq 1$ .
- ▶ We have

$$P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$$

# Generating function

- ▶ Let  $X \in \{0, 1, 2, \dots\}$
- ▶ The (probability) generating function of  $X$  is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k)s^k, \quad s \in \mathbb{R}$$

- ▶ This infinite sum converges (absolutely) for  $|s| \leq 1$ .
- ▶ We have

$$P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$$

- ▶ The pmf can be obtained from the generating function



►  $P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$

- ▶  $P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$
- ▶ Let  $P'_X(s) \triangleq \frac{dP_X(s)}{ds}$  and so on

- ▶  $P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$
- ▶ Let  $P'_X(s) \triangleq \frac{dP_X(s)}{ds}$  and so on
- ▶ We get

$$P'_X(s) = 0 + f_X(1) + f_X(2) 2s + f_X(3) 3s^2 + \dots$$

- ▶  $P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$
- ▶ Let  $P'_X(s) \triangleq \frac{dP_X(s)}{ds}$  and so on
- ▶ We get

$$P'_X(s) = 0 + f_X(1) + f_X(2) 2s + f_X(3) 3s^2 + \dots$$

$$P''_X(s) = 0 + 0 + f_X(2) 2 * 1 + f_X(3) 3 * 2s^1 + \dots$$

- ▶  $P_X(s) = f_X(0) + f_X(1)s + f_X(2)s^2 + f_X(3)s^3 + \dots$
- ▶ Let  $P'_X(s) \triangleq \frac{dP_X(s)}{ds}$  and so on
- ▶ We get

$$P'_X(s) = 0 + f_X(1) + f_X(2) 2s + f_X(3) 3s^2 + \dots$$

$$P''_X(s) = 0 + 0 + f_X(2) 2 * 1 + f_X(3) 3 * 2s^1 + \dots$$

Hence, we get

$$f_X(0) = P_X(0); f_X(1) = \frac{P'_X(0)}{1!}; f_X(2) = \frac{P''_X(0)}{2!}$$

- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k)s^k$

- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k$

$$P'_X(s) = \sum_{k=0}^{\infty} k f_X(k) s^{k-1}$$

- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k$

$$P'_X(s) = \sum_{k=0}^{\infty} k f_X(k) s^{k-1} \Rightarrow P'_X(1) = EX$$



- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k$

$$P'_X(s) = \sum_{k=0}^{\infty} k f_X(k) s^{k-1} \Rightarrow P'_X(1) = EX$$

$$P''_X(s) = \sum_{k=0}^{\infty} k(k-1) f_X(k) s^{k-2}$$

- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k$

$$P'_X(s) = \sum_{k=0}^{\infty} k f_X(k) s^{k-1} \Rightarrow P'_X(1) = EX$$

$$P''_X(s) = \sum_{k=0}^{\infty} k(k-1) f_X(k) s^{k-2} \Rightarrow P''_X(1) = E[X(X-1)]$$

- ▶ The moments (when they exist) can be obtained from the generating function:  $P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k$

$$P'_X(s) = \sum_{k=0}^{\infty} k f_X(k) s^{k-1} \Rightarrow P'_X(1) = EX$$

$$P''_X(s) = \sum_{k=0}^{\infty} k(k-1) f_X(k) s^{k-2} \Rightarrow P''_X(1) = E[X(X-1)]$$

- ▶ For (positive integer valued) discrete random variables, it is more convenient to deal with generating functions than mgf.

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, k = 0, 1, \dots, n$

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, k = 0, 1, \dots, n$

$$P_X(s) = \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k$$

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$

$$\begin{aligned} P_X(s) &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} (sp)^k (1-p)^{n-k} \end{aligned}$$

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, k = 0, 1, \dots, n$

$$\begin{aligned} P_X(s) &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} (sp)^k (1-p)^{n-k} \\ &= (sp + (1-p))^n = (1 + p(s-1))^n \end{aligned}$$

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, k = 0, 1, \dots, n$

$$\begin{aligned} P_X(s) &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} (sp)^k (1-p)^{n-k} \\ &= (sp + (1-p))^n = (1 + p(s-1))^n \end{aligned}$$

► From the above, we get  $P'_X(s) = n(sp + (1-p))^{n-1}p$



## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$

$$\begin{aligned} P_X(s) &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} (sp)^k (1-p)^{n-k} \\ &= (sp + (1-p))^n = (1 + p(s-1))^n \end{aligned}$$

► From the above, we get  $P'_X(s) = n(sp + (1-p))^{n-1}p$

► Thus,

$$EX = P'_X(1) = np;$$

## Example – Generating function for binomial rv

►  $f_X(k) = \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k}, \quad k = 0, 1, \dots, n$

$$\begin{aligned} P_X(s) &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} s^k \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} (sp)^k (1-p)^{n-k} \\ &= (sp + (1-p))^n = (1 + p(s-1))^n \end{aligned}$$

► From the above, we get  $P'_X(s) = n(sp + (1-p))^{n-1}p$

► Thus,

$$EX = P'_X(1) = np; \quad f_X(1) = P'_X(0) = n(1-p)^{n-1}p$$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$
  - ▶  $1 - p \leq 1 - P[X < x]$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$
  - ▶  $1 - p \leq 1 - P[X < x] = 1 - (P[X \leq x] - P[X = x])$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$
  - ▶  $1 - p \leq 1 - P[X < x] = 1 - (P[X \leq x] - P[X = x])$   
 $\Rightarrow 1 - p \leq 1 - F_X(x) + P[X = x]$



- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have

- ▶  $p \leq P[X \leq x] = F_X(x)$
- ▶  $1 - p \leq 1 - P[X < x] = 1 - (P[X \leq x] - P[X = x])$   
 $\Rightarrow 1 - p \leq 1 - F_X(x) + P[X = x]$   
 $\Rightarrow F_X(x) \leq p + P[X = x]$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$
  - ▶  $1 - p \leq 1 - P[X < x] = 1 - (P[X \leq x] - P[X = x])$   
 $\Rightarrow 1 - p \leq 1 - F_X(x) + P[X = x]$   
 $\Rightarrow F_X(x) \leq p + P[X = x]$
- ▶ Thus,  $x$  satisfies (if it is quantile of order  $p$ )

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ Let  $p \in (0, 1)$ . The number  $x \in \Re$  that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad P[X \geq x] \geq 1 - p$$

is called the quantile of order  $p$  or the  $100p^{th}$  percentile of rv  $X$ .

- ▶ Suppose  $x$  is a quantile of order  $p$ . Then we have
  - ▶  $p \leq P[X \leq x] = F_X(x)$
  - ▶  $1 - p \leq 1 - P[X < x] = 1 - (P[X \leq x] - P[X = x])$   
 $\Rightarrow 1 - p \leq 1 - F_X(x) + P[X = x]$   
 $\Rightarrow F_X(x) \leq p + P[X = x]$
- ▶ Thus,  $x$  satisfies (if it is quantile of order  $p$ )

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ Note that for a given  $p$  there can be multiple values for  $x$  to satisfy the above.

- ▶ If  $x$  is a quantile of order  $p$  then

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ If  $x$  is a quantile of order  $p$  then

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ If  $X$  is continuous rv, we need to satisfy  $p = F_X(x)$ .

- ▶ If  $x$  is a quantile of order  $p$  then

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ If  $X$  is continuous rv, we need to satisfy  $p = F_X(x)$ .
- ▶ In general, for a given  $p$ , there may be multiple  $x$  that satisfy the above.

- ▶ If  $x$  is a quantile of order  $p$  then

$$p \leq F_X(x) \leq p + P[X = x]$$


- ▶ If  $X$  is continuous rv, we need to satisfy  $p = F_X(x)$ .
- ▶ In general, for a given  $p$ , there may be multiple  $x$  that satisfy the above.
- ▶ Let us see some examples.

- ▶ Let  $X$  be continuous rv.



- ▶ Let  $X$  be continuous rv.
- ▶ If the df is strictly monotone then  $F_X(x) = p$  would have a unique solution.

- ▶ Let  $X$  be continuous rv.
- ▶ If the df is strictly monotone then  $F_X(x) = p$  would have a unique solution.




- ▶ For continuous rv,  $X$ ,  $F_X$  need not be strictly monotone.

- ▶ For continuous rv,  $X$ ,  $F_X$  need not be strictly monotone.
- ▶ Consider a pdf:  $f_X(x) = 0.5$ ,  $x \in [1, 2] \cup [3, 4]$


- ▶ For continuous rv,  $X$ ,  $F_X$  need not be strictly monotone.
- ▶ Consider a pdf:  $f_X(x) = 0.5$ ,  $x \in [1, 2] \cup [3, 4]$
- ▶ The pdf and the corresponding df are:

- ▶ For continuous rv,  $X$ ,  $F_X$  need not be strictly monotone.
- ▶ Consider a pdf:  $f_X(x) = 0.5$ ,  $x \in [1, 2] \cup [3, 4]$
- ▶ The pdf and the corresponding df are:



- For this df, for  $p = 0.5$ , the quantile of order  $p$  is not unique because there many  $x$  with  $F_X(x) = 0.5$   
But for  $p = 0.75$  it is unique.

- For this df, for  $p = 0.5$ , the quantile of order  $p$  is not unique because there many  $x$  with  $F_X(x) = 0.5$   
But for  $p = 0.75$  it is unique.





- ▶ Let  $X \in \{x_1, x_2, \dots\}$

- ▶ Let  $X \in \{x_1, x_2, \dots\}$
- ▶ Given a  $p$  we want to calculate quantile of order  $p$


- ▶ Let  $X \in \{x_1, x_2, \dots\}$
- ▶ Given a  $p$  we want to calculate quantile of order  $p$
- ▶ Suppose there is a  $x_i$  such that  $F_X(x_i) = p$ .

- ▶ Let  $X \in \{x_1, x_2, \dots\}$
- ▶ Given a  $p$  we want to calculate quantile of order  $p$
- ▶ Suppose there is a  $x_i$  such that  $F_X(x_i) = p$ .
- ▶ Then, for  $x_i \leq x < x_{i+1}$ ,  $F_X(x) = p$

- ▶ Let  $X \in \{x_1, x_2, \dots\}$
- ▶ Given a  $p$  we want to calculate quantile of order  $p$
- ▶ Suppose there is a  $x_i$  such that  $F_X(x_i) = p$ .
- ▶ Then, for  $x_i \leq x < x_{i+1}$ ,  $F_X(x) = p$
- ▶ For  $x_i \leq x \leq x_{i+1}$ , we have  $p \leq F_X(x) \leq p + P[X = x]$

- ▶ Let  $X \in \{x_1, x_2, \dots\}$
- ▶ Given a  $p$  we want to calculate quantile of order  $p$
- ▶ Suppose there is a  $x_i$  such that  $F_X(x_i) = p$ .
- ▶ Then, for  $x_i \leq x < x_{i+1}$ ,  $F_X(x) = p$
- ▶ For  $x_i \leq x \leq x_{i+1}$ , we have  $p \leq F_X(x) \leq p + P[X = x]$
- ▶ So, quantile of order  $p$  is not unique and all such  $x$  qualify.

- This situation is illustrated below



- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .



- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Hence,  $x_i$  is quantile of order  $p$ .

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Hence,  $x_i$  is quantile of order  $p$ .
- ▶ For any  $x < x_i$  we would have  $F_X(x) \leq F_X(x_{i-1}) < p$ .

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Hence,  $x_i$  is quantile of order  $p$ .
- ▶ For any  $x < x_i$  we would have  $F_X(x) \leq F_X(x_{i-1}) < p$ .
- ▶ For any  $x$ , with  $x_i < x < x_{i+1}$  we have  $p + P[X = x_i] = p < F_X(x) = p + \delta_2$ .

- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Hence,  $x_i$  is quantile of order  $p$ .
- ▶ For any  $x < x_i$  we would have  $F_X(x) \leq F_X(x_{i-1}) < p$ .
- ▶ For any  $x$ , with  $x_i < x < x_{i+1}$  we have  
 $p + P[X = x] = p < F_X(x) = p + \delta_2$ .
- ▶ Similarly, for  $x \geq x_{i+1}$  we have  $F_X(x) > p + P[X = x]$ .


- ▶ Now suppose  $p$  is such that  $F_X(x_{i-1}) < p < F_X(x_i)$ .
- ▶ Let  $F_X(x_{i-1}) = p - \delta_1$  and  $F_X(x_i) = p + \delta_2$ . (Note that  $\delta_1, \delta_2 > 0$ )
- ▶ Then  $P[X = x_i] = F_X(x_i) - F_X(x_{i-1}) = \delta_2 + \delta_1$
- ▶ Hence we have

$$p < p + \delta_2 = F_X(x_i) < p + \delta_2 + \delta_1 = p + P[X = x_i]$$

- ▶ Hence,  $x_i$  is quantile of order  $p$ .
- ▶ For any  $x < x_i$  we would have  $F_X(x) \leq F_X(x_{i-1}) < p$ .
- ▶ For any  $x$ , with  $x_i < x < x_{i+1}$  we have  $p + P[X = x] = p < F_X(x) = p + \delta_2$ .
- ▶ Similarly, for  $x \geq x_{i+1}$  we have  $F_X(x) > p + P[X = x]$ .
- ▶ Thus quantile of order  $p$  is unique here.



- This situation is illustrated below



# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.

# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.
- ▶ For a continuous rv, median,  $x$  satisfies:  $F_X(x) = 0.5$ .

# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.
- ▶ For a continuous rv, median,  $x$  satisfies:  $F_X(x) = 0.5$ .
- ▶ For a discrete rv, it satisfies:  
$$0.5 \leq F_X(x) \leq 0.5 + P[X = x].$$

# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.
- ▶ For a continuous rv, median,  $x$  satisfies:  $F_X(x) = 0.5$ .
- ▶ For a discrete rv, it satisfies:  
$$0.5 \leq F_X(x) \leq 0.5 + P[X = x].$$
- ▶ As we saw, median need not be unique.

# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.
- ▶ For a continuous rv, median,  $x$  satisfies:  $F_X(x) = 0.5$ .
- ▶ For a discrete rv, it satisfies:  
 $0.5 \leq F_X(x) \leq 0.5 + P[X = x]$ .
- ▶ As we saw, median need not be unique.
- ▶ Recall that the (standard) Cauchy density is given by

$$f_X(x) = \frac{1}{\pi} \frac{1}{1 + x^2}, \quad -\infty < x < \infty$$

# Median of a distribution

- ▶ For  $p = 0.5$  quantile of order  $p$  is called the median.
- ▶ For a continuous rv, median,  $x$  satisfies:  $F_X(x) = 0.5$ .
- ▶ For a discrete rv, it satisfies:  
 $0.5 \leq F_X(x) \leq 0.5 + P[X = x]$ .
- ▶ As we saw, median need not be unique.
- ▶ Recall that the (standard) Cauchy density is given by

$$f_X(x) = \frac{1}{\pi} \frac{1}{1 + x^2}, \quad -\infty < x < \infty$$

- ▶ One can show that  $\int_{-\infty}^0 f_X(x) dx = 0.5$  and hence the median is at the origin.

- If we want to find  $c$  to minimize  $E[(X - c)^2]$  then the solution is  $c = EX$ .



- ▶ If we want to find  $c$  to minimize  $E[(X - c)^2]$  then the solution is  $c = EX$ .
- ▶ We saw this earlier.

- ▶ If we want to find  $c$  to minimize  $E[(X - c)^2]$  then the solution is  $c = EX$ .
- ▶ We saw this earlier.
- ▶ Suppose we want to find  $c$  to minimize  $E[|(X - c)|]$

- ▶ If we want to find  $c$  to minimize  $E[(X - c)^2]$  then the solution is  $c = EX$ .
- ▶ We saw this earlier.
- ▶ Suppose we want to find  $c$  to minimize  $E[|(X - c)|]$
- ▶ Then we would get  $c$  to be the median.

- ▶ If we want to find  $c$  to minimize  $E[(X - c)^2]$  then the solution is  $c = EX$ .
- ▶ We saw this earlier.
- ▶ Suppose we want to find  $c$  to minimize  $E[|(X - c)|]$
- ▶ Then we would get  $c$  to be the median.  
(Exercise: Show this for discrete and continuous rv)

# Markov Inequality

# Markov Inequality

- ▶ Let  $g : \mathfrak{R} \rightarrow \mathfrak{R}$  be a non-negative function. Then

# Markov Inequality

- ▶ Let  $g : \mathfrak{R} \rightarrow \mathfrak{R}$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

# Markov Inequality

- ▶ Let  $g : \mathfrak{R} \rightarrow \mathfrak{R}$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv



# Markov Inequality

- ▶ Let  $g : \mathfrak{R} \rightarrow \mathfrak{R}$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \end{aligned}$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \\ &\geq \int_{g(x) > c} g(x) f_X(x) dx \end{aligned}$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \\ &\geq \int_{g(x) > c} g(x) f_X(x) dx \quad \text{because } g(x) \geq 0 \end{aligned}$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \\ &\geq \int_{g(x) > c} g(x) f_X(x) dx \quad \text{because } g(x) \geq 0 \\ &\geq c \int_{g(x) > c} f_X(x) dx \end{aligned}$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \\ &\geq \int_{g(x) > c} g(x) f_X(x) dx \quad \text{because } g(x) \geq 0 \\ &\geq c \int_{g(x) > c} f_X(x) dx = c P[g(X) > c] \end{aligned}$$

# Markov Inequality

- ▶ Let  $g : \Re \rightarrow \Re$  be a non-negative function. Then

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ **Proof:** We prove it for continuous rv. Proof is similar for discrete rv

$$\begin{aligned} E[g(X)] &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ &= \int_{g(x) \leq c} g(x) f_X(x) dx + \int_{g(x) > c} g(x) f_X(x) dx \\ &\geq \int_{g(x) > c} g(x) f_X(x) dx \quad \text{because } g(x) \geq 0 \\ &\geq c \int_{g(x) > c} f_X(x) dx = c P[g(X) > c] \end{aligned}$$

$$\text{Thus, } P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$



# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ In all such results an underlying assumption is that the expectation is finite.

# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ In all such results an underlying assumption is that the expectation is finite.
- ▶ Let  $g(x) = |x|^k$  where  $k$  is a positive integer. We have  $g(x) \geq 0, \forall x$ . Let  $c > 0$ .

# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ In all such results an underlying assumption is that the expectation is finite.
- ▶ Let  $g(x) = |x|^k$  where  $k$  is a positive integer. We have  $g(x) \geq 0, \forall x$ . Let  $c > 0$ .
- ▶ We know that  $|x| > c \Rightarrow |x|^k > c^k$  and vice versa.

# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ In all such results an underlying assumption is that the expectation is finite.
- ▶ Let  $g(x) = |x|^k$  where  $k$  is a positive integer. We have  $g(x) \geq 0, \forall x$ . Let  $c > 0$ .
- ▶ We know that  $|x| > c \Rightarrow |x|^k > c^k$  and vice versa.
- ▶ Now we get,

$$P[|X| > c] = P[|X|^k > c^k] \leq \frac{E[|X|^k]}{c^k}$$

# Markov Inequality

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}, \quad (c > 0)$$

- ▶ In all such results an underlying assumption is that the expectation is finite.
- ▶ Let  $g(x) = |x|^k$  where  $k$  is a positive integer. We have  $g(x) \geq 0, \forall x$ . Let  $c > 0$ .
- ▶ We know that  $|x| > c \Rightarrow |x|^k > c^k$  and vice versa.
- ▶ Now we get,

$$P[|X| > c] = P[|X|^k > c^k] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ Markov inequality is often used in this form.

# Chebyshev Inequality

- ▶ Markov Inequality:

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

# Chebyshev Inequality

- ▶ Markov Inequality:

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ Take  $|X|$  as  $|X - EX|$  and take  $k = 2$

# Chebyshev Inequality

- ▶ Markov Inequality:

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ Take  $|X|$  as  $|X - EX|$  and take  $k = 2$

$$P[|X - EX| > c] \leq \frac{E[|X - EX|^2]}{c^2}$$



# Chebyshev Inequality

- ▶ Markov Inequality:

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ Take  $|X|$  as  $|X - EX|$  and take  $k = 2$

$$P[|X - EX| > c] \leq \frac{E[|X - EX|^2]}{c^2} = \frac{\text{Var}(X)}{c^2}$$

# Chebyshev Inequality

- ▶ Markov Inequality:

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ Take  $|X|$  as  $|X - EX|$  and take  $k = 2$

$$P[|X - EX| > c] \leq \frac{E[|X - EX|^2]}{c^2} = \frac{\text{Var}(X)}{c^2}$$

- ▶ This is known as the Chebyshev inequality.

- ▶ The Chebyshev inequality is

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ The Chebyshev inequality is

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ Let  $EX = \mu$  and let  $\text{Var}(X) = \sigma^2$ . Take  $c = k\sigma$

- ▶ The Chebyshev inequality is

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ Let  $EX = \mu$  and let  $\text{Var}(X) = \sigma^2$ . Take  $c = k\sigma$
- ▶ We call,  $\sigma$ , square root of variance, as standard deviation.

- ▶ The Chebyshev inequality is

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ Let  $EX = \mu$  and let  $\text{Var}(X) = \sigma^2$ . Take  $c = k\sigma$
- ▶ We call,  $\sigma$ , square root of variance, as standard deviation.
- ▶ Now, Chebyshev inequality gives us

$$P[|X - \mu| > k\sigma] \leq \frac{\sigma^2}{k^2\sigma^2} = \frac{1}{k^2}$$

- ▶ The Chebyshev inequality is

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ Let  $EX = \mu$  and let  $\text{Var}(X) = \sigma^2$ . Take  $c = k\sigma$
- ▶ We call,  $\sigma$ , square root of variance, as standard deviation.
- ▶ Now, Chebyshev inequality gives us

$$P[|X - \mu| > k\sigma] \leq \frac{\sigma^2}{k^2\sigma^2} = \frac{1}{k^2}$$

- ▶ This is true for all random variables and the RHS above does not depend on the distribution of  $X$ .

- ▶ **Markov inequality:** For a non-negative function,  $g$ ,



- ▶ **Markov inequality:** For a non-negative function,  $g$ ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ **Markov inequality:** For a non-negative function,  $g$ ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ **Markov inequality:** For a non-negative function,  $g$ ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ **Chebyshev inequality**

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ **Markov inequality:** For a non-negative function,  $g$ ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ **Chebyshev inequality**

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ With  $EX = \mu$  and  $\text{Var}(X) = \sigma^2$ , we get

$$P[|X - \mu| > k\sigma] \leq \frac{1}{k^2}$$



# A pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$

# A pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ Each of  $X, Y$  maps  $\Omega$  to  $\mathbb{R}$ .


# A pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ Each of  $X, Y$  maps  $\Omega$  to  $\mathbb{R}$ .
- ▶ We can think of the pair of random variables as a vector-valued function that maps  $\Omega$  to  $\mathbb{R}^2$ .



# A pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ Each of  $X, Y$  maps  $\Omega$  to  $\mathbb{R}$ .
- ▶ We can think of the pair of random variables as a vector-valued function that maps  $\Omega$  to  $\mathbb{R}^2$ .



- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.

- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.
- ▶ That is, by defining the pair of random variables, we essentially create a new probability space with sample space being  $\mathbb{R}^2$ .

- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.
- ▶ That is, by defining the pair of random variables, we essentially create a new probability space with sample space being  $\mathbb{R}^2$ .
- ▶ The events now would be the Borel subsets of  $\mathbb{R}^2$ .

- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.
- ▶ That is, by defining the pair of random variables, we essentially create a new probability space with sample space being  $\mathbb{R}^2$ .
- ▶ The events now would be the Borel subsets of  $\mathbb{R}^2$ .
- ▶ Recall that  $\mathbb{R}^2$  is cartesian product of  $\mathbb{R}$  with itself.

- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.
- ▶ That is, by defining the pair of random variables, we essentially create a new probability space with sample space being  $\mathbb{R}^2$ .
- ▶ The events now would be the Borel subsets of  $\mathbb{R}^2$ .
- ▶ Recall that  $\mathbb{R}^2$  is cartesian product of  $\mathbb{R}$  with itself.
- ▶ So, we can create Borel subsets of  $\mathbb{R}^2$  by cartesian product of Borel subsets of  $\mathbb{R}$ .

- ▶ Just as in the case of a single rv, we can think of the induced probability space for the case of a pair of rv's too.
- ▶ That is, by defining the pair of random variables, we essentially create a new probability space with sample space being  $\mathbb{R}^2$ .
- ▶ The events now would be the Borel subsets of  $\mathbb{R}^2$ .
- ▶ Recall that  $\mathbb{R}^2$  is cartesian product of  $\mathbb{R}$  with itself.
- ▶ So, we can create Borel subsets of  $\mathbb{R}^2$  by cartesian product of Borel subsets of  $\mathbb{R}$ .

$$\mathcal{B}^2 = \sigma(\{B_1 \times B_2 : B_1, B_2 \in \mathcal{B}\})$$


where  $\mathcal{B}$  is the Borel  $\sigma$ -algebra we considered earlier, and  $\mathcal{B}^2$  is the set of Borel sets of  $\mathbb{R}^2$ .

- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.




- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.
- ▶ Let  $I_1, I_2 \subset \mathbb{R}$  be intervals. Then  $I_1 \times I_2 \subset \mathbb{R}^2$  is known as a cylindrical set.

- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.
- ▶ Let  $I_1, I_2 \subset \mathbb{R}$  be intervals. Then  $I_1 \times I_2 \subset \mathbb{R}^2$  is known as a cylindrical set.




- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.
- ▶ Let  $I_1, I_2 \subset \mathbb{R}$  be intervals. Then  $I_1 \times I_2 \subset \mathbb{R}^2$  is known as a cylindrical set.




- ▶  $\mathcal{B}^2$  is the smallest  $\sigma$ -algebra containing all cylindrical sets.

- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.
- ▶ Let  $I_1, I_2 \subset \mathbb{R}$  be intervals. Then  $I_1 \times I_2 \subset \mathbb{R}^2$  is known as a cylindrical set.



- ▶  $\mathcal{B}^2$  is the smallest  $\sigma$ -algebra containing all cylindrical sets.
- ▶ We saw that  $\mathcal{B}$  is also the smallest  $\sigma$ -algebra containing all intervals of the form  $(-\infty, x]$ .

- ▶ Recall that  $\mathcal{B}$  is the smallest  $\sigma$ -algebra containing all intervals.
- ▶ Let  $I_1, I_2 \subset \mathbb{R}$  be intervals. Then  $I_1 \times I_2 \subset \mathbb{R}^2$  is known as a cylindrical set.



- ▶  $\mathcal{B}^2$  is the smallest  $\sigma$ -algebra containing all cylindrical sets.
- ▶ We saw that  $\mathcal{B}$  is also the smallest  $\sigma$ -algebra containing all intervals of the form  $(-\infty, x]$ .
- ▶ Similarly  $\mathcal{B}^2$  is the smallest  $\sigma$ -algebra containing cylindrical sets of the form  $(-\infty, x] \times (-\infty, y]$ .

- ▶ Let  $X, Y$  be random variables on the probability space  $(\Omega, \mathcal{F}, P)$

- ▶ Let  $X, Y$  be random variables on the probability space  $(\Omega, \mathcal{F}, P)$
- ▶ This gives rise to a new probability space  $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$  with  $P_{XY}$  given by

- ▶ Let  $X, Y$  be random variables on the probability space  $(\Omega, \mathcal{F}, P)$
- ▶ This gives rise to a new probability space  $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$  with  $P_{XY}$  given by

$$P_{XY}(B) = P[(X, Y) \in B], \forall B \in \mathcal{B}^2$$



- ▶ Let  $X, Y$  be random variables on the probability space  $(\Omega, \mathcal{F}, P)$
- ▶ This gives rise to a new probability space  $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$  with  $P_{XY}$  given by

$$\begin{aligned} P_{XY}(B) &= P[(X, Y) \in B], \forall B \in \mathcal{B}^2 \\ &= P(\{\omega : (X(\omega), Y(\omega)) \in B\}) \end{aligned}$$

- ▶ Let  $X, Y$  be random variables on the probability space  $(\Omega, \mathcal{F}, P)$
- ▶ This gives rise to a new probability space  $(\mathfrak{R}^2, \mathcal{B}^2, P_{XY})$  with  $P_{XY}$  given by

$$\begin{aligned} P_{XY}(B) &= P[(X, Y) \in B], \forall B \in \mathcal{B}^2 \\ &= P(\{\omega : (X(\omega), Y(\omega)) \in B\}) \end{aligned}$$

- ▶ Recall that for a single rv, the resulting probability space is  $(\mathfrak{R}, \mathcal{B}, P_X)$  with

$$P_X(B) = P[X \in B] = P(\{\omega : X(\omega) \in B\})$$

- ▶ In the case of a single rv, we define a distribution function,  $F_X$  which essentially assigns probability to all intervals of the form  $(-\infty, x]$ .

- ▶ In the case of a single rv, we define a distribution function,  $F_X$  which essentially assigns probability to all intervals of the form  $(-\infty, x]$ .
- ▶ This  $F_X$  uniquely determines  $P_X(B)$  for all Borel sets,  $B$ .

- ▶ In the case of a single rv, we define a distribution function,  $F_X$  which essentially assigns probability to all intervals of the form  $(-\infty, x]$ .
- ▶ This  $F_X$  uniquely determines  $P_X(B)$  for all Borel sets,  $B$ .
- ▶ In a similar manner we define a joint distribution function  $F_{XY}$  for a pair of random variables.

- ▶ In the case of a single rv, we define a distribution function,  $F_X$  which essentially assigns probability to all intervals of the form  $(-\infty, x]$ .
- ▶ This  $F_X$  uniquely determines  $P_X(B)$  for all Borel sets,  $B$ .
- ▶ In a similar manner we define a joint distribution function  $F_{XY}$  for a pair of random variables.
- ▶  $F_{XY}(x, y)$  would be  $P_{XY}((-\infty, x] \times (-\infty, y])$ .

- ▶ In the case of a single rv, we define a distribution function,  $F_X$  which essentially assigns probability to all intervals of the form  $(-\infty, x]$ .
- ▶ This  $F_X$  uniquely determines  $P_X(B)$  for all Borel sets,  $B$ .
- ▶ In a similar manner we define a joint distribution function  $F_{XY}$  for a pair of random variables.
- ▶  $F_{XY}(x, y)$  would be  $P_{XY}((-\infty, x] \times (-\infty, y])$ .
- ▶  $F_{XY}$  fixes the probability of all cylindrical sets of the form  $(-\infty, x] \times (-\infty, y]$  and hence uniquely determines the probability of all Borel sets of  $\mathbb{R}^2$ .

# Joint distribution of a pair of random variables



# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$

# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ The joint distribution function of  $X, Y$  is  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ The joint distribution function of  $X, Y$  is  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ The joint distribution function of  $X, Y$  is  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

$$F_{XY}(x, y) = P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y]))$$

# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ The joint distribution function of  $X, Y$  is  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y])) \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

# Joint distribution of a pair of random variables

- ▶ Let  $X, Y$  be random variables on the same probability space  $(\Omega, \mathcal{F}, P)$
- ▶ The joint distribution function of  $X, Y$  is  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y])) \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

- ▶ The joint distribution function is the probability of the intersection of the events  $[X \leq x]$  and  $[Y \leq y]$ .

# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$   
(These are actually limits:  $\lim_{x \rightarrow -\infty} F_{XY}(x, y) = 0, \forall y$ )



# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$   
(These are actually limits:  $\lim_{x \rightarrow -\infty} F_{XY}(x, y) = 0, \forall y$ )
- ▶  $F_{XY}$  is non-decreasing in each of its arguments

# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$   
(These are actually limits:  $\lim_{x \rightarrow -\infty} F_{XY}(x, y) = 0, \forall y$ )
- ▶  $F_{XY}$  is non-decreasing in each of its arguments
- ▶  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments

# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$   
(These are actually limits:  $\lim_{x \rightarrow -\infty} F_{XY}(x, y) = 0, \forall y$ )
- ▶  $F_{XY}$  is non-decreasing in each of its arguments
- ▶  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments
- ▶ These are straight-forward extensions of single rv case

# Properties of Joint Distribution Function

- ▶ Joint distribution function:

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$   
(These are actually limits:  $\lim_{x \rightarrow -\infty} F_{XY}(x, y) = 0, \forall y$ )
- ▶  $F_{XY}$  is non-decreasing in each of its arguments
- ▶  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments
- ▶ These are straight-forward extensions of single rv case
- ▶ But there is another crucial property satisfied by  $F_{XY}$ .

- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.

- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.
- ▶ Let  $x_1 < x_2$ . Then

$$P[x_1 < X \leq x_2] = F_X(x_2) - F_X(x_1)$$

- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.
- ▶ Let  $x_1 < x_2$ . Then

$$P[x_1 < X \leq x_2] = F_X(x_2) - F_X(x_1)$$

- ▶ The LHS above is a probability.

- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.
- ▶ Let  $x_1 < x_2$ . Then

$$P[x_1 < X \leq x_2] = F_X(x_2) - F_X(x_1)$$

- ▶ The LHS above is a probability.  
The RHS is non-negative because  $F_X$  is non-decreasing.



- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.
- ▶ Let  $x_1 < x_2$ . Then

$$P[x_1 < X \leq x_2] = F_X(x_2) - F_X(x_1)$$

- ▶ The LHS above is a probability.  
The RHS is non-negative because  $F_X$  is non-decreasing.
- ▶ We will now derive a similar expression in the case of two random variables.

- ▶ Recall that, for the case of a single rv, the probability of  $X$  being in any interval is given by the difference of  $F_X$  values at the end points of the interval.
- ▶ Let  $x_1 < x_2$ . Then


$$P[x_1 < X \leq x_2] = F_X(x_2) - F_X(x_1)$$

- ▶ The LHS above is a probability.  
The RHS is non-negative because  $F_X$  is non-decreasing.
- ▶ We will now derive a similar expression in the case of two random variables.
- ▶ Here, the probability we want is that of the pair of rv's being in a cylindrical set.


- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .

- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .

- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .




- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .



$$B \triangleq (-\infty, x_2] \times (-\infty, y_2] = B_1 + (B_2 \cup B_3)$$


- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .



$$B \triangleq (-\infty, x_2] \times (-\infty, y_2] = B_1 + (B_2 \cup B_3)$$

$$B_1 = (x_1, x_2] \times (y_1, y_2]$$


- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .



$$\begin{aligned}
 B &\triangleq (-\infty, x_2] \times (-\infty, y_2] = B_1 + (B_2 \cup B_3) \\
 B_1 &= (x_1, x_2] \times (y_1, y_2] \\
 B_2 &= (-\infty, x_2] \times (-\infty, y_1]
 \end{aligned}$$




- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .



$$\begin{aligned}
 B &\triangleq (-\infty, x_2] \times (-\infty, y_2] = B_1 + (B_2 \cup B_3) \\
 B_1 &= (x_1, x_2] \times (y_1, y_2] \\
 B_2 &= (-\infty, x_2] \times (-\infty, y_1] \\
 B_3 &= (-\infty, x_1] \times (-\infty, y_2]
 \end{aligned}$$

- ▶ Let  $x_1 < x_2$  and  $y_1 < y_2$ . We want  $P[x_1 < X \leq x_2, y_1 < Y \leq y_2]$ .
- ▶ Consider the Borel set  $B = (-\infty, x_2] \times (-\infty, y_2]$ .




$$B \triangleq (-\infty, x_2] \times (-\infty, y_2] = B_1 + (B_2 \cup B_3)$$


$$B_1 = (x_1, x_2] \times (y_1, y_2]$$

$$B_2 = (-\infty, x_2] \times (-\infty, y_1]$$


$$B_3 = (-\infty, x_1] \times (-\infty, y_2]$$

$$B_2 \cap B_3 = (-\infty, x_1] \times (-\infty, y_1]$$







$$P[(X, Y) \in B] = P[X \leq x_2, Y \leq y_2]$$



$$P[(X, Y) \in B] = P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2)$$




$$\begin{aligned}
 P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\
 &= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\
 &= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]
 \end{aligned}$$



$$\begin{aligned}P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\&= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\&= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]\end{aligned}$$


$$P[(X, Y) \in B_2] = P[X \leq x_2, Y \leq y_1] = F_{XY}(x_2, y_1)$$



$$\begin{aligned}
 P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\
 &= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\
 &= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]
 \end{aligned}$$

$$\begin{aligned}
 P[(X, Y) \in B_2] &= P[X \leq x_2, Y \leq y_1] = F_{XY}(x_2, y_1) \\
 P[(X, Y) \in B_3] &= P[X \leq x_1, Y \leq y_2] = F_{XY}(x_1, y_2)
 \end{aligned}$$






$$\begin{aligned}
 P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\
 &= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\
 &= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]
 \end{aligned}$$

$$P[(X, Y) \in B_2] = P[X \leq x_2, Y \leq y_1] = F_{XY}(x_2, y_1)$$

$$P[(X, Y) \in B_3] = P[X \leq x_1, Y \leq y_2] = F_{XY}(x_1, y_2)$$

$$P[(X, Y) \in B_2 \cap B_3] = P[X \leq x_1, Y \leq y_1] = F_{XY}(x_1, y_1)$$




$$\begin{aligned}
 P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\
 &= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\
 &= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]
 \end{aligned}$$

$$P[(X, Y) \in B_2] = P[X \leq x_2, Y \leq y_1] = F_{XY}(x_2, y_1)$$

$$P[(X, Y) \in B_3] = P[X \leq x_1, Y \leq y_2] = F_{XY}(x_1, y_2)$$

$$P[(X, Y) \in B_2 \cap B_3] = P[X \leq x_1, Y \leq y_1] = F_{XY}(x_1, y_1)$$

$$P[(X, Y) \in B_1] = F_{XY}(x_2, y_2) - P[(X, Y) \in (B_2 \cup B_3)]$$



$$\begin{aligned}
 P[(X, Y) \in B] &= P[X \leq x_2, Y \leq y_2] = F_{XY}(x_2, y_2) \\
 &= P[(X, Y) \in B_1 + (B_2 \cup B_3)] \\
 &= P[(X, Y) \in B_1] + P[(X, Y) \in (B_2 \cup B_3)]
 \end{aligned}$$

$$P[(X, Y) \in B_2] = P[X \leq x_2, Y \leq y_1] = F_{XY}(x_2, y_1)$$

$$P[(X, Y) \in B_3] = P[X \leq x_1, Y \leq y_2] = F_{XY}(x_1, y_2)$$

$$P[(X, Y) \in B_2 \cap B_3] = P[X \leq x_1, Y \leq y_1] = F_{XY}(x_1, y_1)$$

$$\begin{aligned}
 P[(X, Y) \in B_1] &= F_{XY}(x_2, y_2) - P[(X, Y) \in (B_2 \cup B_3)] \\
 &= F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1)
 \end{aligned}$$

- ▶ What we showed is the following.

- ▶ What we showed is the following.
- ▶ For  $x_1 < x_2$  and  $y_1 < y_2$

$$\begin{aligned} P[x_1 < X \leq x_2, y_1 < Y \leq y_2] &= F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) \\ &\quad - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \end{aligned}$$

- ▶ What we showed is the following.
- ▶ For  $x_1 < x_2$  and  $y_1 < y_2$

$$P[x_1 < X \leq x_2, y_1 < Y \leq y_2] = F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) \\ - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1)$$

- ▶ This means  $F_{XY}$  should satisfy

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

for all  $x_1 < x_2$  and  $y_1 < y_2$

- ▶ What we showed is the following.
- ▶ For  $x_1 < x_2$  and  $y_1 < y_2$

$$P[x_1 < X \leq x_2, y_1 < Y \leq y_2] = F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) \\ - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1)$$

- ▶ This means  $F_{XY}$  should satisfy

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

for all  $x_1 < x_2$  and  $y_1 < y_2$

- ▶ This is an additional condition that a function has to satisfy to be the joint distribution function of a pair of random variables

# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \Re^2 \rightarrow \Re$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$



# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \Re^2 \rightarrow \Re$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1.  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$

# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1.  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$
2.  $F_{XY}$  is non-decreasing in each of its arguments

# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1.  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$
2.  $F_{XY}$  is non-decreasing in each of its arguments
3.  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments

# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1.  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$
2.  $F_{XY}$  is non-decreasing in each of its arguments
3.  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments
4. For all  $x_1 < x_2$  and  $y_1 < y_2$

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

# Properties of Joint Distribution Function

- ▶ Joint distribution function:  $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1.  $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$   
 $F_{XY}(\infty, \infty) = 1$
2.  $F_{XY}$  is non-decreasing in each of its arguments
3.  $F_{XY}$  is right continuous and has left-hand limits in each of its arguments
4. For all  $x_1 < x_2$  and  $y_1 < y_2$

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

- ▶ Any  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  satisfying the above would be a joint distribution function.