Computational Methods in Physics (PHY 365) FA23

Dr. Muhammad Kamran

Department of Physics

COMSATS University Islamabad

Lab 9

- FX = gradient(F) returns the one-dimensional numerical gradient of vector F.
 - \diamond Here FX corresponds to $\partial F/\partial x$, the differences in "x" (horizontal) direction.
 - ♦ The spacing between points is assumed to be 1.

- FX = gradient(F) returns the one-dimensional numerical gradient of vector F.
 - \diamond Here FX corresponds to $\partial F/\partial x$, the differences in "x" (horizontal) direction.
 - ♦ The spacing between points is assumed to be 1.
- [FX, FY] = gradient(F) returns the x and y components of the two-dimensional numerical gradient of matrix F.

- FX = gradient(F) returns the one-dimensional numerical gradient of vector F.
 - \diamond Here FX corresponds to $\partial F/\partial x$, the differences in "x" (horizontal) direction.
 - ♦ The spacing between points is assumed to be 1.
- [FX, FY] = gradient(F) returns the x and y components of the two-dimensional numerical gradient of matrix F.
- [____] = gradient(F, h) uses h as a uniform spacing between points in each direction.

Problem: Use the gradient function to approximate the first and second derivatives of sin(x).

- Problem: Use the gradient function to approximate the first and second derivatives of sin(x).
- The step size and domain

```
step\_size = 10 ^-3;

x\_vec = -pi : step\_size : pi;
```

- Problem: Use the gradient function to approximate the first and second derivatives of sin(x).
- The step size and domain

```
step\_size = 10 ^-3;

x\_vec = -pi : step\_size : pi;
```

■ The function

$$f_x = \sin(x_vec);$$

- Problem: Use the gradient function to approximate the first and second derivatives of sin(x).
- The step size and domain

```
step\_size = 10 ^-3;

x\_vec = -pi : step\_size : pi;
```


The function $f x = \sin(x \text{ vec});$

■ Calculating the first and second derivatives

```
grad_1 = gradient(f_x, step_size);
grad_2 = gradient(grad_1, step_size);
```

■ Plotting

```
plot(x\_vec \;,\, f\_x \;, `r' \;,\, x\_vec \;,\, grad\_1 \;,\, `b--' \;,\, x\_vec \;,\, grad\_2 \;,\, `m-.')
```



■ The diff function can also be used to differentiate symbolic expressions/functions.

- The diff function can also be used to differentiate symbolic expressions/functions.
- Y = diff(f) differentiates the function "f" with respect to the symbolic scalar variable determined by symvar(f, 1).
- Y = diff(f, var) differentiates the function with respect to the differentiation parameter "var".
- The parameter var can be
 - ♦ a symbolic scalar variable, such as x,
 - \diamond a symbolic function, such as f(x), or
 - \diamond a derivative function, such as diff(f(t), t).

- The diff function can also be used to differentiate symbolic expressions/functions.
- Y = diff(f) differentiates the function "f" with respect to the symbolic scalar variable determined by symvar(f, 1).
- Y = diff(f, var) differentiates the function with respect to the differentiation parameter "var".
- The parameter var can be
 - \diamond a symbolic scalar variable, such as \mathbf{x} ,
 - \diamond a symbolic function, such as f(x), or
 - \diamond a derivative function, such as diff(f(t), t).
- Y = diff(f, var, n) computes the nth derivative of f with respect to var.

■ Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.

- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;

- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;
- $f = 3 * x^4 + 50 * x^2 10;$

- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;
- \blacksquare a1 = diff(f);


- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;
- \blacksquare a1 = diff(f);
- \bullet a2 = diff(f, x);

- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;
- \blacksquare a1 = diff(f);
- \bullet a2 = diff(f, x);
- a3 = diff(f, x, 2);

- Problem: Define a function using MATLAB's symbolic toolbox, and find out its derivatives.
- syms x;
- \blacksquare a1 = diff(f);
- \bullet a2 = diff(f, x);
- \bullet a3 = diff(f, x, 2);
- a4 = diff (a2);

■ Problem: Determine the (symbolic) first and second derivatives of sin(x) using diff function.

■ Problem: Determine the (symbolic) first and second derivatives of sin(x) using diff function.



References

- https://www.mathworks.com/help/matlab/ref/gradient.html
- https://www.mathworks.com/help/symbolic/diff.html
- https://www.mathworks.com/help/symbolic/sym.gradient.html