






1



2



3



4

## 2.1. Probability

- Probability

- Laplace classic definition of probability:

$$P(A) = \frac{\text{Number of outcomes favorable to } A}{\text{Total number of possible outcomes}},$$

- Relative frequency definition of probability:

$$P(A) = \lim_{n \rightarrow \infty} \frac{n_A}{n}$$



5

## 2.1. Probability

- Kolmogorov axiomatic formulation

- $\Omega$ : sample space: set of all experimental outcomes

$$\Omega = \{ \xi_1, \xi_2, \dots, \xi_k, \dots, \xi_n, \dots \}$$

- Event – any subset of  $\Omega$ . Number of subset of sample space :  $2^n$  if  $n < \infty$ .

- $\sigma$ -field  $F$  of subsets of  $\Omega$

- $P$ : a probability measure on the sets in  $F$

- $A$  – any event

- 3 axiom of probability

- (i)  $P(A) \geq 0$  (Probability is a nonnegative number)

- (ii)  $P(\Omega) = 1$  (Probability of the whole set is unity)

- (iii) If  $A \cap B = \emptyset$ , then  $P(A \cup B) = P(A) + P(B)$ .

- $\langle \Omega, F, P \rangle$ : probability model




6

## 2.1. Probability


- Events: A and B

- Mutually exclusive events:  $A \cap B = \emptyset$

- Partition of  $\Omega$ :



$$A_i \cap A_j = \emptyset, \text{ and } \bigcup_{i=1}^n A_i = \Omega$$



$A \cap B = \emptyset$

- Example: experiment of tossing two coins simultaneously

- Elementary events:

$$\xi_1 = (H, H), \xi_2 = (H, T), \xi_3 = (T, H), \xi_4 = (T, T)$$

- The subset  $A = \{ \xi_1, \xi_2, \xi_3 \}$



7

## Probability

- Counting the sample points

- Evaluate is the element of chance associated with the occurrence of certain events when an experiment is performed.

- To solve a probability problem by counting the number of points in the sample space without actually listing each element.

- The fundamental principle of counting - **multiplication rule**:

- if an operation can be performed in  $n_1$  ways, and if for each of these a second operation can be performed in  $n_2$  ways, and for each of the first two a third operation can be performed in  $n_3$  ways, and so forth, then the sequence of  $k$  operations can be performed in  $n_1 n_2 \cdot \dots \cdot n_k$  ways.



8

## 2.1. Probability

- Conditional probability and independence

- $N$  independent trials,
- $N_A, N_B, N_{AB}$  : the number of times events  $A, B$  and  $AB$  occur.
- For large  $N$   $P(A) \approx \frac{N_A}{N}, P(B) \approx \frac{N_B}{N}, P(AB) \approx \frac{N_{AB}}{N}$ .

- Conditional probability:  $P(A|B)$

$$P(A|B) = \frac{N_{AB}}{N_B} = \frac{N_{AB}/N}{N_B/N} = \frac{P(AB)}{P(B)}$$



9

## 2.1. Probability

- Properties of conditional probability

- $P(A|B)$  is nonnegative:

$$P(A|B) = \frac{P(AB)}{P(B)} \geq 0,$$

- $P(\Omega|B) = 1$

$$P(\Omega|B) = \frac{P(\Omega B)}{P(B)} = \frac{P(B)}{P(B)} = 1,$$

- If  $A \cap C = \emptyset$ ,

$$P(A \cup C | B) = P(A|B) + P(C|B),$$



10

## 2.1. Probability

- If  $B \subset A$  then  $P(A|B) = 1$
- If  $A \subset B$  then  $P(A|B) = P(A)$
- Let,  $A_1, A_2, \dots, A_n$  are pair wise disjoint and their union is  $\Omega$ :

$$A_i \cap A_j = \emptyset, \quad \bigcup_{i=1}^n A_i = \Omega.$$

- $B$  is an event

$$P(B) = \sum_{i=1}^n P(BA_i) = \sum_{i=1}^n P(B|A_i)P(A_i).$$



11

## 2.1. Probability

- Independence:  $A$  and  $B$  events

$$P(AB) = P(A)P(B)$$

- If  $A$  and  $B$  are independent events:

$$P(A|B) = P(A)$$

$$P(A|B) = \frac{P(B|A)}{P(B)} \cdot P(A)$$

- Generalized Bayes theorem:

$$P(A_i|B) = \frac{P(B|A_i)P(A_i)}{P(B)} = \frac{P(B|A_i)P(A_i)}{\sum_{i=1}^n P(B|A_i)P(A_i)},$$



12

## 2.1. Probability


- Bayes' theorem interpretation:
  - $P(A)$  represents the a-priori probability of the event  $A$ .
  - The event  $B$  is new knowledge obtained from an experiment.
  - Conditional probability  $P(A|B)$  of  $A$  given  $B$  – a-posteriori probability
  - The new information should be used to improve knowledge of  $A$ .

13

## Probability

### • Examples

- An electrical system consists of four components. The system works if components  $A$  and  $B$  work and either of the components  $C$  or  $D$  works. The reliability (probability of working) of each component is also shown in the Figure. Find the probability that:
  - (a) the entire system works and
  - (b) the component  $C$  does not work, given that the entire system works. Assume that the four components work independently.



14

## 2.1. Probability

### • Example:

- In a box: 6 white and 4 black balls.
- Remove two balls randomly without replacement.
- $P\{\text{the first one is white and the second one is black}\} = ?$
- Question: are events  $W_1$  and  $B_2$  independent?
  - $W_1$  = "first ball removed is white"
  - $B_2$  = "second ball removed is black"

15

## 2.1. Probability

- If, in an experiment, the events  $A_1, A_2, \dots, A_k$  can occur, then

$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \dots P(A_k|A_1 \cap A_2 \cap \dots \cap A_{k-1}).$$

- If the events  $A_1, A_2, \dots, A_k$  are independent, then

$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1)P(A_2) \dots P(A_k).$$

### • Example

- Three cards are drawn in succession, without replacement, from an ordinary deck of playing cards. Find the probability that the event  $A_1 \cap A_2 \cap A_3$  occurs, where  $A_1$  is the event that the first card is a red ace,  $A_2$  is the event that the second card is a 10 or a jack, and  $A_3$  is the event that the third card is greater than 3 but less than 7.

16

## 2.1. Probability

- Example: Two boxes  $B_1$  and  $B_2$  contain 100 and 200 light bulbs respectively. The first box ( $B_1$ ) has 15 defective bulbs and the second - 5. Suppose a box is selected at random and one bulb is picked out.
  - (a) What is the probability that it is defective?
  - (b) Suppose we test the bulb and it is found to be defective. What is the probability that it came from box 1?



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

17

## 2.1. Probability

- Repeated trials, Bernoulli trials
  - Consider  $n$  independent experiments with models  $(\Omega_1, F_1, P_1), (\Omega_2, F_2, P_2), \dots, (\Omega_n, F_n, P_n)$ .
    - Let  $\xi_1 \in \Omega_1, \xi_2 \in \Omega_2, \dots, \xi_n \in \Omega_n$ : elementary events.
    - A joint performance of the  $n$  experiments produces an elementary events  $\omega = (\xi_1, \xi_2, \dots, \xi_n)$ .
    - Consider space  $\Omega = \Omega_1 \times \Omega_2 \times \dots \times \Omega_n : \xi_1 \in \Omega_1, \dots, \xi_n \in \Omega_n$ .
    - Events in combined space  $\Omega$  are of the form  $A_1 \times A_2 \times \dots \times A_n$ .
    - If  $n$  experiments are independent, then  $P(A_1 \times A_2 \times \dots \times A_n) = P(A_1) \times \dots \times P(A_n)$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

18

## 2.1. Probability

- An event  $A$  has probability  $p$  of occurring in a single trial. Find the probability that  $A$  occurs exactly  $k$  times in determined location,  $k \leq n$  in  $n$  trials.

$$\begin{aligned} P_0(\omega) &= P(\{\xi_{i_1}, \xi_{i_2}, \dots, \xi_{i_k}, \dots, \xi_{i_n}\}) = \\ &= P(\{\xi_{i_1}\})P(\{\xi_{i_2}\}) \dots P(\{\xi_{i_k}\}) \dots P(\{\xi_{i_n}\}) = \\ &= \underbrace{P(A)P(A) \dots P(A)}_k \underbrace{P(\bar{A})P(\bar{A}) \dots P(\bar{A})}_{n-k} = p^k q^{n-k}. \end{aligned}$$

- $P\{A \text{ occurs exactly } k \text{ time in } n \text{ trials}\} = C_n^k p^k q^{n-k}$ 
  - Bernoulli formula.



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

19

## 2.1. Probability

- De Moivre – Laplace Theorem
  - Guess that:  $n \rightarrow \infty$  with fixed  $p$ .
  - $k$  is in the neighborhood  $\sqrt{npq}$  of  $np$ .
  - Bernoulli probability estimation:

$$C_n^k p^k q^{n-k} \approx \frac{1}{\sqrt{2\pi npq}} e^{-(k-np)^2 / 2npq}.$$

- Stirling formula for  $n!$  approximation:

$$n! \sim \sqrt{2\pi} n^{n+\frac{1}{2}} e^{-n}$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

20

## 2.1. Probability

- Estimation of Bernoulli formula

$$\binom{n}{k} p^k q^{n-k} = \frac{n!}{(n-k)!k!} p^k q^{n-k},$$

$$\binom{n}{k} p^k q^{n-k} > c_1 \sqrt{\frac{n}{2\pi(n-k)k}} \left(\frac{np}{k}\right)^k \left(\frac{nq}{n-k}\right)^{n-k}$$

$$\binom{n}{k} p^k q^{n-k} < c_2 \sqrt{\frac{n}{2\pi(n-k)k}} \left(\frac{np}{k}\right)^k \left(\frac{nq}{n-k}\right)^{n-k}$$

- Các hằng số  $c_1$  và  $c_2$  khá gần nhau.

$$c_1 = e^{\left\{\frac{1}{12n+1} - \frac{1}{12(n-k)} - \frac{1}{12k}\right\}} \quad c_2 = e^{\left\{\frac{1}{12n} - \frac{1}{12(n-k)+1} - \frac{1}{12k+1}\right\}}.$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

21

## 2.1. Probability

- Example:

- Toss a coin  $n$  times. Probability of getting  $k$  heads in  $n$  trials = ?
- Consider rolling a fair dice eight times. Find the probability that either 3 or 4 shows up five times.
- Suppose 5,000 components are ordered. The probability that a part is defective equals 0.1. What is the probability that the total number of defective parts does not exceed 400 ?



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

22

## 2.2. Random variables

- Definition

- $(\Omega, F, P)$  - probability model for an experiment,
- $X$  - a function that maps every  $\xi \in \Omega$  to a unique point  $x \in R$
- **Random Variable (r.v)**: a finite single valued function that maps the set of all experimental outcomes  $\Omega$  into the set of real numbers  $R$  is said to be a r.v, if the set  $A = \{\xi | X(\xi) \leq x\}$  is an event  $A \subset F$  for every  $x$  in  $R$ .



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

23

## 2.2. Random variables


- Probability distribution function

- Probability distribution function of random variable  $X$ : ( pdf )

$$F_X(x) = P\{\xi | X(\xi) \leq x\} \geq 0$$

- Examples

- $X$  - r.v:  $X(\xi) = c, \xi \in \Omega, F_X(x) = ?$
- $X$  - r.v: coin tossing,  $\Omega = \{H, T\}, X(T)=0, X(H)=1. F_X(x) = ?$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

24

## 2.2. Random variables

- Properties of probability distribution function
  - $F_X(x)$  is a distribution function:
    - $F_X(-\infty) = 0$ ;  $F_X(+\infty) = 1$
  - Pdf is nondecreasing function:
    - if  $x_1 \leq x_2$  then  $F_X(x_1) \leq F_X(x_2)$
  - Pdf is right continuous:  $F_X(x^+) = F_X(x) \forall x$

25

## 2.2. Random variables

- If  $F_X(x_0) = 0$  for some  $x_0$ ,  
then  $F_X(x) = 0 \forall x \leq x_0$
- $P\{X(\xi) > x\} = 1 - F_X(x)$
- If  $x_2 > x_1$ , then  
 $P\{x_1 < X(\xi) \leq x_2\} = F_X(x_2) - F_X(x_1)$
- $P\{X(\xi) = x\} = F_X(x) - F_X(x^-)$

26


## 2.2. Random variables

- Continuous and discrete random variables
  - $X$  – continuous r.v if  $F_X(x)$  is continuous
    - For continuous r.v,  $F_X(x^-) = F_X(x)$  and  $P\{X=x\} = 0$
  - If  $F_X(x) = \text{const}$ , except for a finite number of jump discontinuities, then  $X$  is said to be a discrete-type r.v.
    - If  $x_i$  is such a discontinuity point, then  
 $p_i = P\{X = x_i\} = F_X(x) - F_X(x^-)$

27

## 2.2. Random variables

- Examples
  - A fair coin is tossed twice, and let the r.v  $X$  represent the number of heads. Find  $F_X(x)$



28

## 2.2. Random variables

- Distribution density function

- Derivative of distribution function  $F_X(x)$  is density function  $f_X(x)$  of the r.v.

$$f_X(x) = \frac{dF_X(x)}{dx}.$$

- $F_X(x)$  is monoton nondecreasing function, so that

$$f_X(x) = \frac{dF_X(x)}{dx} = \lim_{\Delta x \rightarrow 0} \frac{F_X(x + \Delta x) - F_X(x)}{\Delta x} \geq 0,$$




VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

29

## 2.2. Random variables

- If r.v is continuous then  $f_X(x)$  is continuous.

- For discrete r.v

$$f_X(x) = \sum p_i \delta(x - x_i),$$


- $x_i$  is jump-discontinuity point function in  $F_X(x)$

$$F_X(x) = \int_{-\infty}^x f_X(u) du.$$

$$F_X(+\infty) = 1, \quad \int_{-\infty}^{+\infty} f_X(x) dx = 1,$$

$$P\{x_1 < X(\xi) \leq x_2\} = F_X(x_2) - F_X(x_1) = \int_{x_1}^{x_2} f_X(x) dx.$$




VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

30

## 2.2. Random variables

$$P\{x_1 < X(\xi) \leq x_2\} = F_X(x_2) - F_X(x_1) = \int_{x_1}^{x_2} f_X(x) dx.$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

31

## 2.2. Random variables

- Some continuous random variables

- Normal (Gaussian) random variables  $X \sim N(\mu, \sigma^2)$

- Density function


$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2 / 2\sigma^2}.$$

- Bell shape curve

- Distribution function

$$F_X(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(y-\mu)^2 / 2\sigma^2} dy = G\left(\frac{x-\mu}{\sigma}\right),$$

$$G(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-y^2 / 2} dy$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG


32



## 2.2. Random variables

- Uniform random variables:

- Density function




$$X \sim U(a, b), \quad a < b,$$

$$f_X(x) = \begin{cases} \frac{1}{b-a}, & a \leq x \leq b, \\ 0, & \text{otherwise.} \end{cases}$$

- Exponential random variables

- Density function

$$f_X(x) = \begin{cases} \frac{1}{\lambda} e^{-x/\lambda}, & x \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

33

## 2.2. Random variables


- Some important discrete random variables

- Bernoulli r.v:  $X$  takes values 0, 1

$$P(X=0) = q, \quad P(X=1) = p.$$

- Binomial r.v  $X \sim B(n, p)$ ,

$$P(X=k) = \binom{n}{k} p^k q^{n-k}, \quad k = 0, 1, \dots, n.$$




VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

34

## Random variables

- Poisson r.v  $X \sim P(\lambda)$ ,

$$P(X=k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots, \infty.$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

35

## 2.3 Some characteristics of random variables

- Mean value

$$\eta_X = \bar{X} = E(X) = \int_{-\infty}^{+\infty} x f_X(x) dx.$$

- For discrete random variable

$$\begin{aligned} \eta_X = \bar{X} = E(X) &= \int x \sum_i p_i \delta(x - x_i) dx = \sum_i x_i p_i \underbrace{\int \delta(x - x_i) dx}_1 \\ &= \sum_i x_i p_i = \sum_i x_i P(X = x_i). \end{aligned}$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

36

## Mean and Variance

- Example: uniform random variable

$$E(X) = \int_a^b \frac{x}{b-a} dx = \frac{1}{b-a} \left. \frac{x^2}{2} \right|_a^b = \frac{b^2 - a^2}{2(b-a)} = \frac{a+b}{2}$$

- Example: exponential random variable

$$E(X) = \int_0^{\infty} \frac{x}{\lambda} e^{-x/\lambda} dx = \lambda \int_0^{\infty} ye^{-y} dy = \lambda,$$

- Example: Poisson random variable

$$\begin{aligned} E(X) &= \sum_{k=0}^{\infty} k P(X=k) = \sum_{k=0}^{\infty} k e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} \sum_{k=1}^{\infty} k \frac{\lambda^k}{k!} \\ &= e^{-\lambda} \sum_{k=1}^{\infty} \frac{\lambda^k}{(k-1)!} = \lambda e^{-\lambda} \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = \lambda e^{-\lambda} e^{\lambda} = \lambda. \end{aligned}$$



37

## 2.3 Some characteristics of random variables

- Variance

- For r.v X with mean  $\mu$

- Variance  $\sigma_X^2 = E[(X - \mu)^2] > 0.$

- Or  $\sigma_X^2 = \int_{-\infty}^{+\infty} (x - \mu)^2 f_X(x) dx > 0.$

- Standard deviation  $\sigma_X = \sqrt{E(X - \mu)^2}$

- Variance and means

$$\begin{aligned} Var(X) = \sigma_X^2 &= E[(X - \mu)^2] = E[X^2 - 2X\mu + \mu^2] = \\ &= E[X^2] - E[2X\mu] + E[\mu^2] = E[X^2] - 2E[X]\mu + \mu^2 \end{aligned}$$

$$= E(X^2) - [E(X)]^2 = \overline{X^2} - \bar{X}^2.$$

38

## 2.3 Some characteristics of random variables

- Example:
- Example: variance of Poisson r.v.

$$\sigma_X^2 = \overline{X^2} - \bar{X}^2 = (\lambda^2 + \lambda) - \lambda^2 = \lambda.$$

- Example: variance of Gaussian r.v.

$$Var(X) = E[(X - \mu)^2] = \int_{-\infty}^{+\infty} (x - \mu)^2 \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2} dx.$$

$$Var(X) = \sigma^2$$



39

## 2.3 Some characteristics of random variables

- Moments

- X – random variables

$$m_n = \overline{X^n} = E(X^n), \quad n \geq 1$$

- Central moments

$$\mu_n = E[(X - \mu)^n]$$

- Relation between moments and central moments

$$\mu_n = E[(X - \mu)^n] = \sum_{k=0}^n C_n^k m_k (-\mu)^{n-k}.$$

- Mean and variance:

$$\mu = m_1, \quad \sigma^2 = \mu_2.$$



40

### 2.3 Some characteristics of random variables





- Generalized moments of X about a

$$E[(X - a)^n]$$

- Absolute moments of X

$$E[|X|^n]$$

- Measure of skewness is  $E[(X - \mu_X)^3]/\sigma_X^3$
- Measure of kurtosis is  $E[(X - \mu_X)^4]/\sigma_X^4$ .



### 2.3 Some characteristics of random variables

- Characteristic function

- Continuous r.v X

$$\Phi_X(\omega) = E(e^{jX\omega}) = \int_{-\infty}^{+\infty} e^{jX\omega} f_X(x) dx.$$

- We have:

$$\Phi_X(0) = 1,$$

- And


- Discrete r.v  $|\Phi_X(\omega)| \leq 1 \quad \forall \omega$

$$\Phi_X(\omega) = \sum_k e^{jk\omega} P(X = k).$$

### 2.4 Chebychev Inequality and Law of Large numbers

- Chebychev Inequality

- Consider an interval of width  $2\varepsilon$  symmetrically centered around its mean  $\mu$



- Chebychev Inequality

$$P(|X - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{\varepsilon^2},$$

### 2.4 Chebychev Inequality and Law of Large numbers

- Weak law and strong law of large numbers

- $X_i$  - independent, identically distributed Bernoulli random variables:

$$P(X_i) = p, \quad P(X_i = 0) = 1 - p = q,$$

- $k = X_1 + X_2 + \dots + X_n$  - number of successes in  $n$  trials
- Weak law of large numbers:


$$P\left\{\left|\frac{k}{n} - p\right| > \varepsilon\right\} \leq \frac{pq}{n\varepsilon^2}.$$

- Strong law of large numbers:

- The ratio  $k/n$  tends to  $p$  not only in probability, but with probability 1

## 2.5. Gaussian distribution

- Gaussian distribution
  - One dimensional normal distribution



45

## 2.5. Gaussian distribution

- Multidimensional Gaussian distribution (  $d$  dimensional )

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left[-\frac{1}{2}(\mathbf{x}-\mu)^T \Sigma^{-1}(\mathbf{x}-\mu)\right]$$

$$\mu \equiv E(\mathbf{x}) = \int \mathbf{x} p(\mathbf{x}) d\mathbf{x}$$

$$\Sigma \equiv E((\mathbf{x}-\mu)(\mathbf{x}-\mu)^T) = \int (\mathbf{x}-\mu)(\mathbf{x}-\mu)^T p(\mathbf{x}) d\mathbf{x}$$

- Where,  $\mathbf{x}$ ,  $\mu$  are  $d$ -dimensional vector
- $\Sigma$  covariance matrix - symmetric and semi-positive defined matrix.
- Consider cases, when  $\Sigma$  positive defined.

$$\mu_i = E(x_i)$$

$$\sigma_{ij} = E((x_i - \mu_i)(x_j - \mu_j))$$

46


## 2.5. Gaussian distribution

- Covariance matrix

$$\Sigma = \begin{bmatrix} \sigma_{00} & & & \\ & \sigma_{11} & \sigma_{ij, i \neq j} & \\ & & \sigma_{22} & \\ & & & \sigma_{33} \end{bmatrix}$$

47

## 2.5. Gaussian distribution



48

## 2.5. Gaussian distribution

- Multidimensional Gaussian distribution is fully defined by  $d+d(d+1)/2$  parameters of mean vector  $\mu$  and covariance matrix  $\Sigma$ .
- Mahalanobis distance:

$$r = (\mathbf{x} - \mu)^T \Sigma^{-1} (\mathbf{x} - \mu)$$



VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

49



50